

RENDICONTI
del
SEMINARIO MATEMATICO
della
UNIVERSITÀ DI PADOVA

D. V. THAMPURAN

Normal neighborhood spaces

Rendiconti del Seminario Matematico della Università di Padova,
tome 45 (1971), p. 95-97

http://www.numdam.org/item?id=RSMUP_1971__45__95_0

© Rendiconti del Seminario Matematico della Università di Padova, 1971, tous droits réservés.

L'accès aux archives de la revue « Rendiconti del Seminario Matematico della Università di Padova » (<http://rendiconti.math.unipd.it/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>

NORMAL NEIGHBORHOOD SPACES

D. V. THAMPURAN *)

The object of this paper is to extend to neighborhood spaces the well-known Urysohn's lemma for topological spaces.

DEFINITION 1. Let X be a set and k a set-valued set-function mapping the power set, of X , to itself. Then (X, k) is said to be a neighborhood space iff,

1. $k\emptyset = \emptyset$
2. $A \subset kA$ for every $A \subset X$ and
3. $kA \subset kB$ if $A \subset B \subset X$.

The neighborhood space (X, k) is said to be directed iff $k(A \cup B) = kA \cup kB$ for all $A, B \subset X$.

For a subset A of X , write $cA = X - A$.

DEFINITION 2. Let (X, k) be a neighborhood space. Take $i = ckc$. Then a set A is said to be a neighborhood of a set B iff $B \subset iA$.

DEFINITION 3. A neighborhood space (X, k) is said to be normal iff $A, B \subset X$ and kA, kB are disjoint imply kA, kB have disjoint neighborhoods.

It is obvious that a neighborhood space (X, k) is normal iff $A, B \subset X$ and $kA \subset iB$ imply there is $C \subset X$ such that $kA \subset iC$ and $kC \subset iB$.

*) Indirizzo dell'A.: Instituto de Matematica, Casilla 114-D, Santiago, Chile.

DEFINITION 4. Let (X, k) , (Y, m) be two neighborhood spaces and f a function from X to Y . Then f is said to be continuous at the point x of X iff the inverse, under f , of every neighborhood of $f(x)$ is a neighborhood of x . We will say f is continuous iff f is continuous at each point of X .

It is easily seen that f is continuous iff $fk \subset mf$.

Let R denote the reals and n the closure function of the usual topology for R . Denote by I the closed unit interval $[0, 1]$ and let p be the restriction of n to I .

LEMMA 1. Let (X, k) be a directed neighborhood space and D a dense subset of the positive reals. For each t in D let $S(t)$ be a subset of X such that

1. $\cup\{S(t) : t \in D\} = X$ and
2. $kS(t) \subset iS(u)$ if $t < u$.

Take $f(x) = \inf\{t : x \in S(t)\}$. Then f is a continuous function from (X, k) to (R, n) .

PROOF. Let $x \in X$. To prove f is continuous it is enough to show that $f(x) < v$ implies $E = \{y : f(y) < v, y \in X\}$ is a neighborhood of x and that $u < f(x)$ implies $F = \{y : f(y) > u, y \in X\}$ is a neighborhood of x .

Now $f(x) < v$ implies there are w, z in D such that $f(x) < w < z < v$. Hence $x \in S(w)$. Also $S(z) \subset E$ since y in $S(z)$ implies $f(y) \leq z < v$. Therefore $x \in iS(z)$ and so E is a neighborhood of x .

Also $u < f(x)$ implies there are r, s in D such that $u < r < s < f(x)$. Then $x \in cS(s)$ since $x \in S(s)$ implies $f(x) \leq s$. Next, $y \in cS(r)$ implies $f(y) \geq r > u$ and so $y \in F$; hence $cS(r) \subset F$. Now $kS(r) \subset iS(s)$ and so $kcS(s) \subset icS(r)$. Hence F is a neighborhood of x .

The next lemma can be proved in the same way as the corresponding part of Urysohn's lemma; for instance we can use the method of proof of Lemma 4 on page 115 of Kelley [1].

LEMMA 2. Let (X, k) be a normal directed neighborhood space and $A, B \subset X$ such that kA, kB are disjoint. Then there is a continuous function f from (X, k) to (I, p) such that f is 0 on kA and 1 on kB .

DEFINITION 5. A directed neighborhood space (X, k) is said to be completely normal iff $A, B \subset X$ and kA, kB are disjoint imply there

is a continuous function f from (X, k) to (I, p) such that f is 0 on kA and 1 on kB .

The next result now easily follows.

THEOREM 1. A directed neighborhood space is normal iff it is completely normal.

Define a neighborhood space for the reals R as follows. For a real number x let $\mathfrak{R}(x)$ be the family of all subsets N of R such that $\{y : y < v\} \subset N$ for some $v > x$ or $\{y : u < y\} \subset N$ for some $u < x$. For a subset A of the reals, let hA be the set of all points x such that each N in $\mathfrak{R}(x)$ intersects A . then (R, h) is a neighborhood space. Let g be the restriction of h to I .

DEFINITION 6. A neighborhood space (X, k) is said to be completely normal iff $A, B \subset X$ and kA, kB are disjoint imply there is a continuous function f from (X, k) to (I, g) such that f is 0 on kA and 1 on kB .

We then have the following result.

THEOREM 2. A neighborhood space is normal iff it is completely normal.

REFERENCES

- [1] KELLEY, J. L.: *General Topology*, Princeton (1968).

Manoscritto pervenuto in redazione l'11 maggio 1970.