

RENDICONTI
del
SEMINARIO MATEMATICO
della
UNIVERSITÀ DI PADOVA

MARCO BIROLI

**An estimate on convergence for an homogenisation
problem for variational inequalities**

Rendiconti del Seminario Matematico della Università di Padova,
tome 58 (1977), p. 9-24

http://www.numdam.org/item?id=RSMUP_1977__58__9_0

© Rendiconti del Seminario Matematico della Università di Padova, 1977, tous droits réservés.

L'accès aux archives de la revue « Rendiconti del Seminario Matematico della Università di Padova » (<http://rendiconti.math.unipd.it/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>

An estimate on convergence for an homogenisation problem for variational inequalities.

MARCO BIROLI (*)

1. - Introduction and results.

Let be $\Omega \subset R^n$ an open bounded set with smooth boundary Γ , $Y = \prod_{i=1}^n [0, y_i]$, $a_{ij}(y) \in L^\infty(Y)$ $i, j = 1, \dots, n$, such that

$$\sum_{i,j=1}^n a_{ij}(y) \xi_i \xi_j \geq \alpha |\xi|^2, \alpha > 0, \text{ a. e. in } Y$$

Let be $\forall u, v \in H_0^1(\Omega)$

$$\langle A^\varepsilon u, v \rangle = \int_{\Omega} \sum_{i,j=1}^n a_{ij} \left(\frac{x}{\varepsilon} \right) \frac{\partial u}{\partial x_j} (x) \frac{\partial v}{\partial x_i} (x) dx$$

(where \langle, \rangle is the duality between $H_0^1(\Omega)$ and $H^{-1}(\Omega)$ and $a_{ij}(y)$ are prolonged to R^n by periodicity) and

$$\langle A u, v \rangle = \int_{\Omega} \sum_{i,j=1}^n q_{ij} \frac{\partial u}{\partial x_j} (x) \frac{\partial v}{\partial x_i} (x) dx$$

where q_{ij} are constants defined as in [3], [9], [10].

(*) Indirizzo dell'A: Istituto di Matematica del Politecnico di Milano - Via Bonardi, 9 Milano.

Lavoro eseguito nell'ambito del gruppo G.N.A.F.A. del CNR.

Let be $f \in H^{-1}(\Omega)$, we have :

Th. 1 — We consider the problems

$$(1,1_\varepsilon) \quad A^\varepsilon u = f$$

$$(1,1) \quad A u = f$$

let be u_ε and u the solution of (1, 1_ε) and (1, 1), we have

$$\lim_{\varepsilon \rightarrow 0}^* u_\varepsilon = u \text{ in } H_0^1(\Omega)$$

Moreover if $a_{ij}(y) \in C^1(R^n)$, $f \in W^{1,p}(\Omega)$, $n < p \leq +\infty$, we have

$$(1,2) \quad \|u_\varepsilon - u\|_{L^\infty} \leq C \|f\|_{W^{1,p}} \varepsilon.$$

The first part of Th. 1 is shown in [7] for the symmetric case and in [8], [12] in the general case by « local energy » method, the second part of Th. 1 is shown by « multiple scales » method in [8] [9] (*). We consider now the problems

$$(1,3_\varepsilon) \quad \begin{aligned} \langle A^\varepsilon u_\varepsilon, v - u_\varepsilon \rangle &\geq \langle f, v - u_\varepsilon \rangle \\ \forall v \in H_0^1(\Omega) &, \quad v \leq \psi \\ u \in H_0^1(\Omega) &, \quad u \geq \psi \end{aligned}$$

$$(1,3) \quad \begin{aligned} \langle A u, v - u \rangle &\geq \langle f, v - u \rangle \\ \forall v \in H_0^1(\Omega) &, \quad v \leq \psi \\ u \in H_0^1(\Omega) &, \quad u \leq \psi \end{aligned}$$

where ψ is a measurable function, such that $\{v \in H_0^1(\Omega), v \leq \psi\} \neq \emptyset$.

Th. 2 Let be u_ε and u solution of (1,3_ε) and (1,3) we have

$$\lim_{\varepsilon \rightarrow 0}^* u_\varepsilon = u \text{ in } H_0^1(\Omega)$$

Moreover if $\psi \in C(\bar{\Omega})$, $f \in L^\infty(\Omega)$

$$\lim_{\varepsilon \rightarrow 0} u_\varepsilon = u \text{ in } C(\bar{\Omega}).$$

(*) Cf. also [1], [2].

The first part of Th. 2 is shown in [11] by the method of « local energy » (cfr. also [5] and [6]), the second part of Th. 2 is shown in [4] by penalisation method and Hölder-estimates. The aim of this work is show an estimate analogous to (1, 2) in the case of variational inequalities.

Th. 3. *Let be $a_{ij}(y) \in C^1(\mathbb{R}^n)$, $f \in L^\infty(\Omega)$, $u_\varepsilon(u)$ the solution of (1, 3 _{ε}) ((1, 3)), α the De Giorgi's exponent, $n \geq 2$*

(I) *If $\|A^\varepsilon \psi\|_{L^\infty}, \|A \psi\|_{L^\infty} \leq C$, we have*

$$\|u_\varepsilon - u\|_{L^\infty} \leq K \varepsilon^{\frac{\alpha}{n-2+3\alpha}}$$

(II) *If $\psi \in W^{1,p}(\Omega)$, $n < p < +\infty$, we have*

$$\|u_\varepsilon - u\|_{L^\infty} \leq K \varepsilon^{\frac{\alpha}{n-2+3\alpha}}$$

The Th. 2 is shown by penalisation and « multiple scales » method. In § 2 we show some preliminary results, in § 3 we study the convergence for the penalised equations and in § 4 we give the proof of Th. 3.

- Rem. 1 (a)* The hypothesis of the part (I) are verified if $\psi = 0$:
 (b) The result of Th. 3 can be generalized to the case $a_{ij}\left(x, \frac{x}{\varepsilon}\right)$
 (c) We don't know if the estimates in Th. 3 are optimal ;

2. – Some preliminary results.

We consider the penalised problem

$$(2,1) \quad A u + \frac{1}{\lambda} (u - \psi)^+ = f, \quad \lambda > 0.$$

LEMMA 1. *The problem (2.1) has a unique solution u^λ ; if $A \psi \in L^\infty(\Omega)$ we have*

$$(2,2) \quad \frac{1}{\lambda} \|(u^\lambda - \psi)^+\|_{L^\infty} \leq \|f\|_{L^\infty} + \|A\psi\|_{L^\infty}$$

$$(2,3) \quad \|A u^\lambda\|_{L^\infty} \leq 2 \|f\|_{L^\infty} + \|A\psi\|_{L^\infty}$$

From (2,1) we have

$$\frac{1}{\lambda^P} \|(u^\lambda - \psi)^+\|_{L^p}^P \leq \frac{1}{\lambda^{P-1}} \|(u^\lambda - \psi)^+\|_{L^p}^{P-1} (\|f\|_{L^p} + \|A\psi\|_{L^p})$$

then

$$(2,4) \quad \frac{1}{\lambda} \|(u^\lambda - \psi)^+\|_{L^p} \leq C_1 + C_2 \|A \psi\|_{L^p}$$

$$(2,5) \quad \|A u^\lambda\|_{L^p} \leq C_1 + C_2 \|A \psi\|_{L^p}$$

From (2,5) we have $u^\lambda \in C(\bar{\Omega})$.

We show now (2,2) « ab absurdum ».

We write

$$(2,6) \quad K = \|f\|_{L^\infty} + \|A \psi\|_{L^\infty}.$$

Let be

$$\Omega' = \left\{ x \mid \frac{1}{\lambda} (u^\lambda - \psi)^+ > K \right\} \neq \emptyset$$

(2,7)

$$\Omega'_\eta = \left\{ x \mid \frac{1}{\lambda} (u^\lambda - \psi)^+ > K + \eta \right\} \neq \emptyset, \quad \eta > 0$$

Ω' and Ω'_η are open set and $\Omega'_\eta \subset \bar{\Omega}'_\eta \subset \Omega'$.

Let be now Ω'' an open set with smooth boundary such that $\bar{\Omega}'_\eta \subset \subset \Omega'' \subset \bar{\Omega}'' \subset \Omega'$.

We have

$$A(u^\lambda - \psi) < 0 \text{ in } \Omega''$$

then from the maximum principle, [13],

$$(2,8) \quad \frac{1}{\lambda} (u^\lambda - \psi)^+ \leq K + \eta \text{ in } \Omega'' \supset \Omega'_\eta$$

From (2,8) we have

$$(2,9) \quad \Omega'_\eta = \emptyset ,$$

then from (2,7) (2,9) we have a contradiction.

From (2,1) (2,2) we have also (2,3).

LEMMA 2. *If $f \in L^\infty(\Omega)$, $\psi \in W^{1,p}(\Omega)$, $n < p \leq +\infty$, we have*

$$(2,10) \quad \|u^\lambda - u\|_{L^\infty} \leq K \lambda^{\frac{\alpha}{n-2(1-\alpha)}}$$

Moreover if $A\psi \in L^\infty(\Omega)$ we have

$$(2,11) \quad \|u^\lambda - u\|_{L^\infty} \leq K \lambda^{\frac{\alpha}{n-2(1-\alpha)}}$$

We show at first (2.11).

We observe that if $A\psi \in L^\infty(\Omega)$ from the lemma 1 we have $\|A u^\lambda\|_{L^\infty}, \|A u\|_{L^\infty} \leq \tilde{C}$ and repeating the calculation of [6] pg. 76-88 or [13] 177-197 we have

$$\|u^\lambda\|_{C^\alpha}, \|u\|_{C^\alpha} \leq C$$

where C depend only on \tilde{C} .

From (1,6_e) (2,1) we have

$$\langle A u^\lambda, u - u^\lambda \rangle \geq \langle f, u - u^\lambda \rangle$$

$$\langle A u, u^\lambda - (u^\lambda - \psi)^+ - u \rangle \geq \langle f, u^\lambda - (u^\lambda - \psi)^+ - u \rangle$$

then

$$(2,12) \quad \|u^\lambda - u\|_{H_0^1} \leq \lambda^{1/2} (C_1 + C_2 \|A\psi\|_{L^\infty})$$

From (2,12) we have

$$(2,13) \quad \|u^\lambda - u\|_{L^{\frac{2n}{n-2}}} \leq \lambda^{1/2} (C'_1 + C'_2 \|A\psi\|_{L^\infty}).$$

We prolongate u^λ and u by 0 and we denote again by u^λ and u these new functions.

From (2,13) we have

$$\int_{B(\bar{x}, \varrho)} |u^\lambda(x) - u(x)| dx \leq \lambda^{1/2} (C'_1 + C'_2 \|A\psi\|_{L^\infty}) |B(\bar{x}, \varrho)|^{1 - \frac{n-2}{2n}}$$

then

$$\begin{aligned} \frac{1}{|B(\bar{x}, \varrho)|} \int_{B(\bar{x}, \varrho)} |u^\lambda(x) - u(x)| dx &\leq \lambda^{1/2} (C'_1 + \\ &+ C'_2 \|A\psi\|_{L^\infty}) |B(\bar{x}, \varrho)|^{-\frac{n-2}{2n}} \leq \lambda^{1/2} (C''_1 + C''_2 \|A\psi\|_{L^\infty}) \varrho^{-\frac{n-2}{2n}} \end{aligned}$$

then there is $\tilde{x} \in B(\bar{x}, \varrho)$ such that

$$(2,14) \quad |u^\lambda(\tilde{x}) - u(\tilde{x})| \leq \lambda^{1/2} (C''_1 + C''_2 \|A\psi\|_{L^\infty}) \varrho^{-\frac{n-2}{2n}}.$$

From (2,14) we have

$$(2,15) \quad |u^\lambda(x) - u(x)| \leq \lambda^{1/2} (C''_1 + C''_2 \|A\psi\|_{L^\infty}) \varrho^{-\frac{n-2}{2n}} + C \varrho^\alpha$$

for $x \in B(\bar{x}, \varrho)$.

We choose $\varrho = \frac{1}{\lambda^{n-2(1-\alpha)}}$ then

$$(2,16) \quad |u^\lambda(x) - u(x)| \leq \overline{\lambda^{n-2(1-\alpha)}} (C''_1 + C''_2 \|A\psi\|_{L^\infty}) + C \lambda^{\frac{\alpha}{n-2(1-\alpha)}}.$$

We show now (2,10).

From the lemma 2.1 of [7] (the hypothesis of symmetry is unnecessary for this lemma) if $\psi \in W^{1,p}(\Omega)$, $n < p \leq +\infty$, there is a sequence $\{\psi_\eta\}$ such that

$$(2,17) \quad \|A \psi_\eta\|_{L^\infty} \leq C_3 \eta^{-1/2}$$

$$(2,18) \quad \|\psi_\eta - \psi\|_{L^\infty} \leq C_3 \eta^{1/2}$$

where C_3 is a constant dependent only on $\|\psi\|_{W^{1,p}}$.

Let be u_η^λ (u_η) the solution of the problem

$$(2,19) \quad A u_\eta^\lambda + \frac{1}{\lambda} (u_\eta^\lambda - \psi_\eta)^+ = f$$

$$(2,20) \quad \begin{aligned} \langle A u_\eta, v - u_\eta \rangle &\geq \langle f, v - u_\eta \rangle \\ \forall v \in H_0^1(\Omega), v &\leq \psi_\eta \\ u_\eta &\in H_0^1(\Omega), u_\eta \leq \psi_\eta. \end{aligned}$$

From [12] pg. 22, part III, we have

$$(2,21) \quad \|u_\eta^\lambda - u^\lambda\|_{L^\infty} \leq C_5 \eta^{1/2}$$

$$(2,22) \quad \|u_\eta - u\|_{L^\infty} \leq C_5 \eta^{1/2}.$$

From (2,16) (2,21) (2,22) we have

$$\begin{aligned} |u^\lambda(x) - u(x)| &\leq |u^\lambda(x) - u_\eta^\lambda(x)| + |u_\eta^\lambda(x) - u_\eta(x)| + |u_\eta(x) - u(x)| \\ &\leq \lambda^{\frac{\alpha}{n-2(1-\alpha)}} (C_1'' + C_2'' \eta^{-1/2}) + C \lambda^{\frac{\alpha}{n-2(1-\alpha)}} + 2 \eta^{1/2} \end{aligned}$$

We choose $\eta = \lambda^{\frac{\alpha}{2[n-2(1-\alpha)]}}$ then

$$|u^\lambda(x) - u(x)| \leq C_6 \lambda^{\frac{\alpha}{2[n-2(1-\alpha)]}}$$

Rem. 1 — We observe that the constants in the lemma 2 depend only on L^∞ — norm of the coefficients of A .

Lemma 3. *We consider the problems*

$$A^\varepsilon u_\varepsilon = f, \quad A u = f$$

for $f \in L^\infty(\Omega)$ we have

$$\|u_\varepsilon - u\|_{L^\infty} \leq C \varepsilon^{1/2}.$$

We consider $\theta = -\Delta^{-1} f$, $\theta \in W^{2,p}(\Omega)$. $\forall p$, for the lemma 2.1 of [7] there is a sequence θ_η such that

$$\|\theta_\eta - \theta\|_{w^{1,\infty}} \leq C_1 \eta^{1/2}$$

$$\|\Delta \theta_\eta\|_{w^{1,\infty}} \leq C_1 \eta^{-1/2}.$$

Let be $f_\eta = -\Delta \theta_\eta$: we have

$$\|f_\eta - f\|_{w^{-1,\infty}} \leq C_2 \eta^{1/2}$$

$$\|f_\eta\|_{w^{1,\infty}} \leq C_2 \eta^{-1/2}.$$

Let be u_ε^η (u^η) the solution of the problem

$$A^\varepsilon u_\varepsilon^\eta = f_\eta \quad (A u^\eta = f_\eta).$$

We have

$$\begin{aligned} \|u_\varepsilon - u\|_{L^\infty} &\leq \|u_\varepsilon - u_\varepsilon^\eta\|_{L^\infty} + \|u_\varepsilon^\eta - u^\eta\|_{L^\infty} + \|u^\eta - u\|_{L^\infty} \leq \\ &\leq 2 C'_1 \eta^{1/2} + C'_2 \varepsilon \eta^{-1/2} \end{aligned}$$

We choose $\eta = \varepsilon$ and we have the result.

3. - The penalised equation.

We suppose for the moment $f = 0$.

We consider the penalised problems

$$(3,1_\varepsilon) \quad A^\varepsilon u_\varepsilon^\lambda + \frac{1}{\lambda} (u_\varepsilon^\lambda - \psi)^+ = 0$$

$$(3.1) \quad A u^\lambda + \frac{1}{\lambda} (u^\lambda - \psi)^+ = 0.$$

Lemma 4. *We have*

$$\|u_\varepsilon^\lambda - u^\lambda\|_{L^\infty} \leq (K_1 + K_2 \lambda^{-1}) \|A \psi\|_{L^p} \varepsilon$$

$$n < p < +\infty.$$

We use for the proof the « multiple scales » method.

Let be $y = \frac{x}{\varepsilon}$ we have, [8] [9],

$$(3.2) \quad A^\varepsilon = \varepsilon^{-2} A_1 + \varepsilon^{-1} A_2 + A_3$$

where

$$A_1 = - \sum_{ij=1}^n a_{ij}(y) \frac{\partial^2}{\partial y_i \partial y_j} - \sum_{ij=1}^n \left(\frac{\partial}{\partial y_i} a_{ij}(y) \right) \frac{\partial}{\partial y_j} =$$

$$= - \sum_{ij=1}^n \frac{\partial}{\partial y_i} \left(a_{ij}(y) \frac{\partial}{\partial y_j} \right)$$

$$A_2 = - \sum_{ij=1}^n a_{ij}(y) \left(\frac{\partial^2}{\partial x_i \partial y_j} + \frac{\partial^2}{\partial x_j \partial y_i} \right) - \sum_{ij=1}^n \left(\frac{\partial}{\partial y_i} a_{ij}(y) \right) \frac{\partial}{\partial x_j} =$$

$$= - \sum_{i,j=1}^n \frac{\partial}{\partial y_j} \left(a_{ij}(y) \frac{\partial}{\partial x_i} \right) - \sum_{i,j=1}^n a_{ij}(y) \frac{\partial^2}{\partial x_i \partial y_j}$$

$$A_3 = - \sum_{i,j=1}^n a_{ij}(y) \frac{\partial^2}{\partial x_i \partial x_j}$$

Let be

$$A_1 \chi^j(y) = \sum_{i=1}^n \frac{\partial}{\partial y_i} (a_{ij}(y))$$

$$\chi^j(y) \text{ Y-periodic, } \int_{\gamma} \chi^j(y) dy = 0$$

we have $\chi^j(y) \in C^2(\bar{\Omega})$.

Let be

$$\chi_1(x; y) = \sum_{i=1}^n \chi^i(y) \frac{\partial u^\lambda}{\partial x_i}; \quad v_\varepsilon = u^\lambda + \varepsilon \chi_1 + \varepsilon^2 \chi_2$$

where $\chi_2(x; y)$ is not given for the moment.

We consider $w = u_\varepsilon^\lambda - v_\varepsilon$,

$$\begin{aligned} A^\varepsilon(w) + \frac{1}{\lambda} (u_\varepsilon^\lambda - \psi)^+ - \frac{1}{\lambda} (u^\lambda + \varepsilon \chi_1 + \varepsilon^2 \chi_2 - \psi)^+ &= \\ &= - (A_1 \chi_2 + A_2 \chi_1 + A_3 u^\lambda + \frac{1}{\lambda} (u^\lambda - \lambda)^+) - \\ &\quad - \varepsilon (A_3 \chi_1 + A_2 \chi_2) - \varepsilon^2 A_3 \chi_2 + \frac{1}{\lambda} (u^\lambda - \psi)^+ \\ &\quad - \frac{1}{\lambda} (u^\lambda + \varepsilon \chi_1 + \varepsilon^2 \chi_2 - \psi)^+ \end{aligned}$$

We choose $\chi_2(x; y)$ such that

$$(3,4) \quad A_1 \chi_2 + A_2 \chi_1 + A_3 u^\lambda + \frac{1}{\lambda} (u^\lambda - \psi)^+ = 0$$

$\chi_2(x; y)$ Y -periodic in y .

We observe that we have

$$\int_Y \left(A_2 \chi_1 + A_3 u^\lambda + \frac{1}{\lambda} (u^\lambda - \psi)^+ \right) dy = 0$$

then the problem (3,4) has a solution.

We observe that

$$\begin{aligned} (3,5) \quad \|A \psi\|_{L^p} \leq C_1 &\Rightarrow \frac{1}{\lambda} \|(u^\lambda - \psi)^+\|_{L^p} \leq C_1 \Rightarrow \\ &\Rightarrow \|u^\lambda\|_{w^{2,p}} \leq C_2 \cdot C_1 \Rightarrow \frac{1}{\lambda} \|(u^\lambda - \psi)^+\|_{w^{1,p}} \leq \frac{C_3 C_1}{\lambda} \\ &\Rightarrow \|u^\lambda\|_{w^{3,p}} \leq \frac{C_3 \cdot C_1}{\lambda} \quad (n < p \leq +\infty). \end{aligned}$$

From (3,5) we have

$$\|\chi_1(x; y)\|_{c^1(Y; w^2, P)} \leq \frac{C_5}{\lambda} \|A\psi\|_{L^P}$$

then

$$(3,6) \quad \|A_3 \chi_1(x; y)\|_{c(Y; L^P)} \leq \frac{C_5}{\lambda} \|A\psi\|_{L^P}$$

$$(3,7) \quad \|A_2 \chi_1(x; y)\|_{c(Y; W^1, P)} \leq \frac{C_7}{\lambda} \|A\psi\|_{L^P}.$$

From (3,4) (3,5) (3,7) we have

$$(3,8) \quad \|\chi_2(x; y)\|_{c^2(Y; w^1, P)} \leq \frac{C_8}{\lambda} \|A\psi\|_{L^P}$$

and, by multiplication for $v \in W^{1, p'}(\Omega)$ $1/p + 1/p' = 1$, we can easily show

$$(3,9) \quad \|\varepsilon A_3 \chi_2\left(x; \frac{x}{\varepsilon}\right)\|_{w^{-1, P}} \leq \frac{C_9}{\lambda} \|A\psi\|_{L^P}.$$

We observe now that

$$(3,10) \quad \begin{aligned} \left\| \frac{1}{\lambda} (u^\lambda - \psi)^+ - \frac{1}{\lambda} (u^\lambda + \varepsilon \chi_1 + \varepsilon^2 \chi_2 - \psi)^+ \right\|_{L^\infty} &\leq \\ &\leq \frac{\varepsilon}{\lambda} \|\chi_1\|_{L^\infty} + \frac{\varepsilon^2}{\lambda} \|\chi_2\|_{L^\infty} \\ &\leq C_{10} \left(\frac{\varepsilon + \varepsilon^2}{\lambda} \right) \|A\psi\|_{L^\infty} \end{aligned}$$

From (3,6) (3,7) (3,8) (3,9) (3,10) we have

$$(3,11) \quad A^\varepsilon(w) + \frac{1}{\lambda} (u^\lambda - \psi)^+ - \frac{1}{\lambda} (u^\lambda + \varepsilon \chi_1 + \varepsilon^2 \chi_2 - \psi)^+ = \theta$$

where

$$\|\theta\|_{w^{-1},P} \leq \left(C_{11} + \frac{C_{12}}{\lambda} \right) \|A\psi\|_{L^P} \varepsilon$$

By the same method used in [10], [13] for the proof of L^∞ – estimate in elliptic problems we have

$$(3,12) \quad \|w\|_{L^\infty} \leq \left(C'_{11} + \frac{C'_{12}}{\lambda} \right) \|A\psi\|_{L^P} \varepsilon$$

then from (3,12)

$$\|u_\varepsilon^\lambda - u^\lambda\|_{L^\infty} \leq \left(C''_1 + \frac{C'_{12}}{\lambda} \right) \|A\psi\|_{L^P} \varepsilon.$$

We consider now the case $f \neq 0$, $f \in L^P$, $n < p \leq +\infty$.
Let be

$$\sigma = A^{-1}f \quad \sigma^\varepsilon = (A^\varepsilon)^{-1}f$$

we have

$$\|\sigma^2 - \sigma\|_{L^\infty} \leq C_{13} \varepsilon^{1/2}$$

We observe that the problems

$$(3,13_\varepsilon) \quad A^\varepsilon u_\varepsilon^\lambda + \frac{1}{\gamma} (u_\varepsilon^\lambda - \psi)^+ = f$$

$$(3,13) \quad Au^\lambda + \frac{1}{\lambda} (u^\lambda - \psi)^+ = f$$

are equivalent to the problems

$$(3,14_\varepsilon) \quad A^\varepsilon v_\varepsilon^\lambda + \frac{1}{\lambda} (v_\varepsilon^\lambda - (\psi - \sigma^\varepsilon))^+ = 0$$

$$(3,14) \quad Av^\lambda + \frac{1}{\lambda} (v^\lambda - (\psi - \sigma))^+ = 0$$

where

$$(3,15) \quad v^\lambda = u^\lambda - \sigma \quad v_\varepsilon^\lambda = u_\varepsilon^\lambda - \sigma^\varepsilon .$$

We consider now the problem

$$(3,14') \quad A^\varepsilon w_\varepsilon^\lambda + \frac{1}{\lambda} (w_\varepsilon^\lambda - (\psi - \sigma))^+ = 0 .$$

From [12] pg 22. part III we have

$$\|v_\varepsilon^\lambda - w_\varepsilon^\lambda\|_{L^\infty} \leq C_{13} \varepsilon^{1/2}$$

and from (3,12) we have

$$\|w_\varepsilon^\lambda - v^\lambda\|_{L^\infty} \leq \left(C_{14} + \frac{C_{15}}{\lambda} \right) (\|A\psi\|_{L^P} + \|f\|_{L^P}) \varepsilon$$

then

$$(3,16) \quad \|u_\varepsilon^\lambda - u^\lambda\|_{L^\infty} \leq (C_{14} + \frac{C_{15}}{\lambda}) (\|A\psi\|_{L^P} + \|f\|_{L^P}) \varepsilon + 3 C_{13} \varepsilon^{1/2} .$$

4. - Prof. of Th. 3.

We consider the problem

$$(4,1_\varepsilon) \quad \langle A^\varepsilon u_\varepsilon, v - u_\varepsilon \rangle \geq \langle f, v - u_\varepsilon \rangle$$

$$\forall v \in H_0^1(\Omega) \quad v \leq \psi$$

$$u_\varepsilon \in H_0^1(\Omega) \quad v_\varepsilon \leq \psi$$

$$(4,1) \quad \langle A u, v - u \rangle \geq \langle f, v - u \rangle$$

$$\forall v \in H_0^1(\Omega) \quad v \leq \psi$$

$$u \in H_0^1(\Omega) \quad u \leq \psi$$

and the corresponding penalised problems

$$(4,2) \quad A^\varepsilon u_\varepsilon^\lambda + \frac{1}{\lambda} (u_\varepsilon^\lambda - \psi)^+ = f$$

$$(4,2) \quad A u^\lambda + \frac{1}{\lambda} (u^\lambda - \psi)^+ = f$$

We suppose for the moment

$$\|A^\varepsilon \psi\|_{L^\infty}, \|A \psi\|_{L^\infty} \leq C.$$

We have

$$\|u_\varepsilon^\lambda - u_\varepsilon\|_{L^\infty} \leq C_1 \lambda^{\frac{1}{n-2(1-\alpha)}}$$

$$\|u^\lambda - u\|_{L^\infty} \leq C_1 \lambda^{\frac{1}{n-2(1-\alpha)}}$$

$$\|u_\varepsilon^\lambda - u^\lambda\|_{L^\infty} \leq \left(C_2 + \frac{C_3}{\lambda}\right) \varepsilon + C_4 \varepsilon^{1/2}$$

then

$$\|u_\varepsilon - u\|_{L^\infty} \leq 2C_1 \lambda^{\frac{1}{n-2(1-\alpha)}} + \left(C_2 + \frac{C_3}{\lambda}\right) \varepsilon + C_4 \varepsilon^{1/2}.$$

We choose $\lambda = \varepsilon^{\frac{n-2+2\alpha}{n-2+3\alpha}}$ then

$$\|u_\varepsilon - u\|_{L^\infty} \leq C_5 \varepsilon^{\frac{\alpha}{n-2+3\alpha}}.$$

Let be now $\psi \in W^{1,p}(\Omega)$, $p > n$; there is $\psi_\varepsilon^\eta, \psi^\eta$ ($\eta > 0$) such that, [7],

$$\|\psi_\varepsilon^\eta - \psi\|_{L^\infty} \leq C_6 \eta^{1/2}$$

$$\|\psi^\eta - \psi\|_{L^\infty} \leq C_6 \eta^{1/2}$$

$$\|A^\varepsilon \psi_\varepsilon^\eta\|_{L^\infty} \leq C_7 \eta^{-1/2}$$

$$\|A \psi^\eta\|_{L^\infty} \leq C_7 \eta^{1/2}.$$

We consider the problems

$$(4,3_\eta) \quad \langle A^\varepsilon u_\varepsilon^\eta, v - u_\varepsilon^\eta \rangle \geq \langle f, v - u_\varepsilon^\eta \rangle$$

$$\forall v \in H_0^1(\Omega), \quad v \leq \psi_\varepsilon^\eta$$

$$u_\varepsilon^\eta \in H_0^1(\Omega), \quad u_\varepsilon^\eta \leq \psi_\varepsilon^\eta$$

$$(4,3) \quad \langle A u^\eta, v - u^\eta \rangle \geq \langle f, v - u^\eta \rangle$$

$$\forall v \in H_0^1(\Omega), \quad v \leq \psi^\eta$$

$$u^\eta \in H_0^1(\Omega), \quad u^\eta \leq \psi^\eta.$$

From the above we have

$$\|u_\varepsilon^\eta - u^\eta\|_{L^\infty} \leq \left(C_8 + \frac{C_9}{\eta^{1/2}} \right) \varepsilon^{\frac{\alpha}{n-2+3\alpha}}$$

$$\|u_\varepsilon^\eta - u_\varepsilon\|_{L^\infty}, \|u^\eta - u\|_{L^\infty} \leq C_7 \eta^{1/2}$$

then

$$\|u_\varepsilon - u\|_{L^\infty} \leq (C_8 + C_{9/\eta^{1/2}}) \varepsilon^{\frac{\alpha}{n-2+3\alpha}} + 2 C_7 \eta^{1/2}.$$

We choose $\eta = \frac{\alpha}{\lambda^{n-2+3\alpha}}$ then

$$\|u_\varepsilon - u\|_{L^\infty} \leq C_{10} \varepsilon^{\frac{\alpha}{2[n-2+3\alpha]}}$$

Rem. 1 — We observe that the dependence on n of the estimate of th. 3 derive from the dependence on n of the estimate in the lemma 2 ; then from (2,13) repeating our calculations we have

$$\|u_\varepsilon - (u + \varepsilon \chi_1)\|_{H_0^1} \leq C \varepsilon^{1/4}$$

$$\text{if } \|A^\varepsilon \psi\|_{L^2}, \|A \psi\|_{L^2} \leq C.$$

REFERENCES

- [1] BABUSKA I., *Solutions of problems with interfaces and singularities*, Int. Fluid Dyn. Appl. Math., April 1974.
- [2] BABUSKA I., *Solution of interference problems by homogenisation*, Inst. Fluid Dyn. Sppl. Math., March 1974.
mès asymptotiques stationnaires, C. R. Acad. Sc. Paris.
- [3] BENSOUSSAN A., LIONS J. L., PAPANICOLAOU G., *Sur quelques phénomènes asymptotiques stationnaires*, C. R. Acad. Sc. Paris.
- [4] BIROLI M., *Sur la G-convergence pour des inéquations quasi-variationnelles*, C. R. Acad. Sc. Paris to appear.
- [5] BOCCARDO L., CAPUZZO-DOLCETTA I., *G-Convergenza per un problema di Dirichlet unilatero*, Boll. U.M.I. 4, 12, 115-123, 1975.
- [6] BOCCARDO L., MARCELLINI P., *Sulla convergenza delle soluzioni di disequazioni quasi-variazionali*, Inst. Mat. Ulisse Dini, Firenze, 1975.
- [7] DE GIORGI, SPAGNOLO S., *Sulla convergenza degli integrali dell'energia per operatori ellittici del 2° ordine*, Boll. UMI, 4, 8, 391-411, 1973.
- [8] LIONS J. L., *Homogenisation*, Cours au College de France 1975-76, Paris.
- [9] LIONS J. L., *Asymptotic behaviour of solution of variational inequalities with highly oscillating coefficients*, Applications of method of Functional Analysis to problems in Mechanics, Lec. Notes in Math., 503, 30-56.
- [10] LADY-ZENSKAIA O. A., URAL'CEVA N. N., *Equations aux dérivées partielles du type elliptique*, Dunod 1968.
- [11] MURAT F., *Sur l'homogenisation d'inéquations elliptiques* Lab. ss. 189 Paris VI, n. 76013.
- [12] PUEL J. P., *Thèse*, 1975 Paris.
- [13] STAMPACCHIA G., *Equations elliptiques du second ordre à coefficients discontinus*, Les presses de l'Univ. de Montreal 1965.
- [14] TARTAR L., *Cours Peccot* 1977, Avril.

Manoscritto pervenuto in redazione il 20 aprile 1977.