

RENDICONTI *del* SEMINARIO MATEMATICO *della* UNIVERSITÀ DI PADOVA

MARTIN HUBER

A simple proof for a theorem of chase

Rendiconti del Seminario Matematico della Università di Padova,
tome 74 (1985), p. 45-49

[<http://www.numdam.org/item?id=RSMUP_1985__74__45_0>](http://www.numdam.org/item?id=RSMUP_1985__74__45_0)

© Rendiconti del Seminario Matematico della Università di Padova, 1985, tous droits réservés.

L'accès aux archives de la revue « Rendiconti del Seminario Matematico della Università di Padova » (<http://rendiconti.math.unipd.it/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

*Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques*
<http://www.numdam.org/>

A Simple Proof for a Theorem of Chase.

MARTIN HUBER (*)

*Dedicated to Professor Laszlo Fuchs
on the occasion of his sixtieth birthday.*

In this note all groups are abelian. A group G is called a *W-group* (Whitehead group) if $\text{Ext}(G, \mathbb{Z}) = 0$. Whitehead's problem asks whether every *W-group* is free. This problem has been solved by Shelah [S1, S2] in the sense that the answer depends on the underlying set theory. The deepest result on Whitehead's problem prior to Shelah's work is Chase's theorem, saying that under $2^{\aleph_0} < 2^{\aleph_1}$ every *W-group* G is *strongly $\aleph_1$ -free* i.e., G is $\aleph_1$ -free and every countable subset of G is contained in a countable $\aleph_1$ -pure subgroup of G . In fact, Chase proved the following somewhat stronger result.

THEOREM [*C*; Thm. 2.3]. Assume $2^{\aleph_0} < 2^{\aleph_1}$. If G is a torsion-free group such that $\text{Ext}(G, \mathbb{Z})$ is torsion, then G is strongly $\aleph_1$ -free.

Among other nontrivial tools, Chase's proof uses the derived functors of the inverse limit. An entirely different proof has been given in [EH]; in there the above theorem is a corollary to a more general result involving the set-theoretic principle «weak diamond». That in the given situation weak diamond is equivalent to $2^{\aleph_0} < 2^{\aleph_1}$ is a nontrivial result of set theory due to Devlin and Shelah [DS].

In this note an alternative proof is given for the above theorem which the author has found a few years ago (cf. p. 167, Remark 2 in [EH]). Since the present proof is both shorter and more elementary

(*) Indirizzo dell'A.: Mathematisches Institut der Universität, Albertstrasse 23 b, D-7800 Freiburg i. Br., Rep. Fed. Tedesca.

than those in [C] or [EH], I think it is well worth being published.

We shall make use of the description of Ext in terms of factor sets. Recall [F; § 49] that $\text{Ext}(G, \mathbb{Z})$ can be identified with $\text{Fact}(G, \mathbb{Z})/\text{Trans}(G, \mathbb{Z})$ where $\text{Fact}(G, \mathbb{Z})$ is the group of factor sets on G to $\mathbb{Z}$ and $\text{Trans}(G, \mathbb{Z})$ is the subgroup of transformation sets. A transformation set is a factor set δg , given by

$$\delta g(x, y) = g(x) - g(x + y) + g(y),$$

where $g: G \rightarrow \mathbb{Z}$ is a map with $g(0) = 0$.

Our proof is divided into three steps, the first of which consists in establishing a lemma.

LEMMA. Let K be a subgroup of H such that $\text{Ext}(H/K, \mathbb{Z})$ contains an element of infinite order. Then there is an element $\tilde{f} \in \text{Fact}(H, \mathbb{Z})$ extending $0 \in \text{Fact}(K, \mathbb{Z})$ such that for all $n > 0$ there is no map $g: H \rightarrow \mathbb{Z}$ for which $g|_K = 0$ and $n\tilde{f} = \delta g$.

PROOF. Let $h \in \text{Fact}(H/K, \mathbb{Z})$ represent an infinite order element of $\text{Ext}(H/K, \mathbb{Z})$. Let $\pi: H \rightarrow H/K$ denote the natural map and define $\tilde{f} \in \text{Fact}(H, \mathbb{Z})$ by

$$\tilde{f}(x, y) = h(\pi x, \pi y).$$

We claim that $\tilde{f}$ satisfies the required conditions. Clearly $\tilde{f}$ extends $0 \in \text{Fact}(K, \mathbb{Z})$. Suppose that there is an $n > 0$ and a function $g: H \rightarrow \mathbb{Z}$ such that $g|_K = 0$ and $n\tilde{f} = \delta g$. Then for each pair $(x, y) \in H \times K$ we have

$$g(x) - g(x + y) = \delta g(x, y) = n\tilde{f}(x, y) = nh(\pi x, 0) = 0.$$

Therefore g is actually a function on H/K i.e., there is a $\bar{g}: H/K \rightarrow \mathbb{Z}$ with $\bar{g}\pi = g$. It follows readily that $nh = \delta\bar{g}$, contradicting the choice of h . $\square$

The second and main task is to prove the following proposition. This is in fact contained in [C; Thm. 1.8]; we notice that the use of factor sets simplifies its proof considerably. By $r_0(A)$ we denote the torsion-free rank of the group A .

PROPOSITION. Let G be a nonzero torsion-free group that does not contain any subgroup H of smaller cardinality such that G/H is $\aleph_1$ -free. Then $r_0(\text{Ext}(G, \mathbb{Z})) = 2^{|G|}$.

PROOF. Let κ denote the cardinality of G . Suppose first that $\kappa = \aleph_0$. By hypothesis G is not free; thus, as is well known, $r_0(\text{Ext}(G, \mathbb{Z})) = 2^{\aleph_0}$ (cf. [C; L. 1.4]). Now suppose that κ is uncountable. The hypothesis enables us to define inductively an increasing chain $\{G_\alpha: \alpha < \kappa\}$ of pure subgroups of G with the following properties:

- (o) $G_0 = 0$;
- (i) $G_\alpha = \bigcup_{\beta < \alpha} G_\beta$ if α is a limit ordinal;
- (ii) for all $\alpha < \kappa$, $G_{\alpha+1}/G_\alpha$ is nonzero and countable;
- (iii) for all $\alpha < \kappa$, $\text{Hom}(G_\alpha, \mathbb{Z}) = 0$.

We have to choose $G_{\alpha+1}$ as a pure subgroup of G which contains G_α such that $G_{\alpha+1}/G_\alpha$ is nonzero, countable, and satisfies $\text{Hom}(G_{\alpha+1}/G_\alpha, \mathbb{Z}) = 0$. That this is possible follows from the facts that G/G_α contains a nonfree countable subgroup and that every countable group is the direct sum of a free group and a group C satisfying $\text{Hom}(C, \mathbb{Z}) = 0$. (For the latter see [F; Cor. 19.3]). For all $\alpha < \kappa$, since G_α is pure in G , it follows that $\text{Ext}(G_{\alpha+1}/G_\alpha, \mathbb{Z})$ contains an element of infinite order.

Let $G' = \bigcup_{\alpha < \kappa} G_\alpha$. We will prove that $r_0(\text{Ext}(G', \mathbb{Z})) = 2^\kappa$. To this end we assign to each sequence $\eta \in 2^\kappa$, $\alpha < \kappa$, a factor set f_η on G_α such that

(a) if ξ is an initial segment of η , then f_η extends f_ξ ;

(b) if $\xi \neq \eta$ have the same domain α , then $f_\xi - f_\eta$ represents an element of $\text{Ext}(G_\alpha, \mathbb{Z})$ of infinite order.

The definition is by induction on the domain of η . Suppose f_η has been defined, where $\eta \in 2^\alpha$. Let $\eta^{\frown}\langle 0 \rangle$, $\eta^{\frown}\langle 1 \rangle$ denote the two extensions of η in $2^{\alpha+1}$. Extend f_η to $f_{\eta^{\frown}\langle 0 \rangle}$ arbitrarily (possible, cf. [HHS; Lemma 1]) and then define $f_{\eta^{\frown}\langle 1 \rangle}$ according to the lemma above, so that there is no $g: G_{\alpha+1} \rightarrow \mathbb{Z}$ for which $g \upharpoonright G_\alpha = 0$ and $n(f_{\eta^{\frown}\langle 1 \rangle} - f_{\eta^{\frown}\langle 0 \rangle}) = \delta g$ for any $n > 0$. We claim that $f_{\eta^{\frown}\langle 1 \rangle} - f_{\eta^{\frown}\langle 0 \rangle}$ represents an element of $\text{Ext}(G_{\alpha+1}, \mathbb{Z})$ of infinite order. Suppose to the contrary that there is a map $h: G_{\alpha+1} \rightarrow \mathbb{Z}$ such that for some $n > 0$, $n(f_{\eta^{\frown}\langle 1 \rangle} - f_{\eta^{\frown}\langle 0 \rangle}) = \delta h$. Then of course $(\delta h) \upharpoonright G_\alpha = 0$; hence $h \upharpoonright G_\alpha$ is a homomorphism. But by definition of G_α , $\text{Hom}(G_\alpha, \mathbb{Z}) = 0$, whence we obtain a contradiction.

It follows that the factor sets f_η , $\eta \in \bigcup_{\alpha < \kappa} 2^\alpha$, satisfy (a) and (b), and in the limit we obtain 2^κ factor sets on G' to $\mathbf{Z}$ which represent pairwise different elements of $\text{Ext}(G', \mathbf{Z})$ modulo torsion. We conclude that

$$2^{|G|} \leq r_0(\text{Ext}(G, \mathbf{Z})) \leq 2^{|G|},$$

the latter inequality being obvious. $\square$

The final step is to deduce the Theorem from the Proposition: Assume that $2^{\aleph_0} < 2^{\aleph_1}$ and let G be an uncountable torsion-free group which is not strongly $\aleph_1$ -free. Thus G contains a countable subgroup H such that for every countable subgroup K containing H , G/K is not $\aleph_1$ -free. Therefore by the proposition we have $r_0(\text{Ext}(G/H, \mathbf{Z})) \geq 2^{\aleph_1}$. Considering exactness of the sequence

$$\text{Hom}(H, \mathbf{Z}) \rightarrow \text{Ext}(G/H, \mathbf{Z}) \rightarrow \text{Ext}(G, \mathbf{Z})$$

we infer that $r_0(\text{Ext}(G, \mathbf{Z})) \geq 2^{\aleph_1}$, since $|\text{Hom}(H, \mathbf{Z})| = 2^{\aleph_0}$ and $2^{\aleph_0} < 2^{\aleph_1}$. Hence, in particular, $\text{Ext}(G, \mathbf{Z})$ is not a torsion group. $\square$

Finally we wish to mention that the Theorem is not true without the hypothesis of $2^{\aleph_0} < 2^{\aleph_1}$. It follows from results in [S1] and [S3] that in a model of Martin's Axiom plus the denial of the Continuum Hypothesis there exist *W-groups* which are not strongly $\aleph_1$ -free.

REFERENCES

- [C] S. CHASE, *On group extensions and a problem of J. H. C. Whitehead*, in *Topics in Abelian Groups*, pp. 173-197, Chicago: Scott, Foresman and Co., 1963.
- [DS] K. DEVLIN - S. SHELAH, *A weak version of $\diamond$ which follows from $2^{\aleph_0} < 2^{\aleph_1}$* , Israel J. Math., **29** (1978), pp. 239-247.
- [EH] P. EKLOF - M. HUBER, *On the rank of Ext*, Math. Z., **174** (1980), pp. 159-185.
- [F] L. FUCHS, *Infinite Abelian Groups*, Vol. I, London, New York, Academic Press, 1970.
- [HHS] H. HILLER - M. HUBER - S. SHELAH, *The structure of $\text{Ext}(A, \mathbf{Z})$ and $V = L$* , Math. Z., **162** (1978), pp. 39-50.

- [S1] S. SHELAH, *Infinite abelian groups. Whitehead problem and some constructions*, Israel J. Math., **18** (1974), pp. 243-256.
- [S2] S. SHELAH, *A compactness theorem for singular cardinals, free algebras, Whitehead problem and transversals*, Israel J. Math., **21** (1975), pp. 319-349.
- [S3] S. SHELAH, *On uncountable abelian groups*, Israel J. Math., **32** (1979), pp. 311-330.

Manoscritto pervenuto in redazione il 9 maggio 1984.