

# RENDICONTI *del* SEMINARIO MATEMATICO *della* UNIVERSITÀ DI PADOVA

JAN AMBROSIEWICZ

## **On the squares of sets of linear groups**

*Rendiconti del Seminario Matematico della Università di Padova*,  
tome 75 (1986), p. 253-256

[http://www.numdam.org/item?id=RSMUP\\_1986\\_\\_75\\_\\_253\\_0](http://www.numdam.org/item?id=RSMUP_1986__75__253_0)

© Rendiconti del Seminario Matematico della Università di Padova, 1986, tous droits réservés.

L'accès aux archives de la revue « Rendiconti del Seminario Matematico della Università di Padova » (<http://rendiconti.math.unipd.it/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

*Article numérisé dans le cadre du programme  
Numérisation de documents anciens mathématiques*  
<http://www.numdam.org/>

## On the Squares of Sets of Linear Groups.

JAN AMBROSIEWICZ (\*)

The covering problem ( $CC \supseteq G$ ,  $C$ -conjugacy class) for the group  $PSL(2, q)$  was described in the paper [1]. In this paper we investigate squares of sets  $K_w = \{g \in G: |g| = w\}$  of elements of the same order of linear groups. We will say that the group  $G$  has the property  $W$  if for each  $w$ , the set  $K_w K_w$  is a subgroup of  $G$  (see [2]). We have proved that the set  $K_2 K_2$  is not a subgroup in any groups  $GL(n, K)$ ,  $SL(n, K)$ ,  $PSL(n, K)$  for  $n \geq 3$ . We have proved, too, that  $K_2 K_2 \leq GL(2, K)$ ,  $K_2 K_2 \leq SL(2, K)$  for  $\text{char } K = 2$  and  $K_2 K_2 \not\leq GL(2, K)$  for  $\text{char } K \neq 2$ . The set  $K_2 K_2$  of the group  $PSL(2, q)$  is described in Theorem 5.

The following Lemma will be useful in the sequel.

**LEMMA 1.** Let  $G$  be a group. An element  $g \in K_2^m$  ( $m \geq 2$ ) if and only if there is an element  $x \in K_2^{m-1}$ ,  $x \neq g^{-1}$  such that  $(gx)^2 = 1$ .

**PROOF.** If  $g = s_1 s_2 \dots s_m$ ,  $s_i \in K_2$  then  $g^{-1} = x g x$  where  $x = s_m s_{m-1} \dots s_2$ . Conversely, if  $(gx)^2 = 1$  and  $x \in K_2^{m-1}$  then  $g = x^{-1}(g^{-1}x^{-1})$  and  $(g^{-1}x^{-1})^2 = g^{-1}x^{-1}g^{-1}x^{-1} = g^{-1}g = 1$ . Since  $g^{-1}x^{-1} \neq 1$  by assumption, hence  $g \in K_2^m$ .

**THEOREM 1.** The groups  $GL(n, K)$ ,  $SL(n, K)$  and  $PSL(n, K)$  with  $n \geq 3$  over any field  $K$  have not property  $W$ .

**PROOF.** Let us consider all possible cases: (i)  $|K| > 3$ , (ii)  $|K| = 2$ , (iii)  $|K| = 3$ .

(\*) Indirizzo dell'A.: Institute of Mathematics ul. Dzierzynskiego 15 m 99, 15-099 Białystok - Polonia.

Ad (i). In this case there is an element  $u$  such that  $u^2 \neq 1$ .

If  $A = \text{diag}[u^{-1}, u, 1]$ ,  $B = \text{diag}[1, u, u^{-1}]$  with  $u^2 \neq 1$  then  $A, B \in K_2K_2$ ,  $AB \notin K_2K_2$  by Lemma 1 ( $m = 2$ ). Using this fact we can see that quasideagonal matrices  $[A, 1, \dots, 1]$ ,  $[B, 1, \dots, 1]$  belong to the set  $K_2K_2$  but their product do not belong to the set  $K_2K_2$ . In the remain cases we act similarly, namely in the case (ii) one can take

$$A = \begin{bmatrix} 1 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix}, \quad B = \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 1 & 1 \end{bmatrix}$$

and in the case (iii)

$$A = \begin{bmatrix} 1 & 0 & 2 \\ 0 & 0 & 2 \\ 0 & 1 & 0 \end{bmatrix}, \quad B = \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 2 & 2 \end{bmatrix}.$$

Since  $A_8 \simeq PSL(4, 2)$ ,  $G_{168} \simeq PSL(3, 2)$  (see [4]) then by Theorem 1 the simple group  $A_8$  and  $G_{168}$  have not property  $W$ .

REMARK 1. If  $X \in K_2 \subset GL(2, K)$  and  $\text{char } K = 2$  then  $\det X = 1$ .

THEOREM 2. If  $\text{char } K = 2$  and  $|K| > 2$  then  $A \in K_2K_2 \subset GL(2, K)$  if and only if  $\det A = 1$  i.e.  $K_2K_2 = SL(2, K)$ .

PROOF. Taking into consideration Corollary 4.7 p. 360 of [3] it is sufficient enough to show our proposition for matrices  $A = N_1 = [0, 1; -a_2, -a_1]$ ,  $A = N_2 = [k_1, 0; 0, k_2]$ .

By Lemma 1

$$(1) \quad N_i \in K_2K_2 \Leftrightarrow \exists_{\substack{X \in K_2 \\ X \neq N_i}} (XN_i = N_i^{-1}X, \det X \neq 0), \quad i = 1, 2.$$

If  $N_i \in K_2K_2$  then  $\det N_i = 1$  by Remark 1. If  $\det N_1 = 1$  then the equation  $XN_1 = N_1^{-1}X$  is satisfied by  $X = [0, 1; 1, 0]$  for  $a_1 \neq 0$  and  $X = [u, u+1; u+1, u]$  ( $u \neq 0, u \neq 1$ ) for  $a_1 = 0$ . Therefore  $N_1 \in K_2K_2$ . The equation  $XN_2 = N_2^{-1}X$  has a solution  $X \in GL(2, K)$  if and only if  $k_1k_2 = 1$  i.e.  $\det N_2 = 1$ . We can take  $X = [0, 1; 1, 0]$  which satisfies other condition of (1).

REMARK 2. If  $|K| = 2$  then  $K_2K_2 \leq GL(2, K) = SL(2, K) = PSL(2, K)$  what one can verify by calculations.

COROLLARY 2.1. If  $\text{char } K = 2$  and  $|K| > 2$  then  $K_2K_2 = PSL(2, K)$ . The proof results from Theorem 2 and from the equality  $SL(2, K) = PSL(2, K)$ .

THEOREM 3. If  $\text{char } K \neq 2$  then  $K_2K_2 \not\leq GL(2, K)$ .  
Indeed, by Lemma 1 and by easy calculations we have

$$[-1, 0; 0, 1], [0, 1; -1, -a_1] \in K_2K_2 (a_1 \neq 0)$$

and

$$[-1, 0; 0, 1] \cdot [0, 1; -1, -a_1] = [0, -1; -1, -a_1] \notin K_2K_2.$$

Theorem 4. The set  $K_2K_2$  is a subgroup of the group  $SL(2, K)$ .

PROOF. If  $\text{char } K = 2$  then Theorem 4 results from Remark 2 and from Theorem 2. If  $\text{char } K \neq 2$  then only the matrix  $[-1, 0; 0, -1]$  has the order 2 what one can verify by calculations.

THEOREM 5. (a) If in the field  $K$  ( $\text{char } K \neq 2$ ) the element  $-1$  is a square then  $K_2K_2 = PSL(2, K)$ .

(b) If in the field  $K$  ( $\text{char } K \neq 2, |K| \neq 3$ ) the element  $-1$  is not a square then  $K_2K_2 \not\leq PSL(2, K)$ .

PROOF. (a) One can verify that  $K_2 = \{[a, b; c, -a], a^2 + bc = -1\}$  in the group  $PSL(2, K)$ . By Lemma 1 we have

$$(2) \quad A \in K_2K_2 \iff \exists_{\substack{T \in K_2 \\ T \neq A}} A^{-1} = T^{-1}AT.$$

Let  $T = [x, y; z, -x], x^2 + yz = -1, A = [a_{11}, a_{12}; a_{21}, a_{22}]$ . The condition (2) is equivalent to the solvability of the system equations

$$(3) \quad \begin{cases} x^2 + yz = -1 \\ a_{21}y + a_{12}z = (a_{22} - a_{11})x \end{cases}$$

or to the solvability of the equation

$$(4) \quad [(a_{11} + a_{22})^2 - 4]x^2 + 4a_{12}a_{21} = u^2 \text{ if } a_{12} \neq 0 \text{ or } a_{21} \neq 0.$$

One can verify that in (2) we can take  $T = [0, y; y^{-1}, 0]$  if  $a_{12} = a_{21} = 0$ ;  $T = [x, (a_{22} - a_{11})a_{21}^{-1}x; 0, -x]$ ,  $x^2 = -1$  if  $a_{21} \neq 0$  and  $T = [x, 0; (a_{22} - a_{11})a_{12}^{-1}x, -x]$ ,  $x^2 = -1$  if  $a_{12} \neq 0$ . Therefore  $K_2K_2 = PSL(2, K)$ .

(b) The matrices  $A = [-1, 1; 1, -2]$ ,  $T = [0, 1; -1, 0]$  fulfil the condition (2). Therefore  $A \in K_2K_2$  and  $A \neq I$ .

If  $A = [0, 1; -1, \pm 2]$  then the equation (4) has no solution and  $A \notin K_2K_2$ . Since  $K_2K_2$  is a normal subset and in this case  $PSL(2, K)$  is a simple group then  $K_2K_2 \not\leq PSL(2, K)$ .

REMARK 3. If  $|K| = 3$  then  $K_2K_2$  is a subgroup of  $PSL(2, K)$  what can be verified by calculations.

COROLLARY 5.1.  $K_2K_2 = A_6$  in the alternating group  $A_6$ .

The proof results from the fact  $A_6 \simeq PSL(2, K)$  (see [4]) and from (a) of Theorem 5.

#### REFERENCES

- [1] J. L. BRENNER - L. CARLITZ, *Covering theorems for finite nonabelian simple groups, III*, Rend. Sem. Mat. Uni. Padova, **55** (1976).
- [2] E. AMBROSIEWICZ, *On groups having the property W*, Demonstratio Mathematica, **10** (1977), pag. 2.
- [3] T. W. HUNGEERFORD, *Algebra*, Springer-Verlag, New York, Heidelberg, Berlin, 1974.
- [4] B. HUPPERT, *Endliche Gruppen I*, Springer-Verlag, Berlin, Heidelberg, New York, 1967.

Manoscritto pervenuto in redazione l'8 dicembre 1984 ed in forma modificata il 26 marzo 1985.