

RENDICONTI *del* SEMINARIO MATEMATICO *della* UNIVERSITÀ DI PADOVA

GIUSEPPE DI FAZIO

Hölder-continuity of solutions for some Schrödinger equations

Rendiconti del Seminario Matematico della Università di Padova,
tome 79 (1988), p. 173-183

http://www.numdam.org/item?id=RSMUP_1988__79__173_0

© Rendiconti del Seminario Matematico della Università di Padova, 1988, tous droits réservés.

L'accès aux archives de la revue « Rendiconti del Seminario Matematico della Università di Padova » (<http://rendiconti.math.unipd.it/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

*Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques*
<http://www.numdam.org/>

Hölder-Continuity of Solutions for Some Schrödinger Equations.

GIUSEPPE DI FAZIO (*)

0. Introduction.

Recently the local regularity properties for solutions of Schrödinger equations of the form

$$(*) \quad Lu \equiv - (a_{ij}u_{x_i})_{x_j} = Vu$$

have been studied by many authors (see e.g. [A-S], [D-M], [C-F-G], [C-F-Z]) allowing V to be a very singular potential, precisely $V \in S$, the Stummel-Kato class (see definition 1.1).

Under this assumption in [C-F-G] was established a Harnack inequality and proved a local continuity result for solutions of (*).

It is easy to see that if Ω is an open bounded set in R^n then $L^p(\Omega) \subseteq S$ for $p > n/2$; hence the result in [C-F-G] generalizes the well known Hölder estimates by Stampacchia [ST], Ladizhenskaia [L-U] etc.

We stress that high integrability of V does not play an essential role.

In fact also the Morrey space $L^{1,\lambda}(\Omega)$ is contained in S for $\lambda > n - 2$ and being in $L^{1,\lambda}(\Omega)$, for any $0 < \lambda < n$, does not imply any extra integrability (see e.g. the examples in [P2]).

In this paper we assume V in $L^{1,\lambda}(\Omega)$ ($\lambda > n - 2$) and prove local hölder-continuity for solutions of (*) hence, in this special situation, we improve the continuity result in [C-F-G].

(*) Indirizzo dell'A.: Via del Canalicchio 9 - 95030 Tremestieri Etneo (Catania), Italy.

Our technique is very close to the one in [C-F-G] heavily relying on the exploitation of well known estimates for the Green function of L .

There is however a technical difficulty.

It is impossible to use the usual C^∞ -approximation for L and V (as in [C-F-G]) because functions in Morrey spaces are not close, in general, to bounded functions in $L^{1,\lambda}(\Omega)$ (see [P1] p. 22 for an example of an $L^{1,\lambda}(\Omega)$ function with distance from $L^\infty(\Omega)$ equal to 1). We overcame this difficulty by developing a representation formula for solutions of (*) that extends classical results on the Green function (see e.g. [ST]).

1. Some function spaces.

Let Ω be an open bounded set of R^n ($n > 3$).

We will need some mild regularity assumption to be satisfied by $\partial\Omega$ e.g.

$$\exists A \in]0, 1[: |\Omega_r(x)| \leq A |B_r(x)| \quad \forall x \in \partial\Omega$$

where $r: 0 < r < \text{diam}(\Omega)$ ⁽¹⁾.

DEFINITION 1.1 (*Stummel-Kato class*). We say that $V: \Omega \rightarrow R$ belongs to the Stummel-Kato class S iff there exists a non decreasing function $\eta(r) > 0$ with $\lim_{r \rightarrow 0} \eta(r) = 0$ such that

$$(1.1) \quad \text{Sup}_{x \in \Omega} \int_{\Omega_r(x)} |V(y)| |x - y|^{2-n} dy \leq \eta(r)$$

Obviously $S \subseteq L^1(\Omega)$.

DEFINITION 1.2 (*Morrey spaces*). $L^{1,\lambda}(\Omega)$ ($0 < \lambda < n$) is the space of functions $f \in L^1(\Omega)$ such that

$$\|f\|_{L^{1,\lambda}(\Omega)} =: \text{Sup}_{\substack{x \in \Omega \\ r > 0}} r^{-\lambda} \int_{\Omega_r(x)} |f(y)| dy < +\infty.$$

⁽¹⁾ $|E|$ denotes the Lebesgue measure of a measurable subset E of R^n :

$$B_r(x) =: \{y \in R^n : |x - y| < r\}; \quad \Omega_r(x) =: \Omega \cap B_r(x).$$

LEMMA 1.1. *If u belongs to $L^{1,\lambda}(\Omega)$ ($n - 2 < \lambda < n$) then u belongs to the Stummel-Kato class and*

$$\int_{\Omega_r(x)} |u(y)| |x - y|^{2-n} dy \leq Cr^{\lambda-n+2} \|u\|_{L^{1,\lambda}(\Omega)}$$

where C depends only on λ and n .

Indeed,

$$\begin{aligned} \int_{\Omega} |u(y)| |x - y|^{2-n} dy &= \sum_{k=0}^{+\infty} \int_{\Omega \cap \{r/2^{k+1} \leq |x-y| < r/2^k\}} |u(y)| |x - y|^{2-n} dy \leq \\ &\leq \sum_{k=0}^{+\infty} (r2^{-k-1})^{2-n} \int_{\Omega_{r/2^k}(x)} |u(y)| dy \leq r^{\lambda-n+2} C \|u\|_{L^{1,\lambda}(\Omega)}. \end{aligned}$$

REMARK 1.1:

$$L^{1,\lambda}(\Omega) \subseteq \mathcal{S} \subseteq L^{1,\mu}(\Omega) \quad \text{where } 0 < \mu \leq n - 2 < \lambda < n.$$

Indeed the inclusion $L^{1,\lambda}(\Omega) \subseteq \mathcal{S}$ is an immediate consequence of Lemma 1.1 and the other inclusion is obvious.

We now recall the definitions of the Sobolev spaces $H^{1,p}(\Omega)$, $H_0^{1,p}(\Omega)$ and $H^{-1,p}(\Omega)$.

DEFINITION 1.3. *We say that u belongs to $H^{1,p}(\Omega)[\widehat{H}_{loc}^{1,p}(\Omega)]$ ($1 < p < +\infty$) iff u ,*

$$\frac{\partial u}{\partial x_i} \in L^p(\Omega)[L_{loc}^p(\Omega)] \quad (i = 1, 2, \dots, n)$$

$H^{1,p}(\Omega)$ is a Banach space under the norm

$$\|u\|_{H^{1,p}(\Omega)} = \|u\|_{L^p(\Omega)} + \sum_{i=1}^n \left\| \frac{\partial u}{\partial x_i} \right\|_{L^p(\Omega)}$$

$H_0^{1,p}(\Omega)$ is the closure of $\mathcal{D}(\Omega)$ with respect to the $H^{1,p}(\Omega)$ norm; $H^{-1,p}(\Omega)$ is the dual space of $H_0^{1,q}(\Omega)$, where $1/p + 1/q = 1$. We have $T \in H^{-1,p}(\Omega)$

iff, $\exists f_i \in L^p(\Omega)$ ($i = 1, 2, \dots, n$) such that $T = \sum_{i=1}^n \partial f_i / \partial x_i$.

2. Green's function and a representation formula.

In the following sections we will consider the operator $L - V$ where L is the divergence form elliptic operator

$$L = - \frac{\partial}{\partial x_j} \left(a_{ij} \frac{\partial}{\partial x_i} \right)$$

satisfying

$$(2.1) \quad \begin{cases} a_{ij} \in L^\infty(\Omega), & a_{ij} = a_{ji} \quad (i, j = 1, 2, \dots, n) \\ \exists \nu > 0: \nu |\xi|^2 < a_{ij} \xi_i \xi_j < \nu^{-1} |\xi|^2 & \forall \xi \in \mathbb{R}^n \end{cases}$$

and V is a function

$$(2.2) \quad V \in L^{1,\lambda}(\Omega) \quad (\lambda > n - 2).$$

DEFINITION 2.1. *We say that $u \in H_{loc}^{1,2}(\Omega)$ is a local weak solution of the equation*

$$Lu = Vu$$

iff

$$(2.3) \quad \int_{\Omega} a_{ij}(x) u_{x_i}(x) \psi_{x_j}(x) dx = \int_{\Omega} V(x) u(x) \psi(x) dx; \quad \forall \psi \in \mathcal{D}(\Omega).$$

Definition 2.1 is meaningful by the inclusion $L^{1,\lambda}(\Omega) \subseteq \mathcal{S}$ and [S] p. 138-140.

We recall that under the weaker hypothesis $V \in \mathcal{S}$ the following regularity result for weak solutions was proven in [C-F-G].

THEOREM 2.1. *There exist two positive constants $C = C(\nu, n)$, $r_0 = r_0(\nu, n, \eta)$ (η from definition 1.1) and a non decreasing function $\omega(r): \lim_{r \rightarrow 0} \omega(r) = 0$ such that, for any local weak solution of $Lu + Vu = 0$ in Ω and for every ball $B_r(x_0): B_{4r}(x_0) \subseteq \Omega$ ($0 < r \leq r_0$) we have:*

$$\text{osc}_{B_r(x_0)} u \leq C \omega(r) \text{Sup}_{B_{3r}(x_0)} |u|.$$

We now define a different class of solutions:

DEFINITION 2.2. *Let L be such that (2.1) holds, let μ be a bounded variation measure in Ω and $T = \sum_{i=1}^n \partial f_i / \partial x_i \in H^{-1,2}(\Omega)$.*

We say that $u \in L^1(\Omega)$ is a very weak solution of the equation

$$Lu = \mu + T$$

if and only if

$$(2.4) \quad \int_{\Omega} u(x) L\varphi(x) dx = \int_{\Omega} \varphi(x) d\mu - \sum_{i=1}^n \int_{\Omega} f_i(x) \frac{\partial \varphi}{\partial x_i} dx$$

for every $\varphi \in H_0^{1,2}(\Omega) \cap C^0(\bar{\Omega})$ such that $L\varphi \in C^0(\bar{\Omega})$. In much the same way as in [ST] it is possible to show

LEMMA 2.1. *Assume μ is a bounded variation measure and $T = \sum_{i=1}^n \partial f_i / \partial x_i \in H^{-1,2}(\Omega)$. If $u \in H_0^{1,2}(\Omega)$ is a weak solution of the equation*

$$Lu = \mu + T$$

i.e.

$$(2.5) \quad \int_{\Omega} a_{ij}(x) u_{x_i}(x) \varphi_{x_j}(x) dx = \int_{\Omega} \varphi(x) d\mu - \sum_{i=1}^n \int_{\Omega} f_i(x) \frac{\partial \varphi}{\partial x_i} dx; \quad \forall \varphi \in H_0^{1,2}(\Omega)$$

then u is the very weak solution of the same equation.

The proof is an easy consequence of the definitions above. We now recall the definition of fundamental solution.

Let $y \in \Omega$ and δ_y the Dirac mass at y .

Consider the equation

$$Lu = \delta_y.$$

We call its (very weak) solution the Green's function relative to the operator L with pole at y and we denote it by $g(x, y)$.

By the definition above the solution $\varphi \in H_0^{1,2}(\Omega) \cap C^0(\bar{\Omega})$ of $L\varphi = \psi$,

where $\psi \in C^0(\Omega)$ is given by the formula

$$\varphi(y) = \int_{\Omega} g(x, y) \psi(x) dx = \langle \psi(x), g(x, y) \rangle.$$

Consider:

$$(2.8) \quad Lu = \mu + T$$

where μ is a bounded variation measure, $T \in H^{-1,p}(\Omega)$ ($p > n$). We have the following

THEOREM 2.2:

$$u(x) = \langle \mu(y), g(x, y) \rangle + \langle T(y), g(x, y) \rangle$$

is the very weak solution of (2.8).

PROOF. We consider only the case $\mu = 0$ (for the case $T = 0$ see [ST] Th. 8.3 p. 227).

We will show that

$$u(x) = \langle T(y), g(x, y) \rangle$$

satisfies:

$$\langle L\psi(x), \langle T(y), g(x, y) \rangle \rangle = \langle T(y), \psi(y) \rangle; \quad \forall \psi \in H_0^{1,2}(\Omega) \cap C^0(\bar{\Omega})$$

such that $L\psi \in C^0(\bar{\Omega})$.

Let

$$T = \sum_{i=1}^n \frac{\partial f_i}{\partial x_i}, \quad \text{where } f_i \in L^p(\Omega), \quad i = 1, 2, \dots, n.$$

Then

$$\langle L\psi(x), \langle T(y), g(x, y) \rangle \rangle = \int_{\Omega} L\psi(x) \left(- \int_{\Omega} \frac{\partial g}{\partial y_i} f_i(y) dy \right) dx.$$

We observe that

$$|L\psi(x) \frac{\partial g}{\partial y_i} f_i(y)| \in L^1(\Omega \times \Omega).$$

Indeed we have:

$$\begin{aligned} \int_{\Omega} \left(\int_{\Omega} |L\psi(x)| \left| \frac{\partial g}{\partial y_i} \right| |f_i(y)| dy \right) dx &= \int_{\Omega} |L\psi(x)| \left(\int_{\Omega} \left| \frac{\partial g}{\partial y_i} \right| |f_i(y)| dy \right) dx \leq \\ &\leq \int_{\Omega} |L\psi(x)| \left\| \frac{\partial g}{\partial y_i} \right\|_{L^{p'}(\Omega)} \|f_i\|_{L^p(\Omega)} dx \leq \max_{\Omega} |L\psi(x)| \|f_i\|_{L^p(\Omega)} \int_{\Omega} \left\| \frac{\partial g}{\partial x_i} \right\|_{L^{p'}(\Omega)} dx. \end{aligned}$$

Then (see [ST] p. 220 (8.6))

$$\int_{\Omega} \left(\int_{\Omega} |L\psi(x)| \left| \frac{\partial g}{\partial y_i} \right| |f_i(y)| dy \right) \leq C \max_{\Omega} |L\psi(x)| \|f_i\|_{L^p(\Omega)}.$$

By Tonelli and Fubini's theorems we have:

$$\begin{aligned} \int_{\Omega} L\psi(x) \left(- \int_{\Omega} \frac{\partial g}{\partial y_i} f_i(y) dy \right) dx &= \int_{\Omega} f_i(y) \left(- \int_{\Omega} \frac{\partial g}{\partial y_i} L\psi(x) dx \right) dy = \\ &= \int_{\Omega} f_i(y) \left(- \frac{\partial}{\partial y_i} \int_{\Omega} g(x, y) L\psi(x) dx \right) dy = \\ &= \int_{\Omega} f_i(y) \left(- \frac{\partial}{\partial y_i} \langle g(x, y), L\psi(x) \rangle \right) = \left\langle \frac{\partial f_i}{\partial y_i}, \langle g(x, y), L\psi(x) \rangle \right\rangle = \\ &= \langle T(x), \psi(x) \rangle. \end{aligned}$$

REMARK 2.1. In the proof above we may differentiate under the integral; i.e.

$$- \int_{\Omega} \frac{\partial g}{\partial y_i} L\psi(x) dx = - \frac{\partial}{\partial y_i} \int_{\Omega} g(x, y) L\psi(x) dx.$$

In fact, for every $\varphi \in \mathcal{D}(\Omega)$ we have, using Fubini's theorem:

$$- \left\langle \frac{\partial}{\partial y_i} \int_{\Omega} g(x, y) L\psi(x) dx, \varphi(y) \right\rangle = \left\langle \int_{\Omega} g(x, y) L\psi(x) dx, \frac{\partial \varphi}{\partial y_i} \right\rangle =$$

$$\begin{aligned}
&= \int_{\Omega} \left(\int_{\Omega} g(x, y) L\psi(x) dx \right) \frac{\partial \varphi}{\partial y_i} dy = - \int_{\Omega} \left(\int_{\Omega} \frac{\partial g}{\partial y_i} \varphi(y) dy \right) L\psi(x) dx = \\
&= - \left\langle \int_{\Omega} \frac{\partial g}{\partial y_i} L\psi(x) dx, \varphi(y) \right\rangle.
\end{aligned}$$

3. Hölder-continuity of local solutions.

We now state the main result of this paper

THEOREM 3.1. *There exist positive numbers $r_0 = r_0(\nu, \|V\|_{1,\lambda}, \lambda, n)$, $\alpha = \alpha(\nu, n)$, $C = C(\nu, n, \|V\|_{1,\lambda}, \lambda)$ such that for any local solution u of $Lu = Vu$ in Ω and for any ball $B_r(x_0)$, with $B_{4r}(x_0) \subseteq \Omega$, $0 < r \leq r_0$ we have*

$$\begin{aligned}
|u(x) - u(x_0)| &\leq C \operatorname{Sup}_{B_{3r}(x_0)} |u| r^{\lambda-n+2} \cdot \\
&\quad \cdot \left(|x - x_0|^{\alpha/2} r^{-\alpha/2} + |x - x_0|^{(\lambda-n+2)/2} r^{-(\lambda-n+2)/2} + \left(\frac{|x - x_0|}{r} \right)^\alpha \right).
\end{aligned}$$

PROOF. Let $V \in L^{1,\lambda}(\Omega)$ and u a local weak solution of $Lu = Vu$ i.e. $u \in H_{\text{loc}}^{1,2}(\Omega)$ such that:

$$(3.1) \quad \int_{\Omega} a_{ij}(x) \frac{\partial u}{\partial x_i} \frac{\partial \psi}{\partial x_j} dx = \int_{\Omega} V(x) \psi(x) dx \quad \forall \psi \in \mathcal{D}(\Omega).$$

Let $\varphi \in \mathcal{D}(\Omega)$. It is easy to see that $u\varphi$ is such that

$$\begin{aligned}
\int_{\Omega} a_{ij}(x) \frac{\partial(u\varphi)}{\partial x_i} \frac{\partial \psi}{\partial x_j} dx &= \int_{\Omega} V(x) u(x) \psi(x) \varphi(x) dx + \\
&\quad + \int_{\Omega} a_{ij}(x) u(x) \frac{\partial \varphi}{\partial x_i} \frac{\partial \psi}{\partial x_j} dx - \int_{\Omega} a_{ij}(x) \frac{\partial u}{\partial x_i} \frac{\partial \varphi}{\partial x_j} \psi(x) dx
\end{aligned}$$

holds.

Therefore, by Lemma 2.1, $u\varphi$ is a very weak solution of

$$L(u\varphi) = V(x)u(x)\varphi(x) - \frac{\partial}{\partial x_j} \left(a_{ij}(x)u(x) \frac{\partial \varphi}{\partial x_i} \right) - a_{ij}(x) \frac{\partial u}{\partial x_i} \frac{\partial \varphi}{\partial x_j}.$$

By Theorem 2.2 we have

$$u(x)\varphi(x) = \int_{\Omega} V(y)u(y)\varphi(y)g(x, y) dy + \int_{\Omega} \frac{\partial g}{\partial y_i} a_{ij}(y)u(y) \frac{\partial \varphi}{\partial y_j} dy - \int_{\Omega} g(x, y)a_{ij}(y) \frac{\partial u}{\partial y_i} \frac{\partial \varphi}{\partial y_j} dy .$$

Now we choose $\varphi \in \mathcal{D}(\Omega)$ such that $0 \leq \varphi \leq 1$, $\varphi(x) = 1$ in $B_{\frac{1}{2}r}(x_0)$, $\text{supp}(\varphi) \subseteq B_{2r}(x_0)$, $|\nabla \varphi| \leq C/r$ where $0 < r \leq r_0$ and r_0 is determined by the local boundedness theorem 1.4 in [C-F-G].

Obviously, for every $x \in B_{2r}(x_0)$ we have:

$$u(x) - u(x_0) = \int_{\Omega} V(y)u(y)\varphi(y)(g(x, y) - g(x_0, y)) dy - \int_{\Omega} (g(x, y) - g(x_0, y))a_{ij}(y) \frac{\partial u}{\partial y_i} \frac{\partial \varphi}{\partial y_j} dy + \int_{\Omega} \left(\left(\frac{\partial g}{\partial y_i} \right)_{(x, y)} - \left(\frac{\partial g}{\partial y_j} \right)_{(x_0, y)} \right) a_{ij}(y)u(y) \frac{\partial \varphi}{\partial y_j} dy = \text{I} - \text{II} + \text{III} .$$

We begin estimating I.

$$\begin{aligned} \text{I} &= \int_{|x-y| > N|x-x_0|} (g(x, y) - g(x_0, y)) V(y)u(y)\varphi(y) dy + \\ &\quad + \int_{|x_0-y| \leq N|x-x_0|} (g(x, y) - g(x_0, y)) V(y)u(y)\varphi(y) dy = A + B . \end{aligned}$$

Where N is a positive number to be fixed later.

To estimate A we use the inequality (see [G-T] p. 200 Th. 8.22 and Harnack's Theorem)

$$|g(x, y) - g(x_0, y)| \leq C(\nu, n) \left(\frac{|x - x_0|}{r} \right)^\alpha g(x_0, y) \leq \leq \frac{C(\nu, n)}{N^\alpha} g(x_0, y) \leq \frac{C(\nu, n)}{N^\alpha |x_0 - y|^{n-2}}$$

hence

$$A \leq \frac{C(v, n)}{N^\alpha} \int_{B_{2r}(x_0)} \frac{|V(y)|}{|x_0 - y|^{n-2}} dy \operatorname{Sup}_{B_{2r}(x_0)} |u|$$

and by Lemma 1.1

$$A \leq \frac{C(\|V\|_{L^{1,\lambda}(\Omega)}, v, n, \lambda)}{N^\alpha} r^{\lambda-n+2} \operatorname{Sup}_{B_{2r}(x_0)} |u|.$$

To estimate B we use Lemma 1.1 and the following bound

$$g(x, y) \leq \frac{C(v, n)}{|x - y|^{n-2}}$$

proven in [L-S-W].

We obtain:

$$|g(x, y) - g(x_0, y)| \leq \frac{C(v, n)}{|x - y|^{n-2}} + \frac{C(v, n)}{|x_0 - y|^{n-2}}$$

and therefore

$$\begin{aligned} B &\leq C(v, n) \int_{|x_0 - y| \leq N|x - x_0|} \frac{|V(y)|}{|x - y|^{n-2}} dy \operatorname{Sup}_{B_{2r}(x_0)} |u| \leq \\ &\leq C(v, n) \int_{|x_0 - y| \leq (N+1)|x - x_0|} \frac{|V(y)|}{|x - y|^{n-2}} dy \leq \\ &\leq C(v, n \|V\|_{L^{1,\lambda}(\Omega)}, \lambda) \operatorname{Sup}_{B_{2r}(x_0)} |u| ((N+1)|x - x_0|)^{\lambda-n+2}. \end{aligned}$$

Now, if we choose $N = (r/|x - x_0|)^{\frac{1}{2}} > 1$ we obtain

$$\begin{aligned} |I| &\leq C(\|V\|_{L^{1,\lambda}(\Omega)}, \lambda, v, n) \operatorname{Sup}_{B_{2r}(x_0)} |u| |x - x_0|^{\alpha/2} r^{\lambda-n+2-\alpha/2} + \\ &+ C(\|V\|_{L^{1,\lambda}(\Omega)}, \lambda, v, n) \operatorname{Sup}_{B_{2r}(x_0)} |u| |x - x_0|^{(\lambda-n+2)/2} r^{(\lambda-n+2)/2}. \end{aligned}$$

Estimating II and III as in [C-F-G] we obtain

$$|II| \leq C(v, n) \left(\frac{|x - x_0|}{r} \right)^\alpha \left(\int_{B_{2r}(x_0)} u(y)^2 dy \right)^{\frac{1}{2}}$$

and

$$|\text{III}| \leq C(\nu, n) \left(\frac{|x - x_0|}{r} \right)^\alpha \left(\int_{B_{2r}(x_0)} u(y)^2 dy \right)^{\frac{1}{2}}.$$

The theorem now follows.

REFERENCES

- [A-S] M. AIZENMAN - B. SIMON, *Brownian motion and Harnack's inequality for Schrödinger operators*, Comm. Pure and Appl. Math., **35** (1982), pp. 209-271.
- [C-F-G] F. CHIARENZA - E. FABES - N. GAROFALO, *Harnack's inequality for Schrödinger operators and the continuity of solutions*, Proc. A.M.S., **98** (1986), pp. 415-425.
- [C-F-Z] M. CRANSTON - E. FABES - Z. ZHAO, *Potential theory for the Schrödinger equation*, Bull. A.M.S., **15** (1986), pp. 213-216.
- [D-M] G. DAL MASO - U. MOSCO, *Wiener criteria and energy decay for relaxed Dirichlet problems*, preprint.
- [G-T] D. GILBARG - N. TRUDINGER, *Elliptic Partial Differential Equation of Second Order* (2nd edition), Springer-Verlag (1983).
- [L-S-W] W. LITTMAN - G. STAMPACCHIA - H. F. WEINBERGER, *Regular points for elliptic equations with discontinuous coefficients*, Ann. Scuola Norm. Sup. di Pisa, S. III, **17** (1963), pp. 45-79.
- [L-U] O. A. LADIZHENSKAIA - N. URALTZEVA, *Linear and quasilinear elliptic equations*, Academic Press (1968).
- [P1] L. C. PICCININI, *Proprietà di inclusione e interpolazione tra spazi di Morrey e loro generalizzazioni* (Tesi di perfezionamento), Pisa (1969).
- [P2] L. C. PICCININI, *Inclusioni tra spazi di Morrey*, Boll. U.M.I., **2** (1969), pp. 95-99.
- [S] M. SCHECHTER, *Spectra of partial differential operators*, North-Holland (1971).
- [ST] G. STAMPACCHIA, *Équations elliptiques du second ordre à coefficients discontinus*, Les Presses de l'Université de Montréal (1965).

Manoscritto pervenuto in redazione il 3 aprile 1987.