

RENDICONTI *del* SEMINARIO MATEMATICO *della* UNIVERSITÀ DI PADOVA

E. OSSANNA

**Comparison between the generalized mean
curvature according to Allard and Federer's
mean curvature measure**

Rendiconti del Seminario Matematico della Università di Padova,
tome 88 (1992), p. 221-227

http://www.numdam.org/item?id=RSMUP_1992__88__221_0

© Rendiconti del Seminario Matematico della Università di Padova, 1992, tous droits réservés.

L'accès aux archives de la revue « Rendiconti del Seminario Matematico della Università di Padova » (<http://rendiconti.math.unipd.it/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>

Comparison between the Generalized Mean Curvature according to Allard and Federer's Mean Curvature Measure.

E. OSSANNA (*)

ABSTRACT - We compare two well known generalized notions of mean curvature for the boundary of a convex body: Federer's mean curvature measure, defined via Steiner's formula, and Allard's generalized mean curvature, which is a vector measure obtained via the first variation of the area. The comparison is got by a suitable approximation lemma for convex sets.

Our purpose is to compare the generalized mean curvature (according to Allard) of a convex body K to its mean curvature measure (according to Federer). We recall that a body K of \mathbf{R}^n is a compact subset of \mathbf{R}^n such that $\overset{\circ}{K} \neq \emptyset$. Our result, which is stated in Theorem 1, follows approximating K by a suitable sequence of regular convex bodies, whose existence is assured by Lemma 2.

After this work was completed we have been informed by Joseph Fu that he has recently obtained [FU], by different methods, a similar (unpublished) result in the context of sets with a generalized unit normal bundle.

First we recall some useful facts.

Let K be a convex body of \mathbf{R}^n , let $p(K, \cdot): \mathbf{R}^n \rightarrow K$ be the nearest-point map for K and define

$$u(K, x) = \frac{p(K, x) - x}{\|p(K, x) - x\|}.$$

If one considers the set $A_\varepsilon(K, E) = \{x \in K_\varepsilon \mid p(K, x) \in E\}$, where E is

(*) Indirizzo dell'A.: Dipartimento di Matematica, Università di Trento, 38050 Povo (Trento), Italy.

Research supported by C.N.R. fellowship at the Department of Mathematics of Trento University.

any Borel subset of \mathbf{R}^n and $K_\varepsilon = \{x \in \mathbf{R}^n \mid d(K, x) < \varepsilon\}$, then [SCH1] there are the so called Federer's *curvature measures* of K , denoted by $\phi_m(K, \cdot)$, $m = 0, \dots, n$ and defined on the Borel sets of \mathbf{R}^n in such a way that

$$H^n(A_\varepsilon(K, E)) = \sum_{m=0}^n \varepsilon^{n-m} \alpha(n-m) \phi_m(K, E),$$

where $\alpha(k) = H^k(\{x \in \mathbf{R}^k \mid \|x\| \leq 1\})$. In particular we call $\phi_{n-2}(K, \cdot)$ the mean curvature measure.

One can also consider the set $M_\varepsilon(K, A) = \{x \in K_\varepsilon \setminus K \mid (p(K, x), u(K, x)) \in A\}$, where A is any Borel subset of $\mathbf{R}^n \times S^{n-1}$. In this case one finds that [SCH2] there are n measures, $\theta_m(K, \cdot)$, $m = 0, \dots, n-1$, defined on the Borel sets of $\mathbf{R}^n \times S^{n-1}$ and called the *generalized curvature measures* of K , such that

$$H^n(M_\varepsilon(K, A)) = \sum_{m=0}^{n-1} \varepsilon^{n-m} \alpha(n-m) \theta_m(K, A).$$

Moreover let \bar{H} be the *generalized mean curvature* of K according to Allard, defined so that

$$\int_{\partial K} \operatorname{div}_{\partial K} X dH^{n-1} = - (n-1) \int_{\partial K} X \cdot d\bar{H}$$

for each vector field $X \in C_0^1(U)$, where U is an open subset of \mathbf{R}^n .

We shall give also the definition of *stratification* of a measure defined on the product of two sets.

LEMMA 1 [SIM]. *Let α be a Radon measure on $\mathbf{R}^n \times S^{n-1}$ and consider the Radon measure σ on \mathbf{R}^n such that $\sigma(A) = \alpha(A \times S^{n-1})$ for each Borel set $A \subset \mathbf{R}^n$. Then for almost all $x \in \mathbf{R}^n$ there is a Radon measure λ_x on S^{n-1} such that for each Borel set $B \subset S^{n-1}$ the function $\lambda_x(B)$ is the density of the measure ρ_B with respect of σ , where $\rho_B(A) = \alpha(A \times B)$. From this decomposition of the measure α one gets*

$$\int_{\mathbf{R}^n \times S^{n-1}} g(x, y) d\alpha = \int_{\mathbf{R}^n} \left(\int_{S^{n-1}} g(x, y) d\lambda_x \right) d\sigma$$

for each function $g(x, y) \in C_0^0(\mathbf{R}^n \times S^{n-1})$.

We call (σ, λ_x) the stratification of the measure α .

THEOREM 1. *Let K be a convex body of \mathbf{R}^n , let \bar{H} be the generalized mean curvature of K and $\phi_{n-2}(K, \cdot)$ the mean curvature measure of K .*

Then for each Borel set $E \subset \mathbf{R}^n$ we have

$$\frac{n-1}{2\pi} \overline{H}(E) = - \int_{\partial K \cap E} \left(\int_{S^{n-1}} y \, d\lambda_x \right) d\phi_{n-2}(K) = - \int_{\partial K \cap E} b_{\lambda_x} \, d\phi_{n-2}(K),$$

where $(\phi_{n-2}(K), \lambda_x)$ is the stratification of the generalized curvature measure $\Theta_{n-2}(K)$ and b_{λ_x} is the barycenter of the measure λ_x .

For the proof of the theorem we need the approximation lemma below. First we recall that [HUT] the oriented varifold associated to an oriented hypersurface M of \mathbf{R}^n is the Radon measure μ on $\mathbf{R}^n \times S^{n-1}$ such that for each function $g \in C_0^0(\mathbf{R}^n \times S^{n-1})$ one has

$$\int_{\mathbf{R}^n \times S^{n-1}} g(x, y) \, d\mu = \int_M g(x, \nu) \, dH^{n-1},$$

where ν is the unit normal field to the surface M .

LEMMA 2. Let K be a convex body of \mathbf{R}^n , then there is a sequence of convex bodies $\{K_j\}_{j \in \mathbf{N}}$ such that $\partial K_j = M_j$ is a smooth surface and such that the following properties hold:

- i) $K_j \xrightarrow{d_H} K$, where d_H stays for Hausdorff distance;
- ii) the oriented varifolds μ_j associated to the surfaces M_j converge weakly to the varifold μ associated to the surface $\partial K = M$.

PROOF. Let K be a convex body of \mathbf{R}^n and M its boundary, suppose that $B_r(0) \subset \overset{\circ}{K}$ and $K \subset B_R(0)$, for suitable $r > 0$ and $R > 0$. Consider the function $u: \mathbf{R}^n \rightarrow \mathbf{R}$ defined in the following way:

$$u(0) = 0, \\ u(x) = \frac{\|x\|}{\|i_M(x)\|} \quad \text{if } x \neq 0,$$

where $i_M(x) = \{tx \mid t \geq 0\} \cap M$. Note that the function i_M is well defined and $i_M(x) \neq 0$ for every $x \in \mathbf{R}^n \setminus \{0\}$.

In this case M consists of the points $x \in \mathbf{R}^n$ such that $u(x) = 1$.

Clearly the function u is convex and lipschitz continuous, moreover the gradient $\nabla u(x)$ exists for almost all x , and where ∇u is defined we have $1/R \leq \|\nabla u(x)\| \leq 1/r$.

Now let $\{\gamma_j\}_{j \in \mathbf{N}}$ be a sequence of mollifiers and define the functions $u_j = u * \gamma_j$. This way we get a sequence of smooth convex functions such that:

- (a) $\{u_j\}_{j \in N}$ converges uniformly to the function u ;
 (b) for every $x \in \mathbf{R}^n$ we have $1/R \leq \|\nabla u_j(x)\| \leq 1/r$.

Now consider the sets $K_j = \{x \in \mathbf{R}^n \mid u_j(x) \leq 1\}$ $j \in N$. One can see easily that the sets K_j are convex and uniformly bounded, hence they are convex bodies.

The statement i) of the lemma follows from (a) after some calculations. Moreover observe that, being K_j convex bodies, convergence i) implies

$$(1) \quad K_j \xrightarrow{L^1} K.$$

From (b) we get that each set $M_j = \{x \in \mathbf{R}^n \mid u_j(x) = 1\}$, boundary of the convex body K_j , is a smooth hypersurface of \mathbf{R}^n .

In what follows we denote by ν the unit outward normal to M (ν is defined almost everywhere) and by ν_j the unit outward normal to M_j .

Now we shall prove statement ii) of the lemma, that is $\mu_j \rightharpoonup \mu$. As a *first step* we consider the vector measures

$$\beta_j = \nu_j(x)(H^{n-1} \llcorner M_j) \quad j \in N, \quad \beta = \nu(x)(H^{n-1} \llcorner M)$$

and their total variation measures

$$|\beta_j| = H^{n-1} \llcorner M_j, \quad |\beta| = H^{n-1} \llcorner M$$

and we shall prove that

$$(2) \quad \beta_j \rightharpoonup \beta,$$

$$(3) \quad |\beta_j| \rightharpoonup |\beta|.$$

To obtain $\beta_j \rightharpoonup \beta$ we use the divergence theorem and the convergence (1), getting for each vector field $X \in C_0^1(\mathbf{R}^n)$

$$(4) \quad \lim_{j \rightarrow +\infty} \int_{\mathbf{R}^n} X \cdot d\beta_j = \int_{\mathbf{R}^n} X \cdot d\beta,$$

and we extend (4) to vector fields belonging to C_0^0 , using the density of C_0^1 in C_0^0 and the uniform boundness of the convex bodies K_j .

Next we show that $|\beta_j| \rightharpoonup |\beta|$. By convergence (2) we get

$$(5) \quad \liminf_{j \rightarrow +\infty} |\beta_j|(A) \geq |\beta|(A) \quad \forall \text{ open } A \subset \mathbf{R}^n,$$

and on the other hand by i) we have

$$(6) \quad |\beta_j|(\mathbf{R}^n) \rightarrow |\beta|(\mathbf{R}^n),$$

in fact, i) implies that for every $\varepsilon > 0$ there are $r > 0$ and $J \in \mathbf{N}$ such that, if we set $\delta = 1 + \varepsilon/r$, it results $\delta^{-1}K \subset K_j \subset \delta K$ for all $j > J$, from which, again for $j > J$, $H^{n-1}(\partial K) \leq H^{n-1}(\partial K_j) \leq H^{n-1}(\delta K)$ follows. Thus, being $|\beta_j|$ and $|\beta|$ finite measures, (3) follows from (5) and (6) [HAL].

The *second step* consists in proving that, if there is a subsequence $\{\mu_{j_h}\}_{h \in \mathbf{N}}$ of the sequence of Radon measures $\{\mu_j\}_{j \in \mathbf{N}}$ and a Radon measure α such that

$$(7) \quad \mu_{j_h} \rightharpoonup \alpha,$$

then $\alpha = \mu$. For this we consider the stratification (σ, λ_x) of the measure α and a function $g \in C_0^0(\mathbf{R}^n)$, then we have

$$\int_{\mathbf{R}^n \times S^{n-1}} g(x) d\alpha = \int_{\mathbf{R}^n} g(x) \left(\int_{S^{n-1}} d\lambda_x \right) d\sigma = \int_{\mathbf{R}^n} g(x) d\sigma,$$

and from (3) and (7) we get

$$\int_{\mathbf{R}^n \times S^{n-1}} g(x) d\alpha = \int_{\mathbf{R}^n} g(x) d|\beta|,$$

which implies

$$(8) \quad \sigma = |\beta|.$$

Moreover since $\beta_j \rightharpoonup \beta$ and $\mu_{j_h} \rightharpoonup \alpha$, between the two measures β and α there is the relationship [RES] $\beta = b(x)\sigma$, where $b(x) = \int_{S^{n-1}} y d\lambda_x$. But

$\beta = \nu(x)(H^{n-1} \llcorner M)$ and $\sigma = |\beta|$, hence $\nu(x)|\beta| = b(x)|\beta|$, which implies $\int_{S^{n-1}} y d\lambda_x = \nu(x)$ for $|\beta|$ -almost all $x \in \mathbf{R}^n$, that is

$$(9) \quad \lambda_x = \delta_{\nu(x)},$$

$|\beta|$ -almost everywhere. Now, using (8) and (9), we can prove $\alpha = \mu$, in fact for each function $g \in C_0^0(\mathbf{R}^n \times S^{n-1})$ we have

$$\begin{aligned} \int_{\mathbf{R}^n \times S^{n-1}} g(x, y) d\alpha &= \int_{\mathbf{R}^n} \left(\int_{S^{n-1}} g(x, y) d\delta_{\nu(x)} \right) d|\beta| = \\ &= \int_{\mathbf{R}^n} g(x, \nu(x)) d|\beta| = \int_{\mathbf{R}^n \times S^{n-1}} g(x, y) d\mu. \end{aligned}$$

Now we end the proof of statement ii) of the lemma. Assume by contradiction there is a function $g \in C_0^0(\mathbf{R}^n \times S^{n-1})$ such that

$\int_{\mathbf{R}^n \times S^{n-1}} g(x, y) d\mu_j$ does not converge to $\int_{\mathbf{R}^n \times S^{n-1}} g(x, y) d\mu$. Then there are a number ε and a subsequence $\{\mu_{j_k}\}_{k \in \mathbf{N}}$ such that

$$(10) \quad \left| \int_{\mathbf{R}^n \times S^{n-1}} g(x, y) d\mu_{j_k} - \int_{\mathbf{R}^n \times S^{n-1}} g(x, y) d\mu \right| > \varepsilon \quad \forall k.$$

On the other hand one has $\mu_{j_k}(\mathbf{R}^n \times S^{n-1}) \leq C \quad \forall k$. Hence there is a further subsequence $\{\mu_{j_{k_h}}\}_{h \in \mathbf{N}}$ that converges weakly to some measure, which, by the second step, must be μ , and this contradicts (10). ■

PROOF OF THEOREM 2. Let $\{K_j\}_{j \in \mathbf{N}}$ be the sequence introduced in the Lemma 2. From the weak convergence of the varifolds μ_j we have

$$\lim_{j \rightarrow +\infty} \int_{\partial K_j} \operatorname{div}_{\partial K_j} X dH^{n-1} = \int_{\partial K} \operatorname{div}_{\partial K} X dH^{n-1},$$

for each vector field $X \in C_0^1(\mathbf{R}^n)$.

Hence, for the definition of generalized mean curvature, we have also

$$(11) \quad \overline{H}_j(x) H^{n-1} \rightharpoonup \overline{H},$$

where \overline{H}_j is the mean curvature vector of the surfaces ∂K_j .

Now if we let E be a Borel set of \mathbf{R}^n such that $\overline{H}(\partial E) = 0$ and $\phi_{n-2}(K, \partial E) = 0$, we have [SCH2]

$$\Theta_{n-2}(K_j) \llcorner E \times S^{n-1} \rightharpoonup \Theta_{n-2}(K) \llcorner E \times S^{n-1},$$

which implies in particular

$$\begin{aligned} \lim_{j \rightarrow +\infty} \int_{\partial K_j \times S^{n-1}} \mathcal{X}_{(E \times S^{n-1})} y d\Theta_{n-2}(K_j) &= \int_{\partial K \times S^{n-1}} \mathcal{X}_{(E \times S^{n-1})} y d\Theta_{n-2}(K) = \\ &= \int_{\partial K \cap E} \int_{S^{n-1}} y d\lambda_x d\phi_{n-2} = \int_{\partial K \cap E} b_{\lambda_x} d\phi_{n-2}(K). \end{aligned}$$

But ∂K_j is a regular surface, thus we have

$$\begin{aligned} \int_{\partial K_j \times S^{n-1}} \mathcal{X}_{(E \times S^{n-1})} y \, d\theta_{n-2}(K_j) &= \frac{n-1}{2\pi} \int_{\partial K_j \cap E} v_j \|\overline{H}_j\| \, dH^{n-1} = \\ &= -\frac{n-1}{2\pi} \int_{\partial K_j \cap E} \overline{H}_j \, dH^{n-1}, \end{aligned}$$

moreover, being $\overline{H}(\partial E) = 0$, convergence (11) implies

$$\lim_{j \rightarrow +\infty} \int_{\partial K_j \cap E} \overline{H}_j \, dH^{n-1} = \overline{H}(E).$$

Hence, for each Borel set $E \subset \mathbf{R}^n$ such that $\overline{H}(\partial E) = 0$ and $\phi_{n-2}(K, \partial E) = 0$, we get

$$\frac{n-1}{2\pi} \overline{H}(E) = - \int_{\partial K \cap E} \left(\int_{S^{n-1}} y \, d\lambda_x \right) d\phi_{n-2}(K) = - \int_{\partial K \cap E} b_{\lambda_x} \, d\phi_{n-2}(K)$$

and this equality holds also for every Borel set $A \subset \mathbf{R}^n$. ■

REFERENCES

- [FU] J. H. G. FU, private communication (1991).
- [HAL] P. R. HALMOS, *Measure Theory*, Springer-Verlag, New York-Berlin (1974).
- [HUT] J. E. HUTCHINSON, *Second fundamental form for varifolds and the existence of surfaces minimizing curvature*, Indiana Univ. Math. J., **35** (1986), pp. 45-71.
- [RES] YU. G. RESHETNYAK, *Weak convergence of completely additive vector functions on a set*, Sibirsk. Mat. Z., **9** (1968), pp. 1386-1394.
- [SCH1] R. SCHNEIDER, *Curvature measures of convex bodies*, Ann. Mat. Pura Appl., (4) **116** (1978), pp. 101- 134.
- [SCH2] R. SCHNEIDER, *Bestimmung konvexer Körper durch Krümmungsmaße*, Comment. Math. Helvetici, **54** (1979), pp. 42-60.
- [SIM] L. SIMON, *Lectures on geometric measure theory*, Proceedings of the Centre for Mathematical Analysis, Australian Nat. Univ. Canberra, Australia, vol. **3** (1984), p. 230.