

RENDICONTI *del* SEMINARIO MATEMATICO *della* UNIVERSITÀ DI PADOVA

BRUNO GABUTTI

GIOVANNA PITTALUGA

LAURA SACRIPANTE

Asymptotics for Meixner polynomials

Rendiconti del Seminario Matematico della Università di Padova,
tome 94 (1995), p. 1-10

http://www.numdam.org/item?id=RSMUP_1995__94__1_0

© Rendiconti del Seminario Matematico della Università di Padova, 1995, tous droits réservés.

L'accès aux archives de la revue « Rendiconti del Seminario Matematico della Università di Padova » (<http://rendiconti.math.unipd.it/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>

Asymptotics for Meixner Polynomials.

BRUNO GABUTTI - GIOVANNA PITTALUGA - LAURA SACRIPANTE (*)

SUMMARY - Asymptotic expansion for Meixner polynomials $m_n(x/(1-c); \alpha+1, c)$ in terms of Laguerre polynomials $L_n^{(\alpha)}(x)$, $c \rightarrow 1$, is derived. Asymptotic approximation and limit relations of the zeros of these polynomials are also given.

1. Introduction and results.

The Meixner polynomials

$$m_n(x; \beta, c) = (\beta)_n {}_2F_1\left(-n, -x; \beta; 1 - \frac{1}{c}\right),$$
$$c \neq 0, \quad \beta \neq 0, -1, -2, \dots,$$

satisfying, for $0 < |c| < 1$, the orthogonality relation

$$\sum_{k=0}^{\infty} m_n(k; \beta, c) m_p(k; \beta, c) \frac{c^k (\beta)_k}{k!} = (1-c)^{-\beta} c^{-n} n! (\beta)_n \delta_{np},$$

are a discrete analogue of the Laguerre polynomials $L_n^{(\beta-1)}(x)$.

These polynomials are of importance for the description of certain Markov processes and in the numerical evaluation of the Hankel transform [3]. Their main properties can be found in Chihara [1], including

(*) Indirizzo degli AA.: University of Turin, Italy.

Work supported by the Consiglio Nazionale delle Ricerche of Italy and by the Ministero dell'Università e della Ricerca Scientifica e Tecnologica of Italy.

the limit relation

$$(1.1) \quad \lim_{c \rightarrow 1} m_n \left(\frac{x}{1-c}; \beta, c \right) = n! L_n^{(\beta-1)}(x),$$

and the recurrence formula

$$(1.2) \quad \begin{aligned} cm_{n+1}(x) &= \\ &= [(c-1)x + (1+c)n + c\beta] m_n(x) - n(n+\beta-1) m_{n-1}(x), \quad n \geq 0. \end{aligned}$$

Now, let $L_n^{(\alpha)}(x)$ be Laguerre polynomials and let $Q_n^{(\alpha)}(x; \lambda)$ be defined by

$$(1.3) \quad Q_n^{(\alpha)}(x; \lambda) = \frac{1}{n!} \left(\frac{\lambda+1}{\lambda-1} \right)^{n/2} m_n \left(-\frac{\lambda x}{2} - \frac{\alpha+1}{2}; \alpha+1, \frac{\lambda+1}{\lambda-1} \right).$$

We notice that, for the transformation formula

$${}_2F_1(a, b; c; z) = (1-z)^{-a} {}_2F_1\left(a, c-b; c; \frac{z}{z-1}\right),$$

the function $Q_n^{(\alpha)}(x; \lambda)$ is even with respect to λ .

We shall assume throughout this paper that the parameter α satisfies the condition $\alpha > -1$.

By putting

$$\frac{x}{1-c} = -\frac{\lambda x}{2} - \frac{\alpha+1}{2} = -\frac{\lambda-1}{2} x \{1 + O(\lambda^{-1})\}, \quad \lambda \rightarrow \infty,$$

from the (1.1) and (1.3) it follows that

$$(1.4) \quad \lim_{\lambda \rightarrow \infty} Q_n^{(\alpha)}(x; \lambda) = L_n^{(\alpha)}(x).$$

If $l_{n,k}^{(\alpha)}$ and $x_{n,k}^{(\alpha, \lambda)}$, $k = 1, 2, \dots, n$, are the zeros of Laguerre and Meixner polynomials (1.3) respectively, that is

$$(1.5) \quad L_n^{(\alpha)}(l_{n,k}^{(\alpha)}) = 0,$$

$$(1.6) \quad Q_n^{(\alpha)}(x_{n,k}^{(\alpha, \lambda)}; \lambda) = 0,$$

then from (1.4) it will be

$$(1.7) \quad \lim_{\lambda \rightarrow \infty} x_{n,k}^{(\alpha, \lambda)} = l_{n,k}^{(\alpha)}, \quad k = 1, 2, \dots, n.$$

In this note we show that the limits (1.4) and (1.7) can be replaced by accurate asymptotic estimates. More precisely, we can prove

THEOREM 1.1. The polynomial $Q_n^{(\alpha)}(x; \lambda)$ defined in (1.3) has the asymptotic representation

$$(1.8) \quad Q_n^{(\alpha)}(x; \lambda) = L_n^{(\alpha)}(x) + \frac{1}{\lambda^2} Q_{n,1}^{(\alpha)}(x) + O\left(\frac{1}{\lambda^4}\right), \quad \lambda \rightarrow \infty,$$

where $L_n^{(\alpha)}(x)$ is the Laguerre polynomial and

$$(1.9) \quad Q_{n,1}^{(\alpha)}(x) = \frac{x}{3} L_n^{(\alpha)}(x) + \\ - \frac{x + \alpha + 1}{2} L_n^{(\alpha+1)}(x) + \frac{x + \alpha + 1}{2} L_n^{(\alpha+2)}(x) - \frac{x}{3} L_n^{(\alpha+3)}(x).$$

THEOREM 1.2. Let $x_{n,k}^{(\alpha,\lambda)}$ and $l_{n,k}^{(\alpha)}$ be the zeros of $Q_n^{(\alpha)}(x; \lambda)$ and $L_n^{(\alpha)}(x)$ respectively. Then

$$(1.10) \quad x_{n,k}^{(\alpha,\lambda)} = l_{n,k}^{(\alpha)} - \frac{1}{6\lambda^2} \left[2l_{n,k}^{(\alpha)} - (2n + \alpha + 1) + \frac{1 - \alpha^2}{l_{n,k}^{(\alpha)}} \right] + O\left(\frac{1}{\lambda^4}\right), \\ k = 1, 2, \dots, n, \quad \lambda \rightarrow \infty.$$

THEOREM 1.3. For the zeros $x_{n,k}^{(\alpha,\lambda)}$ the following limit relations hold

$$(1.11) \quad \lim_{\lambda \rightarrow \infty} \lambda^3 \frac{\partial x_{n,k}^{(\alpha,\lambda)}}{\partial \lambda} = \frac{1}{3} \left[2l_{n,k}^{(\alpha)} - (2n + \alpha + 1) + \frac{1 - \alpha^2}{l_{n,k}^{(\alpha)}} \right],$$

$$(1.12) \quad \lim_{\lambda \rightarrow \infty} \lambda^4 \frac{\partial^2 x_{n,k}^{(\alpha,\lambda)}}{\partial \lambda^2} = - \left[2l_{n,k}^{(\alpha)} - (2n + \alpha + 1) + \frac{1 - \alpha^2}{l_{n,k}^{(\alpha)}} \right].$$

The proofs of the above Theorems are given in the next Sections.

The outlined procedure shows to be efficient in deriving higher order terms in the asymptotic formulas. The required task is, unfortunately, highly manipulative.

2. Proof of Theorem 1.1.

The polynomials $Q_n^{(\alpha)}(x; \lambda)$ satisfy, for $\lambda > 1$, the recurrence formula

$$(2.1) \quad (n+1)Q_{n+1}^{(\alpha)}(x; \lambda) = \frac{\lambda}{\sqrt{\lambda^2 - 1}} (2n + \alpha + 1 - x)Q_n^{(\alpha)}(x; \lambda) - (n + \alpha)Q_{n-1}^{(\alpha)}(x; \lambda),$$

with the initial values

$$(2.2) \quad Q_{-1}^{(\alpha)}(x; \lambda) = 0, \quad Q_0^{(\alpha)}(x; \lambda) = 1.$$

This is an immediate consequence of the Meixner polynomials recurrence formula (1.2). Assume

$$(2.3) \quad Q_n^{(\alpha)}(x; \lambda) = Q_{n,0}^{(\alpha)}(x) + \frac{1}{\lambda^2} Q_{n,1}^{(\alpha)}(x) + \frac{1}{\lambda^4} Q_{n,2}^{(\alpha)}(x) + \dots,$$

and note

$$(2.4) \quad \frac{\lambda}{\sqrt{\lambda^2 - 1}} = \sum_{k=0}^{\infty} \binom{k-1/2}{k} \frac{1}{\lambda^{2k}}, \quad \lambda > 1.$$

Substituting (2.3) and (2.4) into (2.1) and equating the coefficients of the same powers of λ , we find

$$(2.5) \quad (n+1)Q_{n+1,0}^{(\alpha)}(x) = (2n + \alpha + 1 - x)Q_{n,0}^{(\alpha)}(x) - (n + \alpha)Q_{n-1,0}^{(\alpha)}(x),$$

and

$$(2.6) \quad (n+1)Q_{n+1,j}^{(\alpha)}(x) - (2n + \alpha + 1 - x)Q_{n,j}^{(\alpha)}(x) + (n + \alpha)Q_{n-1,j}^{(\alpha)}(x) = \sum_{k=1}^j \binom{k-1/2}{k} (2n + \alpha + 1 - x)Q_{n,j-k}^{(\alpha)}(x),$$

where $j = 1, 2, \dots$.

Moreover we have

$$(2.7) \quad Q_{-1,0}^{(\alpha)}(x) = 0, \quad Q_{0,0}^{(\alpha)}(x) = 1,$$

and

$$(2.8) \quad Q_{-1,j}^{(\alpha)}(x) = Q_{0,j}^{(\alpha)}(x) = 0, \quad j = 1, 2, \dots.$$

From equations (2.5) and (2.7) one recognizes

$$Q_{n,0}^{(\alpha)}(x) = L_n^{(\alpha)}(x), \quad n = 0, 1, \dots$$

The explicit expression of $Q_{n,j}^{(\alpha)}(x)$, $j = 1, 2, \dots$, can be derived by means of the method of generating function [4, p. 35].

In order to determine $Q_{n,1}^{(\alpha)}(x)$, we observe that, from (2.6) and (2.8) with $j = 1$, we have

$$(2.9) \quad \sum_{n=0}^{\infty} (n+1) Q_{n+1,1}^{(\alpha)}(x) z^n - \sum_{n=0}^{\infty} (2n + \alpha + 1 - x) Q_{n,1}^{(\alpha)}(x) z^n + \\ + \sum_{n=0}^{\infty} (n + \alpha) Q_{n-1,1}^{(\alpha)}(x) z^n = \frac{1}{2} \sum_{n=0}^{\infty} (2n + \alpha + 1 - x) Q_{n,0}^{(\alpha)}(x) z^n.$$

Now, setting

$$(2.10) \quad G_1(z) = \sum_{n=0}^{\infty} Q_{n,1}^{(\alpha)}(x) z^n,$$

and

$$(2.11) \quad H_0(z) = \frac{1}{2} \sum_{n=0}^{\infty} (2n + \alpha + 1 - x) Q_{n,0}^{(\alpha)}(x) z^n,$$

by straightforward calculations the (2.9) can be rewritten in the form

$$(2.12) \quad (1-z)^2 G_1'(z) - [(\alpha+1)(1-z) - x] G_1(z) = H_0(z),$$

with

$$(2.13) \quad G_1(0) = 0,$$

where the prime denotes the derivative respect to z .

Using [2, p. 189]

$$(2.14) \quad \sum_{n=0}^{\infty} L_n^{(\alpha)}(x) z^n = (1-z)^{-\alpha-1} \exp\left(\frac{xz}{z-1}\right),$$

and

$$\sum_{n=1}^{\infty} n L_n^{(\alpha)}(x) z^{n-1} = \frac{(\alpha+1)(1-z) - x}{(1-z)^{\alpha+3}} \exp\left(\frac{xz}{z-1}\right),$$

in equation (2.11), we easily obtain

$$H_0(z) = \frac{\alpha + 1 - x - z^2(\alpha + 1 + x)}{2(1 - z)^{\alpha+3}} \exp\left(\frac{xz}{z-1}\right).$$

Thus, from (2.12) and (2.13), it follows

$$G_1(z) = \exp\left(\frac{xz}{z-1}\right).$$

$$\cdot \left[\frac{x}{3(1-z)^{\alpha+1}} - \frac{\alpha+1+x}{2(1-z)^{\alpha+2}} + \frac{\alpha+1+x}{2(1-z)^{\alpha+3}} - \frac{x}{3(1-z)^{\alpha+4}} \right].$$

Then, from (2.10), taking into account (2.14), we find that $Q_{n,1}^{(\alpha)}(x)$ satisfies (1.9). So the expansion (2.3) prove part of estimate (1.8).

To complete the proof, we observe merely that $G_2(z)$, defined by

$$G_2(z) = \sum_{n=0}^{\infty} Q_{n,2}^{(\alpha)}(x) z^n,$$

is analytic near the origin.

In fact, with a procedure perfectly analogous to the one previously used, we can observe that $G_2(z)$ satisfies the non-homogeneous equation

$$(1-z)^2 G_2'(z) - [(\alpha+1)(1-z) - x] G_2(z) = H_1(z),$$

with $G_2(0) = 0$ and

$$H_1(z) = \frac{1}{2} \sum_{n=0}^{\infty} (2n + \alpha + 1 - x) \left[\frac{3}{4} Q_{n,0}^{(\alpha)}(x) + Q_{n,1}^{(\alpha)}(x) \right] z^n.$$

3. Proof of Theorem 1.2.

Suggested by (1.8), we assume that the zeros $x_{n,k}^{(\alpha,\lambda)}$, $k = 1, 2, \dots, n$, of $Q_n^{(\alpha)}(x; \lambda)$ look as

$$(3.1) \quad x_{n,k}^{(\alpha,\lambda)} = l_{n,k}^{(\alpha)} + \frac{1}{\lambda^2} c_1 + \frac{1}{\lambda^4} c_2 + \dots,$$

where $l_{n,k}^{(\alpha)}$ satisfy (1.5).

Putting (3.1) in expansion (1.8) and (1.9) we get

$$(3.2) \quad c_1 = \frac{Q_{n,1}^{(\alpha)}(l_{n,k}^{(\alpha)})}{D_n^{(\alpha)}},$$

where

$$D_n^{(\alpha)} = - \frac{d}{dx} L_n^{(\alpha)}(x) \Big|_{x=l_{n,k}^{(\alpha)}}.$$

The coefficient c_1 may be written in easy form by the repeated use in (1.9) of the following functional relations of Laguerre polynomials, [2, p. 190]

$$(3.3) \quad L_n^{(\alpha+1)}(x) = \frac{1}{x} [(x-n)L_n^{(\alpha)}(x) + (n+\alpha)L_{n-1}^{(\alpha)}(x)],$$

and [2, p. 189]

$$(3.4) \quad x \frac{d}{dx} L_n^{(\alpha)}(x) = nL_n^{(\alpha)}(x) - (n+\alpha)L_{n-1}^{(\alpha)}(x).$$

So, we obtain

$$(3.5) \quad Q_{n,1}^{(\alpha)}(l_{n,k}^{(\alpha)}) = \\ = - \frac{n+\alpha}{6(l_{n,k}^{(\alpha)})^2} L_{n-1}^{(\alpha)}(l_{n,k}^{(\alpha)}) [2(l_{n,k}^{(\alpha)})^2 - (2n+\alpha+1)l_{n,k}^{(\alpha)} + 1 - \alpha^2].$$

Moreover, since (3.4) and (1.5), we have

$$(3.6) \quad D_n^{(\alpha)} = \frac{n+\alpha}{l_{n,k}^{(\alpha)}} L_{n-1}^{(\alpha)}(l_{n,k}^{(\alpha)}).$$

Finally, by means of (3.5), (3.6) and (3.2), Theorem 1.2 is immediately established.

4. Proof of Theorem 1.3.

To prove the first limit relation of the Theorem 1.3, let us set

$$(4.1) \quad R_n^{(\alpha)}(x) = \lim_{\lambda \rightarrow \infty} \frac{\partial Q_n^{(\alpha)}(x; \lambda)}{\partial x},$$

and

$$(4.2) \quad S_n^{(\alpha)}(x) = \lim_{\lambda \rightarrow \infty} \lambda^3 \frac{\partial Q_n^{(\alpha)}(x; \lambda)}{\partial \lambda}.$$

Thus it will be

$$(4.3) \quad \lim_{\lambda \rightarrow \infty} \lambda^3 \frac{\partial x_{n,k}^{(\alpha, \lambda)}}{\partial \lambda} = - \frac{S_n^{(\alpha)}(l_{n,k}^{(\alpha)})}{R_n^{(\alpha)}(l_{n,k}^{(\alpha)})}.$$

To estimate this ratio we establish recurrence formulas for $R_n^{(\alpha)}(x)$ and $S_n^{(\alpha)}(x)$. By differentiating (2.1) and (2.2) with respect to x , we see that $R_n^{(\alpha)}(x)$ satisfies

$$(n+1)R_{n+1}^{(\alpha)}(x) - (2n+\alpha+1-x)R_n^{(\alpha)}(x) + (n+\alpha)R_{n-1}^{(\alpha)}(x) = -L_n^{(\alpha)}(x),$$

with

$$R_{-1}^{(\alpha)}(x) = R_0^{(\alpha)}(x) = 0.$$

In similar manner, by differentiating (2.1) and (2.2) with respect to λ , we find

$$(n+1)S_{n+1}^{(\alpha)}(x) - (2n+\alpha+1-x)S_n^{(\alpha)}(x) + (n+\alpha)S_{n-1}^{(\alpha)}(x) = -(2n+\alpha+1-x)L_n^{(\alpha)}(x),$$

with

$$S_{-1}^{(\alpha)}(x) = S_0^{(\alpha)}(x) = 0.$$

By using the method outlined in Section 2 for the (1.9), we can derive that

$$(4.4) \quad R_n^{(\alpha)}(x) = L_n^{(\alpha)}(x) - L_n^{(\alpha+1)}(x)$$

and

$$(4.5) \quad S_n^{(\alpha)}(x) = -2Q_{n,1}^{(\alpha)}(x),$$

where $Q_{n,1}^{(\alpha)}(x)$ is given by (1.9).

Since (1.5) and (3.3), it follows

$$(4.6) \quad R_n^{(\alpha)}(l_{n,k}^{(\alpha)}) = - \frac{1}{l_{n,k}^{(\alpha)}} (\alpha+n) L_{n-1}^{(\alpha)}(l_{n,k}^{(\alpha)}).$$

So, taking into account (3.5), the limit relation (1.11) can be immediately proved by means of (4.5) and (4.6).

Now, to complete the proof of Theorem 1.3, we first consider that

$$\begin{aligned} \lim_{\lambda \rightarrow \infty} \lambda^4 \frac{\partial^2 x_{n,k}^{(\alpha, \lambda)}}{\partial \lambda^2} &= \\ &= - \frac{T_n^{(\alpha)}(l_{n,k}^{(\alpha)}) R_n^{(\alpha)}(l_{n,k}^{(\alpha)}) - S_n^{(\alpha)}(l_{n,k}^{(\alpha)}) V_n^{(\alpha)}(l_{n,k}^{(\alpha)})}{[R_n^{(\alpha)}(l_{n,k}^{(\alpha)})]^2} + \\ &\quad - \frac{V_n^{(\alpha)}(l_{n,k}^{(\alpha)}) R_n^{(\alpha)}(l_{n,k}^{(\alpha)}) - W_n^{(\alpha)}(l_{n,k}^{(\alpha)}) Z_n^{(\alpha)}(l_{n,k}^{(\alpha)})}{[R_n^{(\alpha)}(l_{n,k}^{(\alpha)})]^2} \lim_{\lambda \rightarrow \infty} \lambda^3 \frac{\partial x_{n,k}^{(\alpha, \lambda)}}{\partial \lambda}, \end{aligned}$$

where $R_n^{(\alpha)}(x)$ and $S_n^{(\alpha)}(x)$ are defined respectively in (4.1) and (4.2) and

$$T_n^{(\alpha)}(x) = \lim_{\lambda \rightarrow \infty} \lambda^4 \frac{\partial^2}{\partial \lambda^2} Q_n^{(\alpha)}(x; \lambda),$$

$$V_n^{(\alpha)}(x) = \lim_{\lambda \rightarrow \infty} \lambda \frac{\partial^2}{\partial x \partial \lambda} Q_n^{(\alpha)}(x; \lambda),$$

$$W_n^{(\alpha)}(x) = \lim_{\lambda \rightarrow \infty} \lambda \frac{\partial}{\partial \lambda} Q_n^{(\alpha)}(x; \lambda),$$

$$Z_n^{(\alpha)}(x) = \lim_{\lambda \rightarrow \infty} \frac{\partial^2}{\partial x^2} Q_n^{(\alpha)}(x; \lambda).$$

By reiterating the previous procedure, we obtain the following results

$$T_n^{(\alpha)}(x) = 6Q_{n,1}^{(\alpha)}(x),$$

$$V_n^{(\alpha)}(x) = W_n^{(\alpha)}(x) = L_n^{(\alpha)}(x),$$

$$Z_n^{(\alpha)}(x) = L_n^{(\alpha+2)}(x) - 2L_n^{(\alpha+1)}(x) + L_n^{(\alpha)}(x),$$

which, remembering (1.5), (3.3), (3.5), (4.3), (4.4) and (4.5), prove the (1.12).

REMARK. In Table 1 we compare the approximations $\bar{x}_{n,k}^{(\alpha, \lambda)}$ obtained by omitting the O -term in the asymptotic representation (1.10) with the exact values $x_{n,k}^{(\alpha, \lambda)}$, for $n = 3$, $\alpha = 2$, and $\lambda = 20, 100$.

TABLE 1.

k	$x_{3,k}^{(2,20)}$	$\bar{x}_{3,k}^{(2,20)}$	$x_{3,k}^{(2,100)}$	$\bar{x}_{3,k}^{(2,100)}$
1	1.5206974	1.5206964	1.5175194542	1.5175194524
2	4.3120309	4.3120301	4.3116010123	4.3116010109
3	9.1672716	9.1672735	9.1708795334	9.1708795366

REFERENCES

- [1] T. S. CHIHARA, *An Introduction to Orthogonal Polynomials*, Gordon & Breach, New York (1978).
- [2] A. ERDÉLYI *et al.*, *Higher Transcendental Functions*, vol. 2, McGraw-Hill Book Co., Inc., New York, N.Y. (1953).
- [3] B. GABUTTI - B. MINETTI, *A New Application of the Discrete Laguerre Polynomials in the Numerical Evaluation of the Hankel Transform of a Strongly Decreasing Even Function*, J. Comput. Phys., **42** (1981), pp. 277-287.
- [4] G. SZEGÖ, *Orthogonal Polynomials*, Colloquium Publications, vol. 23, 4th ed., American Mathematical Society, Providence, R.I. (1975).

Manoscritto pervenuto in redazione il 4 agosto 1993.