

RENDICONTI
del
SEMINARIO MATEMATICO
della
UNIVERSITÀ DI PADOVA

PATRIZIA LONGOBARDI

MERCEDE MAJ

HOWARD SMITH

A note on locally graded groups

Rendiconti del Seminario Matematico della Università di Padova,
tome 94 (1995), p. 275-277

http://www.numdam.org/item?id=RSMUP_1995__94__275_0

© Rendiconti del Seminario Matematico della Università di Padova, 1995, tous droits réservés.

L'accès aux archives de la revue « Rendiconti del Seminario Matematico della Università di Padova » (<http://rendiconti.math.unipd.it/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>

A Note on Locally Graded Groups.

PATRIZIA LONGOBARDI - MERCEDE MAJ - HOWARD SMITH (*)

A group G is locally graded if every nontrivial finitely generated subgroup of G has a nontrivial finite image. The class of locally graded groups is not closed under forming homomorphic images, since free groups (but not all groups) are locally graded. It was shown in [HS] that, if G is locally graded, then G/H is locally graded if, for instance, H is a G -invariant subgroup of the hypercentre of G . A few other results of this kind were established there, but the question as to whether G/H is locally graded if H is abelian was left open. It is not difficult to show that the answer to this question is in the affirmative. Our results are as follows.

THEOREM. *Let G be a locally graded group and H a G -invariant subgroup of the Hirsch-Plotkin radical of G . Then G/H is locally graded.*

COROLLARY. *Suppose G is locally graded and H is a normal subgroup of G . If H is an ascending union of G -invariant subgroups H_λ ($\lambda < \mu$) such that $H_{\lambda+1}/H_\lambda$ is locally nilpotent for each λ , then G/H is locally graded. In particular, if H is soluble then G/H is locally graded.*

(*) Indirizzo degli A.A.: P. LONGOBARDI and M. MAJ: Dipartimento di Matematica, Università di Napoli, Monte S. Angelo, Via Cintia, 80126 Napoli; H. SMITH: Dept. of Mathematics, Bucknell University, Lewisburg, PA 17837, U.S.A.

The proof of our theorem requires the following result.

LEMMA. *Let G be a finitely generated group and suppose that there exists a finitely generated subgroup T of the Hirsch-Plotkin radical H of G such that $G = G' T$. Then $\gamma_j(G)$ is finitely generated, for all $j \geq 1$, and, if c is the nilpotency class of T , then $\gamma_{c+1}(G) = \gamma_{c+i}(G)$ for all $i \geq 1$.*

PROOF. Let $L = \gamma_{c+1}(G)$. Then G/L is nilpotent and $G = G' T$ and so $G = LT$. As G is finitely generated, we may write $G = \langle X, Y \rangle$, for some finite subsets X of L and Y of T , and we may as well assume that Y is closed under forming inverses. Since H is locally nilpotent and normal in G , there is a positive integer k such that $[x, y_1, \dots, y_k] = 1$ for all x in X , y_1, \dots, y_k in Y .

Let $V = \langle X, [x, y_1], [x, y_1, y_2], \dots, [x, y_1, \dots, y_k] : x \in X, y_i \in Y \text{ for all } i \rangle$. Then $\langle X \rangle \leq V$ and $V^y \leq V$ for all $y \in Y$. Since $Y = Y^{-1}$, it follows that V is normal in G . Then $G = VT$ and G/V is nilpotent of class at most c , giving $L \leq V \leq L$. Hence $L = V$, which of course is finitely generated. Since G/L satisfies the maximal condition, all terms of the lower central series of G are thus finitely generated. Finally, since $G = \gamma_{c+2}(G) T$, we see that $G/\gamma_{c+2}(G)$ has class at most c . Thus $L = \gamma_{c+i}(G)$ for all $i \geq 1$.

PROOF OF THE THEOREM. Suppose that the pair (G, H) satisfies the hypotheses of the theorem and suppose, for a contradiction, that G/H is not locally graded. Then, as in the proof of Lemma 1 of [HS], we may assume that G is finitely generated and that G/H is nontrivial but has no nontrivial finite images. This implies that $G = G' H$ and hence that $G = G' T$ for some finitely generated subgroup T of H .

By the lemma, $L = \gamma_{c+1}(G)$ is finitely generated, where c is the nilpotency class of T . Certainly $L \neq 1$, and so it contains a proper normal subgroup N of finite index, and N may be chosen to be normal in G . But now we have $G = HN$ and hence $G/N \cong H/H \cap N$, which is locally nilpotent. Thus G/N is nilpotent. The lemma now gives the contradiction that $L \leq N$.

PROOF OF THE COROLLARY. Assuming the result false, we may choose the ordinal μ to be minimal such that G/H is not locally graded. By the theorem, μ is a limit ordinal. As before, we may now assume that G is finitely generated and that G/H has no nontrivial finite im-

ages. As in the proof of Corollary 6 of [HS], we may further suppose that H is a direct product of G -invariant, finite simple groups. But this of course implies that H is abelian, and the theorem gives us a contradiction.

REFERENCES

- [HS] H. SMITH, *On homomorphic images of locally graded groups*, Rend. Sem. Mat. Univ. Padova, **91** (1994), pp. 53-60.

Manoscritto pervenuto in redazione l'8 aprile 1994
e, in forma revisionata, il 20 ottobre 1994.