

RENDICONTI *del* SEMINARIO MATEMATICO *della* UNIVERSITÀ DI PADOVA

CUI BAOTONG

**Functional differential equations of mixed
type in Banach spaces**

Rendiconti del Seminario Matematico della Università di Padova,
tome 94 (1995), p. 47-54

http://www.numdam.org/item?id=RSMUP_1995__94__47_0

© Rendiconti del Seminario Matematico della Università di Padova, 1995, tous droits réservés.

L'accès aux archives de la revue « Rendiconti del Seminario Matematico della Università di Padova » (<http://rendiconti.math.unipd.it/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>

Functional Differential Equations of Mixed Type in Banach Spaces.

CUI BAOTONG (*)

1. Introduction.

In the last few year the fundamental theory of functional differential equations with a single delay in a Banach space has undergone intensive development (see [1,2]). The fundamental theory of functional differential equations of mixed type, however, is still in a initial stage of development [3-8]. In this paper, we consider the functional differential equation of mixed type of the form

$$(1) \quad x'(t) = f(t, x(t), x(t + \tau(t)), x(t - \tau(t))) \quad \text{a.e. in } (\alpha, \beta),$$

in a Banach space B .

Let $B = (B, \|\cdot\|)$ be a Banach space, $r \geq 0$, $s > 0$ are constants and $-\infty < \alpha < \beta \leq +\infty$. Assume that $f: [\alpha, \beta] \times B \times B \times B \rightarrow B$ satisfies the conditions:

(a) $f(\cdot, x, y, z)$ is strongly measurable for all fixed $x, y, z \in B$ and

there exist $x_0, y_0, z_0 \in B$ such that $\int_{\alpha}^{\beta} \|f(t, x_0, y_0, z_0)\| dt < \infty$;

(b) there exist nonnegative constants a, b and c such that

$$\|f(t, x, y, z) - f(t, \bar{x}, \bar{y}, \bar{z})\| \leq a\|x - \bar{x}\| + b\|y - \bar{y}\| + c\|z - \bar{z}\|,$$

for all $x, y, z, \bar{x}, \bar{y}, \bar{z} \in B$ and $t \in (\alpha, \beta)$; and

(c) $\tau(t)$ is continuous and $r \leq \tau(t) \leq s$.

(*) Indirizzo dell'A.: Department of Mathematics, Binzhou Normal College, Shandong 256604, People's Republic of China.

We consider the equation (1) with two boundary conditions:

$$(2) \quad \begin{cases} x(t) = \phi(t), & t \in [\alpha - s, \alpha], \\ x(t) = \psi(t), & t \in [\beta, \beta + s), \quad \text{if } \beta < \infty, \end{cases}$$

where

$$\phi \in \mathcal{O} = \{ \phi: [\alpha - s, \alpha] \rightarrow B \mid \phi \text{ is Bocher integrable function} \},$$

$$\psi \in \mathfrak{Y} = \{ \psi: [\beta, \beta + s) \rightarrow B \mid \psi \text{ is Bocher integrable function} \},$$

DEFINITION. $x: [\alpha - s, \beta + s) \rightarrow B$ is a solution of the Problem (1) (2), if $x(t)$ is continuous on $[\alpha, \beta)$ and satisfies (1) and (2).

Let

$$\mathcal{A} = \{ x: [\alpha - s, \beta + s) \rightarrow B \mid x(t) \text{ is continuous on } [\alpha, \beta) \text{ and satisfies (2)} \},$$

$$\mathcal{A}_\lambda = \{ x \in \mathcal{A} \mid \sup_{\alpha \leq t < \beta} \exp[-\lambda t] \|x(t)\| < \infty \} \quad \text{for each } \lambda \geq 0.$$

It is easy to see that x is a solution of the problem (1), (2) if and only if x is a fixed point of the operator T , defined on $\mathcal{A}$ by

$$(3) \quad Tx(t) = \begin{cases} \phi(t), & t \in [\alpha - s, \alpha], \\ \psi(t), & t \in [\beta, \beta + s), \quad \text{if } \beta < \infty, \\ \phi(\alpha) + \int_{\alpha}^t f(u, x(u), x(u + \tau(u)), x(u - \tau(u))) du, & t \in (\alpha, \beta). \end{cases}$$

2. Existence and uniqueness.

THEOREM 1 (Existence and uniqueness). Let (a), (b) and (c) hold and $s \leq l = \beta - \alpha$. If there exists a $\lambda \geq 0$ such that

$$(4) \quad D(a, b, c, \lambda) = a \int_0^l e^{-\lambda u} du + b \int_{2r-l}^s e^{\lambda u} du + c \int_{2r-s}^l e^{-\lambda u} du < 1,$$

then, for each $x_0 \in \mathcal{A}_\lambda$ the iterates $T^n x_0$, where $T: \mathcal{A}_\lambda \rightarrow \mathcal{A}_\lambda$ is defined by

(3), converge in the metric ϱ_λ of $\mathcal{C}_\lambda$ defined by

$$(5) \quad \varrho_\lambda(x, \bar{x}) = \sup_{\alpha \leq t < \beta} \{e^{-\lambda t} \|x(t) - \bar{x}(t)\|\}$$

to a solution of the problem (1) (2), which is unique in $\mathcal{C}_\lambda$.

PROOF. Let $x \in \mathcal{C}_\lambda$, then $\sup_{\alpha \leq t < \beta} e^{-\lambda t} \|x(t)\| < \infty$ and $x(t)$ is continuous on $[\alpha, \beta)$ and satisfies (2). Hence $Tx(t)$ is continuous on $[\alpha, \beta)$ and satisfies (2).

From (3), for each $\lambda > 0$ and any $t \in [\alpha, \beta)$, we have that

$$\begin{aligned} e^{-\lambda t} \|Tx(t)\| &\leq \\ &\leq e^{-\lambda t} \left\| Tx(t) - \int_\alpha^\beta f(t, x_0, y_0, z_0) dt \right\| + e^{-\lambda t} \left\| \int_\alpha^\beta f(t, x_0, y_0, z_0) dt \right\| \leq \\ &\leq e^{-\lambda t} \|\phi(\alpha)\| + e^{-\lambda t} \left\| \int_\alpha^t f(u, x(u), x(u + \tau(u)), x(u - \tau(u))) du - \right. \\ &\quad \left. - \int_\alpha^t f(u, x_0, y_0, z_0) du \right\| + \\ &+ e^{-\lambda t} \left\| \int_t^\beta f(u, x_0, y_0, z_0) du \right\| + e^{-\lambda t} \left\| \int_\alpha^\beta f(t, x_0, y_0, z_0) dt \right\| \leq e^{-\lambda t} \|\phi(\alpha)\| + \\ &+ e^{-\lambda t} \left[a \int_\alpha^t \|x(u)\| du + b \int_\alpha^t \|x(u + \tau(u))\| du + c \int_\alpha^t \|x(u - \tau(u))\| du \right] + \\ &+ e^{-\lambda t} (t - \alpha)(a\|x_0\| + b\|y_0\| + c\|z_0\|) + 2e^{-\lambda t} \left\| \int_\alpha^\beta f(t, x_0, y_0, z_0) dt \right\|. \end{aligned}$$

Note that $x \in \mathcal{C}_\lambda$, ϕ and ψ are Bocher integrable functions, so we can obtain that

$$\sup_{\alpha \leq t < \beta} e^{-\lambda t} \|Tx(t)\| < \infty.$$

For $\lambda = 0$, using (a) and $x \in \mathcal{C}_0$, i.e. $\sup_{\alpha \leq t < \beta} \|Tx(t)\| < \infty$, we have $Tx \in \mathcal{C}_0$. So $T(\mathcal{C}_\lambda) \subset \mathcal{C}_\lambda$ for each $\lambda \geq 0$.

Now we define

$$\mathcal{B} = \{y: [\alpha - s, \beta + s] \rightarrow (0, \infty) | y(t) \text{ is continuous in } [\alpha, \beta], \\ y(t) = 0 \text{ for } t \in [\alpha - s, \alpha) \text{ and } t \in [\beta, \beta + s)\}$$

and

$$Wy(t) = 0 \quad \text{for } t \in [\alpha - s, \alpha) \text{ and } t \in [\beta, \beta + s),$$

$$Wy(t) =$$

$$= a \int_{\alpha}^t y(u) du + b \int_{\alpha}^t y(u + \tau(u)) du + c \int_{\alpha}^t y(u - \tau(u)) du \quad \text{for } t \in [\alpha, \beta],$$

then the operator W maps the set $\mathcal{B}$ into itself. We can prove that

$$\|Wy\|_{\lambda} < D(a, b, c, \lambda) \|y\|_{\lambda} \quad \text{for any } y \in \mathcal{B} \text{ and each } \lambda \geq 0,$$

where

$$\|y\|_{\lambda} = \sup_{\alpha \leq t < \beta} \{e^{-\lambda t} y(t)\} \quad \text{for any } y \in \mathcal{B},$$

and

$$\|Tx - T\bar{x}\| \leq W(\|x(t) - \bar{x}(t)\|) \quad \text{for any } x, \bar{x} \in \mathcal{C}_{\lambda} \text{ and } t \in [\alpha, \beta].$$

Thus from (5) we have that

$$\varrho_{\lambda}(Tx, T\bar{x}) \leq D(a, b, c, \lambda) \varrho_{\lambda}(x, \bar{x}), \quad x, \bar{x} \in \mathcal{C}_{\lambda}.$$

Hence T is contractive with respect to ϱ_{λ} by (4). Since $\mathcal{C}_{\lambda}$ is complete in this metric, the Banach's contractive mapping theorem implies that the iterates $T^n x_0, x_0 \in \mathcal{C}_{\lambda}$, converge in $(\mathcal{C}_{\lambda}, \varrho_{\lambda})$ to a unique fixed point of T , i.e. to a unique solution of the problem (1), (2) in $\mathcal{C}_{\lambda}$.

COROLLARY 1. Let (a) and (b) hold, and $\tau(t) = r = \text{const} > 0$. If there exists $\lambda \geq 0$ such that

$$(6) \quad a \int_0^l e^{-\lambda u} du + b \int_{2r-l}^r e^{\lambda u} du + c \int_r^l e^{-\lambda u} du < 1,$$

then the problem (1), (2) has an unique solution in $\mathcal{C}_{\lambda}$.

EXAMPLE. Consider the Lecornu's equations [4,6]

$$(7) \quad x'(t) = ax(t+1) + bx(t-1)$$

where a and b are constants and $\tau(t) = 1$. The conditions (a) and (b) are verified. By the Corollary 1 and (6), if there exists a $\lambda \geq 0$ such that

$$|\alpha| \int_{\alpha-l}^1 e^{\lambda u} du + |b| \int_1^l e^{-\lambda u} du < 1, \text{ then the problem (7), (2) has a unique}$$

solution in $\mathcal{C}_\lambda$ for $r = s = 1$.

COROLLARY 2. Let the hypotheses of the Theorem 1 hold and $\beta < \infty$, then the problem (1), (2) has a bounded solution on $[\alpha, \beta)$.

PROOF. Since $\beta < \infty$, then $\mathcal{C}_\lambda = \mathcal{C}_0$ for all $\lambda \geq 0$. Hence the problem (1), (2) has a unique solution in $\mathcal{C}_0$ by the Theorem 1, and the solution is bounded.

3. Dependence on the boundary equations.

THEOREM 2 (Dependence). Let all conditions of the Theorem 1 hold. Suppose that $\{\phi_n\} \subset \mathcal{O}$, $\{\psi_n\} \subset \mathfrak{I}$,

$$\|\phi_n - \phi\| \stackrel{\text{def}}{=} \int_{\alpha-s}^{\alpha} \|\phi_n(t) - \phi(t)\| dt \rightarrow 0, \quad n \rightarrow \infty,$$

$$\|\psi_n - \psi\| \stackrel{\text{def}}{=} \int_{\beta}^{\beta+s} \|\psi_n(t) - \psi(t)\| dt \rightarrow 0, \quad n \rightarrow \infty,$$

and $\phi_n(\alpha) \rightarrow \phi(\alpha)$ as $n \rightarrow \infty$. Let $x_n(t)$ be the solution of the equation (1) with the boundary conditions

$$(8)_n \quad \begin{cases} x_n(t) = \phi_n(t), & t \in [\alpha - s, \alpha], \\ x_n(t) = \psi_n(t), & t \in [\beta, \beta + s), \quad \text{if } \beta < \infty, \end{cases}$$

$n = 1, 2, \dots$. Then the sequence $\{x_n(t)\}$ of the solutions of the problem (1), $(8)_n$ ($n = 1, 2, \dots$) has the following properties:

- (i) if $\beta < \infty$, then $x_n(t) \rightarrow x(t)$ as $n \rightarrow \infty$ uniformly on $[\alpha, \beta)$;
- (ii) if $\beta = \infty$, then $x_n(t) \rightarrow x(t)$ as $n \rightarrow \infty$ uniformly on the compact intervals of $[\alpha, \beta)$.

PROOF. Let

$$\mathcal{A}_n = \{x: [\alpha - s, \beta + s] \rightarrow B \mid x(t) \text{ is continuous on } [\alpha, \beta]$$

and satisfies (8)_n \}

$$\mathcal{A}_{n\lambda} \equiv \{x \in \mathcal{A}_n \mid \sup e^{-\lambda t} \|x(t)\| < \infty\} \quad \text{for each } \lambda \geq 0,$$

$n = 1, 2, \dots$, and we define the operators T_n :

$$(9)_n \quad T_n x_n(t) = \begin{cases} \phi_n(t), & t \in [\alpha - s, \alpha], \\ \psi_n(t), & t \in [\beta, \beta + s), \quad \text{if } \beta < \infty, \\ \phi_n(\alpha) + \int_{\alpha}^t (u \cdot x_n(u), x_n(u + \tau(u)), x_n(u - \tau)) du, & t \in [\alpha, \beta), \end{cases}$$

for $n = 1, 2, \dots$. Similarly to the proof of the Theorem 1, we can prove that $T_n(\mathcal{A}_{n\lambda}) \subset \mathcal{A}_{n\lambda}$ and the operator T_n has a unique fixed point for each n . Hence for each n the problem (1), (6)_n has a unique solution in $\mathcal{A}_{n\lambda}$.

Let $x \in \mathcal{A}_\lambda$ be the fixed point of T and $x_n \in \mathcal{A}_{n\lambda}$ be the fixed point of T_n for each n , then we have that

$$\varrho_\lambda(x, x_n) \leq M_n e^{-\lambda \alpha} / (1 - D(a, b, c, \lambda)) \quad \text{for all } n,$$

where $M_n = \text{Const.} \geq 0$ and $\lim_{n \rightarrow \infty} M_n = 0$. Thus from $\lim_{n \rightarrow \infty} \varrho_\lambda(x, x_n) = 0$, it is easy to prove our results.

4. Remarks.

REMARK 1.. The above results can be extended naturally to the problem involving several arguments

$$(10) \quad x'(t) = f(t, x(t), x(t + \tau_1(t)), x(t + \tau_2(t)), \dots, x(t + \tau_m(t)),$$

$$x(t - \tau_1(t)), x(t - \tau_2(t)), \dots, x(t - \tau_m(t))) \quad \text{a.e. in } (\alpha, \beta),$$

where $\tau_1(t) > 0$ ($i = 1, 2, \dots, m$) are continuous, and the exist nonnegative constants r_i and s_i ($i = 1, 2, \dots, m$) such that $0 \leq r_i \leq \tau_i(t) \leq s_i$ and

also to functional differential equation of mixed type of the form

$$x'(t) = f(t, x(t), x(t + \tau_1(t)), x(t + \tau_2(t)), \dots, x(t + \tau_p(t)), \\ x(t - h_1(t)), x(t - h_2(t)), \dots, x(t - h_q(t))) \quad \text{a.e. in } (\alpha, \beta),$$

where p and q are positive numbers, $\tau_i(t)$ and $h_j(t)$ are nonnegative continuous functions.

REMARK 2. The equations

$$(11) \quad x'(t) = f(t, x(t), x(g(t)t), x(h(t)t)) \quad \text{a.e. in } (\alpha, \infty),$$

and

$$x'(t) = f(t, x(t), x(g_1(t)t), x(g_2(t)t), \dots, x(g_p(t)t), \\ x(h_1(t)t), x(h_2(t)t), \dots, x(h_q(t)t)) \quad \text{a.e. in } (\alpha, \infty),$$

where g, h, g_i ($i = 1, 2, \dots, p$) and h_j ($j = 1, 2, \dots, q$) are continuous, and $g > 1, g_i > 1, 0 < h < 1, 0 < h_j < 1$ for each i and j , are of type (10) too.

In fact, let $t = e^y$, $X(y) = x(e^y)$, then we have that from (11)

$$X'(y) = x'(e^y)e^y = e^y f(e^y, x(e^y), x(e^{y + \log g(e^y)}), x(e^{y + \log h(e^y)})) \stackrel{\text{def}}{=} \\ \stackrel{\text{def}}{=} F(y, X(y), X(y + \log g(e^y)), X(y + \log h(e^y))) \quad \text{a.e.}$$

where $\log(e^y) > 0, \log h(e^y) < 0$.

Acknowledgment. The author is grateful to the referee for helpful suggestions.

REFERENCES

- [1] V. LAKSHMIKANTHAM - S. LEELA, *Nonlinear Differential Equations in Abstract Spaces*, Pergamon Press, New York (1981).
- [2] F. J. BURKOWSKI, *Existence and uniqueness theorems for differential equations with deviating arguments of mixed type*, in *Delay and Functional Differential Equations and Their Applications* (Edited by Klaus Schmitt), Academic Press, New York (1972), pp. 309-311.
- [3] CUI BAOTONG - XIA DAFENG, *Existence and uniqueness of analytic solutions of functional differential equations*, *Fuyang Shiyuan Xuebao*, 2 (1989), pp. 87-93.

- [4] ZHENG ZUXIU, *Functional Differential Equations*, Xi'an Jiaotong University Press (1989).
- [5] CUI BAOTONG, *Asymptotic stability of the solutions of nonlinear functional differential equations of neutral type*, J. Math. Res. & Exposition, **11** (1991), p. 423.
- [6] J. HALE, *Theory of Functional Differential Equations*, Springer-Verlag, New York (1977).
- [7] CUI BAOTONG, *Asymptotic stability of large systems of neutral type*, Matematica Applicata, **3**(3) (1989), pp. 81-83.
- [8] CUI BAOTONG, *Oscillation criteria for n -th order delay differential equations*, Ann. Diff. Eqs., **7** (1991), pp. 1-4.

Manoscritto pervenuto in redazione il 12 novembre 1991
e, in forma revisionata, il 17 gennaio 1994.