

RENDICONTI *del* SEMINARIO MATEMATICO *della* UNIVERSITÀ DI PADOVA

A. MOHAMMADI HASSANABADI

Residually finite PSP-groups

Rendiconti del Seminario Matematico della Università di Padova,
tome 101 (1999), p. 59-61

http://www.numdam.org/item?id=RSMUP_1999__101__59_0

© Rendiconti del Seminario Matematico della Università di Padova, 1999, tous droits réservés.

L'accès aux archives de la revue « Rendiconti del Seminario Matematico della Università di Padova » (<http://rendiconti.math.unipd.it/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

*Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques*
<http://www.numdam.org/>

Residually Finite PSP-Groups.

A. MOHAMMADI HASSANABADI (*)

ABSTRACT - Let G be a group. If there exists an integer $n > 1$ such that for each n -tuple (H_1, \dots, H_n) of subgroups of G there is a non-identity permutation σ in S_n such that the two sets $H_1 \dots H_n$ and $H_{\sigma(1)} \dots H_{\sigma(n)}$ are equal, then G is said to have the property of permutable subgroup products. This is a note to show that a finitely generated residually finite group has the above property if and only if it is finite-by-abelian.

1. - Introduction.

Groups with the property of permutable subgroup products are denoted by PSP. If $n > 1$ is an integer then G is a PSP_n -group if for each n -tuple (H_1, \dots, H_n) of subgroups of G there is a permutation $\sigma \neq 1$ in S_n such that $H_1 \dots H_n = H_{\sigma(1)} \dots H_{\sigma(n)}$. Thus PSP is the union of the classes PSP_n ; $n = 2, 3, \dots$

It was proved in [1] that a finitely generated residually soluble group G is a PSP-group if and only if G is finite-by-abelian. But the question(**) whether finitely generated residually finite groups have the same property was left open. The result of this note is the following which answers the above question affirmatively.

(*) Indirizzo dell'A.: Department of Mathematics, University of Isfahan, Isfahan, Iran.

(**) The author thanks Professor Akbar Rhemtulla, of the Department of Mathematical Sciences, University of Alberta for drawing his attention to this question. He also would like to thank the Referee for his valuable suggestions.

THEOREM. *Suppose that G is a finitely generated residually finite group. Then G is a PSP-group if and only if G is finite-by-abelian.*

2. – Proof.

We first prove the following Lemma in which we use the notation of [2] for twisted wreath products. We recall the definition of twisted wreath product (compare Neumann [2]). Let A, B be finite groups and H a subgroup of B , and suppose that a faithful action σ of H on A is given. We may form the direct product F of $|B:H|$ copies of A , and define a faithful action of B on F in such a way that B permutes transitively the copies of A and H acts on the first of these copies according to the action of σ . The split extension of F by B with this action is the twisted wreath product of A by B with respect to H ; it will be denoted by $\text{Atwr}(H)B$ and reference to the action σ will be suppressed. The subgroup F is called the base group of the twisted wreath product. If $H = 1$, the group becomes the standard wreath product $\text{Awr}B$ of A by B . We are concerned with the case in which B is cyclic and A is an elementary abelian p -group for some prime p , and H acts faithfully and irreducibly on A .

LEMMA. *Let $G = \text{Atwr}(H)B$ be the twisted wreath product of a finite elementary abelian group A and a finite cyclic group B with respect to a subgroup H of B , where H acts on A irreducibly. If G is a PSP_n -group, $n > 1$ and $|B:H| = m$, then $m < 7^{2n}$.*

PROOF. Let $B = \langle b \rangle$ so that a right transversal to H in B can be chosen to be $T = \{1, b, b^2, \dots, b^{m-1}\}$. Let $G = BF$, where $F = \{f: T \rightarrow A\} = A^T$. For each $i, i = 0, 1, \dots, m-1$, put $A_i = \{f \in F | f(b^j) = 1 \text{ for all } j \neq i\}$. Then for $f \in A_0, f^b(b^i) = 1$ for all $i \neq 1$, since for each $t \in T, f^b(t) = f(t^{b^{-1}})^{h(b,t)}$ where $t^{b^{-1}} = (tb^{-1})^\tau, h(b,t) = (tb^{-1})^{\tau-1}$, and τ is the mapping of B to T that maps elements of B onto their coset representatives in T . Thus for $t = b^i, 0 < i < m$, we have $t^{b^{-1}} = (b^i b^{-1})^\tau = (b^{i-1})^\tau = b^{i-1}$, so that $f^b(t) = f(t^{b^{-1}})^{h(b,t)} = f(b^{i-1})^{h(b,t)} = f(b^{i-1})^{(b^{i-1})^{\tau-1}} = 1$, if $i \neq 1$. Therefore $A_1 = A_0^b$. Similarly $A_i^b = A_{i+1}$ for $i = 1, 2, \dots, m-1$ and $A_m = A_0$.

Now the proof follows from Lemma 2.1 of [1], since the conditions of that Lemma are satisfied. We include the statement of Lemma 2.1 of [1] for convenience.

LEMMA 2.1 of [1]. *Let G be the split extension of a finite abelian group F of prime exponent p and a finite cyclic group $\langle b \rangle$. Suppose that F is the direct sum $A_0 \oplus \dots \oplus A_{m-1}$ of m subgroups each normalized by $\langle b^m \rangle$ and $A_i^b = A_{i+1}$ for all $i = 1, 2, \dots, m-1$ and $A_m = A_0$. If G is a PSP $_n$ -group, $n > 1$ then $m < 7^{2n}$.*

PROOF OF THE THEOREM. Let G be a finitely generated residually finite PSP-group. By the Lemma and Theorem 4 of [4], G is soluble-by-finite. By the main theorem of [1] G is finite-by-abelian-by-finite. We can assume that there is a torsion-free abelian normal subgroup A with G/A finite.

Let $g \in G$. Then g acts nilpotently on A by [3], Lemma 2. Since $|gA|$ is finite, it follows that $[A, g] = 1$. Hence $A \leq Z(G)$, $G/Z(G)$ is finite and G' is finite, as required.

The converse follows from the proof of the main result of [3].

REFERENCES

- [1] P. LONGOBARDI - M. MAJ - A. H. RHEMTULLA, *Residually solvable PSP-groups*, Boll. Un. Mat. Ital., (7) 7-B (1993), pp. 253-261.
- [2] B. H. NEUMANN, *Twisted wreath product of groups*, Arch. Math. (Basel), 14 (1963), pp. 1-6.
- [3] A. H. RHEMTULLA - A. R. WEISS, *Groups with permutable subgroup products*, *Proceeding of the 1987 Singapore Conference in Group Theory*, Walter der Gruyter, Berlin, New York.
- [4] J. S. WILSON, *Two-generator conditions for residually finite groups*, Bull. London Math. Soc., 23 (1991), pp. 239-248.

Manoscritto pervenuto in redazione il 9 aprile 1997.