

SÉMINAIRE D'ANALYSE FONCTIONNELLE ÉCOLE POLYTECHNIQUE

T. STARBIRD

Enflo operators of L^1

Séminaire d'analyse fonctionnelle (Polytechnique) (1975-1976), exp. n° 24, p. 1-5

<http://www.numdam.org/item?id=SAF_1975-1976___A19_0>

© Séminaire Maurey-Schwartz
(École Polytechnique), 1975-1976, tous droits réservés.

L'accès aux archives du séminaire d'analyse fonctionnelle implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>

ÉCOLE POLYTECHNIQUE

CENTRE DE MATHÉMATIQUES

PLATEAU DE PALAISEAU - 91120 PALAISEAU

Téléphone : 941.81.60 - Poste N°

Télex : ECOLEX 091596 F

S E M I N A I R E M A U R E Y - S C H W A R T Z 1 9 7 5 - 1 9 7 6

ENFLO OPERATORS ON L^1

par T. STARBIRD
(Berkeley)

We present here a summary of some results obtained by P. Enflo and the author. The complete proofs can be found in [1] or [6]. We study a class of operators on L^1 called Enflo operators. The definition of an Enflo operator is given below. Theorem 1 gives a characterization of Enflo operators which could serve as an alternate definition. Theorem 2 is the main structural result concerning Enflo operators. Several corollaries are then derived.

We deal with $L^1 = L^1([0,1],\mu)$, the Banach space of equivalence classes of Lebesgue integrable real-valued functions defined on $[0,1]$. μ is Lebesgue measure. An "isomorphism" is a linear homeomorphism into. An "isomorph of L^1 " is a Banach space isomorphic (i.e. linearly homeomorphic) to L^1 . If two Banach spaces X and Y are isomorphic, we write $X \sim Y$. "Subspace" will mean "closed linear subspace". If E is a (Lebesgue) measurable subset of $[0,1]$, we write $|E|$ for $\mu(E)$. χ_E denotes the characteristic function of E , that is,

$$\chi_E(t) = \begin{cases} 1 & \text{if } t \in E \\ 0 & \text{if } t \notin E \end{cases} .$$

Definitions :

(1) A collection $\{F_i^n : i = 1, 2, \dots, 2^n ; n = 0, 1, \dots\}$ of measurable subsets of $[0,1]$ is called a tree if

- (a) $|F_1^0| > 0$;
- (b) for any n and i , $1 \leq i \leq 2^n$,

$$F_{2i-1}^{n+1} \cup F_{2i}^{n+1} = F_i^n ;$$

- (c) $F_i^n \cap F_j^n = \emptyset$ whenever $i \neq j$, $1 \leq i, j \leq 2^n$;
- and (d) $\max_{1 \leq i \leq 2^n} |F_i^n| \rightarrow 0$ as $n \rightarrow \infty$.

(2) A collection $\{F_i^n : i = 1, 2, \dots, M_n ; n = 0, 1, \dots\}$ of measurable subsets of $[0,1]$ is called a bush if

- (a) $M_0 = 1$ and $|F_1^0| > 0$,
- (b) for any n and i , $1 \leq i \leq M_{n+1}$, the set F_i^{n+1} is contained in some F_j^n ;
- (c) for each n , the collection $\{F_i^n : i = 1, \dots, M_n\}$ forms a partition of F_1^0 ;

and (d) $\max_{1 \leq i \leq M_n} |F_i^n| \rightarrow 0$ as $n \rightarrow \infty$.

(3) Let $T: L^1 \rightarrow L^1$ be a bounded linear operator. T is called an Enflo operator if there exist $\delta > 0$ and a bush (E_i^n) , $i = 1, \dots, M_n$; $n = 0, 1, \dots$ of subsets of $[0, 1]$ such that

$$\frac{1}{|E_1^0|} \int \max_{1 \leq i \leq M_n} |T(\chi_{E_i^n})| > \delta$$

for each n . If T is an Enflo operator and $\lambda > 0$, T is called an Enflo operator of constant λ if

$$\lambda \leq \sup \lim_{n \rightarrow \infty} \frac{1}{|E_1^0|} \int \max_{1 \leq i \leq M_n} |T(\chi_{E_i^n})| ,$$

where the supremum is taken over all bushes (E_i^n) in which $|E_1^0| > 0$. (It can be shown that actually for any such bush, $\lim_{n \rightarrow \infty} \frac{1}{|E_1^0|} \int \max_{1 \leq i \leq M_n} |T(\chi_{E_i^n})|$ exists.)

Our first major result is

Theorem 1 : Let $T: L^1 \rightarrow L^1$ be a bounded linear operator. T is an Enflo operator if and only if there exists a subspace Y of L^1 with Y isomorphic to L^1 and with $T|_Y$ an isomorphism (into).

The proof of Theorem 1 depends upon the next Theorem :

Theorem 2 : Suppose T is an Enflo operator of constant λ , and $0 < \varepsilon < 1/2$. Then there exist a measurable subset E_0 of $[0, 1]$ and a σ -algebra \mathcal{Q} of subsets of E_0 such that

1. $\mu|_{\mathcal{Q}}$ is purely non-atomic, and hence $L^1(\mu|_{\mathcal{Q}})$ is isometric to L^1 (see [2]) ;
2. $T|_{L^1(\mu|_{\mathcal{Q}})}$ is an isomorphism ; for $f \in L^1(\mu|_{\mathcal{Q}})$,

$$\|Tf\| \geq \left(\frac{1 - \varepsilon}{1 + \varepsilon} \right)^2 \lambda \|f\| ;$$

3. The image $T(L^1(\mu|_{\mathcal{Q}}))$ is complemented ; it is the range of a projection of norm at most

$$\frac{\|T\|}{\lambda} \left(\frac{1 + \varepsilon}{1 - \varepsilon} \right)^2 .$$

In fact, there is a tree (A_i^n) , $i = 1, \dots, 2^n$; $n = 0, 1, \dots$ of measurable subsets of $E_0 = A_1^0$ with

$$(a) \quad (1 - \varepsilon) \frac{|E_0|}{2^n} \leq |A_i^n| \leq (1 + \varepsilon) \frac{|E_0|}{2^n}$$

and there is a tree (F_i^n) , $i = 1, \dots, 2^n$; $n = 0, 1, \dots$ of measurable subsets of a set $F_1^0 \subset [0, 1]$ such that for each n and i , $1 \leq i \leq 2^n$,

$$(b) \quad (1 - \varepsilon)\lambda \frac{|E_0|}{2^n} \leq \int_{F_1^0} |T\chi_{A_i^n}| < (1 + \varepsilon) \int_{F_i^n} |T\chi_{A_i^n}| .$$

Remark : Conclusions 1, 2, and 3 follow, using facts about relative disjointness (see [5]), from the existence of the trees (A_i^n) and (F_i^n) satisfying (a) and (b).

Remark : Note that Theorem 2, conclusions 1 and 2, asserts a strong form of the direct implication claimed in Theorem 1.

Corollary 1 : Let $Z = L^1$. If X is a subspace of Z and $X \sim L^1$, then there exists a subspace Y of X with $Y \sim L^1$ and Y complemented in Z .

Proof : Let $T : L^1 \rightarrow Z$ be any isomorphism onto X , apply Theorem 1 and then Theorem 2.

Q.E.D.

A consequence of Corollary 1 together with the decomposition method of Pełczyński [4] is that if a complemented subspace X of L^1 contains a subspace isomorphic to L^1 , then $X \sim L^1$.

Before stating the next corollary, we make the following

Remark : If $T_1 + T_2$ is an Enflo operator, then either T_1 or T_2 must be an Enflo operator. For

$$(*) \quad \int \max_i |(T_1 + T_2)E_i^n| \leq \int \max_i |T_1 E_i^n| + \int \max_i |T_2 E_i^n|$$

for each n and for any bush (E_i^n) . If neither T_1 nor T_2 is an Enflo operator, then as $n \rightarrow \infty$ the limit of the right-hand side, and hence the left-hand side, of (*) is 0. Since this is true for all bushes, $T_1 + T_2$ cannot be an Enflo operator.

Now we can state the next corollary, an alternate proof of which is given in [3].

Corollary 2 : L^1 is primary ; i.e., if $L^1 \sim X \oplus Y$, then either $X \sim L^1$ or $Y \sim L^1$ (or both).

Proof : Consider X and Y as complementary subspaces in L^1 , with projections P onto X and $I-P$ onto Y . Then, since these two operators sum to the identity operator, which is certainly an Enflo operator, one of them (let us say P) is an Enflo operator.

Hence by theorem 2 the range of P , which is X , must contain a complemented subspace isomorphic to L^1 . Pełczyński's decomposition method [4] then implies that $X \sim L^1$.

Q.E.D.

Corollary 3 : Let $T: L^1 \rightarrow L^1$ be a bounded linear operator. If there exists a subspace Y isomorphic to L^1 with $T|_Y$ an isomorphism, then there exists a subspace Z isometric to L^1 with $T|_Z$ an isomorphism.

Proof : Combine Theorems 1 and 2.

Q.E.D.

Recall that if $T: L^1 \rightarrow L^1$ is a bounded linear operator, its absolute value, $|T|$, is the operator on L^1 defined for $f \geq 0$, $f \in L^1$ by

$$(|T|f)(t) = \sup \left\{ \sum_{i=1}^m |Tf_i(t)| \mid \sum_{i=1}^m f_i = f, f_i \geq 0 \right\}$$

for all t in $[0,1]$, and defined for general $f \in L^1$ by linearity, writing f as the difference of two non-negative functions.

Proposition : Let $T: L^1 \rightarrow L^1$ be a bounded linear operator. T is an Enflo operator if and only if $|T|$ is an Enflo operator.

BIBLIOGRAPHIE

- [1] P. Enflo and T. Starbird, Operators on L^1 which are isomorphisms on subspaces isomorphic to L^1 , in preparation.
- [2] D. Maharam, On homogeneous measure algebras, Proc. Nat. Acad. Sci. USA, 28 (1942), 108-111.
- [3] B. Maurey, Sous-espaces complémentés de L^p , d'après P. Enflo, Séminaire Maurey-Schwartz 1974-75, No 3.
- [4] A. Pełczyński, Projections in certain Banach spaces, Studia Math. 19 (1960), 209-228.
- [5] H.P. Rosenthal, On relatively disjoint families of measures, with some applications to Banach space theory, Studia Math. 37 (1970), 13-36.
- [6] T. Starbird, Subspaces of L^1 containing L^1 , Ph. D. dissertation, University of California at Berkeley, June, 1976.
