

SÉMINAIRE N. BOURBAKI

FRIEDRICH HIRZEBRUCH

Singularities and exotic spheres

Séminaire N. Bourbaki, 1968, exp. n° 314, p. 13-32

<http://www.numdam.org/item?id=SB_1966-1968__10__13_0>

© Association des collaborateurs de Nicolas Bourbaki, 1968, tous droits réservés.

L'accès aux archives du séminaire Bourbaki (<http://www.bourbaki.ens.fr/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

SINGULARITIES AND EXOTIC SPHERES

by Friedrich HIRZEBRUCH

BRIESKORN has proved [4] that the n -dimensional affine algebraic variety $z_0^3 + z_1^2 + \dots + z_n^2 = 0$ (n odd, $n \geq 1$) is a topological manifold though the variety has an isolated singular point (which is normal for $n \geq 2$). Such a phenomenon cannot occur for normal singularities of 2-dimensional varieties, as was shown by MUMFORD ([12], [6]). BRIESKORN's result stimulated further research on the topology of isolated singularities (BRIESKORN [5], MILNOR [11] and the speaker [5], [7]). BRIESKORN [5] uses the paper of F. PHAM [14], whereas the speaker studied certain singularities from the point of view of transformation groups using results of BREDON ([2], [3]), W.C. HSIANG and W.Y. HSIANG [8] and JÄNICH [9].

§ 1. The integral homology of some affine hypersurfaces.

PHAM [14] studies the non-singular subvariety $V_a = V(a_0, a_1, \dots, a_n)$ of \mathbb{C}^{n+1} given by

$$z_0^{a_0} + z_1^{a_1} + \dots + z_n^{a_n} = 1 \quad (n \geq 0),$$

where $a = (a_0, \dots, a_n)$ consists of integers $a_j \geq 2$.

Let G_{a_j} be the cyclic group of order a_j multiplicatively written and generated by w_j . Define the group $G_a = G_{a_0} \times G_{a_1} \times \dots \times G_{a_n}$ and put $\varepsilon_j = \exp(2\pi i/a_j)$.

Then $w_0^{k_0} w_1^{k_1} \dots w_n^{k_n}$ is an element of G_a whereas $\varepsilon_0^{k_0} \varepsilon_1^{k_1} \dots \varepsilon_n^{k_n}$ is a complex number. G_a operates on V_a by

$$w_0^{k_0} \dots w_n^{k_n} (z_0, \dots, z_n) = (\varepsilon_0^{k_0} z_0, \dots, \varepsilon_n^{k_n} z_n).$$

Let \hat{G}_{a_j} be the group of a_j -th roots of unity and $x \mapsto \hat{x}$ the isomorphism $G_{a_j} \rightarrow \hat{G}_{a_j}$ given by $w_j \mapsto \varepsilon_j = \hat{w}_j$.

PHAM considers the following subspace U_a of V_a

$$U_a = \{z \mid z \in V_a \text{ and } z_j^{a_j} \text{ real } \cong 0 \text{ for } j = 0, \dots, n\}$$

LEMMA.- The subspace U_a is a deformation retract of V_a by a deformation compatible with the operations of G_a .

For the proof see PHAM [14], p. 338.

U_a can also be described by the conditions

$$z_j = u_j |z_j| \quad \text{with } u_j \in \hat{G}_{a_j} \quad (j = 0, \dots, n).$$

Put $|z_j|^{a_j} = t_j$. Then U_a becomes the space of $(n+1)$ -tuples of complex numbers

$$t_0 u_0 \oplus t_1 u_1 \oplus \dots \oplus t_n u_n$$

with

$$u_j \in \hat{G}_{a_j}, \quad t_j \geq 0, \quad \sum_{j=0}^n t_j = 1$$

Thus U_a can be identified with the join $G_{a_0} * G_{a_1} * \dots * G_{a_n}$ of the finite sets G_{a_j} (see MILNOR [10]).

LEMMA 2.1 in [10] states in particular that the reduced integral homology groups of the join $A * B$ of two spaces A, B without torsion are given by a canonical isomorphism

$$\tilde{H}_{r+1}(A * B) \cong \sum_{i+j=r} \tilde{H}_i(A) \otimes \tilde{H}_j(B),$$

whereas LEMMA 2.2 in [10] shows that $A * B$ is simply connected provided B is arcwise connected and A is any non-vacuous space. These properties of the join together with its associativity imply

THEOREM. The subvariety V_a of \mathbb{C}^{n+1} is $(n-1)$ -connected. Moreover

$$(1) \quad \tilde{H}_n(V_a) \cong \tilde{H}_0(G_{a_0}) \otimes \tilde{H}_0(G_{a_1}) \otimes \dots \otimes \tilde{H}_0(G_{a_n}).$$

This is a free abelian group of rank $r = \prod (a_j - 1)$.

The isomorphism (1) is compatible with the operations of G_a .

All other reduced integral homology groups of V_a vanish.

It can be shown that V_a has the homotopy type of a connected union $S^n \vee \dots \vee S^n$ of r spheres of dimension n .

The identification of U_a with a join was explained to the speaker by MILNOR.

$U_a = G_{a_0} * G_{a_1} * \dots * G_{a_n}$ is an n -dimensional simplicial complex which has an n -simplex for each element of G_a . The n -simplex belonging to the unit of G_a is denoted by e . All other n -simplices are obtained from e by operations of G_a . Thus we have for the n -dimensional simplicial chain group

$$(2) \quad C_n(U_a) = J_a e$$

where J_a is the group ring of G_a . The homology group $\tilde{H}_n(U_a) = \tilde{H}_n(V_a)$ is an additive subgroup of $J_a e = C_n(U_a) \cong J_a$.

The face operator ∂_j commutes with all operations of G_a on $C_n(U_a)$ and furthermore satisfies $\partial_j = w_j \partial_j$. Therefore

$$(3) \quad h = (1-w_0)(1-w_1)\dots(1-w_n) e$$

is a cycle. Thus $h \in \tilde{H}_n(U_a)$. It follows easily that $\tilde{H}_n(V_a) = J_a h$. This yields the

THEOREM. The map $w \rightarrow wh \ (w \in G_a)$ induces an isomorphism

$$J_a/I_a \cong \tilde{H}_n(V_a) = J_a \cdot h$$

where $I_a \subset J_a$ is the annihilator ideal of h which is generated by the elements

$$1 + w_j + w_j^2 + \dots + w_j^{a_j-1}, \quad (j = 0, \dots, n).$$

Therefore $w_0^{k_0} w_1^{k_1} \dots w_n^{k_n} h$ (where $0 \leq k_j \leq a_j-2, j = 0, \dots, n$) is a basis of $\tilde{H}_n(V_a)$.

We recall that $\tilde{H}_n(V_a)$ is the integral singular homology group (of course with compact support). V_a is a $2n$ -dimensional oriented manifold without boundary (non-compact for $n \geq 1$). Therefore the bilinear intersection form S is well defined over $\tilde{H}_n(V_a)$. It is symmetric for n even, skew-symmetric for n odd. It is compatible with the operations of G_a .

PHAM ([14], p.358) constructs an n -dimensional cycle \tilde{h} in V_a which is homologous to h and intersects U_a exactly in two interior points of the simplices e and $w_0 w_1 \dots w_n e$ (sign questions have to be observed). In this way he obtains (using the G_a -invariance of S) the following result, reformulated somewhat for our purposes.

THEOREM. Put $\eta = (1-w_0) \dots (1-w_n)$. The bilinear form S over J_a $\eta \cong \tilde{H}_n(V_a)$ is given by

$$S(x\eta, y\eta) = f(\bar{y} \cdot x\eta), \quad (x, y \in J_a),$$

where $f : J_a \rightarrow \mathbb{Z}$ is the additive homomorphism with

$$f(1) = -f(w_0 \dots w_n) = (-1)^{\frac{n(n-1)}{2}}$$

$$f(w) = 0 \text{ for } w \in G_a, \quad w \neq 1, \quad w \neq w_0 \dots w_n,$$

and where $y \mapsto \bar{y}$ is the ring automorphism of the group ring J_a induced by $w \mapsto w^{-1} \ (w \in G_a)$.

§ 2. The quadratic form of V_a .

Let G be a finite abelian group, $J(G)$ its group ring. The ring automorphism of $J(G)$ induced by $g \mapsto g^{-1}$ ($g \in G$) is denoted by $x \mapsto \bar{x}$ ($x \in J(G)$). Give an element $\eta \in J(G)$ and a function $f : G \rightarrow \mathbb{Z}$. The additive homomorphism $J(G) \rightarrow \mathbb{Z}$ induced by f is also called f . Put $\hat{f} = \sum_{w \in G} f(w)w$. We assume

$$a) f(\bar{x}\eta) = f(x\eta) \text{ for all } x \in J(G), [\text{equivalently } \hat{f}\bar{\eta} = \bar{\hat{f}}\eta]$$

or

$$b) f(\bar{x}\eta) = -f(x\eta) \text{ for all } x \in J(G), [\text{equivalently } \hat{f}\bar{\eta} = -\bar{\hat{f}}\eta].$$

The bilinear form S over the lattice $J(G)\eta$ defined by

$$S(x\eta, y\eta) = f(\bar{y}x\eta), (x, y \in J(G)),$$

is symmetric in case a), skew symmetric in case b). Since S is a form with integral coefficients, its determinant is well-defined. The signature

$$\tau(S) = \tau^+(S) - \tau^-(S), \text{ case a),}$$

is the number $\tau^+(S)$ of positive minus the number $\tau^-(S)$ of negative diagonal entries in a diagonalisation of S over \mathbb{R} . Let χ run through the characters of G .

LEMMA. With the preceeding assumptions

$$\pm \det S = \prod_{\chi(\eta) \neq 0} \chi(\hat{f}). \text{ order of the torsion subgroup of } J(G)/J(G)\eta$$

and in case a)

$$\tau^+(S) = \text{number of characters } \chi \text{ with } \chi(\hat{f}\bar{\eta}) > 0$$

$$\tau^-(S) = \text{number of characters } \chi \text{ with } \chi(\hat{f}\bar{\eta}) < 0.$$

The proof is an exercise as in [1], p. 444.

The lemma and the last theorem of § 1 imply for the affine hypersurface

$$V_a = V(a_0, \dots, a_n) \text{ the}$$

THEOREM. Let S be the intersection form of V_a . Then

$$(1) \quad \pm \det S = \prod_{1 \leq k_j \leq a_j - 1} (1 - \varepsilon_0^{k_0} \varepsilon_1^{k_1} \dots \varepsilon_n^{k_n})$$

where $\varepsilon_j = \exp(2\pi i/a_j)$. For n even, we have

$$\tau^+(S) = \text{number of } (n+1)\text{-tuples of integers } (x_0, \dots, x_n), 0 < x_j < a_j,$$

$$\text{with } 0 < \sum_{j=0}^n \frac{x_j}{a_j} < 1 \pmod{2\mathbb{Z}}$$

(2)

$$\tau^-(S) = \text{number of } (n+1)\text{-tuples of integers } (x_0, \dots, x_n), 0 < x_j < a_j,$$

$$\text{with } -1 < \sum_{j=0}^n \frac{x_j}{a_j} < 0 \pmod{2\mathbb{Z}}.$$

See [5] for details.

REMARK. The intersection form S of $V(a_0, \dots, a_n)$ with $n \equiv 0 \pmod{2}$ is
even, i.e. $S(x, x) \equiv 0 \pmod{2}$ for $x \in \tilde{H}_n(V_n)$. Therefore, by a well-known
theorem, $\det S = \pm 1$ implies $\tau^+(S) - \tau^-(S) = \tau(S) \equiv 0 \pmod{8}$.

Following MILNOR we introduce for $a = (a_0, \dots, a_n)$ the graph $\Gamma(a)$:
 $\Gamma(a)$ has the $(n+1)$ vertices a_0, \dots, a_n . Two of them (say a_i, a_j) are
joined by an edge if and only if the greatest common divisor (a_i, a_j) is
greater than 1. Then we have [5]

LEMMA. det S as given in the preceeding theorem equals ± 1 if and only if
 $\Gamma(a)$ satisfies

- a) $\Gamma(a)$ has at least two isolated points, or,
- b) it has one isolated point and at least one connectedness component K
with an odd number of vertices such that $(a_i, a_j) = 2$ for
 $a_i, a_j \in K$ $(i \neq j)$.

Now suppose n even and $a = (a_0, \dots, a_n) = (p, q, 2, \dots, 2)$ with p, q odd
and $(p, q) = 1$. Then $\det S = \pm 1$ and

$$(3) \quad (-1)^{n/2} \cdot \tau(S) = \frac{(p-1)(q-1)}{2} + 2(N_{p,q} + N_{q,p}),$$

where $N_{p,q}$ is the number of $q \cdot x$ ($1 \leq x \leq \frac{p-1}{2}$) whose remainder mod p of smallest absolute value is negative. This follows from the preceding theorem. Observe that by the above remark $\tau(S)$ is divisible by 4 (even by 8) and that this is related to one of the proofs of the quadratic reciprocity law ([1], p. 450).

In particular, for n even and $(a_0, \dots, a_n) = (3, 6k-1, 2, \dots, 2)$ the signature $\tau(S)$ equals $(-1)^{n/2} \cdot 8k$.

§ 3. Exotic spheres.

A k -dimensional compact oriented differentiable manifold is called a k -sphere if it is homeomorphic to the k -dimensional standard sphere. A k -sphere not diffeomorphic to the standard k -sphere is said to be exotic. The first exotic sphere was discovered by MILNOR in 1956. Two k -spheres are called equivalent if there exists an orientation preserving diffeomorphism between them. The equivalence classes of k -spheres constitute for $k \geq 5$ a finite abelian group Θ_k under the connected sum operation. Θ_k contains the subgroup bP_{k+1} of those k -spheres which bound a parallelizable manifold. bP_{4m} ($m \geq 2$) is cyclic of order

$$2^{2m-2} (2^{2m-1} - 1) \text{ numerator } \left(\frac{4B_m}{m} \right),$$

where B_m is the m -th Bernoulli number. Let ξ_m be a generator of bP_{4m} . If a $(4m-1)$ -sphere Σ bounds a parallelizable manifold B of dimension $4m$, then the signature $\tau(B)$ of the intersection form of B is divisible by 8 and

$$(1) \quad \Sigma = + \frac{\tau(B)}{8} \xi_m$$

(g_m should be chosen in such a way that we have always the plus-sign in (1)).

For $m = 2$ and 4 we have

$$bP_8 = \Theta_7 = \mathbb{Z}_{28}, \quad bP_{12} = \Theta_{11} = \mathbb{Z}_{992}.$$

All these results are due to MILNOR-KERVAIRE. The group bP_{2n} (n odd, $n \geq 3$) is either 0 or \mathbb{Z}_2 . It contains only the standard sphere and the KERVAIRE sphere (obtained by plumbing two copies of the tangent bundle of S^n). It is known that bP_{2n} is \mathbb{Z}_2 (equivalently that the KERVAIRE sphere is exotic) if $n \equiv 1 \pmod{4}$ and $n \geq 5$ (E. BROWN-F. PETERSON).

Let $V_a^0 = V^0(a_0, a_1, \dots, a_n) \subset \mathbb{C}^{n+1}$ (where $a_j \geq 2$) be defined by

$$\frac{a_0}{z_0^2} + \frac{a_1}{z_1^2} + \dots + \frac{a_n}{z_n^2} = 0.$$

This affine variety has exactly one singular point, namely the origin of \mathbb{C}^{n+1} .

Let

$$S^{2n+1} = \{z \in \mathbb{C}^{n+1} \mid \sum_{j=0}^n z_j \bar{z}_j = 1\}.$$

Then $\Sigma_a = \Sigma(a_0, \dots, a_n) = V_a^0 \cap S^{2n+1}$ is a compact oriented differentiable manifold (without boundary) of dimension $2n-1$.

THEOREM. Let $n \geq 3$. Then Σ_a is $(n-2)$ -connected. It is a $(2n-1)$ -sphere if and only if the graph $\Gamma(a)$ defined in § 2 satisfies the condition a) or b). If Σ_a is a $(2n-1)$ -sphere, then it belongs to bP_{2n} . If, moreover, $n = 2m$, then

$$\Sigma_a = \frac{\tau}{8} g_m,$$

where $\tau = \tau^+ - \tau^-$ and τ^+, τ^- are as in § 2 (2). In particular

$$\sum_{i=0}^{2m} z_i \bar{z}_i = 1$$

$$z_0^3 + z_1^{6k-1} + z_2^2 + \dots + z_{2m}^2 = 0$$

is a $(4m-1)$ -sphere embedded in $S^{4m+1} \subset \mathbb{C}^{2m+1}$ which represents the element $(-1)^m k \cdot g_m \in bP_{4m}$. Example : For $m = 2$ and $k = 1, \dots, 28$ we get the 28 classes of 7-spheres, for $m = 3$ and $k = 1, \dots, 992$ the 992 classes of 11-spheres.

COROLLARY. The affine variety $V^0(a_0, \dots, a_n)$, $n \geq 3$, is a topological manifold if and only if the graph $\Gamma(a)$ satisfies a) or b) of § 2.

For this theorem and for the case n odd see BRIESKORN [5].

Proof. If we remove from V_a^0 the points with $z_n = 0$ we get a space \tilde{V}_a whose fundamental group has $\pi_1(V_a - \{0\}) \cong \pi_1(\Sigma_a)$ as homomorphic image. \tilde{V}_a is fibred over \mathbb{C}^* with $V(a_0, \dots, a_{n-1})$ as fibre which is simply-connected. Thus $\pi_1(\tilde{V}_a) \cong \mathbb{Z}$ and $\pi_1(\Sigma_a)$ is commutative. Because of this and by SMALE-POINCARÉ we have to study only the homology of Σ_a .

Let $V_a^\varepsilon \subset \mathbb{C}^{n+1}$ be the affine variety

$$z_0^{a_0} + z_1^{a_1} + \dots + z_n^{a_n} = \varepsilon$$

($V_a = V_a^1$). Let D^{2n+2} be the full ball in \mathbb{C}^{n+1} with center 0 and radius 1 and S^{2n+1} , as before, its boundary. Σ_a is diffeomorphic to $\Sigma_a^\varepsilon = S^{2n+1} \cap V_a^\varepsilon$ for $\varepsilon > 0$ and small. It is the boundary of $B_a^\varepsilon = D^{2n+2} \cap V_a^\varepsilon$ whose interior (for ε small) is diffeomorphic to V_a^ε and V_a . The exact homology sequence of the pair $(B_a^\varepsilon, V_a^\varepsilon)$ shows that Σ_a is $(n-2)$ -connected. Using POINCARÉ duality we get the exact sequence

$$0 \rightarrow H_n(\Sigma_a) \rightarrow H_n(V_a) \xrightarrow{\sigma} \text{Hom}(H_n(V_a), \mathbb{Z}) \rightarrow H_{n-1}(\Sigma_a) \rightarrow 0$$

where the homomorphism σ is given by the bilinear intersection form S of V_a (see § 2). This determines $H^*(\Sigma_a)$ completely : $H_n(\Sigma_a) = 0$ if and only

if $\det S \neq 0$. If $\det S \neq 0$, then $|\det S|$ equals the order of $H_{n-1}(\Sigma_a)$.

The manifold B_a^ε is parallelizable since its normal bundle is trivial.

This finishes the proof in view of § 2.

§ 4. Manifolds with actions of the orthogonal group.

$O(n)$ denotes the real orthogonal group with $O(m) \subset O(n)$, $m < n$, by

$$A \mapsto \begin{pmatrix} 1 & 0 \\ 0 & A \end{pmatrix}, \quad (A \in O(m), 1 = \text{unit of } O(n-m)).$$

Let X be a compact differentiable manifold of dimension $2n-1$ on which $O(n)$ acts differentiably ($n \geq 2$). Suppose each isotropy group is conjugate to $O(n-2)$ or $O(n-1)$. Then the orbits are either Stiefel manifolds $O(n)/O(n-2)$ (of dimension $2n-3$) or spheres $O(n)/O(n-1)$ (of dimension $n-1$). Suppose that the 2-dimensional representation of an isotropy group of type $O(n-2)$ normal to the orbit is trivial whereas the n -dimensional representation of an isotropy group of type $O(n-1)$ normal to the orbit is the 1-dimensional trivial representation plus the standard representation of $O(n-1)$. Under these assumptions the orbit space is a compact 2-dimensional manifold X' with boundary, the interior points of X' corresponding to orbits of type $O(n)/O(n-2)$, the boundary points of X' to the orbits of type $O(n)/O(n-1)$. Suppose finally that X' is the 2-dimensional disk.

It follows from the classification theorems of [8] and [9] that the classes of manifolds X with the above properties under equivariant diffeomorphisms are in one-to-one correspondence with the non-negative integers. We let $W^{2n-1}(d)$ be the $(2n-1)$ -dimensional $O(n)$ -manifold corresponding to the integer $d \geq 0$. The fixed point set of $O(n-2)$ in $W^{2n-1}(d)$ is a 3-dimensional $O(2)$ -manifold, namely $W^3(d)$, which by ([9], § 5, Korollar 6) is the lens

space $L(d,1)$. Thus in order to determine the d associated to a given $O(n)$ -manifold of our type we just have to look at the integral homology group H_1 of the fixed point set of $O(n-2)$. $W^{2n-1}(0)$ is $S^n \times S^{n-1}$, the manifold $W^{2n-1}(1)$ is S^{2n-1} , the actions of $O(n)$ are easily constructed. Consider for $d \geq 2$ the manifold $\Sigma(d,2,\dots,2)$ in \mathbb{C}^{n+1} given by

$$(1) \quad \begin{aligned} z_0^d + z_1^2 + \dots + z_n^2 &= 0 \\ \sum_{i=0}^n z_i \bar{z}_i &= 1 \end{aligned}$$

(see § 3). It is easy to check that this is an $O(n)$ -manifold satisfying all our assumptions. The operation of $A \in O(n)$ on (z_0, z_1, \dots, z_n) is, of course, given by applying the real orthogonal matrix $A \in O(n)$ on the complex vector (z_1, \dots, z_n) leaving z_0 untouched. The fixed point set of $O(n-2)$ is $\Sigma(d,2,2)$ which is $L(d,1)$, see [6].

THEOREM. The $O(n)$ -manifold $\Sigma(d,2,\dots,2)$ given by (1) is equivariantly diffeomorphic with $W^{2n-1}(d)$, $n \geq 2$. It can also be obtained by equivariant plumbing of $d-1$ copies of the tangent bundle of S^n along the graph A_{d-1}

$$\bullet \text{---} \bullet \text{---} \dots \text{---} \bullet \quad d-1 \text{ vertices} .$$

For the proof it suffices to establish the $O(n)$ -action on the manifold obtained by plumbing and check all properties :

$O(n)$ acts on S^n and on the unit tangent bundle of S^n . Since the action of $O(n)$ on S^n has two fixed points the plumbing can be done equivariantly. The fixed point set of $O(n-2)$ is the manifold obtained by plumbing $d-1$ tangent bundles of S^2 which is well-known to be $L(d,1)$, (see [6], resolution of the singularity of $z_0^d + z_1^2 + z_2^2 = 0$).

The above theorem gives another method to calculate the homology of $\Sigma(d, 2, \dots, 2)$ and to prove that $\Sigma(d, 2, \dots, 2)$ for d odd and an odd number of 2's is a sphere. In particular, $\Sigma(3, 2, 2, 2, 2, 2)$ is the exotic 9-dimensional Kervaire sphere (see § 3). The calculation of the ARF invariant of the A_{d-1} -plumbing shows more generally that

$$\Sigma(d, 2, \dots, 2), \quad (d \text{ odd, an odd number of } 2\text{'s})$$

is the standard sphere for $d \equiv +1 \pmod{8}$ and the Kervaire sphere for $d \equiv +3 \pmod{8}$, in agreement with a more general result in [5].

REMARKS. The $O(n)$ -manifold $W^{2n-1}(d)$ coincides with Bredon's manifolds M_k^{2n-1} for $d = 2k+1$, see Bredon [3]. $\Sigma(3, 2, 2, 2)$ is the standard 5-sphere (since $\theta_5 = 0$). Therefore S^5 admits a differentiable involution α with the lens space $L(3, 1)$ as fixed point set and a diffeomorphism β of period 3 with the real projective 3-space as fixed point set. Compare [3]. α and β are defined on $\Sigma(3, 2, 2, 2)$ given by (1) as follows

$$\alpha(z_0, z_1, z_2, z_3) = (z_0, z_1, z_2, -z_3)$$

$$\beta(z_0, z_1, z_2, z_3) = (\varepsilon z_0, z_1, z_2, z_3), \quad \text{where } \varepsilon = \exp(2\pi i/3).$$

Many more such examples of "exotic" involutions etc. which are not differentially equivalent to orthogonal involutions etc. can be constructed.

§ 5. Manifolds associated to knots.

Let X be a compact differentiable manifold of dimension $2n-1$ on which $O(n-1)$ acts differentiably ($n \geq 3$). Suppose each isotropy group is conjugate to $O(n-3)$ or $O(n-2)$ or is $O(n-1)$. Then the orbits are either Stiefel manifolds $O(n-1)/O(n-3)$ (of dimension $2n-5$) or spheres $O(n-1)/O(n-2)$ (of dimension $n-2$) or points (fixed points of the whole action). The

representations of the isotropy groups $O(n-3)$, $O(n-2)$ and $O(n-1)$ respectively normal to the orbit are supposed to be the 4-dimensional trivial representation, the 3-dimensional trivial plus the standard representation of $O(n-2)$, the 1-dimensional trivial plus the sum of two copies of the standard representation of $O(n-1)$. The orbit space X' is then a 4-dimensional manifold with boundary. We suppose that X' is the 4-dimensional disk D^4 .

Then the points of the interior of D^4 correspond to Stiefel-manifold-orbits, the points of $\partial D^4 = S^3$ to the other orbits. The set F of fixed points corresponds to a 1-dimensional submanifold of S^3 , also called F .

We suppose F non-empty and connected, it is then a knot in S^3 . We shall call an $O(n-1)$ -manifold of dimension $2n-1$ a "knot manifold" if all the above conditions are satisfied.

Let K be the set of isomorphism classes of differentiable knots (i.e. isomorphism classes of pairs (S^3, F) - F a compact connected 1-dimensional submanifold - under diffeomorphisms of S^3). For the following theorem see JÄNICH ([9], § 6), compare also W.C. HSIANG and W.Y. HSIANG [8].


THEOREM. For any $n \geq 3$ there is a one-to-one correspondence

$$\kappa_n : K \rightarrow \mathfrak{K}_{2n-1} ,$$

where \mathfrak{K}_{2n-1} is the set of isomorphism classes of $(2n-1)$ -dimensional knot manifolds under equivariant diffeomorphisms. κ_n^{-1} associates to a knot manifold the knot F considered above.

REMARK. The 2-fold branched covering of S^3 along a knot F is an $O(1)$ -manifold which will be denoted by $\kappa_2(F)$.

If we plumb 8 copies of the tangent bundles of S^n ($n \geq 1$) according to the tree E_8



we get a $(2n-1)$ -dimensional manifold $M^{2n-1}(E_8)$. For $n=2$ this is S^3/G , where G is the binary pentagondodecahedral group [6]. For n odd, $M^{2n-1}(E_8)$ is the standard sphere, as the ARF invariant shows. For $n = 2m \geq 4$, the manifold $M^{4m-1}(E_8)$ is an exotic sphere, it is the famous MILNOR sphere which represents the generator $\pm g_m$ of bP_{4m} (see § 3).

$M^{2n-1}(E_8)$ admits an action of $O(n-1)$ as follows: $O(n-1)$ operates as subgroup of $O(n+1)$ on S^n and thus on the unit tangent bundle of S^n . The action on S^n leaves a great circle fixed.

When plumbing the eight copies of the tangent bundle we put the center of the plumbing operation always on this great circle; (for one copy, corresponding to the central vertex of the E_8 -tree, we need three such centers, therefore, we cannot have an action of $O(n)$, which has only 2 fixed points on S^n .) Then the action of $O(n-1)$ on each copy of the tangent bundle is compatible with the plumbing and extends to an action of $O(n-1)$ on $M^{2n-1}(E_8)$ which, for $n \geq 3$, becomes a knot manifold as can be checked. The resulting knot can be seen on a picture attached at the end of this lecture. The speaker had convinced himself that this is the torus knot $t(3,5)$, but ZIESCHANG and VOGT showed him a better proof. This implies the

THEOREM. Suppose $n \geq 3$. Then $\kappa_n(t(3,5))$ is equivariantly diffeomorphic to $M^{2n-1}(E_8)$ with the $O(n-1)$ -action defined by equivariant plumbing. (This is still true for $n=2$, see Remark above).

We now consider the manifold $\Sigma(p, q, 2, 2, \dots, 2) \subset \mathbb{C}^{n+1}$ given by the equations (see § 3)

$$z_0^p + z_1^q + z_2^2 + \dots + z_n^2 = 0$$

$$\sum_{i=0}^n z_i \bar{z}_i = 1 \quad (n \geq 3).$$

This is an $O(n-1)$ -manifold, the action being defined similarly as in § 4. Suppose $(p, q) = 1$. Then it can be shown that $\Sigma(p, q, 2, 2, \dots, 2)$ is a knot manifold : It is $\kappa_n(t(p, q))$ where $t(p, q)$ is the torus knot. Therefore, by the preceding theorem we have an equivariant diffeomorphism

$$M^{2n-1}(E_8) \cong \Sigma(3, 5, \underbrace{2, \dots, 2}_{n-1}).$$

This gives a different proof (based on the classification of knot manifolds) that $\Sigma(3, 5, \underbrace{2, \dots, 2}_{2m-1})$ represents for $m \geq 2$ a generator of bP_{4m} .

(compare § 3).

§ 6. A theorem on knot manifolds.

Let F be a knot in S^3 . Then the signature $\tau(F)$ can be defined in the following way which MILNOR explained to the speaker in a letter. MILNOR also considers higher dimensional cases. We cite from his letter, but restrict to classical knots :

Let X be the complement of an open tubular neighbourhood of F in S^3 .

Then the cohomology

$$H^* = H^*(\hat{X}, \partial\hat{X}; \mathbb{R})$$

where \hat{X} is the infinite cyclic covering of X , satisfies Poincaré duality just as if \hat{X} were a 2-dimensional manifold bounded by F .

In particular the pairing

$$U : H^1 \otimes H^1 \rightarrow H^2 \simeq \mathbb{R}$$

is non-degenerate. Let t denote a generator for the group of covering transformations of \hat{X} . Then for $a, b \in H^1$ the pairing

$$\langle a, b \rangle = a \cup t^*b + b \cup t^*a$$

is symmetric and non-degenerate. Hence, the signature

$$\tau^+(F) - \tau^-(F) = \tau(F) \text{ is defined.}$$

There exist earlier definitions of the signature by MURASUGI [13] and TROTTER [17]. The signature is a cobordism invariant of the knot. A cobordism invariant mod 2 was introduced by ROBERTELLO [15] inspired by an earlier paper of KERVAIRE-MILNOR. Let F be a knot and Δ its Alexander polynomial, then the ROBERTELLO invariant $c(F)$ is an integer mod 2, namely

$$c(F) = 0, \text{ if } \Delta(-1) \equiv \pm 1 \pmod{8}$$

$$c(F) = 1, \text{ if } \Delta(-1) \equiv \pm 3 \pmod{8}$$

We recall that the first integral homology group of $\kappa_2(F)$, the 2-fold branched covering of the knot F (see a remark in § 5), is always finite, its order is odd, and equals up to sign the determinant of F . We have $\pm \det F = \Delta(-1)$.

THEOREM. Let F be a knot, then $\kappa_n(F)$, $n \geq 2$, is the boundary of a parallelizable manifold. For n odd, $\kappa_n(F)$ is homeomorphic to S^{2n-1} and thus represents an element of bP_{2n} , it is the standard sphere if

$c(F) = 0$, the Kervaire sphere if $c(F) = 1$. If $n = 2m$, then $\kappa_{2m}(F)$ is
 $(2m-2)$ -connected and $H_{2m-1}(\kappa_{2m}(F), \mathbb{Z}) \simeq H_1(\kappa_2(F), \mathbb{Z})$. For $m \geq 2$ it is
homeomorphic to S^{4m-1} if and only if $\det F = +1$. Then $\kappa_{2m}(F)$ represents
(up to sign) an element of bP_{4m} which is $\pm \frac{\tau(F)}{8} \cdot g_m$ (see § 3).

The proof uses an equivariant handlebody construction starting out from a Seifert surface [16] spanned in the knot F . For simplicity, not out of necessity, we have disregarded orientation questions in § 5 and § 6.

REMARK. § 2(3) gives up to sign a formula for the signature of the torus knot $t(p,q)$, (p,q) odd with $(p,q) = 1$.

BIBLIOGRAPHY
=====

- [1] S. BOREWICZ und I. ŠAFAREVIČ.- Zahlentheorie, Birkhäuser Verlag, Basel und Stuttgart 1966.
- [2] G. BREDON - Transformation groups on spheres with two type of orbits, Topology 3 (1965), p. 103-113.
- [3] G. BREDON - Examples of differentiable group actions, Topology 3 (1965), p. 115-122.
- [4] E. BRIESKORN - Examples of singular normal complex spaces which are topological manifolds, Proc. Nat. Acad. Sci. USA 55 (1966), p. 1395-1397.
- [5] E. BRIESKORN - Beispiele zur Differentialtopologie von Singularitäten, Inventiones mathematicae, 2 (1966), p. 1-14.
- [6] F. HIRZEBRUCH - The topology of normal singularities of an algebraic surface (d'après un article de D. Mumford), Séminaire Bourbaki 15ème année, 1962/63, n° 250.
- [7] F. HIRZEBRUCH - $O(n)$ -Mannigfaltigkeiten... in Vorbereitung.
- [8] W. HSIANG und W.-Y. HSIANG - Some results on differentiable actions, Bull. Amer. Math. Soc. 72 (1966), p. 134-138.
- [9] K. JÄNICH - Differenzierbare Mannigfaltigkeiten mit Rand als Orbiträume differenzierbarer G -Mannigfaltigkeiten ohne Rand, to appear in Topology.
- [10] J. MILNOR - Construction of universal bundles II, Annals of Math. 63 (1956), p. 430-436.
- [11] J. MILNOR - On isolated singularities of hypersurfaces, mimeographed Princeton 1966.
- [12] D. MUMFORD - The topology of normal singularities of an algebraic surface and a criterion for simplicity, Paris, Presses universitaires de France, 1961 (Institut des Hautes Études Scientifiques, Publications mathématiques, 9).

- [13] K. MURASUGI - On a certain numerical invariant of link types, Trans. Americ. Math. Soc. 117 (1965), p. 387-422.
- [14] F. PHAM - Formules de Picard-Lefschetz généralisées et ramification des intégrales, Bull. Soc. math. France 93 (1965), p. 333-367.
- [15] R. ROBERTELLO - An invariant of knot cobordism, Comm. Pure Appl. Math. 18 (1965), p. 543-555.
- [16] H. SEIFERT - Über das Geschlecht von Knoten, Math. Annalen 110 (1934), p. 571-592.
- [17] H.F. TROTTER - Homology of group systems with applications to knot theory, Annals of Math. 76 (1962), p. 464-498.

ERRATUM

Page 314-07. Ligne 4 du bas, au lieu de "Let g_m be a generator of bp_{4m} ." lire: "Let g_m be the Milnor generator of bp_{4m} , see p. 314-14."

