

SÉMINAIRE N. BOURBAKI

MICHAEL SCHNEIDER

Holomorphic vector bundles on $\mathbb{P}_n$

Séminaire N. Bourbaki, 1980, exp. n° 530, p. 80-102

<http://www.numdam.org/item?id=SB_1978-1979__21__80_0>

© Association des collaborateurs de Nicolas Bourbaki, 1980, tous droits réservés.

L'accès aux archives du séminaire Bourbaki (<http://www.bourbaki.ens.fr/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

HOLOMORPHIC VECTOR BUNDLES ON $\mathbb{P}_n$

by Michael SCHNEIDER

The classification of holomorphic (= algebraic) vector bundles on complex projective space $\mathbb{P}_n$ could be tried along the following lines :

- I) Classify the topological complex vector bundles on $\mathbb{P}_n$.
- II) Determine which topological bundles admit an analytic structure.
- III) Classify for fixed topological bundle all possible analytic structures.

This is a survey of some of the main results concerning I) - III) as well as a guide to the literature. We included only a few open problems. But in fact most of the work has still to be done.

Notation. - No distinction will be made between holomorphic vector bundles and locally free coherent analytic sheaves. $\mathcal{O}(1)$ is the line bundle having a holomorphic section vanishing precisely on a hyperplane. $E(k) := E \otimes \mathcal{O}(1)^{\otimes k}$,

$h^i(\mathbb{P}_n, E) := \dim_{\mathbb{C}} H^i(\mathbb{P}_n, E)$ for a vector bundle E on $\mathbb{P}_n$. The total Chern class of E will be denoted by $c(E) = 1 + c_1(E) + \dots + c_r(E)$. The Chern classes $c_i(E) \in H^{2i}(\mathbb{P}_n, \mathbb{Z}) \simeq \mathbb{Z}$ will be regarded mostly as integers. The holomorphic tangent bundle of $\mathbb{P}_n$ will be denoted by $T_{\mathbb{P}_n}$.

1. Topological classification

Let $\text{Vect}_{\text{top}}^r(\mathbb{P}_n)$ be the isomorphism classes of topological complex vector bundles of rank r on $\mathbb{P}_n$. It is well known that $\text{Vect}_{\text{top}}^r(\mathbb{P}_n) \simeq \text{Vect}_{\text{top}}^n(\mathbb{P}_n)$ for all $r \geq n$.

Schwarzenberger [53] noticed that the Chern classes of $E \in \text{Vect}_{\text{top}}^r(\mathbb{P}_n)$ satisfy the condition

$$(S_n) \quad \sum_{i=1}^r \binom{\delta_i}{k} \in \mathbb{Z} \quad \text{for } 2 \leq k \leq n.$$

Here the δ_i are as usual related to the Chern class of E by

$$c(E) = \prod_{i=1}^r (1 + \delta_i).$$

The conditions (S_n) for $r = 2$ are as follows :

(S₂) no condition

(S₃) $c_1 c_2 \equiv 0 \pmod{2}$

(S₄) $c_2(c_2 + 1 - 3c_1 - 2c_1^2) \equiv 0 \pmod{12}$

(S₅) is equivalent to (S₄).

For $r = 3$ one gets for instance (S₃): $c_3 \equiv c_1 c_2 \pmod{2}$.

A. Thomas [60] proved that the Schwarzenberger condition (S_n) classifies stable bundles on $\mathbb{P}_n$ i.e.

$$\text{Vect}_{\text{top}}^n(\mathbb{P}_n) \simeq \{(c_1, \dots, c_n) \in \mathbb{Z}^n : (c_1, \dots, c_n) \text{ satisfy } (S_n)\}.$$

For $\mathbb{P}_2$ this gives

$$\text{Vect}_{\text{top}}^r(\mathbb{P}_2) \simeq \mathbb{Z} \times \mathbb{Z} \quad \text{for } r \geq 2.$$

For $\mathbb{P}_3$ there remains the classification of 2-bundles. This has been done by Atiyah and Rees [2]. They showed that for c_1, c_2 with $c_1 c_2 \equiv 0 \pmod{2}$ and c_1 odd there exists exactly one 2-bundle with these c_i as Chern classes. For c_1 even there are exactly two 2-bundles with these c_i as Chern classes. These two bundles are distinguished by a certain mod 2 invariant α .

On $\mathbb{P}_4$ there remains the classification of bundles of rank 2 and 3. Switzer [55], complementing the results of Atiyah and Rees, showed

$$\text{Vect}_{\text{top}}^2(\mathbb{P}_4) \simeq \{(c_1, c_2) \in \mathbb{Z} \times \mathbb{Z} : (S_4) \text{ is true}\}.$$

Switzer [55] recently pushed the classification of 2-bundles up to $\mathbb{P}_6$. As a sample let us state his results on $\mathbb{P}_5$ because this is the first case where not all c_1, c_2 satisfying the Schwarzenberger conditions arise as the Chern classes of a vector bundle of rank 2. Set $\Delta = \frac{c_1^2 - 4c_2}{4}$. Then for c_1, c_2 satisfying

(S₅) there exists at least one 2-bundle with these c_i as Chern classes if c_1 is odd or if c_1 is even and $\Delta^2(\Delta - 1) \equiv 0 \pmod{24}$ (if c_1 is even and $\Delta^2(\Delta - 1) \not\equiv 0 \pmod{24}$ there is no 2-bundle with these c_i as Chern classes). For $c_2 \not\equiv c_1^2 \pmod{3}$ there exists exactly one 2-bundle and for $c_2 \equiv c_1^2 \pmod{3}$ there are exactly three 2-bundles.

2. Construction of holomorphic vector bundles on $\mathbb{P}_n$

In this section we will give some general procedures to construct holomorphic bundles. These will be applied to show that all topological vector bundles on $\mathbb{P}_n$, $n \leq 3$, admit an analytic structure.

Let us start by recalling that all line bundles on $\mathbb{P}_n$ are of the form $\mathcal{O}(k)$, $k \in \mathbb{Z}$. To convince the reader that the difficulties arise only if rank and dimension are bigger than 1 we include a short proof of the fact that all holomorphic vector bundles on $\mathbb{P}_1$ split into line bundles (see [19]).

THEOREM (Grothendieck [21]).- Any holomorphic vector bundle E on $\mathbb{P}_1$ is of the form $E = \mathcal{O}(a_1) \oplus \dots \oplus \mathcal{O}(a_r)$.

Proof. The proof is by induction on $r = \text{rk } E$. We may assume $r \geq 2$. Choose $k \in \mathbb{Z}$ minimal with $H^0(E(k)) \neq 0$ (k exists by Serre's results on the cohomology of coherent sheaves on $\mathbb{P}_n$). We may assume $k = 0$. Any nonzero $\sigma \in H^0(E)$ has zeroes only in codimension 2. Hence a nonzero $\sigma \in H^0(E)$ gives a trivial line subbundle of E

$$(*) \quad 0 \rightarrow \mathcal{O} \xrightarrow{\sigma} E \rightarrow F \rightarrow 0.$$

By induction we have $F \simeq \mathcal{O}(a_2) \oplus \dots \oplus \mathcal{O}(a_r)$. From $(*)$ one gets the exact sequence

$$\rightarrow H^0(E(-1)) \rightarrow H^0(F(-1)) \rightarrow H^1(\mathcal{O}(-1)) = 0.$$

This shows $H^0(F(-1)) = 0$ and therefore $a_i \leq 0$ for all i . The obstruction to split $(*)$ lies in $H^1(F^*) = \bigoplus_i H^1(\mathcal{O}(-a_i)) = 0$, since $a_i \leq 0$ for all i .

Hence $(*)$ splits and we get

$$E \simeq \mathcal{O} \oplus \mathcal{O}(a_2) \oplus \dots \oplus \mathcal{O}(a_r).$$

2.1. Vector bundles of rank $n-1$ on $\mathbb{P}_n$

Tango [58] constructed indecomposable holomorphic $(n-1)$ -bundles on $\mathbb{P}_n$ for each $n \geq 3$ using the following generalization of a general position argument of Serre's.

PROPOSITION 2.1.- Let E be a holomorphic vector bundle on $\mathbb{P}_n$ generated by global sections. If $c_i(E) = 0$ for some $i \leq r = \text{rk } E$ then E has a trivial subbundle of rank $r - i + 1$.

COROLLARY 1.- For $n \geq 3$ there is an indecomposable $(n-1)$ -bundle on $\mathbb{P}_n$.

Proof. $\Omega^1(2)$ is generated by global sections. Let

$$\varphi : H^0(\mathbb{P}_n, \Omega^1(2)) \times \mathbb{P}_n \rightarrow \Omega^1(2)$$

be the canonical surjection and put $E = (\ker \varphi)^*$. One calculates $c_n(E) = 0$. Hence E has a trivial subbundle such that the quotient F is of rank $n-1$. The indecompos-

bility of F can be proved by inspecting its cohomology groups.

COROLLARY 2.- For n odd there is a $(n-1)$ -bundle N on $\mathbb{P}_n$ with Chern class

$$c(N) = 1 + h^2 + h^4 + \dots + h^{n-1}.$$

Here $h = c_1(\mathcal{O}(1))$ is the canonical generator of $H^2(\mathbb{P}_n, \mathbb{Z})$.

Proof. $\Omega^1(2)$ is generated by global sections and $c_n(\Omega^1(2)) = 0$ for n odd. This shows the existence of a trivial line subbundle of $\Omega^1(2)$. This gives a surjection

$$T(-1) \longrightarrow \mathcal{O}(1).$$

Let N be the kernel of this map. Then

$$\begin{aligned} c(N) &= c(T(-1))(1+h)^{-1} \\ &= (1-h)^{-1}(1+h)^{-1} \\ &= 1 + h^2 + h^4 + \dots + h^{n-1}. \end{aligned}$$

Remarks.- 1) N is the Null-correlation bundle.

2) The tangent bundle $T_{\mathbb{P}_n}$ is indecomposable.

3) Maruyama [38] has shown that for each $r > n$ there exist indecomposable r -bundles on $\mathbb{P}_n$ if $n \geq 2$.

2.2. Subvarieties of $\mathbb{P}_n$ of codimension 2 and holomorphic vector bundles of rank 2

In this section we will explain the connection of locally complete intersection subvarieties of codimension 2 and holomorphic bundles of rank 2. This correspondence essentially goes back to Serre [49] and has been rediscovered and reformulated many times [28], [9], [18], [23], [25]. Here we follow mainly Hartshorne's presentation.

Let E be a holomorphic 2-bundle on $\mathbb{P}_n$ and suppose E has a holomorphic section σ vanishing in codimension 2 only (this can always be achieved by replacing E by $E(k)$ with $k \in \mathbb{Z}$ minimal with respect to $H^0(E(k)) \neq 0$). Then $Y = \{\sigma = 0\}$ is of codimension 2 and locally a complete intersection. Y is in general neither reduced nor irreducible. The Koszul complex of σ is

$$0 \longrightarrow \det E^* \longrightarrow E^* \longrightarrow J_Y \longrightarrow 0.$$

This implies

$$E^*|_Y \simeq J/J^2.$$

Hence E is an extension of the normal bundle $N_{Y|\mathbb{P}_n} = (J/J^2)^*$ of Y in $\mathbb{P}_n$ to the whole of $\mathbb{P}_n$. Inserting

$$E^* \simeq E \otimes \det E^*$$

into the Koszul complex gives.

$$0 \longrightarrow 0 \xrightarrow{\sigma} E \longrightarrow J_Y \otimes \det E \longrightarrow 0.$$

It is clear that

$$c_2(E) = \text{dual of } Y.$$

Hence $c_2(E) = \deg Y$.

The interesting point is the reversal of this procedure. Take a locally complete intersection $Y \subset \mathbb{P}_n$ of codimension 2. We would like to construct a 2-bundle E together with a $\sigma \in H^0(\mathbb{P}_n, E)$ giving $Y = \{\sigma = 0\}$. By what we have seen it is natural to try getting E^* as extension of J_Y by some line bundle.

PROPOSITION 2.2.1.- Let Y be a locally complete intersection of codimension 2 in $\mathbb{P}_n$, $n \geq 3$. Assume that $\det N_Y|_{\mathbb{P}_n} \simeq \mathcal{O}_Y(k)$. Then there exists a holomorphic 2-bundle E on $\mathbb{P}_n$ with a holomorphic section $\sigma \in H^0(\mathbb{P}_n, E)$ such that

$$Y = \{\sigma = 0\}.$$

In particular $c_1(E) = k$, $c_2(E) = \deg Y$.

Proof. The extensions of J_Y by $\mathcal{O}(-k)$ are classified by $\text{Ext}_{\mathcal{O}}^1(J_Y, \mathcal{O}(-k))$. The exact sequence

$$\begin{aligned} 0 \rightarrow H^1(\mathbb{P}_n, \underline{\text{Hom}}(J_Y, \mathcal{O}(-k))) &\rightarrow \text{Ext}_{\mathcal{O}}^1(J_Y, \mathcal{O}(-k)) \rightarrow H^0(\mathbb{P}_n, \underline{\text{Ext}}_{\mathcal{O}}^1(J_Y, \mathcal{O}(-k))) \rightarrow \\ &\rightarrow H^2(\mathbb{P}_n, \underline{\text{Hom}}(J_Y, \mathcal{O}(-k))) \end{aligned}$$

gives for $n \geq 3$ an isomorphism

$$\text{Ext}_{\mathcal{O}}^1(J_Y, \mathcal{O}(-k)) \xrightarrow{\sim} H^0(\mathbb{P}_n, \underline{\text{Ext}}_{\mathcal{O}}^1(J_Y, \mathcal{O}(-k)))$$

since $\underline{\text{Hom}}(J_Y, \mathcal{O}(-k)) = \mathcal{O}(-k)$ and $H^i(\mathbb{P}_n, \mathcal{O}(-k)) = 0$ for $1 \leq i \leq n-1$ and all $k \in \mathbb{Z}$. Using

$$\begin{aligned} \underline{\text{Ext}}_{\mathcal{O}}^1(J_Y, \mathcal{O}(-k)) &\xrightarrow{\sim} \underline{\text{Ext}}^2(\mathcal{O}_Y, \mathcal{O}(-k)) \\ &\simeq \underline{\text{Ext}}^2(\mathcal{O}_Y, \mathcal{O}(-n-1)) \otimes \mathcal{O}(-k+n+1) \\ &\simeq \omega_Y \otimes \mathcal{O}_Y(-k+n+1) \quad \text{see [22]} \\ &\simeq \mathcal{O}_Y(-n-1) \otimes \det N \otimes \mathcal{O}_Y(-k+n+1) \\ &\simeq \mathcal{O}_Y, \end{aligned}$$

one finally gets an isomorphism

$$\text{Ext}_{\mathcal{O}}^1(J_Y, \mathcal{O}(-k)) \xrightarrow{\sim} H^0(Y, \mathcal{O}_Y).$$

The canonical section ξ in $H^0(Y, \mathcal{O}_Y)$ therefore gives an extension

$$0 \rightarrow \mathcal{O}(-k) \rightarrow \mathcal{F} \rightarrow J_Y \rightarrow 0$$

of J_Y by $\mathcal{O}(-k)$ through a coherent sheaf. Since ξ locally generates each stalk of $\underline{\text{Ext}}_{\mathcal{O}}^1(J_Y, \mathcal{O}(-k))$ it follows from [49] that $\mathcal{F}$ is locally free. $E := \mathcal{F}^*$ is the desired bundle.

Remarks. - 1) Barth, Larsen and Ogus [36], [45] have shown that $\text{Pic}(\mathbb{P}_n) \xrightarrow{\sim} \text{Pic}(Y)$ for $n \geq 6$ and nonsingular Y . Thus each nonsingular submanifold $Y \subset \mathbb{P}_n$, $n \geq 6$, of codimension 2 gives a holomorphic vector bundle of rank 2 on $\mathbb{P}_n$.

2) The above construction does not work without further considerations on $\mathbb{P}_2$. But if $k \leq 2$ the group $H^2(\mathbb{P}_2, \mathcal{O}(-k))$ still vanishes and the proposition 2.2.1 remains valid in that case. For arbitrary k see [51], [18].

Let us apply this proposition to produce many holomorphic 2-bundles on $\mathbb{P}_2$ and $\mathbb{P}_3$.

Examples.

1) Take Y to be the union of d simple points in $\mathbb{P}_2$. Then $\det N_{Y|\mathbb{P}_2} = \mathcal{O}_Y(2)$ and we get a holomorphic 2-bundle E on $\mathbb{P}_2$ with $c_1 = 2$ and $c_2 = d$. This shows the existence of 2-bundles with $c_1 = 0$, $c_2 \geq 0$.

2) Take Y to be the union of d disjoint lines in $\mathbb{P}_3$. Then $\det N_{Y|\mathbb{P}_3} = \mathcal{O}_Y(2)$ and we get a 2-bundle with $c_1 = 2$, $c_2 = d$. Normalizing gives $c_1 = 0$, $c_2 \geq 0$ arbitrary.

3) Take Y to be the union of r disjoint nonsingular conics in $\mathbb{P}_3$. Then $\det N_{Y|\mathbb{P}_3} \simeq \mathcal{O}_Y(3)$ and we get a 2-bundle with $c_1 = 3$, $c_2 = 2r$. This shows the existence of 2-bundles with $c_1 = -1$, $c_2 \geq 0$ even.

4) Horrocks [28]

Let $p \geq 2$ be an integer and $m_1, \dots, m_r \in \mathbb{Z}$ with $0 < m_i < p$. Choose r disjoint lines $L_i \subset \mathbb{P}_3$ and give them a nilpotent structure through

$J_{L_i} = (x^{m_i}, y^{p-m_i})$. Here x, y are equations for L_i . Take Y to be the union

of these fattened lines. Then $\det N_{Y|\mathbb{P}_3} \simeq \mathcal{O}_Y(p)$ and we get a 2-bundle with

$$c_1 = p, \quad c_2 = \sum_{i=1}^r m_i(p - m_i).$$

A short calculation shows that all $c_1, c_2 \in \mathbb{Z}$ with $c_1 c_2 \equiv 0 \pmod{2}$ are of this form (modulo twisting). Therefore all c_1, c_2 with $c_1 c_2 \equiv 0 \pmod{2}$ are the Chern classes of a holomorphic 2-bundle on $\mathbb{P}_3$.

Atiyah and Rees [2] showed that for a holomorphic 2-bundle E with even c_1 the α -invariant can be given by

$$\alpha(E) = h^0(E_{\text{norm}}(-2)) + h^2(E_{\text{norm}}(-2)) \pmod{2}.$$

Here E_{norm} denotes $E(-c_1/2)$ for c_1 even and $E((-c_1+1)/2)$ for c_1 odd.

Note that $h^2(E_{\text{norm}}(-2)) = h^1(E_{\text{norm}}(-2))$ by Serre-duality.

It takes some arithmetic [2] to show that by the above Horrocks construction one can achieve both values of α . This implies

$$\text{Vect}_{\text{hol}}^2(\mathbb{P}_3) \longrightarrow \text{Vect}_{\text{top}}^2(\mathbb{P}_3)$$

is surjective.

5) Take Y to be the disjoint union of a plane nonsingular cubic curve and a nonsingular elliptic space curve of degree d . Y gives a 2-bundle on $\mathbb{P}_3$ with Chern classes $c_1 = 4$, $c_2 = d + 3$. A short calculation shows $\alpha = 1$. Normalizing one gets the invariants

$$c_1 = 0, \quad c_2 = d + 1, \quad \alpha = 1.$$

Note that in Example 2) one has $\alpha = 0$.

6) Horrocks, Mumford [32]

These authors show the existence of a 2-bundle on $\mathbb{P}_4$ which comes from an abelian surface $Y \subset \mathbb{P}_4$. Suppose you have shown the embedding of an abelian surface Y into $\mathbb{P}_4$. The exact sequence

$$0 \longrightarrow \mathcal{O}_Y^2 \longrightarrow T_{\mathbb{P}_4}|_Y \longrightarrow N_{Y|\mathbb{P}_4} \longrightarrow 0$$

gives

$$\det N_{Y|\mathbb{P}_4} = \mathcal{O}_Y(5) \quad \text{and} \quad \deg Y = 10.$$

Hence we get a 2-bundle with $c_1 = 5$, $c_2 = 10$. This is essentially the only known indecomposable 2-bundle on $\mathbb{P}_4$.

Problem 1. Are there any holomorphic 2-bundles on $\mathbb{P}_n$, $n \geq 5$, which do not split into line bundles?

Let us close this section by some remarks on the connection of 3-bundles on $\mathbb{P}_n$ and locally complete intersections $Y \subset \mathbb{P}_n$ of codimension 2.

PROPOSITION 2.2.2 (Van de Ven, Vogelaar [64]).- Let Y be a locally complete intersection of codimension 2 in $\mathbb{P}_n$, $n \geq 3$. Suppose there is a holomorphic line bundle L on Y together with holomorphic sections $\sigma_1, \sigma_2 \in H^0(Y, L)$ such that $\{\sigma_1 = 0\} \cap \{\sigma_2 = 0\} = \emptyset$. If furthermore $\det N_{Y|\mathbb{P}_n} \otimes L^* \simeq \mathcal{O}_Y(k)$ then there is a holomorphic 3-bundle E on $\mathbb{P}_n$ with

$$c_1(E) = k, \quad c_2(E) = \deg Y, \quad c_3(E) = \deg(\sigma_1 = 0).$$

Remark.- One gets E as an extension

$$0 \longrightarrow \mathcal{O}^2 \longrightarrow E \longrightarrow J_Y(k) \longrightarrow 0.$$

As an application it is shown that all $c_1, c_2, c_3 \in \mathbb{Z}$ with $c_3 \equiv c_1 c_2 \pmod{2}$ occur as the Chern classes of a holomorphic 3-bundle on $\mathbb{P}_3$. Combining with 4) one obtains the surjectivity of the map

$$\text{Vect}_{\text{hol}}^r(\mathbb{P}_3) \longrightarrow \text{Vect}_{\text{top}}^r(\mathbb{P}_3)$$

for all r .

2.3. Monads

The description of holomorphic vector bundles on $\mathbb{P}_n$ by monads is due to Horrocks [27], [29], [31] and was recently put into a general frame by Beilinson [11]. In specific cases they have been studied by Barth, Hulek, Drinfeld and Manin [5], [8], [33], [12].

DEFINITION 2.3.1.- A monad is a complex of holomorphic vector bundles

$$0 \longrightarrow A \xrightarrow{a} B \xrightarrow{b} C \longrightarrow 0$$

which is exact except possibly at B .

Remark.- $E := \text{Ker } b / \text{Im } a$ is a holomorphic vector bundle with

$\text{rk } E = \text{rk } B - \text{rk } A - \text{rk } C$ and Chern class

$$c(E) = c(B) c(A)^{-1} c(C)^{-1}.$$

The following version of the Beilinson construction I learned from Verdier.

THEOREM 2.3.2 (Beilinson [11]).- Let E be a holomorphic vector bundle on $\mathbb{P}_n$. There exists a spectral sequence with

$$\begin{aligned} E_1^{pq} &= H^q(\mathbb{P}_n, E \otimes \Omega^p(-p)) \otimes \mathcal{O}(p), \\ E_\infty^{pq} &= 0 \quad \text{for } p + q \neq 0 \end{aligned}$$

and a filtration of E whose associated graded module is $\bigoplus_p E_\infty^{p, -p}$.

Proof. Let $\mathbb{P}_n = \mathbb{P}(V)$, V a complex vector space of dimension $n+1$. Consider the canonical exact sequence

$$0 \longrightarrow \mathcal{O}(-1) \longrightarrow \mathbb{P}(V) \times V \longrightarrow \mathcal{Q} \longrightarrow 0.$$

Here $\mathcal{Q} = T(-1)$ and $H^0(\mathbb{P}_n, \mathcal{Q}) = V$. On $\mathbb{P}_n \times \mathbb{P}_n$ we look at

$\mathcal{Q} \boxtimes \mathcal{O}(1) := \text{pr}_1^* \mathcal{Q} \otimes \text{pr}_2^* \mathcal{O}(1)$. There is a canonical section

$\sigma \in H^0(\mathbb{P}_n \times \mathbb{P}_n, \mathcal{Q} \boxtimes \mathcal{O}(1)) = V \otimes V^*$ corresponding to id_V . This section vanishes precisely and transversally at the diagonal Δ of $\mathbb{P}_n \times \mathbb{P}_n$. Hence we have the Koszul complex

$$0 \longrightarrow \Omega^n(n) \boxtimes \mathcal{O}(-n) \longrightarrow \dots \longrightarrow \Omega^1(1) \boxtimes \mathcal{O}(-1) \longrightarrow \mathcal{O}_{\mathbb{P}_n \times \mathbb{P}_n} \xrightarrow{\sigma} \mathcal{O}_\Delta \longrightarrow 0.$$

This gives

$$R^i \text{pr}_{2*}(C^\bullet \otimes \text{pr}_1^* E) = \begin{cases} 0 & \text{for } i \neq 0 \\ E & \text{for } i = 0, \end{cases}$$

where $C^v = \Omega^{-v}(-v) \boxtimes \mathcal{O}(v)$ for $v \leq 0$ and $C^v = 0$ for $v > 0$. The spectral sequence for the hypercohomology of pr_{2*} now gives the result.

Remark.- Interchanging pr_1 with pr_2 in the above proof gives a spectral sequence with

$$E_1^{pq} = H^q(\mathbb{P}_n, E(p)) \otimes \Omega^{-p}(-p)$$

satisfying the same properties as the one in the theorem.

Applications (compare [8] and [31] for a different approach)

1) Let E be a holomorphic r -bundle on $\mathbb{P}_2$ with $H^0(\mathbb{P}_2, E(-1)) = H^0(\mathbb{P}_2, E^*(-1)) = 0$. Then E is the cohomology of a monad

$$H^1(E(-2)) \otimes \mathcal{O}(-1) \rightarrow H^1(E \otimes \Omega^1) \otimes \mathcal{O} \rightarrow H^1(E(-1)) \otimes \mathcal{O}(1).$$

If $c_1(E) = 0$, then $h^1(E(-2)) = h^1(E(-1)) = c_2(E)$ by Riemann-Roch. In case E is orthogonal or symplectic (i.e. we have a nondegenerate symmetric or skew bilinear form on E), one can give the bundles in terms of linear algebra. Let H and K be complex vector spaces of dimension n and $2n + r$. K should be equipped with an orthogonal or symplectic nondegenerate form. $GL(H) \times O(K)$ acts on the linear mappings $L(H, K)$ by

$$(f, g) \cdot \varphi = g \varphi f^{-1}.$$

Using the above description of bundles by monads it is easy to show that the isomorphism classes of orthogonal (symplectic) holomorphic r -bundles on $\mathbb{P}_2 = \mathbb{P}(V)$ with $H^0(\mathbb{P}_2, E(-1)) = 0$ and $c_2(E) = n$ correspond one to one to the orbits of $GL(H) \times O(K)$ on the set of all linear maps $\alpha : V \rightarrow L(H, K)$ with

- (i) $\alpha(v)$ is injective for all $v \neq 0$
- (ii) $\alpha(v)(H)$ is for all $v \in V$ a totally isotropic subspace of K .

Remark.- $H^0(E) = 0$ is equivalent to the surjectivity of the map $H \otimes V \rightarrow K$ induced by α .

2) Let E be a holomorphic r -bundle on $\mathbb{P}_2 = \mathbb{P}(V)$ with $H^0(\mathbb{P}_2, E) = H^0(\mathbb{P}_2, E^*(-1)) = 0$. Then E comes from a monad

$$H^1(E(-2)) \otimes \mathcal{O}(-1) \xrightarrow{a} H^1(E(-1)) \otimes \Omega^1(1) \xrightarrow{b} H^1(E) \otimes \mathcal{O}.$$

One can make explicit the maps a and b [37]:

for $z \in V^* = \Gamma(\mathbb{P}_2, \mathcal{O}(1))$ denote the maps

$$H^1(E(-2)) \rightarrow H^1(E(-1)) \quad \text{and} \quad H^1(E(-1)) \rightarrow H^1(E)$$

given by the multiplication with z by $\alpha(z)$ and $\beta(z)$. At the point $x \in \mathbb{P}_2$ the map a is given by

$$(z' \wedge z'') \otimes h \rightarrow z'' \otimes \alpha(z')h - z' \otimes \alpha(z'')h.$$

Here $z', z'' \in \Omega^1(1)_x$ (note that $\mathcal{O}(-1) = \det \Omega^1(1)$). The map b is given at $x \in \mathbb{P}_2$ by

$$z \otimes k \mapsto \beta(z)k.$$

The injectivity of α is equivalent to :

for each nonzero $h \in H^1(E(-2))$ the map $z \mapsto \alpha(z)h$ from V^* to $H^1(E(-1))$ has rank at least 2 .

Now let E be of rank 2 and $c_1(E) = -1$. Serre-duality gives a symmetric nondegenerate form on $H^1(E(-1))$ and an isomorphism $H^1(E(-2))^* \simeq H^1(E)$. In this case $\beta(z) = \alpha(z)^t$, $z \in V^*$. From this one can deduce as in 1) a bijective correspondence (see [37]) between the isomorphism classes of holomorphic 2-bundles E on $\mathbb{P}_2$ with $c_1(E) = -1$, $H^0(E) = 0$, $c_2(E) = n$ and the orbits of $GL(H) \times O(K)$ on the set of all linear maps $\alpha : V^* \rightarrow L(H, K)$ satisfying

- (i) $\alpha(z')^t \alpha(z'') = \alpha(z'')^t \alpha(z')$ for $z' , z'' \in V^*$
- (ii) the map $z \mapsto \alpha(z)h$ from V^* to K is for all nonzero $h \in H$ of rank at least 2 .

Here H and K are complex vector spaces of dimension $n - 1$ and n . Furthermore K is equipped with a nondegenerate symmetric bilinear form.

The case $c_1(E) = 0$ is different. Here Serre-duality gives

$$H^1(\mathbb{P}_2, E(-2))^* \simeq H^1(\mathbb{P}_2, E(-1))$$

and for $z \in V^*$ the map

$$\alpha(z) : H^1(E(-2)) \rightarrow H^1(E(-2))^*$$

is symmetric. It takes some work (see [5], [37]) to show that the isomorphism classes of 2-bundles E with $c_1(E) = 0$, $H^0(E) = 0$ and $c_2(E) = n$ are in bijective correspondence with the orbits of $GL(H)$ acting on the set of all linear maps $\alpha : V^* \rightarrow S^2 H^*$ satisfying

- (i) the map $z \mapsto \alpha(z)h$ from V^* to H^* is for all nonzero $h \in H$ of rank at least 2
- (iii) there is a base (z_0, z_1, z_2) of V^* such that $\alpha(z_0)$ is invertible and the map $H \rightarrow H^*$ given by $\alpha(z_1)\alpha(z_0)^{-1}\alpha(z_2) - \alpha(z_2)\alpha(z_0)^{-1}\alpha(z_1)$ is of rank 2 .

Here H is a complex vector space of dimension n (≥ 2) . Monads of this type have been used by Barth [5] to classify stable 2-bundles on $\mathbb{P}_2$ with $c_1 = 0$.

3) Let E be a holomorphic r -bundle on $\mathbb{P}_3$ with $H^0(E(-1)) = 0$, $H^1(E(-2)) = 0$ ("instanton condition"), $E \simeq E^*$ and $c_2(E) = n$. Then E comes from a monad

$$H^1(E(-3) \otimes T) \otimes \mathcal{O}(-1) \rightarrow H^1(E \otimes \Omega^1) \otimes \mathcal{O} \rightarrow H^1(E(-1)) \otimes \mathcal{O}(1) .$$

In particular this shows that $H^1(\mathbb{P}_3, E(-v)) = 0$ for all $v \geq 2$. Using the notation of the first application one gets in the same way a bijection between isomorphism classes of orthogonal (symplectic) r -bundles on $\mathbb{P}_3 = \mathbb{P}(V)$ satisfying the

conditions $H^0(E(-1)) = 0$, $H^1(E(-2)) = 0$, $c_2(E) = n$ and the orbits of $GL(H) \times O(K)$ acting on the linear maps $\alpha: V \rightarrow L(H, K)$ with

- (i) $\alpha(v): H \rightarrow K$ is injective for all $v \neq 0$
- (ii) $\alpha(v)(H)$ is for all $v \in V$ a totally isotropic subspace of K .

Remark. - The condition $H^0(P_3, E) = 0$ is equivalent to the surjectivity of the map $H \otimes V \rightarrow K$ induced by α .

Monads of this type have been used to describe instantons [1], [22].

4) Let E be a holomorphic r -bundle on P_3 with $H^0(E) = H^1(E(-2)) = 0$ and $E \simeq E^*$. Then E comes from a monad

$$H^2(E(-3)) \otimes \mathcal{O}(-1) \rightarrow H^1(E(-1)) \otimes \Omega^1(1) \rightarrow H^1(E) \otimes \mathcal{O}.$$

3. Stable bundles

DEFINITION 3.1.- A holomorphic r -bundle E on P_n is said to be stable if for all proper coherent subsheaves $\mathcal{F}$ of E of rank s we have the inequality

$$\frac{c_1(\mathcal{F})}{s} < \frac{c_1(E)}{r}.$$

If we have only " $\leq$ " instead of " $<$ " then E is called semi-stable. A bundle which is not semi-stable is usually called unstable.

Remarks. - 1) This definition is due to Mumford and Takemoto [56]. Recently Gieseker [17] suggested a slightly different definition. He calls E stable if

$$\frac{p_{\mathcal{F}}(m)}{s} < \frac{p_E(m)}{r}$$

for $m \gg 0$. Here $p_{\mathcal{F}}(m) = \chi(P_n, \mathcal{F}(m))$ is the Hilbert polynomial of $\mathcal{F}$. With this definition one generally gets more stable but fewer semi-stable bundles than before.

2) It is straightforward [56] that stable bundles E are always simple, i.e. $H^0(E^* \otimes E) = \mathbb{C}$, and therefore indecomposable.

3) T_{P_n} is stable [35].

PROPOSITION 3.2 [4].- The stable 2-bundles on P_n are precisely the simple ones.

Proof. Assume E to be simple. We can choose $k \in \mathbb{Z}$ minimal with $H^0(E(k)) \neq 0$. Take a nonzero $\sigma \in H^0(E(k))$ and put $Y = \{\sigma = 0\}$. Y is of codimension 2 and we get an exact sequence

$$0 \rightarrow \mathcal{O} \xrightarrow{\sigma} E(k) \rightarrow J_Y(c_1(E) + 2k) \rightarrow 0.$$

If $c_1 + 2k \leq 0$ we get a "non-trivial" endomorphism of $E(k)$ by composing

$$E(k) \rightarrow J_Y(c_1 + 2k) \hookrightarrow \mathcal{O}(c_1 + 2k) \hookrightarrow \mathcal{O} \xrightarrow{\sigma} E(k).$$

Hence $c_1 + 2k > 0$.

Now let $\mathcal{O}(\ell)$ be a subsheaf of E . By minimality of k we get $-\ell \geq k$ and therefore $\ell < c_1/2$. This shows the stability of E .

Remark.— It is easy to see that a 2-bundle E on $\mathbb{P}_n$ is stable if and only if $H^0(\mathbb{P}_n, E_{\text{norm}}) = 0$. For 3-bundles with $c_1 = 0$ stability is equivalent to $H^0(E) = H^0(E^*) = 0$.

Problem 2. Give a similar criterion of stability for bundles of higher rank.

Schwarzenberger has shown [52] that Riemann-Roch implies that the Chern classes of a stable 2-bundle on $\mathbb{P}_2$ have to satisfy $c_1^2 - 4c_2 < 0$ (for a semi-stable 2-bundle one has $c_1^2 - 4c_2 \leq 0$). In fact $c_1^2 - 4c_2 = -4$ cannot occur for a stable 2-bundle on $\mathbb{P}_2$ [38].

It is a general fact, proved by Maruyama [43], that the restriction of a semi-stable r -bundle on $\mathbb{P}_n$, $r < n$, to a general hyperplane is semi-stable again (Barth [4] showed the same to be true for stable 2-bundles on $\mathbb{P}_n$, $n \geq 3$, with the exception of the Null-correlation bundle). Hence for a semi-stable 2-bundle E on $\mathbb{P}_n$ we have

$$c_1^2 - 4c_2 \leq 0$$

and for stable 2-bundles one necessarily has

$$c_1^2 - 4c_2 < 0.$$

Problem 3. Determine similar necessary conditions for stable (semi-stable) holomorphic bundles of higher rank.

We show next how stability of a 2-bundle E on $\mathbb{P}_n$ coming from a locally complete intersection $Y \subset \mathbb{P}_n$ of codimension 2 is reflected by Y .

Let $Y \subset \mathbb{P}_n$, $n \geq 2$, be a locally complete intersection of codimension 2 and $\det N_{Y|\mathbb{P}_n} = \mathcal{O}_Y(k)$. Then we can find an extension E of $N_{Y|\mathbb{P}_n}$ as in 2.2.1.

PROPOSITION 3.3 (see [25]).— E is stable if and only if $k > 0$ and Y is not contained in any hypersurface of degree $d \leq k/2$.

Proof. We have an exact sequence

$$0 \rightarrow \mathcal{O} \rightarrow E \rightarrow J_Y(k) \rightarrow 0.$$

If E is stable then $c_1(E) = k > 0$.

Assume k to be even. The sequence

$$0 \rightarrow \mathcal{O}(-k/2) \rightarrow E_{\text{norm}} \rightarrow J_Y(k/2) \rightarrow 0$$

gives

$$H^0(E_{\text{norm}}) \xrightarrow{\sim} H^0(J_Y(k/2)).$$

Stability of E is equivalent to $H^0(E_{\text{norm}}) = 0$. Therefore $H^0(J_Y(k/2)) = 0$, which is equivalent to the fact that Y is not contained in any hypersurface of degree $\leq k/2$. Assume on the other hand $k > 0$ and $H^0(J_Y(k/2)) = 0$. This gives $H^0(E_{\text{norm}}) = 0$ which is the stability of E . The case c_1 odd is treated in a similar way.

Using this criterion we re-examine the examples of 2.1.

Examples.- 1) If E comes from d simple points in $\mathbb{P}_2$, E is stable if and only if the points do not all lie on a line. This shows the existence of stable 2-bundles on $\mathbb{P}_2$ with $c_1 = 0$, $c_2 \geq 2$.

2) If E comes from d disjoint lines in $\mathbb{P}_3$ then E is stable if and only if these lines are not contained in a plane. This is the case for $d \geq 2$. This gives stable 2-bundles on $\mathbb{P}_3$ with $c_1 = 0$, $c_2 \geq 1$, $\alpha = 0$.

3) For bundles coming from disjoint nonsingular conics we have the same result as in 2). One gets stable bundles with $c_1 = -1$, $c_2 \geq 2$ even.

4) If E comes from a plane cubic and a disjoint elliptic curve of degree d then E is stable if $d \geq 4$. This gives stable bundles on $\mathbb{P}_3$ with $c_1 = 0$, $c_2 \geq 5$, $\alpha = 1$.

5) The 2-bundle of Horrocks and Mumford on $\mathbb{P}_4$ is stable since an abelian surface Y can neither lie in some $\mathbb{P}_3$ (because of $\pi_1(Y) \neq 0$) nor in some hyperquadric Q (consider normal bundles).

Here we wish to draw the attention of the reader to an example of a stable 3-bundle on $\mathbb{P}_5$ constructed by Horrocks [30] using representation theory.

Let us close this section by giving the following

Conjecture.- Each 2-bundle on $\mathbb{P}_n$, $n \geq 5$, which is not stable is a direct sum of line bundles.

In [20] a "proof" for this was given even for $n \geq 4$. Unfortunately there is a gap in that paper.

The conjecture has nice consequences [20], [50] :

1) Each topological 2-bundle on $\mathbb{P}_n$, $n \geq 5$, which is not the direct sum of two line bundles and satisfying $c_1^2 - 4c_2 \geq 0$ cannot have an analytic structure. By [46],

[54], [55] there are many topological 2-bundles with $c_1^2 - 4c_2 \geq 0$ and which do not split.

2) Each holomorphic 2-bundle on $\mathbb{P}_5$ which can be extended topologically to $\mathbb{P}_n$, n arbitrarily large, is the direct sum of line bundles.

This sharpened the theorem of Barth and Van de Ven [9] on Babylonian vector bundles (see also [48], [61]).

3) Each nonsingular submanifold $Y \subset \mathbb{P}_n$ of codimension 2 is a complete intersection if $n \geq 6$ and $n \geq \frac{1}{3}\sqrt{\deg(Y)} + 1$. This would improve some results in [3].

One can even show, for example, that a nonsingular 4-dimensional submanifold $Y \subset \mathbb{P}_6$ is a complete intersection if $\deg Y \leq 514$.

4) Furthermore one could improve the results of Barth and Van de Ven in [10].

4. Moduli of stable bundles

Sofar we commented the points I and II of the introduction. To deal with III one would like to introduce on the set of isomorphism classes of stable holomorphic r -bundles on $\mathbb{P}_n$ with fixed topological type a "good" analytic structure.

Consider the functor

$$\Sigma(c_1, \dots, c_r) : \underline{\text{An}} \longrightarrow \underline{\text{Ens}}$$

from analytic spaces to sets given by

$$\Sigma(c_1, \dots, c_r)(S) := \{ \text{bundles } E \text{ on } \mathbb{P}_n \times S \text{ of fixed rank with } E(s) \text{ stable and} \\ c_i(E(s)) = c_i \text{ for } i = 1, \dots, r \text{ and } s \in S \} / \sim .$$

Here $E_2 \sim E_1$ if $E_2 \cong \text{pr}_S^*(L) \otimes E_1$ for a holomorphic line bundle L on S .

Σ is contravariant in an obvious way.

DEFINITION 4.1.- $M = M(c_1, \dots, c_r) \in \underline{\text{An}}$ is a coarse moduli space for $\Sigma(c_1, \dots, c_r)$ if there is a morphism of functors

$$\Sigma \longrightarrow \text{Hom}(-, M)$$

with

$$\Sigma(\text{pt}) \xrightarrow{\sim} M .$$

Furthermore M should be minimal with respect to these properties, i.e. if N is another analytic space satisfying the above then there should be a unique morphism $M \rightarrow N$ making the diagram

$$\begin{array}{ccc} \Sigma & \longrightarrow & \text{Hom}(-, M) \\ & \searrow & \swarrow \\ & \text{Hom}(-, N) & \end{array}$$

commutative.

If a coarse moduli space exists one has put in a functorial way an analytic structure onto the stable bundles on $\mathbb{P}_n$ with fixed Chern classes and fixed rank.

If M represents Σ then M is said to be a fine moduli space. This is equivalent to the existence of a universal family over $M \times \mathbb{P}_n$.

It seems much easier to construct a coarse moduli space M in the analytic category than to do it in the algebraic category. In the algebraic category the existence was proved by Maruyama [39], [40], [41] by using Mumford's geometric invariant approach. Maruyama could not show that M is always of finite type. For $n = 2$ and arbitrary rank this was shown to be true by Gieseker [17]. For arbitrary n and rank ≤ 4 it was verified recently by Maruyama [43].

These authors also study compactifications of M and it turns out that one has not only to admit semi-stable bundles but also semi-stable torsion free coherent sheaves.

Our object here is only to mention some specific results for the moduli spaces M of bundles over $\mathbb{P}_2$ and $\mathbb{P}_3$.

By deformation theory the Zariski tangent space of M at m is $H^1(\text{End}(E))$ if E is the bundle corresponding to m . If $H^2(\text{End}(E)) = 0$ then M is smooth at m . In particular the moduli spaces of stable bundles on $\mathbb{P}_2$ are nonsingular. By Riemann-Roch we get

$$\dim M_{\mathbb{P}_2}(c_1, c_2, r) = (1-r)c_1^2 + 2rc_2 - r^2 + 1.$$

For rank 2 we get

$$\dim M_{\mathbb{P}_2}(c_1, c_2) = 4c_2 - c_1^2 - 3.$$

Let us summarize the properties of $M_{\mathbb{P}_2}(c_1, c_2)$.

THEOREM 4.2.- $M_{\mathbb{P}_2}(c_1, c_2)$ is a smooth, quasi-projective manifold of dimension $4c_2 - c_1^2 - 3$. M is connected and rational. M is a fine moduli space if and only if $4c_2 - c_1^2 \neq 0$ (8).

Remarks.- The rationality and connectedness was proved by Barth [5] for c_1 even and by Hulek [33] for c_1 odd using monads. Maruyama [42] showed that M is connected, unirational (and in some cases rational) and that M is a fine moduli space if $4c_2 - c_1^2 \neq 0$ (8). Le Potier [37] proved the nonexistence of a universal family for $4c_2 - c_1^2 \equiv 0$ (8) using monads. He showed that in this case there are topological obstructions to the existence of the universal family. In doing this he calculated

$$\pi_1(M(0, c_2)) = \begin{cases} \mathbb{Z}/3\mathbb{Z} & \text{for } c_2 = 2 \\ 0 & \text{otherwise} \end{cases}$$

$$\pi_2(M(0, c_2)) = \begin{cases} \mathbb{Z}/2\mathbb{Z} & \text{for } c_2 = 2 \\ \mathbb{Z}/2\mathbb{Z} \oplus \mathbb{Z} & \text{for } c_2 \geq 2, \quad c_2 \text{ even} \\ \mathbb{Z} & \text{for } c_2 \text{ odd.} \end{cases}$$

To conclude this section we give the simplest examples of moduli spaces on $\mathbb{P}_2$ and $\mathbb{P}_3$ which can be deduced quickly from the description of bundles by monads.

Examples. - 1) $M_{\mathbb{P}_2}(-1, 1) = \{\Omega^1(1)\}$.

This follows immediately from Application 2 of 2.3.

2) $M_{\mathbb{P}_2}(-1, 2) = S^2\mathbb{P}_2 \setminus \Delta$ (see [37]).

The application 2 of 2.3 shows that

$$M(-1, 2) = \{\alpha : V^* \rightarrow \mathbb{C}^2 \text{ linear and surjective}\} \text{ modulo the action of } \mathbb{C}^* \times O(\mathbb{C}^2).$$

Here $\mathbb{C}^2$ is equipped with a nondegenerate symmetric bilinear form. A linear algebraic calculation identifies the righthand side to $(\mathbb{P}(V) \times \mathbb{P}(V)) \setminus \Delta$ modulo $\mathbb{Z}/2\mathbb{Z}$. This finally gives $M(-1, 2) \simeq S^2\mathbb{P}_2 \setminus \Delta$.

3) $M_{\mathbb{P}_2}(0, 2) = \{\text{nonsingular conics in } \mathbb{P}_2\}$, [5].

By application 2 of 2.3 one has

$$M(0, 2) = \text{Isom}(V^*, S^2H^*) / \text{GL}(H).$$

H is of dimension 2. Let $C := \{q \in S^2(H^*) : \det q = 0\}$; for $\alpha \in \text{Isom}(V^*, S^2H^*)$ the inverse image $\alpha^{-1}(C)$ will be a nonsingular conic. α' , $\alpha \in \text{Isom}(V^*, S^2H^*)$ with $\alpha'^{-1}(C) = \alpha'^{-1}(C)$ differ by an automorphism $\gamma \in \text{Aut}(S^2H^*)$ with $\gamma(C) = C$. But these γ 's come from automorphisms of H . This proves our claim.

4) $M_{\mathbb{P}_3}(0, 1) = \text{PGL}(3, \mathbb{C}) / \text{Sp}(2, \mathbb{C})$ (see [4]).

By application 3 of 2.3 we have

$$M(0, 1) = \text{Isom}(\mathbb{C}^4, \mathbb{C}^4) / \mathbb{C}^* \times \text{Sp}(2, \mathbb{C}) = \text{PGL}(3, \mathbb{C}) / \text{Sp}(2, \mathbb{C}).$$

In particular $\text{PGL}(3)$ operates transitively on $M(0, 1)$. The Null-correlation bundle belongs to $M(0, 1)$.

Hartshorne [25] gives a description of $M_{\mathbb{P}_3}(0, 2)$. In particular $M_{\mathbb{P}_3}(0, 2)$ is still connected. For $c_2 \geq 3$ the space $M_{\mathbb{P}_3}(0, c_2)$ will be divided into 2 components by the α -invariant. The following example due to Barth and Hulek [8] (see also [25]) shows that $M_{\mathbb{P}_3}(0, c_2)$ is reducible if c_2 is odd and at least 5.

Consider the monad

$$\mathcal{O}(-m-1) \xrightarrow{a} \mathcal{O}(m) \oplus \mathcal{O} \oplus \mathcal{O} \oplus \mathcal{O}(-m) \xrightarrow{b} \mathcal{O}(m+1)$$

on $\mathbb{P}_3$. The map $a \in H^0(\mathcal{O}(2m+1) \oplus \mathcal{O}(m+1) \oplus \mathcal{O}(m+1) \oplus \mathcal{O}(1))$ has to be chosen such that the a_i have no common zero. On $\mathcal{O}(m) \oplus \mathcal{O} \oplus \mathcal{O} \oplus \mathcal{O}(-m)$ take the symplectic form

$$q = \begin{pmatrix} & & & 1 \\ & 0 & & \\ & & 1 & \\ -1 & & & 0 \end{pmatrix}$$

and put $b = a^t$.

The stable 2-bundles defined by these monads have Chern classes $c_1 = 0$, $c_2 = 2m + 1$.

This family of bundles depends effectively on

$$\#a's - \dim(\mathbb{C}^* \times \mathcal{O}(q))$$

parameters (compare 2.3).

One checks that $\dim \mathcal{O}(q) = 4 + 2\binom{m+3}{3} + \binom{2m+3}{3}$ and thus gets that the family depends on

$$3m^2 + 10m + 8$$

parameters.

For $m \geq 2$ this number is bigger than $16m + 5 = 8c_2 - 3$ which is the dimension of the Zariski-open smooth part of bundles E with $H^2(\mathbb{P}_3, \text{End}(E)) = 0$.

Questions. - 1) Are $M_{\mathbb{P}_3}(0,3)$ and $M_{\mathbb{P}_3}(0,4)$ nonsingular and do they have only two components (given by α)?

2) What can be said about $M(0, c_2)$, c_2^- even?

3) Is the Zariski-open part of mathematical instanton bundles of $M_{\mathbb{P}_3}(0, c_2)$, i.e. the bundles E with $H^1(\mathbb{P}_3, E(-2)) = 0$, nonsingular?

5. Jumping lines and uniform bundles

If E is a holomorphic r -bundle on $\mathbb{P}_n$ the restriction of E to a projective line $L \subset \mathbb{P}_n$ is by the theorem of Grothendieck of the form

$$E|_L \simeq \mathcal{O}(a_1) \oplus \dots \oplus \mathcal{O}(a_r).$$

The integers a_i depend on L but are the same for the general line L . Lines for which $E|_L$ is different from the generic form are called jumping lines. The set of jumping lines will be denoted by $S(E)$. It is a closed analytic subset of $\text{Gr}(1, n)$.

One of the main tools in studying stable 2-bundles on $\mathbb{P}_n$ is the theorem of Grauert and Müllich [18], [4].

THEOREM 5.1.- For a stable normalized 2-bundle E on $\mathbb{P}_n$ the restriction of E to the general line is

$$E|L \simeq \begin{cases} \mathcal{O} \oplus \mathcal{O} & \text{for } c_1 = 0 \\ \mathcal{O} \oplus \mathcal{O}(-1) & \text{for } c_1 = -1. \end{cases}$$

To study stable bundles of higher rank it would be desirable to solve the following

Problem 4. Let E be a stable r -bundle on $\mathbb{P}_n$. Is it true that for the general line L one has

$$E|L \simeq \mathcal{O}(a_1) \oplus \dots \oplus \mathcal{O}(a_r).$$

with $a_1 \geq a_2 \geq \dots \geq a_r$, $a_{i-1} - a_i \leq 1$ for $i = 2, \dots, r$?

For $r = 2$ it is true by the Grauert-Müllich theorem. For $r = 3$ and $n = 2$ it is true by [43].

For stable 2-bundles E with c_1 even one can say more about $S(E)$. The Grauert-Müllich theorem implies for a normalized stable 2-bundle E on $\mathbb{P}_n$:

$$S(E) = \{L : H^0(L, E(-1)|L) \neq 0\}.$$

Suppose now $n = 2$ and $c_1 = 0$. The exact sequence

$$0 \rightarrow H^0(E(-1)|L) \rightarrow H^1(E(-2)) \xrightarrow{\alpha(L)} H^1(E(-1))$$

shows that

$$S(E) = \{L \in \mathbb{P}_2^* : \det \alpha(L) = 0\},$$

because $h^1(E(-2)) = h^1(E(-1)) = c_2(E)$. Hence $S(E)$ is a curve of degree $c_2(E)$. Barth [4] has shown that this remains true if $n > 2$, i.e. $S(E)$ is a divisor of degree $c_2(E)$ in $\text{Gr}(1, n)$.

For c_1 odd $S(E)$ is not a hypersurface. For example look at $E \in M_{\mathbb{P}_2}(-1, 2) = S^2\mathbb{P}_2 \setminus \Delta$. If E corresponds to 2 different points $p_1, p_2 \in \mathbb{P}_2$ then there is only one jumping line: the line containing p_1 and p_2 . In order to associate geometric objects to $M_{\mathbb{P}_2}(-1, c_2)$ Hulek [33] gives the following

DEFINITION 5.2.- Let E be a normalized 2-bundle on $\mathbb{P}_2$. A line $L \subset \mathbb{P}_2$ is called a jumping line of the second kind if $H^0(E|L^2) \neq 0$. Here L^2 denotes the first infinitesimal neighborhood of L in $\mathbb{P}_2$.

Hulek shows that for stable 2-bundles on $\mathbb{P}_2$ with $c_1 = -1$ the set $C(E)$ of jumping lines of the second kind is a curve in $\mathbb{P}_2^*$ of degree $2c_2(E) - 2$. Furthermore

$$S(E) \subset \text{Sing } C(E)$$

and in general one has equality.

Holomorphic bundles E on $\mathbb{P}_n$ with $S(E) = \emptyset$ are called uniform. Van de Ven [63] showed that a uniform 2-bundle on $\mathbb{P}_n$ either splits into line bundles or is of the form $T_{\mathbb{P}_2}(k)$, $k \in \mathbb{Z}$. This was generalized by Sato [47] to r -bundles on $\mathbb{P}_n$ with $r \leq n$. Elencwajg [14] proved that uniform 3-bundles E on $\mathbb{P}_2$ (and therefore on $\mathbb{P}_n$ for all n by Sato's result) are homogeneous, i.e. $\sigma^*E \simeq E$ for all $\sigma \in \text{PGL}(n)$. This gave much evidence to the old conjecture [51] that uniform bundles of arbitrary rank on $\mathbb{P}_n$ are homogeneous.

Recently Elencwajg [15] gave an example of a uniform 4-bundle on $\mathbb{P}_2$ which is not homogeneous. In fact he uses a monad of the type described in application 2 of 2.3.

Problem 5. Does every uniform unstable bundle on $\mathbb{P}_n$ split?

For rank two this is true (and easy to see).

Finally we recommend to consult a recent problem list (26 problems) on vector bundles on $\mathbb{P}_n$ compiled by Hartshorne [26]. There one can especially find many problems related to instantons which we have almost completely neglected due to limited space and knowledge.

BIBLIOGRAPHY

- [1] M. F. ATIYAH, N. J. HITCHIN, V. G. DRINFELD, Ju. MANIN - Construction of instantons, Physics Letters, 65 A (1978), 185-187.
- [2] M. F. ATIYAH, E. REES - Vector bundles on projective 3-space, Inventiones math., 35 (1976), 131-153.
- [3] W. BARTH - Submanifolds of low codimension in projective space, Proc. I.C.M. Vancouver (1975), 409-413.
- [4] W. BARTH - Some properties of stable rank -2 vector bundles on $\mathbb{P}_n$, Math. Ann., 226 (1977), 125-150.
- [5] W. BARTH - Moduli of vector bundles on the projective plane, Inventiones math. 42 (1977), 63-91.
- [6] W. BARTH - Stable vector bundles on $\mathbb{P}_3$, some experimental data, Preprint 1978.
- [7] W. BARTH, G. ELENCAWAG - Concernant la cohomologie des fibrés algébriques stables sur $\mathbb{P}_n(\mathbb{C})$. In Variétés Analytiques Compactes, Nice 1977, Lecture Notes in Math., 683 (1978), 1-24.
- [8] W. BARTH, K. HULEK - Monads and moduli of vector bundles, Manuscripta math. 25(1978), 323-347.
- [9] W. BARTH, A. VAN DE VEN - A decomposability criterion for algebraic 2-bundles on projective spaces, Inventiones Math., 25 (1974), 91-106.
- [10] W. BARTH, A. VAN DE VEN - On the geometry in codimension 2 of Grassmann manifolds, In classification of algebraic varieties and compact complex manifolds, Lecture Notes in Math., 412 (1974), 1-35.
- [11] A. BEILINSON - Coherent sheaves on $\mathbb{P}^N$ and problems of linear algebra, J. of functional analysis and its applications, 12 (1978), 68-69.
- [12] V. G. DRINFELD, Ju. MANIN - On locally free sheaves over $\mathbb{P}_3$ connected with Yang-Mills fields, Uspechi, 33 (1978), 165-166.
- [13] V. G. DRINFELD, Ju. MANIN - J. Functional Analysis and its applications, 12(1978), 78-79.
- [14] G. ELENCAWAG - Les fibrés uniformes de rang 3 sur $\mathbb{P}_2(\mathbb{C})$ sont homogènes, Math. Ann., 231 (1978), 217-227.
- [15] G. ELENCAWAG - Des fibrés uniformes non homogènes, Preprint 1978, Nice.
- [16] D. FERRAND - Courbes gauches et fibrés de rang 2, C. R. Acad. Sci. Paris, 281 (1975), A 345-347.
- [17] D. GIESEKER - On the moduli of vector bundles on an algebraic surface, Ann. of Math., 106 (1977), 45-60.
- [18] H. GRAUERT, G. MÜLICH - Vektorbündel vom Rang 2 über dem n-dimensionalen komplex-projektiven Raum, Manuscripta math., 16 (1975), 75-100.

- [19] H. GRAUERT, R. REMMERT - Zur Spaltung lokal-freier Garben über Riemannschen Flächen, Math. Z., 144 (1975), 35-43.
- [20] H. GRAUERT, M. SCHNEIDER - Komplexe Unterräume und holomorphe Vektorraumbündel vom Rang zwei, Math. Ann., 230 (1977), 75-90.
- [21] A. GROTHENDIECK - Sur la classification des fibrés holomorphes sur la sphère de Riemann, Amer. J. Math., 79 (1956), 121-138.
- [22] A. GROTHENDIECK - Théorèmes de dualité pour les faisceaux algébriques cohérents, Séminaire Bourbaki, Exposé 149, Mai 1957, W. A. Benjamin-Addison-Wesley, New York, 1966.
- [23] R. HARTSHORNE - Varieties of small codimension in projective space, Bull. Amer. Math. Soc., 80 (1974), 1017-1032.
- [24] R. HARTSHORNE - Stable vector bundles and instantons, Comm. Math. Phys., 59 (1978), 1-15.
- [25] R. HARTSHORNE - Stable vector bundles of rank 2 on $\mathbb{P}^3$, Math. Ann., 238(1978), 229-280.
- [26] R. HARTSHORNE - Algebraic vector bundles on projective spaces: a problem list, Preprint 1978.
- [27] G. HORROCKS - Vector bundles on the punctured spectrum of a local ring, Proc. London Math. Soc. (3), 14 (1964), 689-713.
- [28] G. HORROCKS - A construction for locally free sheaves, Topology, 7 (1968), 117-120.
- [29] G. HORROCKS - Letter to Mumford (1971).
- [30] G. HORROCKS - Examples of rank three vector bundles on five-dimensional projective space, J. London Math. Soc. (2), 18 (1978), 15-27.
- [31] G. HORROCKS - Construction of bundles on $\mathbb{P}^n$, Preprint 1978.
- [32] G. HORROCKS, D. MUMFORD - A rank 2 vector bundle on $\mathbb{P}^4$ with 15,000 symmetries, Topology, 12 (1973), 63-81.
- [33] K. HULEK - Talk at Oberwolfach 1978.
- [34] W. HULSBERGEN - Vector bundles on the complex projective plane, Thesis, Leiden (1976).
- [35] S. G. LANGTON - Valuative criteria for families of vector bundles on algebraic varieties, Ann. of Math. (2), 101 (1975), 88-110.
- [36] M. E. LARSEN - On the topology of complex projective manifolds, Inventiones math., 19 (1973), 251-260.
- [37] J. LE POTIER - Fibrés stables de rang 2 sur $\mathbb{P}_2(\mathbb{C})$, Preprint 1978, Paris VII.
- [38] M. MARUYAMA - On a family of algebraic vector bundles, Number theory, algebraic geometry and commutative algebra, in honor of Y. Akizuki, Kinokuniya, Tokyo (1973), 95-146.

- [39] M. MARUYAMA - Stable vector bundles on an algebraic surface, Nagoya Math. J., 58 (1975), 25-68.
- [40] M. MARUYAMA - Openness of a family of torsion free sheaves, J. Math. Kyoto Univ., 16 (1976), 627-637.
- [41] M. MARUYAMA - Moduli of stable sheaves, I. J. Math. Kyoto Univ., 17 (1977), 91-126.
- [42] M. MARUYAMA - Moduli of stable sheaves, II. J. Math. Kyoto Univ., to appear.
- [43] M. MARUYAMA - Boundedness of semi-stable sheaves of small ranks, Preprint 1978.
- [44] G. MÜLICH - Familien holomorpher Vektorraumbündel über $\mathbb{P}_1$ und unzerlegbare holomorphe 2-Bündel über der projektiven Ebene, Thesis, Göttingen, 1974.
- [45] A. OGUS - On the formal neighborhood of a subvariety of projective space, Amer. J. Math., 97 (1976), 1085-1107.
- [46] E. REES - Some rank two bundles on $\mathbb{P}_n(\mathbb{C})$ whose Chern classes vanish, In Variétés analytiques Compactes, Nice 1977, Lecture Notes in Math., 683 (1978), 25-28.
- [47] E. SATO - Uniform vector bundles on a projective space, J. Math. Soc. Japan, 28 (1976), 123-132.
- [48] E. SATO - On the decomposability of infinitely extendable vector bundles on projective spaces and Grassmann varieties, J. Math. Kyoto Univ., 17 (1977), 127-150.
- [49] J.-P. SERRE - Sur les modules projectifs, Séminaire Dubreil-Pisot, (1960/61), Exposé 2.
- [50] M. SCHNEIDER - Mannigfaltigkeiten der Codimension zwei im projektiven Raum, Private Notes.
- [51] R. L. E. SCHWARZENBERGER - Vector bundles on algebraic surfaces, Proc. London Math. Soc. (3), 11 (1961), 601-622.
- [52] R. L. E. SCHWARZENBERGER - Vector bundles on the projective plane, Proc. London Math. Soc. (3), 11 (1961), 623-640.
- [53] R. L. E. SCHWARZENBERGER - Appendix I of Topological Methods in Algebraic Geometry by F. Hirzebruch, Springer 1966.
- [54] L. SMITH - Complex 2-plane bundles over $\mathbb{CP}(n)$, Manuscripta Math., 24 (1978), 221-228.
- [55] R. SWITZER - Complex 2-plane bundles over complex projective space, Preprint 1978, Göttingen.
- [56] F. TAKEMOTO - Stable vector bundles on algebraic surfaces, Nagoya Math. J. 47(1972), 29-48.

- [57] H. TANGO - On $(n-1)$ -dimensional projective spaces contained in the Grassmann variety $Gr(n,1)$, J. Math. Kyoto Univ., 14 (1974), 415-460.
- [58] H. TANGO - An example of indecomposable vector bundle of rank $n-1$ on P^n , J. Math. Kyoto Univ., 16(1976), 137-141.
- [59] H. TANGO - On morphisms from projective space P^n to the Grassmann variety $Gr(n,d)$, J. Math. Kyoto Univ., 16(1976), 201-207.
- [60] A. THOMAS - Almost complex structures on complex projective spaces, Trans. Amer. Math. Soc., 193 (1974), 123-132.
- [61] A. N. TJURIN - Finite dimensional vector bundles over infinite varieties, Math. U.S.S.R. Izvestija 10 (1976), 1187-1204.
- [62] G. TRAUTMANN - Moduli for bundles on P_n , Preprint 1978, Kaiserslautern.
- [63] A. VAN DE VEN - On uniform vector bundles, Math. Ann., 195(1972), 245-248.
- [64] A. VAN DE VEN, J. VOGELAAR - A construction for vector bundles of rank 3, Preprint 1978, Leiden.