

SÉMINAIRE ÉQUATIONS AUX DÉRIVÉES PARTIELLES – ÉCOLE POLYTECHNIQUE

R. BEALS

C. FEFFERMAN

Classes of spatially inhomogeneous pseudodifferential operators

Séminaire Équations aux dérivées partielles (Polytechnique) (1972-1973), exp. n° 26,
p. 1-6

http://www.numdam.org/item?id=SEDP_1972-1973____A27_0

© Séminaire Équations aux dérivées partielles (Polytechnique)
(École Polytechnique), 1972-1973, tous droits réservés.

L'accès aux archives du séminaire Équations aux dérivées partielles (<http://sedp.cedram.org>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>

ECOLE POLYTECHNIQUE
CENTRE DE MATHÉMATIQUES
17, rue Descartes
75230 Paris Cedex 05

S E M I N A I R E G O U L A O U I C - S C H W A R T Z 1 9 7 2 - 1 9 7 3

CLASSES OF SPATIALLY INHOMOGENEOUS

PSEUDODIFFERENTIAL OPERATORS

by R. BEALS and C. FEFFERMAN

"Reprinted from the Proceedings of the National Academy U. S. A."

Abstract. One can obtain sharp information on a pseudodifferential operator $p(x, D)$ by embedding the symbol p in a symbolic calculus specially designed to reflect the behavior of p . We sketch the development of symbolic calculi arising in this connection, and use our results to give simple proofs of the sharp Gårding inequality and of sufficiency of Nirenberg-Treves' condition (P) for local solvability of equations of principal type.

I. A Calculus of Symbols.

Let ϕ and φ be positive smooth functions on $\mathbb{R}^n \times \mathbb{R}^r$ satisfying the inequalities

$$(1) \quad c \leq \phi(x, \xi) \leq C(1 + |\xi|^2)^{1/2}, \quad c(1 + |\xi|^2)^{-1/2} \leq \varphi(x, \xi) \leq C.$$

$$(2) \quad \phi\varphi \geq c.$$

$$(3) \quad |D_x \varphi| \leq C, \quad |D_\xi \phi| \leq C, \quad |D_x \phi(x, \xi)| \leq C \frac{\phi(x, \xi)}{\varphi(x, \xi)},$$

$$|D_\xi \varphi(x, \xi)| \leq C \frac{\varphi(x, \xi)}{\phi(x, \xi)}.$$

$$(4) \quad c \frac{\phi(x, \xi)}{\varphi(x, \xi)} \leq \frac{\phi(y, \eta)}{\varphi(y, \eta)} \leq C \frac{\phi(x, \xi)}{\varphi(x, \xi)} \quad \text{if} \quad |\xi - \eta| < \frac{1}{2} |\xi|.$$

For real numbers M and m we define the class $S_{\phi, \varphi}^{M, m}$ of symbols to consist of all smooth functions $a(x, \xi)$ on $\mathbb{R}^n \times \mathbb{R}^n$ which satisfy the conditions $|D_x^\alpha D_\xi^\beta a(x, \xi)| \leq C_{\alpha\beta} \phi^{M-|\beta|}(x, \xi) \varphi^{m-|\alpha|}(x, \xi)$ for all multi-indices α and β .

Example: Take $\phi = (1 + |\xi|^2)^{\rho/2}$ and $\varphi = (1 + |\xi|^2)^{-\delta/2}$ with $0 \leq \delta \leq \rho \leq 1$ ($\delta \neq 1$). Then $S_{\phi, \varphi}^{M, m}$ is essentially Hörmander's class $S_{\rho, \delta}^{Mp - m\delta}$ (see [1]).

Corresponding to $a(x, \xi) \in S_{\phi, \varphi}^{M, m}$ we define an operator $a(x, D)$ from the Schwartz class \mathcal{S} to itself by the standard formula

$$a(x, D) u(x) = \int_{\mathbb{R}^n} e^{ix \cdot \xi} a(x, \xi) \hat{u}(\xi) d\xi,$$

where \wedge denotes the Fourier transform.

Theorem 1. Let $a \in S_{\phi, \varphi}^{M, n}$ and $b \in S_{\phi, \varphi}^{M', m'}$ be symbols. Then

(A) The operator $a(x, D) \circ b(x, D)$ defined on \mathcal{S} is the pseudodifferential operator arising from a symbol $a \circ b \in S_{\phi, \varphi}^{M+M', m+m'}$. Moreover, $a \circ b$ has the asymptotic expansion $a \circ b \sim \sum_{\alpha} \frac{1}{\alpha!} \partial_\xi^\alpha a(x, \xi) \cdot D_x^\alpha b(x, \xi)$ in the sense that for any $N > 0$,

$$a \circ b - \sum_{|\alpha| < N} \frac{1}{\alpha!} \partial_\xi^\alpha a \cdot D_x^\alpha b \in S_{\phi, \varphi}^{M+M' - N, m+m' - N}$$

(B) Similarly, the adjoint operator $a^*(x, D)$ restricted to \mathcal{S} is the pseudodifferential operator arising from a symbol $a^\#(x, \xi) \in S_{\phi, \varphi}^{M, m}$. Moreover

$a^\#$ has the asymptotic expansion $a^\# \sim \sum_{\alpha} \frac{1}{\alpha!} \partial_\xi^\alpha D_x^\alpha \bar{a}(x, \xi)$, i.e.,

$$a^\# - \sum_{|\alpha| < N} \frac{1}{\alpha!} \partial_\xi^\alpha D_x^\alpha \bar{a} \in S_{\phi, \varphi}^{M-N, m-N}.$$

Theorem 2. If $a(x, \xi)$ is a symbol in $S_{\phi, \varphi}^{0,0}$, then $a(x, D)$ is bounded on L^2 .

Remarks: 1. Theorem 2 leads to an extension of itself in which one defines analogues of the Sobolev spaces H^s relative to ϕ and φ .

2. If, in addition to (1)-(4) one assumes the estimate

$$(5) \quad \phi(x, \xi) \geq c |\xi| \varphi(x, \xi),$$

then one can prove a theorem on the behavior of operators in $S_{\phi, \varphi}^{M, m}$ under smooth changes of co-ordinates.

We forgo writing down the explicit results.

Our proofs of theorems 1 and 2 rely heavily on ideas from a recent paper of A. P. Calderón and R. Vaillancourt [2].

II. Applications. We begin with a simple proof of the sharp Gårding inequality [3], which asserts that $\operatorname{Re}(a(x, D)u, u) \geq -C \|u\|^2$ for all $u \in L^2$, whenever $a(x, \xi)$ is a non-negative classical first-order symbol. Clearly, the estimate is unaffected (only the constant C changes) if we replace a by $a+1$, so we may and shall assume that $a(x, \xi) \geq 1$. Now set $\phi(x, \xi) = (1 + |\xi|^2)^{1/4} a^{1/2}(x, \xi)$ and $\varphi(x, \xi) = (1 + |\xi|^2)^{-1/4} a^{1/2}(x, \xi)$ and observe that inequalities (1)-(4) hold, ^{and} that $a^{1/2}(x, \xi)$ belongs to $S_{\phi, \varphi}^{1/2, 1/2}$. Thus for $T = a^{1/2}(x, D)$ we have $T^*T = a(x, D) +$ a pseudo-differential operator with symbol in $S_{\phi, \varphi}^{0,0}$ by Theorem 1; and we know that the error

term is bounded, by Theorem 2. That is, $a(x, D) = T^* T +$ a bounded error, from which the sharp Gårding inequality is obvious. Note that we could have replaced T by $T' = (a^{1/4}(x, D)) \circ (a^{1/4}(x, D))$ to obtain a positive self-adjoint pseudo-differential operator whose symbol differs from $a(x, D)$ only by a bounded error. A slight generalization of this procedure produces approximate square roots of operators with non-negative symbols in $S_{\phi, \varphi}^{1,1}$ classes; the square roots have symbols in $S_{\Psi, \psi}^{1/2, 1/2}$ for suitable Ψ, ψ .

Next we sketch a new proof of sufficiency of Nirenberg-Treves' condition (P) for local solvability of partial differential equations of principal type. In their paper [4], Nirenberg and Treves reduced the whole problem to proving the following result: Lemma NT: Let $p_t(x, \xi)$ be a first-order classical symbol depending smoothly on the real parameter t ; and suppose that for each fixed (x, ξ) , the function $t \mapsto p_t(x, \xi)$ does not change sign. Then the corresponding operator $p_t(x, D)$ may be written in the form $p_t(x, D) = A_t B + C_t$, where B is a fixed (unbounded) self-adjoint operator, A_t is self-adjoint, bounded and non-negative, and C_t is a bounded error.

Ideally, one should prove the lemma simply by writing the symbol $p_t(x, \xi)$ in the form

(*) $p_t = a_t \cdot b + c_t$, where $a_t \geq 0$ and a_t, b, c_t are classical symbols of order $0, 1, 0$ respectively.

If p_t could be so expressed, the conclusion of Lemma NT would follow instantly from the classical symbolic calculus and sharp Gårding inequality. Nirenberg and Treves carried this out in [6] for the case of real-analytic symbols p_t , but unfortunately p_t cannot in general be written in the form (*) using classical symbols. (The unpublished counterexample is due to Mather.) In our original solution [5] to the sufficiency problem, the authors circumvented direct use of Lemma NT by a complicated argument. Now, however, we can use $S_{\phi, \varphi}^{M, m}$ classes to give a rather simple direct proof of Lemma NT: we simply pick the proper ϕ and φ to enable us to write $p_t(x, \xi)$ in the form (*) with a_t, b, c_t in $S_{\phi, \varphi}^{0,0}$, $S_{\phi, \varphi}^{1,1}$, $S_{\phi, \varphi}^{0,0}$ respectively. Essentially, we can take

$$\varphi(x, \xi) = (1 + |\xi|^2)^{-1/4} + \sup_t \left\{ |D_{\xi} p_t(x, \xi)| + \frac{1}{|\xi|} |D_x p_t(x, \xi)| \right\} + \sup\{\delta > 0 \mid p_t(y, \eta)$$

does not change sign as a function of (t, y, η) in the region

$$|y-x| < \delta, \quad |\xi-\eta| < \delta |\xi| \}, \text{ and } \phi(x, \xi) = (1 + |\xi|^2)^{1/2} \varphi(x, \xi).$$

The construction of a_t, b, c_t is not hard, and can be made without mentioning the elaborate Whitney decompositions of [5], though the latter still lurk in the background.

Detailed proofs of the results mentioned in this note will appear in [6].

*This research was partially supported by NSF grants GP28148 and GP28271.

1. Hörmander, L., "Pseudo-differential operators and hypo-elliptic equations," Amer. Math. Soc. Proc. Symp. in Pure Math., vol. X, 138-183 (1967).
 2. Calderón, A.P., and R. Vaillancourt, "A class of bounded pseudo-differential operators," Proc. Nat. Acad. Sci. U.S.A. ,
 3. Hörmander, L., "Pseudo-differential operators and non-elliptic boundary value problems, Ann. Math. 83, 129-209 (1966).
 4. Nirenberg, L., and F. Treves, " On local solvability of linear partial differential equations, Part II," Comm. Pure Appl. Math. 23, 459-509 (1970).
 5. Beals, R., and C. Fefferman, "On local solvability of linear partial differential equations," Ann. Math.
 6. Beals, R., and C. Fefferman,
-