

SÉMINAIRE ÉQUATIONS AUX DÉRIVÉES PARTIELLES – ÉCOLE POLYTECHNIQUE

J. J. KOHN

Regularity of the $\bar{\partial}$ -Neumann problem

Séminaire Équations aux dérivées partielles (Polytechnique) (1980-1981), exp. n° 19,
p. 1-7

http://www.numdam.org/item?id=SEDP_1980-1981____A21_0

© Séminaire Équations aux dérivées partielles (Polytechnique)
(École Polytechnique), 1980-1981, tous droits réservés.

L'accès aux archives du séminaire Équations aux dérivées partielles (<http://sedp.cedram.org>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>

ÉCOLE POLYTECHNIQUE

CENTRE DE MATHÉMATIQUES

91128 PALAISEAU CEDEX - FRANCE

Tél. (6) 941.82.00 - Poste N°

Télex : ECOLEX 691 596 F

S E M I N A I R E G O U L A O U I C - M E Y E R - S C H W A R T Z 1 9 8 0 - 1 9 8 1

REGULARITY OF THE $\bar{\partial}$ -NEUMANN PROBLEM

by

J.J. KOHN

Let $\Omega \subset \mathbb{C}^n$ be a bounded domain with a smooth boundary $b\Omega$. If z_1, \dots, z_n are coordinates in \mathbb{C}^n we set $x_j = \text{Re}(z_j), y_j = \text{Im}(z_j)$ and, when u is a differentiable function, we define :

$$(1) \quad \begin{aligned} u_{z_j} &= \frac{1}{2} (u_{x_j} - \sqrt{-1} u_{y_j}) \\ u_{\bar{z}_j} &= \frac{1}{2} (u_{x_j} + \sqrt{+1} u_{y_j}) . \end{aligned}$$

Given $\alpha_1, \dots, \alpha_n$ on Ω we will study the system

$$(2) \quad u_{\bar{z}_j} = \alpha_j \text{ for } j = 1, \dots, n.$$

We will assume that the α_j satisfy the compatibility conditions

$$(3) \quad \alpha_j \bar{z}_k = \alpha_k \bar{z}_j .$$

We shall also assume that $b\Omega$ is pseudo-convex, that is if r is a real C^∞ function defined in a neighborhood of $b\Omega$ such that $dr \neq 0$, $r < 0$ in Ω and $r > 0$ outside of $\bar{\Omega}$, if $P \in b\Omega$ and if $(\zeta_1, \dots, \zeta_n) \in \mathbb{C}^n$ satisfy

$$(4) \quad \sum r_{z_i}(P) \zeta_i = 0$$

then we have

$$(5) \quad \sum r_{z_i \bar{z}_j}(P) \zeta_i \bar{\zeta}_j \geq 0.$$

Definition : If $u \in L_2(\Omega)$ we define the singular support of u , denoted by $s.s.(u)$, to be the subset of $\bar{\Omega}$ defined as follows. The point $P \notin s.s.(u)$ if and only if there exists a neighborhood U of P such that the restriction of u to $U \cap \bar{\Omega}$ is in $C^\infty(U \cap \bar{\Omega})$.

This lecture will be concerned with the following two problems.

Problem I. : Given $\alpha_j \in L_2(\Omega)$, $j = 1, \dots, n$ satisfying (3) does

there exist a $u \in L_2(\Omega)$ such that

$$(5) \quad \text{s.s.}(u) \subset \bigcup_j \text{s.s.}(\alpha_j) \quad ?$$

Problem II. : Let $\mathcal{H}(\Omega)$ denote the subspace of $L_2(\Omega)$ consisting of holomorphic functions. Let u be the unique solution of (2) such that $u \perp \mathcal{H}(\Omega)$. Does u satisfy (5) ?

Observe that the system (2) is elliptic thus $\Omega \cap \text{s.s.}(u)$ is contained in the singular support of the α_j . Hence problem I is only interesting for points in $b\Omega$. It is easy to construct a holomorphic function h which is singular at $b\Omega$, then if u satisfies (2) so does $u + h$, so that (5) cannot hold for all solutions of (2). In case the right side of (5) is empty we have the following result.

Theorem : If Ω is pseudo-convex and if $\alpha_j \in C^\infty(\bar{\Omega})$, $j = 1, \dots, n$ satisfying (3), then there exists $u \in C^\infty(\bar{\Omega})$ which satisfies (2).

It is at present unknown whether, under the hypotheses of the above theorem, the solution of (2) which is orthogonal to $\mathcal{H}(\Omega)$ is also in $C^\infty(\bar{\Omega})$. One of the reasons for the solution orthogonal to $\mathcal{H}(\Omega)$ is because the Bergman projection $B : L_2(\Omega) \rightarrow \mathcal{H}(\Omega)$ can be expressed in terms of it. Namely, if $f \in L_2(\Omega)$, if the $f_{z_j} \in L_2(\Omega)$ and if $u \perp \mathcal{H}(\Omega)$ satisfies $u_{z_j} = f_{z_j}$ for $j = 1, \dots, n$ then clearly Bf , the orthogonal projection of f in $\mathcal{H}(\Omega)$, is given by :

$$(6) \quad Bf = f - u.$$

This (5) implies $\text{s.s.}B(f) \subset \text{s.s.}(f)$. By a result of Bell and Ligocka regularity properties of B imply boundary regularity of biholomorphic maps.

It is known that the solution of Problem I depends on the geometry of Ω . The following results of Catlin show this .

Theorem (Catlin) : If Ω is pseudo-convex and if V is a complex-analytic curve contained in $b\Omega$ then there exists α_j satisfying (3) such that (5) is not satisfied for any solution of (2).

Theorem (Catlin) : There exists a pseudo-convex domain $\Omega \subset \mathbb{C}^3$ such that $b\Omega$ does not contain any complex curves and such that there exist α_j as above, so that (5) does not hold.

The $\bar{\partial}$ -Neumann problem enables one to express the solution of II in terms of a boundary value problem. In particular we have the notion of subellipticity of this problem defined as follows.

Definition : The $\bar{\partial}$ -Neumann problem on Ω is subelliptic at $x_0 \in b\Omega$ if there exists a neighborhood U of x_0 and constants $\varepsilon > 0$, $c > 0$ such that for all n-tuples $(\varphi_1, \dots, \varphi_n)$ with $\varphi_j \in C^\infty(U \cap \bar{\Omega})$ and

$$(7) \quad \sum r_{z_j} \varphi_j = 0 \text{ on } b\Omega$$

we have

$$(8) \quad \sum \|\varphi_j\|_\varepsilon^2 \leq c \left(\sum_1^n \|\varphi_{jz_k} - \varphi_{kz_j}\|^2 + \|\sum_1^n \varphi_j z_j\|^2 \right),$$

where $\|\cdot\|_\varepsilon$ denotes the ε -Sobolev-norm.

The quadratic form defined by the left side of (8) is connected with the solution u of II as follows. If $(\alpha_1, \dots, \alpha_n)$ satisfies (3) and $(\varphi_1, \dots, \varphi_n)$ with $\varphi_j \in C^\infty(\bar{\Omega})$ satisfies (7) and

$$(9) \quad \sum_{j,k} (\varphi_{j\bar{z}_k} - \varphi_{k\bar{z}_j}, \psi_{j\bar{z}_k} - \psi_{k\bar{z}_j}) + \left(\sum_j \varphi_j z_j, \sum_k \psi_k \bar{z}_k \right) = \sum (\alpha_j, \psi_j),$$

for all (ψ_1, \dots, ψ_n) with $\psi_j \in C^\infty(\bar{\Omega})$ and $\sum r_{z_j} \psi_j = 0$ on $b\Omega$. Then

$$(10) \quad u = - \sum_j \varphi_j z_j$$

is the unique solution of (2) which is orthogonal to $\mathcal{K}(\Omega)$.

Proposition : If the $\bar{\partial}$ -Neumann is subelliptic at $x_0 \in b\Omega$, if $(\alpha_1, \dots, \alpha_n)$ satisfies (3) and if $\alpha_j \in H_{loc}^s(x_0)$ then $u \in H_{loc}^{s+1}(x_0)$, where $u \perp \mathcal{K}(\Omega)$ and u satisfies (2). In particular we have

$$(11) \quad U \cap \text{s.s.}(u) \subset U \cap \bigcup_j \text{s.s.}(\alpha_j).$$

Hence if the $\bar{\partial}$ -Neumann problem is subelliptic at all points in $b\Omega$ then II is settled affirmatively. From Catlin's result it then follows that subellipticity cannot hold at $x_0 \in b\Omega$ if there is a complex-analytic curve through x_0 contained in $b\Omega$. Catlin also obtained a quantitative measure of the dependence of subellipticity on how close curves through x_0 can be to $b\Omega$.

Theorem (Catlin) : Suppose Ω is pseudo-convex, $x_0 \in b\Omega$ and (8) holds for some $\varepsilon > 0$. Suppose further that there is a complex-analytic curve V with $x_0 \in V$, which has the property that there exists a neighborhood U of x_0 and constants $C > 0$, $\eta > 0$ such that

$$(12) \quad |r(z)| \leq C|z - x_0|^\eta$$

for all $z \in U \cap V$. Then $\varepsilon \leq \frac{1}{\eta}$.

Definition : Suppose $x_0 \in b\Omega$, let $\mathcal{V}(x_0) = \mathcal{V}$ denote the set of germs of complex analytic curves through x_0 . The order of x_0 denoted by $\mathcal{O}(x_0)$ is defined by

$$(13) \quad \mathcal{O}(x_0) = \sup_{V \in \mathcal{V}} \mathcal{O}(x_0, V),$$

where $\mathcal{O}(x_0, V)$, the order of contact of V to $b\Omega$ at x_0 , is defined by

$$(14) \quad \mathcal{O}(x_0, V) = \sup\{\eta \mid \exists \text{ a neighborhood } U \text{ of } x_0 \text{ and } C > 0 \text{ so that (12) holds for all } z \in U \cap V\}.$$

Denoting by \mathcal{W} the set of germs of non-singular complex-analytic curves at x_0 we define $\text{reg } \mathcal{O}(x_0)$ the regular order of x_0 by

$$(15) \quad \text{reg } \mathcal{O}(x_0, V) = \sup_{V \in \mathcal{W}} \mathcal{O}(x_0, V).$$

It is clear that for $n = 1$ we have $\mathcal{O}(x_0) = \text{reg } \mathcal{O}(x_0) = 1$ and that if $n > 1$ then $\mathcal{O}(x_0) \geq \text{reg } \mathcal{O}(x_0) \geq 2$. Furthermore, it is easy to show that for $n = 2$ and Ω pseudo-convex we have $\mathcal{O}(x_0) = \text{reg } \mathcal{O}(x_0)$. However for $\Omega \subset \mathbb{C}^3$, we have the following example (studied by Bloom and Graham), let Ω be given by

$$(16) \quad r(z) = \text{Re}(z_3) + |z_1^2 - z_2^3|^2 + \exp(-1/|z|^2)$$

then for $x_0 \neq 0$ we have $\mathcal{O}(x_0) = \text{reg } \mathcal{O}(x_0) = 2$ and when $x_0 = 0$, we have $\text{reg } \mathcal{O}(0) = 6$ and $\mathcal{O}(0) = \infty$.

D'Angelo has shown that for pseudo-convex domains if $\text{reg } \mathcal{O}(x_0) \leq 4$ then $\mathcal{O}(x_0) = \text{reg } \mathcal{O}(x_0)$.

Returning to subellipticity for pseudo-convex domains the convex of Catlin's theorem holds if $n \leq 2$ and if $\mathcal{O}(x_0) = 2$, that is in those cases when $\mathcal{O}(x_0) < \infty$, then subellipticity holds at x_0 for all $\varepsilon \leq \frac{1}{\mathcal{O}(x_0)}$. A recent example of d'Angelo shows that this is not true in general. Let $\Omega \subset \mathbb{C}^n$ for $n \geq 3$ defined by :

$$(17) \quad r(z) = \text{Re}(z_n) + \sum_{j=1}^{n-2} |z_j^m - z_n z_{j+1}|^2 + |z_{n-1}^m|^2,$$

for this domain $\mathcal{O}(x_0) = 2m$ when $x_0 = (0, \dots, 0)$, $\mathcal{O}(x_0) = 2$ when $z_{n-1}(x_0) \neq 0$ and $\mathcal{O}(x_0) = (2m)^n$ when $x_0 = (0, \dots, 0, i\delta)$ whenever $\delta \in \mathbb{R} - \{0\}$. Catlin shows that subellipticity at $x_0 = (0, \dots, 0)$ holds for $\varepsilon \leq (2m)^{-n}$ but does not hold for $\varepsilon > (2m)^{-n}$. Using the phenomena involved in d'Angelo's example Catlin has constructed for each ε_0 , with $0 < \varepsilon_0 \leq \frac{1}{4}$, a domain such that subellipticity holds with $\varepsilon = \varepsilon_0$ but does not hold for $\varepsilon > \varepsilon_0$. Furthermore he constructs a domain such that subellipticity holds for all $\varepsilon < \varepsilon_0$ but does not hold for $\varepsilon = \varepsilon_0$.

The major problem now is whether $\mathcal{O}(x_0) < \infty$ implies subellipticity on pseudo-convex domain. Here we will briefly describe the sufficient conditions that are known.

Definition : If $x_0 \in \text{b } \Omega$ we define a succession of ideals $I_1(x_0) \subset I_2(x_0) \subset \dots \subset I_k(x_0) \subset C^\infty(x_0)$, where $C^\infty(x_0)$ denotes the ring of germs of C^∞ functions at x_0 . We define by

$$(18) \quad \lambda = \det \begin{pmatrix} 0 & r_{z_1} & \dots & r_{z_n} \\ r_{\bar{z}_1} & r_{z_1 \bar{z}_1} & \dots & r_{z_n \bar{z}_1} \\ \vdots & \vdots & & \vdots \\ r_{\bar{z}_n} & r_{z_1 \bar{z}_n} & \dots & r_{z_n \bar{z}_n} \end{pmatrix}$$

Let $I_1(x_0) = \sqrt{\mathbb{R}(\lambda)}$, where (λ) denotes the ideal generated by λ and $\sqrt{\mathbb{R}}$ is the real radical defined as follows. If $J \subset C^\infty(x_0)$ is an ideal then $\sqrt{\mathbb{R} J} = \{f \in C^\infty(x_0) \mid \exists m \text{ and } g \in J \text{ with } |f|^m \leq |g|\}$. To define $I_k(x_0)$ consider for any $f^{(1)}, \dots, f^{(n+1)} \in I_{k-1}(x_0)$ the $(n+1) \times 2(n+1)$ matrix

$$(19) \quad \begin{pmatrix} 0 & r_{z_1} & \dots & r_{z_n} \\ r_{\bar{z}_1} & r_{z_1 \bar{z}_1} & \dots & r_{z_n \bar{z}_1} \\ \vdots & \vdots & & \\ r_{\bar{z}_n} & r_{z_1 \bar{z}_n} & \dots & r_{z_n \bar{z}_n} \\ 0 & f_{z_1}^{(1)} & \dots & f_{z_n}^{(1)} \\ 0 & f_{z_1}^{(n+1)} & \dots & f_{z_n}^{(n+1)} \end{pmatrix}$$

Let $D^k(f^{(1)}, \dots, f^{(n+1)})$ denote the set of all determinants of $(n+1) \times (n+1)$ minors of the above matrix. Then we define $I_k(x_0)$ by

$$(20) \quad I_k(x_0) = \sqrt{\mathbb{R}(I_{k-1}(x_0), \cup D^k(f^{(1)}, \dots, f^{(n+1)}))},$$

where the union is taken over all $(n+1)$ -tuples in $I_{k-1}(x_0)$.

Theorem : If Ω is pseudo-convex and if $1 \in I_k(x_0)$ the subellipticity holds at x_0 .

In case r is real-analytic in a neighborhood of x_0 it can be shown (using a result of Diederich and Fornæss) that for pseudo-convex domain the condition $1 \in I_k(x_0)$ for some k is equivalent to the non-existence of complex-analytic curves through x_0 which lie in $b\Omega$.

Recent result of d'Angelo give a new analysis of the notion of order of x_0 and in many cases give sharp bounds for the $\mathcal{O}(x)$ in terms of $\mathcal{O}(x_0)$ when x is near x_0 .

References : The recent results of Catlin and d'Angelo are not yet published. For the other references and a more detailed survey see : "Several complex variables from the point of view of linear partial differential equations" by J.J. Kohn in Proc. Conf. on PDE and Diff. Geom. 1980, Beijing. Acad. Sinica.

