SÉMINAIRE DE PROBABILITÉS (STRASBOURG)

KAI LAI CHUNG

A simple proof of Doob's convergence theorem

Séminaire de probabilités (Strasbourg), tome 5 (1971), p. 76

http://www.numdam.org/item?id=SPS_1971__5__76_0

© Springer-Verlag, Berlin Heidelberg New York, 1971, tous droits réservés.

L'accès aux archives du séminaire de probabilités (Strasbourg) (<http://portail.mathdoc.fr/SemProba/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>

A SIMPLE PROOF OF DOOB'S CONVERGENCE THEOREM

by K.L. CHUNG

Doob's version of the fundamental convergence theorem of potential theory asserts that if (f_n) is a decreasing sequence of excessive function and f is the supermedian function $\inf_n f_n$, then the set where f differs from $\hat{f}$ (its regularized function) is semi-polar. Many beautiful proofs of this result are available in the literature. Here is a trivial one.

By truncation we can assume f_1 is bounded. Set $A = \{f \geq \hat{f} + \varepsilon\}$, $\varepsilon > 0$. Since f is fine u.s.c., $\hat{f}$ fine continuous, A is finely closed. Calling P_A the usual réduite operator, we have

$$f \geq P_A f \geq P_A(\hat{f} + \varepsilon) = P_A \hat{f} + P_A \varepsilon$$

(the first $\geq$ follows from $f_n \geq P_A f_n$, the second $\geq$ from the fact that the measures $P_A(x, \cdot)$ are carried by A). Applying the semi-group

$$P_t f \geq P_t P_A \hat{f} + P_t P_A \varepsilon$$

As $t \rightarrow 0$, we get

$$\hat{f} \geq P_A \hat{f} + P_A \varepsilon$$

At a point x regular for A , this implies the absurd inequality $\hat{f}(x) \geq \hat{f}(x) + \varepsilon$. Thus A has no regular point, and $\{f > \hat{f}\} = \bigcup \{f \geq \hat{f} + \frac{1}{n}\}$ is semi-polar.

The same proof shows that if f is nearly Borel, positive and such that $f \geq P_K f$ for every compact set K (it is well known that f then is supermedian), then $\{f > \hat{f}\}$ is semi-polar. One must just take care to apply the above reasoning, not to A (which isn't known to be finely closed) but to $K \subset A$, compact. If x is regular for A one chooses an increasing sequence K_n of compact subsets of A such that $T_{K_n} \downarrow 0$ P^x -a.s. and gets the same contradiction as above.

As a matter of fact the latter version, given very recently by Gettoor and Rao, is the one that I stumbled into. On going over the above proof with Sieveking he pointed out the even shorter cut to the original Doob version.