SÉMINAIRE DE PROBABILITÉS (STRASBOURG)

JIA-AN YAN

A comparison theorem for semimartingales and its applications

Séminaire de probabilités (Strasbourg), tome 20 (1986), p. 349-351

http://www.numdam.org/item?id=SPS_1986__20__349_0

© Springer-Verlag, Berlin Heidelberg New York, 1986, tous droits réservés.

L'accès aux archives du séminaire de probabilités (Strasbourg) (<http://portail.mathdoc.fr/SemProba/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>

A COMPARISON THEOREM FOR SEMIMARTINGALES

AND ITS APPLICATIONS

by YAN Jia-an

We work on a filtered probability space $(\Omega, \mathbb{F}, P; (\mathbb{F}_t))$ satisfying the usual conditions. Let X be a semimartingale such that $\sum_{0 \leq s \leq t} |\Delta X_s| < \infty$ for $t < \infty$ (as usual, we allow an evanescent exceptional set in our inequalities without mentioning it): this is the class of semimartingales for which Yor (Astérisque 52-53, Temps Locaux, p.23-35) has shown the existence of local times $L_t^a(X)$ continuous in t , and cadlag in a . On the other hand, X has a unique decomposition

$$X = X_0 + M + A$$

where M is a continuous local martingale, and A is of finite variation. We denote by A^c the continuous part of A .

LEMMA 1. Assume the following conditions

$$(i) L^0(X) = 0 \quad (ii) \int_0^\cdot I_{\{X_{s-} > 0\}} dA_s^c \leq 0 \quad (iii) \Delta X \leq 0.$$

Then we have $X \leq 0$ on the set $\{X_0 \leq 0\}$.

Proof. We have from Tanaka-Meyer's formula

$$X_t^+ = X_0^+ + \sum_{0 \leq s \leq t} I_{\{X_{s-} \leq 0\}} X_s^+ + \frac{1}{2} L_t^0 + \sum_{0 \leq s \leq t} I_{\{X_{s-} > 0\}} X_s^- + \int_0^t I_{\{X_{s-} > 0\}} dX_s$$

On $\{X_0 \leq 0\}$ the first term vanishes. The second one vanishes because of (iii) and the third one because of (i). Therefore on $\{X_0 \leq 0\}$

$$X_t^+ = \sum_{0 \leq s \leq t} I_{\{X_{s-} > 0\}} (X_s^- + \Delta X_s) + \int_0^t I_{\{X_{s-} > 0\}} (dM_s + dA_s^c)$$

We have $X^- + \Delta X = X^+ - X_- \leq 0$ on $\{X_- > 0\}$ by (iii) and $\int_0^t I_{\{X_{s-} > 0\}} dA_s^c \leq 0$ by (ii). Therefore

$$\int_0^t I_{\{X_{s-} > 0\}} dM_s \geq 0 \text{ on } \{X_0 \leq 0\}$$

Since this is a continuous local martingale starting from 0, it must be equal to 0, and from this we deduce $X_t^+ \leq 0$, and finally $X \leq 0$.

We apply this lemma to a generalization of the comparison lemma given by Ikeda-Watanabe ([1], p.352). One might extract from the proof a slightly more general version of lemma 1, but we shall not give it explicitly.

THEOREM 1. Let X^1, X^2 be solutions of two stochastic differentials equations

$$X_t^i = X_0^i + \int_0^t \sigma(s, X_{s-}^i) dM_s + \int_0^t b^i(s, X_{s-}^i) dB_s + \int_0^t c^i(s, X_{s-}^i) dC_s \quad (i=1,2)$$

where M is a continuous local martingale, B is a continuous increasing process and C an increasing process (B and C adapted). We assume

- $\sigma(s, x)$ is Borel measurable, $|\sigma(s, x) - \sigma(s, y)| \leq \rho(|x - y|)$, where ρ is an increasing function on $\mathbb{R}_+$ such that $\int_0^{+\infty} \rho^{-2}(u) du = +\infty$.
- $b^i(s, x), c^i(s, x)$ are continuous on $\mathbb{R}_+ \times \mathbb{R}$ given the product of the right topology on $\mathbb{R}_+$ and the ordinary topology on $\mathbb{R}$.
- $b^1(s, x) < b^2(s, x)$ and $c^1(s, x) < c^2(s, x)$
- $x < y \Rightarrow c^1(s, x) \leq c^2(s, y)$.

Then we have $X^1 \leq X^2$ on the set $\{X_0^1 \leq X_0^2\}$.

Proof. We may assume $X_0^1 \leq X_0^2$ everywhere. Consider the stopping time

$$T = \inf\{t > 0 : X_t^1 - X_t^2 > 0\}$$

We assume $P\{T < \infty\} > 0$ and derive a contradiction. First of all, we have $X_{T-}^1 \geq X_{T-}^2$ (on $\{T < \infty\}$), and $X_{T-}^1 \leq X_{T-}^2$ on $\{0 < T < \infty\}$. We cannot have $X_{T-}^1 < X_{T-}^2$ on $\{0 < T < \infty\}$, because $\Delta X_{T-}^1 \leq \Delta X_{T-}^2$ (last hypothesis) would then imply $X_{T-}^1 < X_{T-}^2$. Therefore $X_{T-}^1 = X_{T-}^2$ on $\{0 < T < \infty\}$. On $T=0$, we have by convention $X_{T-}^1 = X_{T-}^2$, and it is clear that $X_{T-}^1 = X_{T-}^2$ on this set.

Let X be the semimartingale $(X^1 - X^2)_{T+t}$ on $\{T < \infty\}$, relative to the family $(\mathbb{F}_{T+t})$. From the above, we have $X_0 = 0$. X belongs to the class of semimartingales considered at the beginning, and we set $X = M + A$ as before. There is an interval $[0, U(\omega)[$ on which $\Delta X \leq 0$, $\int_0^t \mathbb{I}_{\{X_{s-} > 0\}} dA_s^C \leq 0$, due to the third hypothesis, and the right continuity of $b^i(T+s, X_{T+s-}^i)$, $c^i(T+s, X_{T+s-}^i)$. Finally, the first hypothesis will imply, exactly as in LeGall's paper [2], that $L^0(X) = 0$ (this is the key point of the proof).

Then we apply lemma 1, not to X , but to X stopped at U -, where

$$U = \inf\{t > 0 : \Delta X_t > 0 \text{ or } \int_0^t \mathbb{I}_{\{X_{s-} > 0\}} dA_s^C > 0\}$$

which is a.s. $> T$ due to the above: we deduce that $X \leq 0$ on $[0, U[$, which contradicts the definition of T .

REMARKS. 1) The first hypothesis can be weakened as

- $\sigma(s, x)$ is Borel measurable, and for any x there is a $\delta(x) > 0$ such that $|\sigma(s, x) - \sigma(s, y)| \leq \rho(|x - y|)$ for $y \in [x - \delta(x), x + \delta(x)]$.

In fact, if we set $V = \inf\{t > 0 : |\sigma(t, X_{t-}^1) - \sigma(t, X_{t-}^2)| > \rho(|X_{t-}^1 - X_{t-}^2|)\}$ $L^0(X^{V-}) = 0$ and we may apply lemma 1 to $X^{(U \wedge V)-}$.

2) As we mentioned, the key point of the proof is to check $L^0(X)=0$, and we deduced this from our first hypothesis as in [2]. Similar conditions ensuring that $L^0(X)=0$ (see [2], Corollaire 1.2) will lead to the same conclusion $X^1 \leq X^2$.

Similarly, we can prove the following theorem.

THEOREM 2. Let X^i be solutions of the following stochastic differential equations

$$\begin{aligned} X_t^i = X_0^i + \int_0^t \sigma(s, X_{s-}^i) dW_s + \int_0^t b^i(s, X_{s-}^i) ds + \int_0^t \int_{U_0} f(s, X_{s-}^i, u) \hat{N}_p(ds, du) \\ + \int_0^t \int_{U \setminus U_0} g^i(s, X_{s-}^i, u) N_p(ds, du) . \end{aligned}$$

Here (W_t) is a Wiener process, N_p is the counting measure of a quasi-left continuous point process p on a standard measurable space U , $U_0 \subset U$ is a measurable subset such that $E[N_p(t, U \setminus U_0)] < \infty$ for t finite, and $\hat{N}_p = N_p - \tilde{N}_p$ ($\sim$ denoting compensation as usual).

We may assert that $X^1 \leq X^2$ on $\{X_0^1 \leq X_0^2\}$ if the following hypotheses are satisfied :

- σ and b^i are as in the preceding theorem.
- f^i, g^i are measurable functions on $\mathbb{R}_+ \times \mathbb{R} \times U$, and for any fixed $u \in U$, $f^i(s, x, u)$ and $g^i(s, x, u)$ are continuous on $\mathbb{R}_+ \times \mathbb{R}$ in the same topology as in theorem 1.
- $(x \leq y) \Rightarrow (f^1(s, x, u) \leq f^2(s, y, u) \text{ and } g^1(s, x, u) \leq g^2(s, y, u))$.

REFERENCES.

- [1]. IKEDA (N.) and WATANABE (S.). Stochastic differential equations and diffusion processes. North Holland, Kodansha, 1981.
- [2]. LE GALL (J.F.). Applications du temps local aux équations différentielles stochastiques unidimensionnelles. Sémin. Prob. XVII, Lect. Notes in M. 986, p. 15-31. Springer-Verlag 1982.

Institute of Applied Mathematics
Academia Sinica
Beijing, China.