

SÉMINAIRE DE PROBABILITÉS (STRASBOURG)

KALYANAPURAM RANGACHARI PARTHASARATHY **A generalized Biane process**

Séminaire de probabilités (Strasbourg), tome 24 (1990), p. 345-348

http://www.numdam.org/item?id=SPS_1990__24__345_0

© Springer-Verlag, Berlin Heidelberg New York, 1990, tous droits réservés.

L'accès aux archives du séminaire de probabilités (Strasbourg) (<http://portail.mathdoc.fr/SemProba/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

A Generalised Biane Process

K. R. Parthasarathy

Indian Statistical Institute, 7 S. J. S. Sansanwal Marg, New Delhi 110 016

Using the methods of quantum probability in a toy Fock space as outlined in [2] and the theory of finite dimensional representations of the 3-dimensional simple Lie algebra $sl(2, \mathbb{C})$ Ph. Biane [1] constructed a quantum Markov chain in discrete time and derived a classical Markov chain which is the discrete time quantum analogue of a classical Bessel process. Here exploiting the Peter-Weyl theory of representations of compact groups we extend Biane's construction to an arbitrary compact group G and derive a classical Markov chain whose state space is the space $\Gamma(G)$ of all characters of irreducible representations of G .

Let G be a compact second countable topological group and let $g \rightarrow L_g$ denote its left regular representation in the complex Hilbert space $L_2(G)$ of all square integrable functions on G with respect to its normalised Haar measure. Let $\mathcal{W}(G)$ denote the W^* algebra generated by the family $\{L_g, g \in G\}$ and let $\mathcal{Z}(G)$ be its centre. Denote by $\Gamma(G)$ the countable set of all characters of irreducible unitary representations of G . For any χ let U^χ be an irreducible unitary representation of G with character χ and dimension $d(\chi)$. If $\chi_1, \chi_2 \in \Gamma(G)$ the tensor product $U^{\chi_1} \otimes U^{\chi_2}$ decomposes into a direct sum of irreducible representations. We shall denote by $m(\chi_1, \chi_2; \chi)$ the multiplicity with which the type U^χ appears in such a decomposition of $U^{\chi_1} \otimes U^{\chi_2}$. Define

$$p_{\chi_1, \chi_2}^\chi = \frac{m(\chi, \chi_1; \chi_2) d(\chi_2)}{d(\chi) d(\chi_1)}. \quad (1)$$

Then

$$\sum_{\chi_2 \in \Gamma(G)} p_{\chi_1, \chi_2}^\chi = 1 \quad \text{for each } \chi, \chi_1 \in \Gamma(G).$$

In other words, for every fixed $\chi \in \Gamma(G)$ the matrix $P^\chi = ((p_{\chi_1, \chi_2}^\chi))$ is a stochastic matrix over the state space $\Gamma(G)$. In each row of P^χ all but a finite number of entries are 0 and each entry is rational. Inspired by Biane's construction in [1] we shall now combine the Peter-Weyl theorem and the methods of quantum probability in order to realise explicitly a Markov chain with transition probability matrix P^χ in the state space $\Gamma(G)$.

As a special case consider $G = SU_2$. Let χ_n denote the character of the "unique" irreducible unitary representation of G of dimension n . By (1) and

Clebsch-Gordon formula

$$p_{\chi_i, \chi_j}^{\chi_2} = \begin{cases} \frac{i-1}{2i} & \text{if } j = i-1, \\ \frac{i+1}{2i} & \text{if } j = i+1, \\ 0 & \text{otherwise.} \end{cases}$$

This is the case covered by Biane [1].

To realise our goal we recall that by Peter-Weyl theorem $L_2(G)$ admits the Plancherel decomposition:

$$L_2(G) = \oplus_{\chi \in \Gamma(G)} \mathcal{H}_\chi$$

where $\dim \mathcal{H}_\chi = d(\chi)^2$, L_g leaves each \mathcal{H}_χ invariant and $L_g|_{\mathcal{H}_\chi}$ is a direct sum of $d(\chi)$ copies of the representation U^χ . If π_χ denotes the orthogonal projection onto the component \mathcal{H}_χ then

$$\pi_\chi = d(\chi)^{-1} \int_G \chi(g) L_g dg \tag{2}$$

thanks to Schur orthogonality relations. Furthermore the abelian W^* algebra $\mathcal{Z}(G)$ is generated by the family $\{\pi_\chi, \chi \in \Gamma(G)\}$.

Fix $\chi_0 \in \Gamma(G)$. Let U^{χ_0} act in the Hilbert space \mathcal{H} . Denote by ρ the density matrix $d(\chi_0)^{-1} I$ in \mathcal{H} . Fix a positive integer N and consider the Hilbert space $\mathcal{H}^{\otimes N} = \mathcal{H} \otimes \dots \otimes \mathcal{H}$ where the tensor product is taken N -fold. Denote by $\mathcal{B}_{[n]}$ the W^* algebra generated by all operators of the form $X_1 \otimes \dots \otimes X_n \otimes I \otimes \dots \otimes I$ where X_i are bounded operators in \mathcal{H} . Then $\mathcal{B}_{[1]} \subset \mathcal{B}_{[2]} \subset \dots \subset \mathcal{B}_{[N]} = \mathcal{B}$ yields a finite filtration in \mathcal{B} with conditional expectation maps $E_{[n]} : \mathcal{B} \rightarrow \mathcal{B}_{[n]}$, $1 \leq n \leq N$, defined by

$$E_{[n]} X_1 \otimes \dots \otimes X_N = \left(\prod_{i=n+1}^N \text{tr } \rho X_i \right) X_1 \otimes \dots \otimes X_n \otimes I \otimes \dots \otimes I$$

for all $X_i \in \mathcal{B}(\mathcal{H})$ and linear extension. Thanks to Peter-Weyl theorem there exists a unique identity preserving and continuous $*$ homomorphism $j_n : \mathcal{W}(G) \rightarrow \mathcal{B}_{[n]}$ satisfying

$$j_n(L_g) = U_g^{\chi_0} \otimes \dots \otimes U_g^{\chi_0} \otimes I \otimes \dots \otimes I \text{ for all } g \in G \tag{3}$$

where $U_g^{\chi_0}$ appears n -fold and I , $(N - n)$ -fold. Then

$$E_{[n-1]} j_n(L_g) = j_{n-1}(d(\chi_0)^{-1} \chi_0(g) L_g) \text{ for all } g \in G. \tag{4}$$

Thus there exists a completely positive map $T : \mathcal{W}(G) \rightarrow \mathcal{W}(G)$ satisfying

$$E_{[n-1]} j_n(X) = j_{n-1}(T(X)), \quad X \in \mathcal{W}(G), \tag{5}$$

$$T(L_g) = d(\chi_0)^{-1} \chi_0(g) L_g \text{ for all } g \in G. \tag{6}$$

From (2) and Schur orthogonality relations it now follows that

$$\begin{aligned}
 T(\pi_\chi) &= [d(\chi_0)d(\chi)]^{-1} \int \chi_0(g)\chi(g)L_g dg \\
 &= \sum_{\chi' \in \Gamma(G)} [d(\chi_0)d(\chi)]^{-1} d(\chi') m(\chi_0, \chi; \chi') \pi_{\chi'} \\
 &= \sum_{\chi' \in \Gamma(G)} p_{\chi, \chi'}^{\chi_0} \pi_{\chi'} .
 \end{aligned}
 \tag{7}$$

We now establish the following lemma.

Lemma 1. For any $m < n$, $Z \in \mathcal{Z}(G)$, $X \in \mathcal{W}(G)$

$$[j_m(Z), j_n(X)] = 0 . \tag{8}$$

In particular, the family $\{j_m(Z), m = 1, 2, \dots, N, Z \in \mathcal{Z}(G)\}$ is commutative.

Proof. We have for any $g, h \in G$,

$$\begin{aligned}
 [j_m(L_g), j_n(L_h)] &= \underbrace{[U_g^{\chi_0} \otimes \dots \otimes U_g^{\chi_0} \otimes I \otimes \dots \otimes I]}_{m\text{-fold}}, \underbrace{[U_h^{\chi_0} \otimes \dots \otimes U_h^{\chi_0} \otimes I \otimes \dots \otimes I]}_{n\text{-fold}} \\
 &= j_m([L_g, L_h]) \underbrace{I \otimes \dots \otimes I}_{m\text{-fold}} \underbrace{U_h^{\chi_0} \otimes \dots \otimes U_h^{\chi_0}}_{(n-m)\text{-fold}} \underbrace{\otimes I \otimes \dots \otimes I}_{(N-n)\text{-fold}} .
 \end{aligned}$$

Since Z can be approximated by linear combinations of $L_g, g \in G$, it follows that $[j_m(Z), j_n(L_h)] = 0$. Since X can be approximated by linear combinations of L_h we have (8). The second part is immediate.

From (3), (5) - (7) and Lemma 1 we have the following theorem.

Theorem 2. For any fixed $\chi_0 \in \Gamma(G)$ let the W^* homomorphisms $j_n : \mathcal{W}(G) \rightarrow \mathcal{B}$, $1 \leq n \leq N$, be defined by (3). In the state $\rho^{\otimes N}$, where $\rho = d(\chi_0)^{-1}I$, the sequence $\{j_n, 1 \leq n \leq N\}$ is a quantum Markov chain in the sense of Accardi-Frigerio-Lewis with transition operator $T : \mathcal{W}(G) \rightarrow \mathcal{W}(G)$ satisfying $T(L_g) = d(\chi_0)^{-1}\chi_0(g)L_g$ for all $g \in G$. T leaves the centre $\mathcal{Z}(G)$ of $\mathcal{W}(G)$ invariant. The family $\{j_n(Z), Z \in \mathcal{Z}(G), 1 \leq n \leq N\}$ is commutative. In the state $\rho^{\otimes N}$ the sequence $\{j_n|_{\mathcal{Z}(G)}, 1 \leq n \leq N\}$ is a classical Markov chain with state space $\Gamma(G)$ and transition probability matrix $P^{\chi_0} = ((p_{\chi, \chi'}^{\chi_0}))$, $\chi, \chi' \in \Gamma(G)$, defined by (1).

Remark. We can replace the W^* algebra $\mathcal{W}(G)$ by the $*$ unital algebra $\mathcal{U}(G)$ of left invariant differential operators on G if G is a compact connected Lie group. In such a case $\mathcal{Z}(G)$ can be replaced by the centre $z(G)$ of $\mathcal{U}(G)$. If we choose this infinitesimal description, put $G = SU_2$ and choose χ_0 as the character of the unique 2-dimensional irreducible unitary representation of G , then we obtain Biane's example in [1].

References

- [1] Ph. Biane: Marches de Bernoulli quantiques, Université de Paris VII, preprint, 1989.
- [2] P. A. Meyer: Eléments de probabilités quantiques, Séminaire de Probabilités XX, LNM Springer 1204, pp. 186–312, 1986.