SÉMINAIRE DE PROBABILITÉS (STRASBOURG)

JEAN-CLAUDE GRUET

ZHAN SHI

On the Spitzer and Chung laws of the iterated logarithm for brownian motion

Séminaire de probabilités (Strasbourg), tome 29 (1995), p. 237-247

<http://www.numdam.org/item?id=SPS_1995__29__237_0>

© Springer-Verlag, Berlin Heidelberg New York, 1995, tous droits réservés.

L'accès aux archives du séminaire de probabilités (Strasbourg) (<http://portail.mathdoc.fr/SemProba/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

ON THE SPITZER AND CHUNG LAWS OF THE ITERATED LOGARITHM FOR BROWNIAN MOTION

by

J.-C. Gruet⁽¹⁾ and Z. Shi⁽²⁾

(1) *Laboratoire de Probabilités, Université Paris VI, 4 Place Jussieu, 75252 Paris Cedex 05, France & Université de Reims, UFR Sciences, 51062 Reims Cedex, France*

(2) *L.S.T.A., Université Paris VI, 4 Place Jussieu, 75252 Paris Cedex 05, France*

1. Introduction.

Let $\{W(t); t \geq 0\}$ be a two-dimensional Brownian motion. It is well-known that the Brownian path is almost surely dense in the plane, but never hits a given point at positive time. A natural question is thus to study the rate with which the small values of $\|W - x\|$ (the symbol “ $\|\cdot\|$ ” denoting the usual Euclidean modulus) approach 0 for any $x \in \mathbb{R}^2$. Without loss of generality, we assume $W(0) = (1, 0)$ and $x = (0, 0)$. Let

$$X(t) = \inf_{0 \leq s \leq t} \|W(s)\|, \quad t > 0.$$

The following celebrated Spitzer (1958) integral test characterizes the lower functions of X :

Theorem A (Spitzer 1958). *For any non-decreasing function $f > 1$,*

$$\mathbb{P}\left[X(t) < \frac{t^{1/2}}{f(t)}, \text{ i.o.}\right] = \begin{cases} 0 \\ 1 \end{cases} \iff \int \frac{dt}{t \log f(t)} \begin{cases} < \infty \\ = \infty \end{cases}.$$

Here and in the sequel, “i.o.” stands for “infinitely often” as t tends to infinity.

Spitzer's Theorem A answers the how-small-are-the-small-values-of- $\|W\|$ question. We propose to study the corresponding "how big" problem for the small values. Our Theorem 1, stated as follows, provides a characterization of the upper functions of X .

Theorem 1. *If $g > 1$ is non-decreasing, then*

$$\mathbb{P}\left[X(t) > \exp\left(-\frac{\log t}{g(t)}\right), \text{ i.o.}\right] = \begin{cases} 0 \\ 1 \end{cases} \iff \int_1^\infty \frac{dt}{t g(t) \log t} \begin{cases} < \infty \\ = \infty \end{cases}.$$

Theorems A and 1 together give an accurate description of the almost sure asymptotic behaviours of X . For example, it is immediately seen from the above theorems that

$$\mathbb{P}\left[X(t) > \exp\left(-\frac{\log t}{(\log \log t)^a}\right), \text{ i.o.}\right] = \begin{cases} 0 & \text{if } a > 1; \\ 1 & \text{otherwise.} \end{cases}$$

$$\mathbb{P}\left[X(t) < \exp\left(-(\log t)(\log \log t)^a\right), \text{ i.o.}\right] = \begin{cases} 0 & \text{if } a > 1; \\ 1 & \text{otherwise.} \end{cases}$$

What about the lower functions of the big values of Brownian motion? Let us recall the classical Chung (1948) integral test for linear Brownian motion.

Theorem B (Chung 1948). *Let B be a real-valued Brownian motion. For every non-decreasing function $h > 0$ such that $t^{-1/2}h(t)$ is non-increasing, we have*

$$\mathbb{P}\left[\sup_{0 \leq s \leq t} |B(s)| < \frac{t^{1/2}}{h(t)}, \text{ i.o.}\right] = \begin{cases} 0 \\ 1 \end{cases} \iff \int_1^\infty \frac{dt}{t} h^2(t) \exp\left(-\frac{\pi^2}{8} h^2(t)\right) \begin{cases} < \infty \\ = \infty \end{cases}.$$

Chung's Theorem B was obtained for linear Brownian motion. The following natural question was raised by Révész (1990, p.195): for the planar Brownian motion W , what can be said on the liminf behaviour of $\sup_{0 \leq s \leq t} \|W(s)\|$?

The same question can be asked for a Brownian motion of any dimension. Let $\{V(t); t \geq 0\}$ denote a d -dimensional Brownian motion, and let

$$Y(t) = \sup_{0 \leq s \leq t} \|V(s)\|, \quad t > 0.$$

Our answer to the problem is the following

Theorem 2. *Let $d \geq 1$ and let $h > 0$ be a non-decreasing function. Then*

$$\mathbb{P}\left[Y(t) < \frac{t^{1/2}}{h(t)}, \text{ i.o.}\right] = \begin{cases} 0 \\ 1 \end{cases} \iff \int_1^\infty dt \frac{h^2(t)}{t} \exp\left(-\frac{j_\nu^2}{2} h^2(t)\right) \begin{cases} < \infty \\ = \infty \end{cases},$$

where j_ν denotes the smallest positive zero of the Bessel function $J_\nu(x)$ of index $\nu \equiv (d-2)/2$.

Remarks. (i) Since $j_{-1/2} = \pi/2$, Theorem B is a special case of the above result.

(ii) An interesting feature in Theorem 2 is that we do not suppose $t^2/h(t)$ to be non-decreasing. Thus the latter condition can be removed from Chung's Theorem B.

(iii) As usual, Theorem 2 has a "local" version for t tending to 0, of which the statement and proof are omitted.

Corollary 1. *We have, for $d \geq 1$,*

$$(1.1) \quad \liminf_{t \rightarrow \infty} \left(\frac{\log \log t}{t} \right)^{1/2} Y(t) = \frac{j_\nu}{2^{1/2}} \quad \text{a.s.},$$

with rate of convergence

$$\liminf_{t \rightarrow \infty} \frac{(\log \log t)^{3/2}}{t^{1/2} \log \log t} \left(Y(t) - \frac{j_\nu}{2^{1/2}} \frac{t^{1/2}}{(\log \log t)^{1/2}} \right) = -\frac{j_\nu}{2^{1/2}}, \quad \text{a.s.}$$

Remark. The LIL (1.1) was previously obtained by Lévy (1953) for $d = 2$ and by Ciesielski & Taylor (1962) for any dimension d .

Theorem 1 is proved in Section 2, and Theorem 2 in Section 3.

2. The proof of Theorem 1.

Let W be as before a Brownian motion in the plane, starting from $(1, 0)$. Define

$$H(x) = \inf\{t > 0 : \|W(t)\| = x\}, \quad 0 < x \leq 1,$$

the first hitting time of $\|W\|$ at level x . Obviously the process $y \mapsto H(1/y)$ (for $y \geq 1$) is increasing, and it has independent increments by using the strong Markov property of $\|W\|$. Since

$$H(2^{-n}) = \sum_{k=1}^n \left(H(2^{-k}) - H(2^{-(k-1)}) \right),$$

using Brownian scaling gives

$$H(2^{-n}) = \sum_{k=1}^n 2^{-2(k-1)} \xi_k,$$

where $(\xi_k)_{k \geq 1}$ is an i.i.d. sequence of random variables having the same law as $H(1/2)$. Consequently,

$$(2.1) \quad 2^{-2(n-1)} \sum_{k=1}^n \xi_k \leq H(2^{-n}) \leq \sum_{k=1}^n \xi_k.$$

Let us first establish a preliminary result for the partial sum of (ξ_k) :

Lemma 1. *Let $\{\Lambda(t); t \geq 0\}$ be a subordinator, and assume that $\Lambda(1)$ has the same law as $H(1/2)$. Then for any function $f > 1$ such that $f(t)/t$ is non-decreasing, we have*

$$\limsup_{t \rightarrow \infty} \frac{\Lambda(t)}{f(t)} = \begin{cases} 0 \\ \infty \end{cases}, \quad \text{a.s.} \quad \Longleftrightarrow \quad \int^{\infty} \frac{dt}{\log f(t)} \begin{cases} < \infty \\ = \infty \end{cases}.$$

Proof of Lemma 1. The Laplace transform of $H(1/2)$ is well-known (see Kent (1978 Theorem 3.1)):

$$\mathbb{E} \exp(-\lambda H(1/2)) = \frac{K_0(\sqrt{2\lambda})}{K_0(\sqrt{\lambda/2})}, \quad \forall \lambda > 0,$$

where K_0 is the modified Bessel function. Recall that $\Lambda(1)$ has the same law as $H(1/2)$. Write

$$\mathbb{E} \exp(-\lambda \Lambda(1)) = \exp(-\Psi(\lambda)).$$

Thus

$$\Psi(\lambda) = \log K_0(\sqrt{\lambda/2}) - \log K_0(\sqrt{2\lambda}).$$

Using elementary asymptotics of K_0 , we immediately arrive at the following estimate

$$\Psi(\lambda) - \lambda\Psi'(\lambda) \sim \frac{2\log 2}{\log(1/\lambda)}, \quad \lambda \rightarrow 0,$$

(the usual symbol “ $a(x) \sim b(x)$ ” ($x \rightarrow x_0$) means $\lim_{x \rightarrow x_0} a(x)/b(x) = 1$). Now the statement of Lemma 1 follows by applying a general result for subordinators (see for example Fristedt (1974 Theorem 6.1)) which tells that $\limsup_{t \rightarrow \infty} \Lambda(t)/f(t) = 0$ or ∞ (almost surely) according as

$$\int^{\infty} \left(\Psi\left(\frac{1}{f(t)}\right) - \frac{1}{f(t)} \Psi'\left(\frac{1}{f(t)}\right) \right) dt$$

converges or diverges. □

Lemma 2. *If $h > 1$ is a non-decreasing function with $\int^{\infty} dt/h(t) = \infty$, then*

$$\int^{\infty} \frac{dt}{t + h(t)} = \infty.$$

Proof of Lemma 2. The proof is briefly sketched, since it involves only elementary computations. Set $\mathcal{A} = \{t : h(t) \leq t\}$ and $\mathcal{B} = \{t : h(t) > t\}$. Obviously, we have

$$\frac{1}{t + h(t)} \geq \frac{1}{2} \left(\frac{1}{t} \mathbf{1}_{\mathcal{A}}(t) + \frac{1}{h(t)} \mathbf{1}_{\mathcal{B}}(t) \right).$$

Assume $\int^{\infty} \mathbf{1}_{\mathcal{A}}(t)(dt/t) < \infty$. We only have to show $\int^{\infty} \mathbf{1}_{\mathcal{B}}(t)(dt/h(t)) = \infty$. Write $F_{\mathcal{A}}(t) \equiv \int_1^t \mathbf{1}_{\mathcal{A}}(s)ds$. Using integration by parts for $\int \mathbf{1}_{\mathcal{A}}(s)(ds/s)$, it is seen that $t \mapsto F_{\mathcal{A}}(t)/t$ is a Cauchy family for $t > 1$. Thus $\lim_{t \rightarrow \infty} F_{\mathcal{A}}(t)/t$ exists. If $\lim_{t \rightarrow \infty} F_{\mathcal{A}}(t)/t > 0$, then $\int_1^t F_{\mathcal{A}}(s)(ds/s^2)$ would diverge, which contradicts the convergence of $\int_1^{\infty} \mathbf{1}_{\mathcal{A}}(s)(ds/s)$ (the latter is obviously greater than $\int_1^t F_{\mathcal{A}}(s)(ds/s^2)$ by integration by parts). Consequently, $\lim_{t \rightarrow \infty} F_{\mathcal{A}}(t)/t = 0$. Thus $\int_1^t \mathbf{1}_{\mathcal{B}}(s)ds \geq t/2$ for sufficiently large t . Again using integration by parts, we obtain

$$\int_1^t \mathbf{1}_{\mathcal{B}}(t) \frac{ds}{h(s)} \geq \frac{1}{2} \int_1^t \frac{ds}{h(s)} + \text{a finite term},$$

which diverges as t tends to infinity. Lemma 2 is proved. □

Proof of Theorem 1. Pick $0 < x < 1$, and let us write $2^{-n} \leq x \leq 2^{-(n-1)}$ (which means $(n-1)\log 2 \leq \log(1/x) \leq n\log 2$). Then $H(2^{-(n-1)}) \leq H(x) \leq H(2^{-n})$. Using (2.1), we have

$$(2.2) \quad x^2 \Lambda\left(\log(1/x)/\log 2 - 1\right) \leq H(x) \leq \Lambda\left(\log(1/x)/\log 2 + 1\right).$$

First, we show the following integral test for H :

$$(2.3) \quad \limsup_{x \rightarrow 0^+} \frac{H(x)}{f(x)} = \begin{cases} 0 \\ \infty \end{cases}, \quad \text{a.s.} \iff \int_{0^+} \frac{dx}{x \log f(x)} \begin{cases} < \infty \\ = \infty \end{cases},$$

for any function $f > 1$ such that $\log f(x)/\log(1/x)$ is non-increasing. Indeed, assume that $\int_{0^+} dx/(x \log f(x))$ converges. Define $\hat{f}(t) = f(2^{-t/2})$. Then $\hat{f}(t)/t$ is non-decreasing, with $\int^\infty dt/\log \hat{f}(t) < \infty$. By Lemma 1, we have $\limsup_{t \rightarrow \infty} \Lambda(t)/\hat{f}(t) = 0$, with probability 1. Thus $\limsup_{t \rightarrow \infty} \Lambda(t)/f(2^{-(t-1)}) = 0$. Using the second part of (2.2), this implies $\limsup_{x \rightarrow 0^+} H(x)/f(x) = 0$. It remains to verify the divergent half of (2.3). Suppose $\int_{0^+} dx/(x \log f(x)) = \infty$. Then $\int^\infty dt/\log f(2^{-2(t+1)})$ diverges as well. According to Lemma 2, this implies

$$\int^\infty \frac{dt}{\log \tilde{f}(t)} = \infty,$$

for $\tilde{f}(t) \equiv 2^{2(t+1)} f(2^{-2(t+1)})$. Applying Lemma 1 gives $\limsup_{t \rightarrow \infty} \Lambda(t)/\tilde{f}(t) = \infty$, which, by means of the first part of (2.2), yields $\limsup_{x \rightarrow 0^+} H(x)/f(x) = \infty$. Hence (2.3) is proved. By noting $[H(x) > t] = [X(t) > x]$ (for any $0 < x < 1$ and $t > 0$), several lines of standard calculation readily confirm that the integral test (2.3) is equivalent to that in Theorem 1. $\square$

3. The proof of Theorem 2.

In this section, V denotes a d -dimensional Brownian motion, which, without loss of generality, is assumed to start from 0. Let $H(x) = \inf\{t > 0 : \|V(t)\| = x\}$ (for $x > 0$). The proof of Theorem 2 is essentially based on the following exact density function of $H(1)$ due to Ciesielski & Taylor (1962):

$$\mathbb{P}[H(1) \in dt]/dt = \frac{1}{2^\nu \Gamma(\nu+1)} \sum_{n=1}^{\infty} \frac{j_{\nu,n}^{\nu+1}}{J_{\nu+1}(j_{\nu,n})} \exp\left(-\frac{j_{\nu,n}^2}{2} t\right), \quad t > 0,$$

where $\nu = (d-2)/2$, and $0 < j_{\nu,1} < j_{\nu,2} < \dots$ are the positive zeros of the Bessel function J_ν (and of course $J_{\nu+1}$ denotes the Bessel function of index $\nu+1$). Let Y be as before the supremum process of $\|V\|$. By Brownian scaling, we have, for any $x > 0$,

$$\begin{aligned}\mathbb{P}[Y(1) < x] &= \mathbb{P}[H(x) > 1] = \mathbb{P}[H(1) > 1/x^2] \\ &= \frac{2^{1-\nu}}{\Gamma(\nu+1)} \sum_{n=1}^{\infty} \frac{1}{j_{\nu,n}^{1-\nu} J_{\nu+1}(j_{\nu,n})} \exp\left(-\frac{j_{\nu,n}^2}{2x^2}\right),\end{aligned}$$

which implies

$$(3.1) \quad \mathbb{P}[Y(1) < x] \sim \frac{2^{1-\nu}}{\Gamma(\nu+1)j_{\nu,1}^{1-\nu}J_{\nu+1}(j_{\nu,1})} \exp\left(-\frac{j_{\nu,1}^2}{2x^2}\right), \quad \text{as } x \rightarrow 0,$$

(recall that $j_\nu \equiv j_{\nu,1}$ is the smallest positive zero of J_ν). Write in the sequel $\rho \equiv j_\nu^2/2$.

Let $h > 0$ be a non-decreasing function. In the rest of the note, generic constants will be denoted by K_i ($1 \leq i \leq 9$).

We begin with the convergent part of Theorem 2, which is an immediate consequence of the tail estimation (3.1). Indeed, pick a sufficiently large initial value t_0 and define the sequence $(t_k)_{k \geq 1}$ by $t_{k+1} = (1 + h^{-2}(t_k))t_k$ for $k = 0, 1, 2, \dots$, and write $h_k \equiv h(t_k)$ for notational convenience. Obviously t_k increases to infinity (as $k \rightarrow \infty$). Assume that $\int^\infty (dt/t)h^2(t)\exp(-\rho h^2(t))$ converges. This implies, by several lines of elementary calculation, the convergence of $\sum_k \exp(-\rho h_k^2)$. From (3.1) and scaling it follows that

$$\begin{aligned}\mathbb{P}\left[Y(t_k) < \frac{t_k^{1/2}}{h_k - 1/h_k}\right] &= \mathbb{P}\left[Y(1) < \frac{1}{h_k - 1/h_k}\right] \\ &\leq K_1 \exp(-\rho(h_k - 1/h_k)^2) \\ &\leq K_2 \exp(-\rho h_k^2),\end{aligned}$$

which sums. According to Borel-Cantelli lemma, (almost surely) for large k , $Y(t_k) \geq t_k^{1/2}/(h_k - 1/h_k)$. Let $t \in [t_k, t_{k+1}]$. Then by our construction of (t_k) ,

$$Y(t) \geq Y(t_k) \geq \frac{t_k^{1/2}}{h_k - 1/h_k} = \frac{t_{k+1}^{1/2}}{(1 + 1/h_k^2)(h_k - 1/h_k)} \geq \frac{t_{k+1}^{1/2}}{h_k} \geq \frac{t^{1/2}}{h(t)},$$

which yields the convergent part of Theorem 2.

It remains to show the divergent part. Let h be such that

$$(3.2) \quad \int^\infty \frac{dt}{t} h^2(t) \exp(-\rho h^2(t)) = \infty.$$

In view of (1.1), we assume without loss of generality that

$$(3.3) \quad \frac{(\log \log t)^{1/2}}{(2\rho)^{1/2}} \leq h(t) \leq \frac{(2 \log \log t)^{1/2}}{\rho^{1/2}}.$$

Define $t_k = \exp(k/\log k)$ (for $k \geq k_0$) and write as before $h_k \equiv h(t_k)$. In what follows, we only deal with the index k tending ultimately to infinity. Therefore our results, sometimes without further mention, are to be understood for sufficiently large k 's. Obviously (3.2) is equivalent to the following

$$(3.4) \quad \sum_k \exp(-\rho h_k^2) = \infty.$$

Using (3.3) gives

$$(3.5) \quad \frac{(\log k)^{1/2}}{(3\rho)^{1/2}} \leq h_k \leq \log k.$$

Fix an $\varepsilon > 0$, then

$$(3.6) \quad k^{-1/2} h_k^2 \leq \frac{\varepsilon}{\rho},$$

(for $k \geq k_0$). Consider the measurable event $A_k = \{Y(t_k) < t_k^{1/2}/h_k\}$. From (3.1) it follows that for $k \geq k_0$,

$$(3.7) \quad \mathbb{P}(A_k) = \mathbb{P}\left[Y(1) < \frac{1}{h_k}\right] \geq (1 - \varepsilon) \frac{2^{1-\nu}}{\Gamma(\nu+1) j_\nu^{1-\nu} J_{\nu+1}(j_\nu)} \exp(-\rho h_k^2),$$

which, by means of (3.4), yields

$$(3.8) \quad \sum_k \mathbb{P}(A_k) = \infty.$$

Let $k < \ell$. Since V has independent and stationary increments, we have

$$\begin{aligned} \mathbb{P}(A_k A_\ell) &= \mathbb{P}\left[\sup_{0 \leq t \leq t_k} \|V(t)\| < \frac{t_k^{1/2}}{h_k}, \sup_{0 \leq t \leq t_\ell} \|V(t)\| < \frac{t_\ell^{1/2}}{h_\ell}\right] \\ &\leq \mathbb{P}(A_k) \sup_{\|x\| \leq t_k^{1/2}/h_k} \mathbb{P}\left[\sup_{0 \leq t \leq t_\ell - t_k} \|V(t) + x\| < \frac{t_\ell^{1/2}}{h_\ell}\right]. \end{aligned}$$

Using a general property of Gaussian measures (see for example Ledoux & Talagrand (1991 p.73)), it follows that

$$\begin{aligned} \mathbb{P}(A_k A_\ell) &\leq \mathbb{P}(A_k) \mathbb{P}\left[\sup_{0 \leq t \leq t_\ell - t_k} \|V(t)\| < \frac{t_\ell^{1/2}}{h_\ell}\right] \\ (3.9) \quad &= \mathbb{P}(A_k) \mathbb{P}\left[Y(1) < \frac{t_\ell^{1/2}}{(t_\ell - t_k)^{1/2} h_\ell}\right]. \end{aligned}$$

For every $n > k_0$, define

$$\mathcal{E}(n) = \{k_0 \leq k < \ell \leq n : \ell - k \leq (\log k)^3\},$$

$$\mathcal{F}(n) = \{k_0 \leq k < \ell \leq n : \ell - k > (\log k)^3\}.$$

It is seen that when $k < \ell < k + (\log k)^3$,

$$\begin{aligned} \frac{\ell}{\log \ell} - \frac{k}{\log k} &= \frac{\ell \log k - k \log \ell}{\log k \log \ell} \\ &= \frac{(\ell - k) \log k - k \log(1 + (\ell - k)/k)}{\log k \log \ell} \\ &\sim \frac{\ell - k}{\log k}, \quad (k \rightarrow \infty), \end{aligned}$$

which implies

$$\frac{t_k}{t_\ell} \leq \exp\left(-\frac{\ell - k}{2 \log k}\right).$$

Let $(k, \ell) \in \mathcal{E}(n)$. From the above estimate it follows that

$$\frac{t_\ell^{1/2}}{(t_\ell - t_k)^{1/2}} \leq \left(1 - \exp\left(-\frac{\ell - k}{2 \log k}\right)\right)^{-1/2} \leq \left[K_3 \min\left((\ell - k)/\log k, 1\right)\right]^{-1/2},$$

which, by means of (3.5), yields

$$\frac{t_\ell^{1/2}}{(t_\ell - t_k)^{1/2} h_\ell} \leq \frac{t_\ell^{1/2}}{(t_\ell - t_k)^{1/2} h_k} \leq \left[K_4 \min(\ell - k, \log k)\right]^{-1/2}.$$

From (3.9) and (3.1) it follows that

$$\mathbb{P}(A_k A_\ell) \leq K_5 \mathbb{P}(A_k) \exp(-K_6(\ell - k)) + K_5 \mathbb{P}(A_k) k^{-K_6}.$$

Obviously,

$$\begin{aligned} \sum_{\ell > k} \exp(-K_6(\ell - k)) &\leq K_7, \\ \sum_{k < \ell < k + (\log k)^3} k^{-K_6} &\leq k^{-K_6} (\log k)^3 \leq K_8. \end{aligned}$$

Therefore,

$$(3.10) \quad \sum_{(k, \ell) \in \mathcal{E}(n)} \mathbb{P}(A_k A_\ell) \leq K_9 \sum_{k=k_0}^n \mathbb{P}(A_k).$$

Now let $(k, \ell) \in \mathcal{F}(n)$. In this case, $\ell - (\log \ell)^2 \geq k + (\log k)^3 - (\log(k + (\log k)^3))^2 > k$, thus $\ell - k > (\log \ell)^2$. Since

$$\frac{\ell}{\log \ell} - \frac{k}{\log k} = \frac{(\ell - k) \log k - k \log(1 + (\ell - k)/k)}{\log k \log \ell} \sim \frac{\ell - k}{\log \ell},$$

we have

$$\frac{t_\ell^{1/2}}{(t_\ell - t_k)^{1/2} h_\ell} \leq \frac{1}{[1 - \exp(-(\ell - k)/2 \log \ell)]^{1/2} h_\ell} \leq \frac{1}{(1 - \ell^{-1/2})^{1/2} h_\ell}.$$

By means of (3.9), (3.1), (3.6) and (3.7), this implies

$$\begin{aligned} \mathbb{P}(A_k A_\ell) &\leq \mathbb{P}(A_k)(1 + \varepsilon) \exp(\rho \ell^{-1/2} h_\ell^2) \frac{2^{1-\nu}}{\Gamma(\nu + 1) j_\nu^{1-\nu} J_{\nu+1}(j_\nu)} \exp(-\rho h_k^2) \\ &\leq (1 + 3\varepsilon) e^\varepsilon \mathbb{P}(A_k) \mathbb{P}(A_\ell). \end{aligned}$$

Combining the above estimate together with (3.10) and (3.8) yields

$$\liminf_{n \rightarrow \infty} \sum_{k=k_0}^n \sum_{\ell=k_0}^n \mathbb{P}(A_k A_\ell) \Big/ \left(\sum_{k=k_0}^n \mathbb{P}(A_k) \right)^2 \leq 1.$$

According to a well-known version of Borel-Cantelli lemma (see for example Révész (1990 p.28)), we have $\mathbb{P}(A_k, \text{i.o.}) = 1$. The proof of the divergent part of Theorem 2 is completed. $\square$

Acknowledgement. We are grateful to Davar Khoshnevisan for helpful comments on the first draft of this work.

REFERENCES

- Chung, K.L. (1948): On the maximum partial sums of sequences of independent random variables. *Trans. Amer. Math. Soc.* 64, 205-233.
- Ciesielski, Z. & Taylor, S.J. (1962). First passage times and sojourn times for Brownian motion in space and the exact Hausdorff measure of the sample path. *Trans. Amer. Math. Soc.* 103 434-450.
- Fristedt, B.E. (1974): Sample functions of stochastic processes with stationary independent increments. In: *Advances in Probability 3* (Eds.: P. Ney & S. Port) pp. 241-396. Dekker, New York.
- Itô, K. & McKean, H.P. (1965): *Diffusion Processes and their Sample Paths*. Springer, Berlin.

- Ledoux, M. & Talagrand, M. (1991): *Probability in Banach Spaces: Isoperimetry and Processes*. Springer, Berlin.
- Lévy, P. (1953): La mesure de Hausdorff de la courbe du mouvement brownien. *Giornale dell'Istituto Italiano degli Attuari* 16 1-37.
- Révész, P. (1990): *Random Walk in Random and Non-Random Environments*. World Scientific, Singapore.
- Revuz, D. & Yor, M. (1994): *Continuous Martingales and Brownian Motion*. 2nd Edition. Springer, Berlin.
- Spitzer, F. (1958): Some theorems concerning 2-dimensional Brownian motion. *Trans. Amer. Math. Soc.* 87 187-197.