SÉMINAIRE DE PROBABILITÉS (STRASBOURG)

YUKUANG CHIU

The multiplicity of stochastic processes

Séminaire de probabilités (Strasbourg), tome 31 (1997), p. 207-215

[<http://www.numdam.org/item?id=SPS_1997__31__207_0>](http://www.numdam.org/item?id=SPS_1997__31__207_0)

© Springer-Verlag, Berlin Heidelberg New York, 1997, tous droits réservés.

L'accès aux archives du séminaire de probabilités (Strasbourg) (<http://portail.mathdoc.fr/SemProba/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>

The Multiplicity of Stochastic Processes

Yukuang Chiu

Department of Mathematics, 0112
University of California, San Diego
La Jolla, CA 92093-0112, USA

This paper studies the multiplicity of non-Gaussian, non-infinitely divisible and non-stationary processes associated with the “chaos” space of N. Wiener [12], and for each positive integer N and for $N = \infty$, constructs a process of multiplicity N . The examination of multiplicity of a process has been of interest to many authors such as H. Cramér [2,3,4], T. Hida [5,6], K. Itô [7] and G. Kallianpur and V. Mandrekar [10].

Our approach here begins with a classical, well known theorem on a separable Hilbert space.

Let U_t , ($t \in R$) be a one parameter group of unitary operators acting on a separable Hilbert space H , and let E_λ be its spectral measure, i.e.,

$$U_t = \int_{-\infty}^{\infty} e^{it\lambda} dE_\lambda.$$

Then there exists a sequence $\{f_n\}$ of elements in H , which will be referred to as cyclic vectors, such that the Hilbert space H can be Hellinger-Hahn [9] decomposed into a direct sum

$$H = \sum_{n \geq 1} \oplus H_n,$$

where

$$\begin{aligned} H_n &= \left\{ \int_{-\infty}^{\infty} g(\lambda) dE_\lambda f_n; g \in L^2(R, \mu_n) \right\} \\ &= \text{linear span of } \{U_t f_n; -\infty < t < \infty\}, \end{aligned}$$

which will be referred to as a cyclic subspace of H with f_n , with the notation

$$d\mu_n(\lambda) = \|dE_\lambda f_n\|^2,$$

we further have

$$d\mu \stackrel{\text{def}}{=} d\mu_1 \gg d\mu_2 \gg \cdots,$$

where $d\mu \gg d\nu$ means that the measure $d\mu$ is absolutely continuous with respect to the measure $d\nu$. The type of the measure sequence $\{d\mu_n\}$ is invariant with respect to the choice of $\{f_n\}$'s. This is to say that if $H = \sum_{n \geq 1} \oplus H'_n$ is another decomposition with H'_n a cyclic subspace with cyclic vector f'_n , then

$$d\mu_n \sim d\mu'_n (\text{equivalence}), n = 1, 2, \dots,$$

where $d\mu'_n(\lambda) = \|dE_\lambda f'_n\|^2$.

Denote the support for $d\mu_n$ by Λ_n . The integer $m(\lambda) = \max\{n; \lambda \in \Lambda_n\}$ is referred to as the multiplicity of λ , and the pair $\{d\mu, m\}$, the spectral type of U_t . The spectral type of U_t is said to be σ -Lebesgue if $d\mu$ is equivalent to Lebesgue measure and if $m(\lambda) \equiv \infty$; and that of U_t is said to be simple Lebesgue if $d\mu$ is equivalent to Lebesgue measure and if $m(\lambda) \equiv 1$.

As a further consequence of Hellinger-Hahn decomposition, we have that if U_t and U'_t are one parameter groups of unitary operators acting on $\mathbf{H}$ and $\mathbf{H}'$ respectively, and if they are unitary equivalent, i.e., if there exists an isometry V of $\mathbf{H}$ onto $\mathbf{H}'$ such that $U'_t = VU_tV^{-1}$, then the associated measure sequences $\{d\mu_n\}$ and $\{d\mu'_n\}$ are of the same type. Conversely, if these two sequences are of the same type, then we can construct an isometry between $\mathbf{H}$ and $\mathbf{H}'$ such that $\{U_t\}$ and $\{U'_t\}$ are unitary equivalent. In other words, the sequence $\{d\mu_n\}$ is unitary invariant.

Example Define θ_t to be the transform of $L^2(R)$

$$\theta_t : \begin{array}{ccc} L^2(R) & \rightarrow & L^2(R) \\ F(\cdot) & \rightarrow & F(\cdot - t). \end{array}$$

Then θ_t consists of a one parameter group of unitary operators on $L^2(R)$, and its spectral type is simple Lebesgue.

To see this, let us write

$$\tau(u) = \begin{cases} e^u, & u < 0 \\ 0, & u \geq 0 \end{cases}.$$

Then

$$\hat{\tau}(\lambda) \stackrel{\text{def}}{=} \frac{1}{\sqrt{2\pi}} \int_{-\infty}^0 e^u e^{i\lambda u} du = \frac{1}{\sqrt{2\pi}} \frac{1}{1 + i\lambda}.$$

Since the Fourier transform is topologically isomorphic on $L^2(R)$ by Plancherel's theorem [13], it follows that

$$\text{linear span } \{e^{i\lambda t} \hat{\tau}(\lambda), t \in R\} = \text{linear span } \{\theta_t \tau(\cdot), t \in R\} = L^2(R).$$

Thus $L^2(R)$ itself turns out to be a cyclic space with cyclic vector τ . Now

$$\begin{aligned} \left(\int_{-\infty}^{\infty} e^{it\lambda} dE_{\lambda} \tau \right)(u) &= \tau(u - t) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{-i\lambda(u-t)} \hat{\tau}(\lambda) d\lambda, \\ \int_{-\infty}^{\infty} e^{it\lambda} \|dE_{\lambda} \tau\|^2 &= \int_{-\infty}^{\infty} e^{it\lambda} |\hat{\tau}(\lambda)|^2 d\lambda. \end{aligned}$$

Hence

$$d\mu(\lambda) = \|dE_{\lambda} \tau\|^2 = \frac{1}{2\pi} \frac{1}{1 + \lambda^2} d\lambda.$$

This shows that the measure $d\mu$ is equivalent to Lebesgue measure and $m(\lambda) \equiv 1$, and consequently simple Lebesgue.

In the sequel, let U_t be the one parameter group of unitary operators induced by Brownian motion flow T_t on $L^2(\mathbf{B})$ [5], i.e., the collection of all variables measurable with respect to the σ -field generated by Brownian motion $\mathbf{B}$ with finite variances. First, we look at the spectral type of U_t .

To begin with, let $L^2(\mathbf{B}) = \sum_{n=0}^{\infty} \oplus \mathcal{H}_n$ be the Wiener-Itô decomposition [8, 12] of $L^2(\mathbf{B})$. It is well known that each $\mathcal{H}_n$, which consists of an U_t -invariant subspace, is topologically isomorphic to $\sqrt{n!} \tilde{L}^2(R^n)$ (via $\mathcal{J}$ -transformation [5]), where $\tilde{L}^2(R^n)$ denotes all the symmetric functions of $L^2(R^n)$, and that each element in $\mathcal{H}_n$ can be expressed as an n -multiple Wiener integral. Without ambiguity, we still write U_t to be the restriction of U_t on $\mathcal{H}_n$. We then have

Theorem For each $n \geq 2$, the spectral type of U_t on $\mathcal{H}_n$ is σ -Lebesgue.

To prove this, we introduce a unitary isometry V_t of U_t . Since spectral type is unitary invariant, the investigation of spectral type of U_t may be reduced to a search for that of V_t . Now let us put

$$L_{nc}^2 = L^2((u_1, u_2, \dots, u_n) \in R^n; u_1 \leq u_2 \leq \dots \leq u_n)$$

and define $\mathcal{C}$:

$$\mathcal{C}: \begin{array}{ccc} \hat{L}^2(R^n) & \rightarrow & \sqrt{n!} L_{nc}^2 \\ F(u_1, u_2, \dots, u_n) & \rightarrow & F(u_{\pi(1)}, u_{\pi(2)}, \dots, u_{\pi(n)}), \end{array}$$

where π is a permutation of $\{1, 2, \dots, n\}$ such that $u_{\pi(1)} \leq u_{\pi(2)} \leq \dots \leq u_{\pi(n)}$. Obviously $\mathcal{C}$ defines an isometric mapping from $\hat{L}^2(R^n)$ to $\sqrt{n!} L_{nc}^2$. Further let

$$A_n = \begin{pmatrix} \frac{1}{n} & \frac{1}{n} & \dots & \frac{1}{n} & \frac{1}{n} \\ -1 & 1 & \dots & 0 & 0 \\ \dots & \dots & \dots & \dots & \dots \\ 0 & 0 & \dots & 1 & 0 \\ 0 & 0 & \dots & -1 & 1 \end{pmatrix}$$

and define $\mathcal{E}$:

$$\mathcal{E}: \begin{array}{ccc} L_{nc}^2 & \rightarrow & L^2(R \times R_+^{n-1}) \\ F(u_1, u_2, \dots, u_n) & \rightarrow & G(v_1, v_2, \dots, v_n), \end{array}$$

where $R_+ = [0, \infty)$ and

$$\begin{pmatrix} v_1 \\ v_2 \\ \vdots \\ v_n \end{pmatrix} = A_n \begin{pmatrix} u_1 \\ u_2 \\ \vdots \\ u_n \end{pmatrix}.$$

Then again we verify that $\mathcal{E}$ defines an isometric mapping from L_{nc}^2 to $L^2(R \times R_+^{n-1})$. Hence if

$$V_t \stackrel{\text{def}}{=} (\mathcal{E} \cdot \mathcal{C} \cdot \mathcal{J})^{-1} U_t (\mathcal{E} \cdot \mathcal{C} \cdot \mathcal{J}),$$

then $\{V_t, t \in R\}$ consists of a one parameter group of unitary operators on $L^2(R \times R_+^{n-1})$. As a matter of fact, with the diagram

$$\begin{array}{ccc} & \mathcal{H}_n & \xrightarrow{\mathcal{J}} \sqrt{n!} \hat{L}^2(R^n) \\ \mathcal{E} \cdot \mathcal{C} \cdot \mathcal{J}: & \uparrow & \downarrow \mathcal{C} \\ & n! L^2(R \times R_+^{n-1}) & \xleftarrow{\mathcal{E}} n! L_{nc}^2 \end{array}$$

in mind, we see that if

$$\mathcal{E} \cdot \mathcal{C} \cdot \mathcal{J}: \begin{array}{ccc} \mathcal{H}_n & \rightarrow & n! L^2(R \times R_+^{n-1}) \\ \varphi & \rightarrow & n! G(v_1, v_2, \dots, v_n), \end{array}$$

then

$$\begin{aligned} (1) \quad U_t \varphi &\rightarrow n! G(v_1 - t, v_2, \dots, v_n) \\ (2) \quad &= n! (V_t G)(v_1, v_2, \dots, v_n). \end{aligned}$$

To see the spectral type of V_t on $L^2(R \times R_+^{n-1})$, we decompose $L^2(R \times R_+^{n-1})$ into a direct sum by means of a complete orthonormal basis $\{\eta_n; n \geq 0\}$ of $L^2(R_+)$:

$$(3) \quad L^2(R \times R_+^{n-1}) = \sum_{k_2, \dots, k_n \geq 0} \oplus L_{k_2, \dots, k_n},$$

where

$$L_{k_2, \dots, k_n} = \{f(v_1) \otimes \eta_{k_2}(v_2) \otimes \dots \otimes \eta_{k_n}(v_n); f \in L^2(R)\},$$

and $\otimes$ means tensor product. Such η_n 's may be taken, for example as the Laguerre functions. Apparently, the subspace $L_{k_2, \dots, k_n}$ of $L^2(R \times R_+^{n-1})$ by (1) and (2) is V_t invariant, and the spectral type of V_t on each $L_{k_2, \dots, k_n}$, as seen in the example, is simple Lebesgue. Combining this with (3), we have proven that the spectral type of V_t on $L^2(R \times R_+^{n-1})$ is σ -Lebesgue.

Here, let us note that if we put

$$X_{k_2, \dots, k_n}(t) = (\mathcal{E} \cdot \mathcal{C} \cdot \mathcal{J})^{-1}(\tau(v_1 - t)\eta_{k_2}(v_2) \dots \eta_{k_n}(v_n))$$

then $X_{k_2, \dots, k_n}(t)$ may be expressed as a stochastic integral

$$\begin{aligned} X_{k_2, \dots, k_n}(t) &= \int_{-\infty}^t dB(u_n) \int_{-\infty}^{u_n} \eta_{k_n}(u_n - u_{n-1}) dB(u_{n-1}) \times \dots \times \\ &\int_{-\infty}^{u_3} \eta_{k_3}(u_3 - u_2) dB(u_2) \times \int_{-\infty}^{u_2} \tau\left(\frac{u_1 + \dots + u_n}{n} - t\right) \eta_{k_2}(u_2 - u_1) dB(u_1). \end{aligned}$$

Hence if we put

$$\mathcal{H}_n(X_{k_2, \dots, k_n}) = (\mathcal{E} \cdot \mathcal{C} \cdot \mathcal{J})^{-1} L_{k_2, \dots, k_n},$$

then

$$\mathcal{H}_n = \sum_{k_2, \dots, k_n \geq 0} \oplus \mathcal{H}_n(X_{k_2, \dots, k_n}).$$

This is the decomposition of $\mathcal{H}_n$ corresponding to that of $L^2(R \times R_+^{n-1})$.

Further, if we notice that the expectations in $\mathcal{H}_n$ correspond to the multiple integrations in $L^2(R \times R_+^{n-1})$, then we can immediately compute, for example

$$E[(X_{k_2, \dots, k_n}(t) - X_{k_2, \dots, k_n}(s))^2] = \int_{-\infty}^{\max\{t, s\}} (\tau(u - t) - \tau(u - s))^2 du,$$

and

$$(4) \quad E[X_{k_2, \dots, k_n}(t) X_{k_2, \dots, k_n}(s)] = \frac{1}{2} e^{-|t-s|}.$$

In the case where $n = 2$, which is of particular interest, we will write

$$X_n(t) = \int_{-\infty}^t dB(u_2) \int_{-\infty}^{u_2} \tau\left(\frac{u_1 + u_2}{2} - t\right) \eta_n(u_2 - u_1) dB(u_1).$$

We now focus on the multiplicity of a process $X(t) \in \mathcal{H}_2$:

$$X(t) \stackrel{\text{def}}{=} \sum_{n=0}^{\infty} F(t)^n X_n(t),$$

where $F(t)$ on R is an absolutely continuous function with (i) $0 < F(t) \leq \delta < 1$.

Theorem If $F(t)$ further satisfies the conditions (ii) the derivative F' of F is in $L^1(R)$; (iii) for any open interval (a, b) ,

$$\int_a^b F'^2 dt = +\infty,$$

then the multiplicity of $X(t)$ is infinity.

The proof will be done by constructing another process $Y(t)$ which is both canonically represented by Brownian motion and has the same reproducing kernel Hilbert space as that of $X(t)$. Consequently, the determination of the multiplicity for process $X(t)$ may be reduced to that for $Y(t)$.

Before constructing $Y(t)$, let us first find a process $T(t)$ such that $T(t)$ can be canonically represented by Brownian motion, and that $T(t)$ shares the same covariance with $X_n(t)$. Since the covariance of $X_n(t)$ is given by (4), it follows from N. Wiener [11] that such a process must be Ornstein-Uhlenbeck process

$$T(t) = \int_{-\infty}^t e^{-(t-u)} dB(u).$$

Let us prepare a sequence of independent Brownian motions on R : $B_0, B_1, B_2, \dots$, and let

$$Y_n(t) = \int_{-\infty}^t e^{-(t-u)} dB_n(u).$$

Then a process $Y(t)$ defined as

$$Y(t) = \sum_{n=0}^{\infty} F(t)^n Y_n(t)$$

shares the same reproducing kernel Hilbert space as that of $X(t)$. Hence the multiplicity of $Y(t)$ equals that of $X(t)$.

To say that the multiplicity of $Y(t)$ is infinity, it suffices to show by T. Hida [5,6] that the representation of $Y(t)$ is canonical, i.e., fix $T \in R$, for $n = 0, 1, 2, \dots$, take $f_n \in L^2((-\infty, T])$ such that

$$\sum_{n=0}^{\infty} \int_{-\infty}^T |f_n(t)|^2 dt < \infty$$

and let

$$g_n(t) = \int_{-\infty}^{\min(t, T)} e^{-(t-u)} f_n(u) du.$$

We then have to show that if

$$h_0(t) := \sum_{n=0}^{\infty} F(t)^n g_n(t) = 0,$$

then $f_n = 0$ in $L^2((-\infty, T])$, $n = 0, 1, 2, \dots$. For this purpose, let

$$\begin{aligned} h_k(t) &= \sum_{n=k}^{\infty} n(n-1) \cdots (n-k+1) F(t)^{n-k} g_n(t), \quad k \geq 1 \\ l_k(t) &= \sum_{n=k}^{\infty} n(n-1) \cdots (n-k+1) F(t)^{n-k} g'_n(t), \quad k \geq 1 \\ l_0(t) &= \sum_{n=0}^{\infty} F(t)^n g'_n(t). \end{aligned}$$

It is clear that for all k ,

$$l_k(t) \in L_{loc}^2(R), \quad h_k(t) \in C(R),$$

where $L_{loc}^2(R)$ and $C(R)$ denote all the locally L^2 integrable functions and all the continuous functions on R respectively. It then follows, by mathematical induction and hypotheses on F that

$$\begin{aligned} h'_0(t) = l_0(t) + F'(t)h_1(t) = 0 &\implies h_1(t) = 0 \\ h'_1(t) = l_1(t) + F'(t)h_2(t) = 0 &\implies h_2(t) = 0 \\ &\dots \dots \dots \\ h'_k(t) = l_k(t) + F'(t)h_{k+1}(t) = 0 &\implies h_{k+1}(t) = 0 \\ &\dots \dots \dots \end{aligned}$$

In matrix form,

$$A_t \cdot \begin{pmatrix} g_0(t) \\ g_1(t) \\ \vdots \\ g_n(t) \\ \vdots \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 0 \\ \vdots \end{pmatrix},$$

where

$$A_t = \begin{pmatrix} 1 & F(t) & F(t)^2 & F(t)^3 & \dots & \binom{n}{0}F(t)^n & \dots & \dots \\ 0 & 1 & 2F(t) & 3F(t)^2 & \dots & \binom{n}{1}F(t)^{n-1} & \dots & \dots \\ 0 & 0 & 1 & 3F(t) & \dots & \binom{n}{2}F(t)^{n-2} & \dots & \dots \\ 0 & 0 & 0 & 1 & \dots & \binom{n}{3}F(t)^{n-3} & \dots & \dots \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ \dots & \dots & \dots & \dots & \dots & \binom{n}{n-1}F(t) & \dots & \dots \\ 0 & 0 & 0 & 0 & \dots & 1 & \dots & \dots \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots \end{pmatrix}.$$

On the other hand, if we let

$$B_t = \begin{pmatrix} 1 & -F(t) & F(t)^2 & -F(t)^3 & \dots & (-1)^n \binom{n}{0}F(t)^n & \dots & \dots \\ 0 & 1 & -2F(t) & 3F(t)^2 & \dots & (-1)^{n-1} \binom{n}{1}F(t)^{n-1} & \dots & \dots \\ 0 & 0 & 1 & -3F(t) & \dots & (-1)^{n-2} \binom{n}{2}F(t)^{n-2} & \dots & \dots \\ 0 & 0 & 0 & 1 & \dots & (-1)^{n-3} \binom{n}{3}F(t)^{n-3} & \dots & \dots \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ \dots & \dots & \dots & \dots & \dots & -\binom{n}{n-1}F(t) & \dots & \dots \\ 0 & 0 & 0 & 0 & \dots & 1 & \dots & \dots \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots \end{pmatrix},$$

then $B_t A_t = A_t B_t$ turns out to be an infinite unit matrix. This results in $g_n(t) = 0$ and hence $f_n = 0, n = 0, 1, 2, \dots$. The proof of the theorem is thus completed.

As a consequence of the approach, we may easily prove that for each positive integer N , the multiplicity of a process defined as

$$X(t) = \sum_{n=0}^{N-1} F(t)^n X_n(t)$$

is exactly N .

The argument for this follows if, in the proof, we define $Y(t)$ as

$$Y(t) = \sum_{n=0}^{N-1} F(t)^n B_n(t)$$

and A_t as

$$A_t = \begin{pmatrix} 1 & F(t) & F(t)^2 & F(t)^3 & \dots & \binom{N-1}{0} F(t)^{N-1} \\ 0 & 1 & 2F(t) & 3F(t)^2 & \dots & \binom{N-1}{1} F(t)^{N-2} \\ 0 & 0 & 1 & 3F(t) & \dots & \binom{N-1}{2} F(t)^{N-3} \\ 0 & 0 & 0 & 1 & \dots & \binom{N-1}{3} F(t)^{N-4} \\ \dots & \dots & \dots & \dots & \dots & \dots \\ \dots & \dots & \dots & \dots & \dots & \binom{N-1}{N-2} F(t) \\ 0 & 0 & 0 & 0 & \dots & 1 \end{pmatrix}$$

and B_t as

$$B_t = \begin{pmatrix} 1 & -F(t) & F(t)^2 & -F(t)^3 & \dots & (-1)^{N-1} \binom{N-1}{0} F(t)^{N-1} \\ 0 & 1 & -2F(t) & 3F(t)^2 & \dots & (-1)^{N-2} \binom{N-1}{1} F(t)^{N-2} \\ 0 & 0 & 1 & -3F(t) & \dots & (-1)^{N-3} \binom{N-1}{2} F(t)^{N-3} \\ 0 & 0 & 0 & 1 & \dots & (-1)^{N-4} \binom{N-1}{3} F(t)^{N-4} \\ \dots & \dots & \dots & \dots & \dots & \dots \\ \dots & \dots & \dots & \dots & \dots & -\binom{N-1}{N-2} F(t) \\ 0 & 0 & 0 & 0 & \dots & 1 \end{pmatrix}.$$

Finally, we need to demonstrate the existence of the function F . The construction will be done by using the Monotone Convergence Theorem.

Notation: Let $f(t)$ be a function locally symmetric at $t = x$ and let $N(x)$ denote the local support of f at x and $|N(x)|$ denote the Lebesgue measure of the support $N(x)$.

We first proceed to construct a sequence of functions $s_n(t)$, $n = 1, 2, \dots$ as follows.

$s_1(t)$: (i) symmetric about y-axis, (ii) locally symmetric at $t = \frac{n}{2}$, $n = 1, 2, \dots$ and $|N(\frac{n}{2})| \leq \frac{1}{2^2}$, and (iii) $0 < \int_{N(\frac{n}{2})} s_1(t) dt \leq \frac{1}{2} \frac{\delta}{2^{2+n}}$ and $\int_{N(\frac{n}{2})} s_1^2(t) dt = +\infty$, $n = 1, 2, \dots$;
 $s_2(t)$: (i) symmetric about y-axis, (ii) locally symmetric at $t = \frac{n}{2^2}$, $n = 1, 3, 5, \dots$, and $|N(\frac{n}{2^2})| \leq \frac{1}{2^3}$, and (iii) $0 < \int_{N(\frac{n}{2^2})} s_2(t) dt \leq \frac{1}{2} \frac{\delta}{2^{3+n}}$ and $\int_{N(\frac{n}{2^2})} s_2^2(t) dt = +\infty$, $n = 1, 2, \dots$. In general, for $k \geq 3$, we similarly construct $s_k(t)$ as $s_k(t)$: (i) symmetric about y-axis, (ii) locally symmetric at $t = \frac{n}{2^k}$, $n = 1, 3, 5, \dots$, and $|N(\frac{n}{2^k})| \leq \frac{1}{2^{k+1}}$, and (iii) $0 < \int_{N(\frac{n}{2^k})} s_k(t) dt \leq \frac{1}{2} \frac{\delta}{2^{k+1+n}}$ and $\int_{N(\frac{n}{2^k})} s_k^2(t) dt = +\infty$, $n = 1, 2, \dots$.

Now, let us consider the sum

$$S_n(t) \stackrel{\text{def}}{=} \sum_{k=1}^n s_k(t).$$

Since we obviously have

$$0 \leq S_1(t) \leq S_2(t) \leq \dots,$$

and

$$0 < \lim_{n \rightarrow \infty} \int_{-\infty}^{\infty} S_n(t) dt \leq \sum_{k=1}^{\infty} 2^{-k} \delta = \delta,$$

it follows from the Monotone Convergence Theorem that

$$S(t) \stackrel{\text{def}}{=} \lim_{n \rightarrow \infty} S_n(t)$$

exists for almost all t . Now define function F as

$$F(t) = \int_{-\infty}^t S(u) du.$$

We may easily verify that the function F satisfies the conditions as in the theorem.

Acknowledgements

I thank Prof. T. Hida of Nagoya University, Japan for his academic supervision. As his 70th birthday anniversary draws close, I wish him in good health. I thank Drs Fred Wright and Chuck Berry at University of California, San Diego for their encouragement and support.

References

- [1] N. Aronszajn, Theory of reproducing kernels. Trans. Amer. Math. Soc. Vol. 68, 337-404 (1950).
- [2] H. Cramér, Stochastic Processes as Curves in Hilbert Space. Theory Probability Appl., 9(2), 169-179 (1964).
- [3] H. Cramér, A Contribution to the Multiplicity Theory of Stochastic Processes. Proc. Fifth Berkeley Symp. Stat. Appl. Probability, II, 215-221 (1965).
- [4] H. Cramér, Structural and Statistical Problems for a Class of Stochastic Processes. The First Samuel Stanley Wilks Lecture at Princeton University, March 17 1970, 1-30 (1971).
- [5] T. Hida, Brownian Motion. Springer-Verlag (1980).
- [6] T. Hida, Canonical Representations of Gaussian Processes and Their Applications. mem. College Sci., Univ. Kyoto, A33 (1), 109-155 (1960).
- [7] K. Itô, Spectral Type of The Shift Transformation of Differential Processes With Stationary Increments. Trans. Amer. Math. Soc., Vol. 81, No. 2, 253-263 (1956).
- [8] K. Itô, Multiple Wiener integral. J. Math. Soc. Japan 3, 157-169 (1951).
- [9] S. Itô, On Hellinger-Hahn's Theorem. (In Japanese) Sūgaku, vol. 5, no. 2, 90-91 (1953).

- [10] G. Kallianpur and V. Mandrekar, On the Connection between Multiplicity Theory and O. Hanner's Time Domain Analysis of Weakly Stationary Stochastic Processes. Univ. North Carolina Monograph Ser. Probability Stat., No. 3, 385-396 (1970).
- [11] N. Wiener, Time Series. M.I.T. (1949).
- [12] N. Wiener, Nonlinear Problems in Random Theory. M.I.T. (1958).
- [13] N. Wiener, The Fourier Integral and Certain of Its Applications. Dover Publications. INC., New York (1958).