

SÉMINAIRE DE PROBABILITÉS (STRASBOURG)

AZZOUZ DERMOUNE

OCTAVE MOUTSINGA

Generalized variational principles

Séminaire de probabilités (Strasbourg), tome 36 (2002), p. 183-193

<http://www.numdam.org/item?id=SPS_2002__36__183_0>

© Springer-Verlag, Berlin Heidelberg New York, 2002, tous droits réservés.

L'accès aux archives du séminaire de probabilités (Strasbourg) (<http://portail.mathdoc.fr/SemProba/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

GENERALIZED VARIATIONAL PRINCIPLES

Azzouz Dermoune

Octave Moutsinga

Université des Sciences et Technologies de Lille
Laboratoire de Statistique et Probabilités
F.R.E. CNRS 2222
59655 Villeneuve d'Ascq cédex, France

Abstract

In [7] Weinan E, Y. G. Rykov, and Y. G. Sinai have introduced a generalized variational principles in order to give a weak solution of the pressureless gas equations with initial velocity u_0 and distribution of masses given by a probability measure P . The aim of this work is to connect these generalized variational principles at each time $t > 0$ with the convex hull of any primitive of the map $m \in (0, 1) \rightarrow F^{-1}(m) + tu_0(F^{-1}(m))$. Here F is the distribution function of the probability measure P and F^{-1} is its inverse. The latter convex hull is also used to obtain the solutions of the scalar conservation law and the Hamilton-Jacobi equation associated with the pressureless gas equations.

1 Motivation

First we recall the definition of the system of conservation law studied in [7].

Definition. Let u_0 be a continuous and bounded real function. Let $(P_t : t \geq 0)$ be a family of probability measures on $\mathbf{R}$, weakly continuous with respect to t . For each $t \geq 0$, let $I_t = u(t, \cdot)P_t$ be a measure absolutely continuous with respect to P_t . $(P_t, I_t, u(\cdot, t) : t \geq 0)$ is a weak solution of the pressureless gas equations

$$\begin{cases} \frac{\partial P_t}{\partial t} + \frac{\partial I_t}{\partial x} = 0 \\ \frac{\partial I_t}{\partial t} + \frac{\partial (u I_t)}{\partial x} = 0 \\ P_t \rightarrow P, u P_t \rightarrow u_0 P \quad \text{weakly,} \end{cases}$$

if, for any $f \in C_0^1(\mathbf{R})$, the space of C^1 -functions on $\mathbf{R}$ with compact support, and $0 < t_1 < t_2$,

$$\int f(x) dP_{t_2}(x) - \int f(x) dP_{t_1}(x) = \int_{t_1}^{t_2} \int f'(x) dI_t(x) dt,$$

and

$$\int f(x) dI_{t_2}(x) - \int f(x) dI_{t_1}(x) = \int_{t_1}^{t_2} \int f'(x) u(x, t) dI_t(x) dt.$$

Weinan E, Rykov, Sinai in [7] have constructed a weak solution for the pressureless gas equations using sticky particle dynamics. Each particle is indexed by its initial position $x \in \mathbf{R}$, initial velocity $u_0(x)$, and the mass of the set of particles $(-\infty, x]$ is equal to $F(x) := P(-\infty, x]$. Before collision $x + tu_0(x)$ is the position of the particle x at time t . At collisions the colliding particles stick and form a new massive particle. The mass and the velocity of this new particle are given by the laws of conservation of mass and momentum. The method used in [7] is based on the construction of a partition ξ_t of $\mathbf{R}$, which divides the particles into ordered intervals (called clusters), so that each group of particles initially located in an interval $[\alpha, \beta] \in \xi_t$ are glued to a single one before or at time t , and different clusters are at different locations at time t . Each element $[\alpha, \beta]$ of ξ_t is then completely determined by its endpoints α, β . These endpoints are characterized by the following generalized variational principles, denoted in the sequel by (GVP).

(GVP1) α is the left endpoint of an element of ξ_t iff

$$\frac{\int_{[y_1, \alpha]} [\eta + tu_0(\eta)] dP(\eta)}{P[y_1, \alpha]} < \frac{\int_{[\alpha, y_2]} [\eta + tu_0(\eta)] dP(\eta)}{P[\alpha, y_2]},$$

for all $y_1 < \alpha < y_2$.

(GVP2) β is the right endpoint of an element of ξ_t iff

$$\frac{\int_{(y_1, \beta]} [\eta + tu_0(\eta)] dP(\eta)}{P(y_1, \beta]} < \frac{\int_{(\beta, y_2]} [\eta + tu_0(\eta)] dP(\eta)}{P(\beta, y_2]} y,$$

for all $y_1 < \beta < y_2$.

Having $(\xi_t : t \geq 0)$, Weinan E, Rykov, Sinai in [7] have defined the forward flow map associated to pressureless gas equations as follows:

$$\varphi(t, x, P, u_0) = \frac{\int_{[\alpha, \beta]} [\eta + tu_0(\eta)] dP(\eta)}{P[\alpha, \beta]}, \quad (1)$$

where $[\alpha, \beta]$ is the unique element of ξ_t which contains x . They showed that $(P_t = P \circ \varphi^{-1}(t, \cdot, P, u_0), I_t = u(\cdot, t)P_t : t \geq 0)$ is a weak solution of the pressureless gas equations. Here

$$u(x, t) = \frac{\int_{[\alpha, \beta]} u_0(\eta) dP(\eta)}{P[\alpha, \beta]}. \quad (2)$$

In the other hand Dermoune [3, 4, 5] has constructed, for any probability distribution P , a process $(X_t, t \geq 0)$ describing trajectories of sticky particle dynamics with initial velocity u_0 , and masses distributed following the probability P . This process is solution of the non-linear stochastic differential equation

$$dX_t = \mathbf{E}[u_0(X_0) | X_t] dt, \quad \mathcal{L}(X_0) = P, \quad (3)$$

and

$$X_t = \mathbf{E}[X_0 + tu_0(X_0) | X_t], \forall t \geq 0. \quad (4)$$

Using a simple proof based on the formula of change of variables he showed that $(P_t = \mathcal{L}(X_t), I_t = u(\cdot, t)P_t)_{t \geq 0}$ is a weak solution of the pressureless gas equations. Here $u(x, t) = \mathbf{E}[u_0(X_0) | X_t = x]$. The formula (4) gives the relation between the trajectories $(X_t, t \geq 0)$ of sticky particles and the trajectories $(X_0 + tu_0(X_0), t \geq 0)$ of the particles without any interaction.

Now we make some remarks on the (GVP). If P is continuous then the partition ξ_t given by (GVP1) and (GVP2) is ambiguous. In fact, in this case, every left endpoint is also a right endpoint. Our aim here is to clarify this point and to give a precise definition of ξ_t .

The idea in our work is to index each particle by $m \in F(\mathbf{R})$, with initial position $F^{-1}(m)$ (**Fig. 1**), velocity $u_0(F^{-1}(m))$, and the mass of the set of particles $(0, m]$ (i.e. initially located in $(-\infty, F^{-1}(m)]$) is equal to m . Before collision $F^{-1}(m) + tu_0(F^{-1}(m))$ is the position of the particle m at time t (**Fig. 2**). After the collision the map $m \in (0, 1) \rightarrow F^{-1}(m) + tu_0(F^{-1}(m))$ oscillates, and now the positions of massive particles are given by the derivative of the convex hull $H(\cdot, t)$ of any primitive of the map $m \in (0, 1) \rightarrow F^{-1}(m) + tu_0(F^{-1}(m))$. The right inverse

$$M^*(x, t) := \inf\{m \in (0, 1) : \partial_m H(m, t) > x\}$$

is the mass at time t of the set of clusters located in $(-\infty, x]$, and the left inverse

$$M_*(x, t) := \sup\{m \in (0, 1) : \partial_m H(m, t) < x\},$$

is the mass of the set of clusters located in $(-\infty, x)$ (see **Fig. 3**). The velocity at time t of the cluster located at x is given by

$$u(x, t) = \frac{\int_{M_*(x, t)}^{M^*(x, t)} u_0(F^{-1}(m)) dm}{M^*(x, t) - M_*(x, t)},$$

and the family $(\partial_x M^*(x, t), u(x, t) \partial_x M^*(x, t), u(x, t))_{t \geq 0}$ is a weak solution of our pressureless gas equations.




Fig. 1. The particle indexed by m is initially located at $F^{-1}(m)$.

Fig. 2. Before collision $F^{-1}(m) + tu_0(F^{-1}(m))$ is the position at time t of the particle indexed by m . The colliding particles before or at time t are represented by the graphic regions where the map $m \mapsto F^{-1}(m) + tu_0(F^{-1}(m))$ oscillates.

Fig. 3. The ordinate x of a point (m, x) of the graph represents a location at time t of a cluster. The length $M^* - M_*$ of an horizontal segment belonging to the graph is the mass of a cluster at time t .

In [2] *Brenier* and *Grenier* have established a connection between the pressureless gas equations and the following scalar conservation law :

$$\partial_t M + \partial_x (A(M)) = 0, \quad M(x, 0) = F(x), \quad (5)$$

where A is any primitive of $m \in (0, 1) \rightarrow u_0(F^{-1}(m))$. In our work we show that the entropy solution of (5) is given by the map $(x, t) \rightarrow M^*(x, t)$, defined above.

Plan of the paper. In Section 2 we give a precise definition of the (GVP). Section 3 is consecrated to the study of the extreme points of the convex hull of

the map $z \in (0, 1) \rightarrow \int_{\frac{1}{2}}^z [F^{-1}(m) + tu_0(F^{-1}(m))] dm$ as a function of the position of the clusters. We end our work by Section 4 which contains the connection between the (GVP), the scalar conservation law (5) and the Hamilton-Jacobi equation

$$\partial_t \Psi(x, t) + A(\Psi(x, t)) = 0 ,$$

with initial condition $\Psi(\cdot, 0)$ is any primitive of F .

2 A precise definition of the (GVP)

The support $\text{supp}(P)$ of P is defined by

$$\text{supp}(P) = \{a \in \mathbf{R} : P(a - \varepsilon, a + \varepsilon) > 0, \forall \varepsilon > 0\} .$$

Let $S_- = \{a \in \mathbf{R} : P(a - \varepsilon, a] > 0, \forall \varepsilon > 0\}$, and $S_+ = \{a \in \mathbf{R} : P[a, a + \varepsilon) > 0, \forall \varepsilon > 0\}$. Then $\text{supp}(P) = S_+ \cup S_-$. For $z \in (0, 1]$ we define

$$F^{-1}(z) = \inf\{a : F(a) \geq z\} ,$$

where $F(a) = P(-\infty, a]$. We have $F(F^{-1}(z)) = z$ if $z \in F(\mathbf{R})$, and if $F(x - 0) < z \leq F(x)$ for some $x \in \mathbf{R}$, then $F(F^{-1}(z)) = F(x)$. It is easy to show that F^{-1} is one to one from $F(\mathbf{R})$ into S_- .

Let us consider, for each fixed $t > 0$, a primitive of the map $m \in (0, 1) \rightarrow F^{-1}(m) + tu_0(F^{-1}(m))$, for example

$$\Phi(z, t) = \int_{\frac{1}{2}}^z [F^{-1}(m) + tu_0(F^{-1}(m))] dm .$$

We denote by λ the Lebesgue measure on $[0, 1]$. It follows from the equality $\lambda \circ (F^{-1})^{-1} = P$ that for all $\alpha, \beta \in \mathbf{R}$,

$$\Phi(F(\beta), t) - \Phi(F(\alpha), t) = \int_{(\alpha, \beta]} [\eta + tu_0(\eta)] dP(\eta) .$$

We suppose that $\text{supp}(P)$ is bounded or $\int_0^x \eta dP(\eta) \rightarrow +\infty$ as $|x| \rightarrow +\infty$. Under this assumption the convex hull $H(\cdot, t)$ of the map $\Phi(\cdot, t)$ is well defined.

Now we construct a partition ξ_t of $\text{supp}(P)$ as follows. An element of ξ_t has the form $[\alpha, \beta] \cap \text{supp}(P)$. The endpoints α, β are given by the extreme points of $H(z, t)$. More precisely let

$$F_l^{-1}(z) = \inf\{a : F(a) > z\} , \quad \text{and} \quad F_r^{-1}(z) = \sup\{a : F(a) < z\} .$$

If m is an extreme point of $H(\cdot, t)$, then $F_r^{-1}(m)$ a right endpoint of an element $G_1 \in \xi_t$ and $F_l^{-1}(m)$ is a left endpoint of an element $G_2 \in \xi_t$. The group of particles G_1 is just on the left of the group G_2 . We can show that a right endpoint of ξ_t belongs to S_- , and a left endpoint belongs to S_+ .

Now we give a detailed description of endpoints of ξ_t from extreme points of $H(\cdot, t)$, see **Fig.4** for a graphic illustration. We distinguish four cases.

A) m is isolated in the set of extreme points. Namely, there exist $p_1 < p_2$ and z_1, z_2 two extremes points such that

$$\frac{\Phi(t, m) - \Phi(t, z_1)}{m - z_1} = p_1 < p_2 = \frac{\Phi(t, z_2) - \Phi(t, m)}{z_2 - m},$$

and

$$\frac{\Phi(t, m) - \Phi(t, z')}{m - z'} \leq p_1 < p_2 \leq \frac{\Phi(t, z'') - \Phi(t, m)}{z'' - m}, \text{ for all } z' < m < z'', \quad (6)$$

and there is no extreme point in $(z_1, m) \cup (m, z_2)$. We have two cases.

i) $F_r^{-1}(m) = F_l^{-1}(m)$. In this case there exist two sequences (z_1^n) and (z_2^n) in $F(\mathbf{R})$ such that $z_1^n \rightarrow m - 0$ and $z_2^n \rightarrow m + 0$. Using the continuity of u_0 in (6) we get

$$\alpha + tu_0(\alpha) \leq p_1 < p_2 \leq \alpha + tu_0(\alpha),$$

which is absurd.

ii) So necessarily $F_r^{-1}(m) < F_l^{-1}(m)$. $F_r^{-1}(m)$ is the right endpoint of a cluster with mass $m - z_1$ and $F_l^{-1}(m)$ is the left endpoint of a cluster with mass $z_2 - m$.

B) m is isolated from the right. Namely there exists $z_2 > m$ an extreme point such that (m, z_2) does not contain any extreme point, but for any $z' < m$, (z', m) contains an extreme point. In this case we have two situations.

- i) $F_r^{-1}(m) = F_l^{-1}(m) := \alpha$. This situation implies that α is the left endpoint of the cluster with mass $z_2 - m$. And any interval $(\alpha - \varepsilon, \alpha)$ contains a cluster.
- ii) $F_r^{-1}(m) < F_l^{-1}(m)$. This situation implies that $\{F_r^{-1}(m)\} \in \xi_t$. The cluster with mass $z_2 - m$ has $F_l^{-1}(m)$ as a left endpoint. Any interval $(F_r^{-1}(m) - \varepsilon, F_r^{-1}(m))$ contains a cluster.

C) z is isolated from the left. Namely there exists $z_1 < z$ an extreme point such that (z_1, z) does not contain any extreme point, but for any $z < z'$, (z, z') contains an extreme point.

- i) $F_r^{-1}(m) = F_l^{-1}(m) := \alpha$. This situation implies that α is the right endpoint of the cluster with mass $m - z_1$. Any interval $(\alpha, \alpha + \varepsilon)$ contains a cluster.
- ii) $F_r^{-1}(m) < F_l^{-1}(m)$. Then $F_r^{-1}(m)$ is a right endpoint of a cluster of mass $m - z_1$ and $\{F_l^{-1}(m)\} \in \xi_t$.

D) For all $z_1 < m < z_2$ there is an extreme point in (z_1, m) and an extreme point in (m, z_2) . Then necessarily $F_r^{-1}(m) = F_l^{-1}(m) := \alpha$. In this case the particle α did not meet any other particle during the interval $[0, t]$, and any interval $(\alpha - \varepsilon, \alpha)$ or $(\alpha, \alpha + \varepsilon)$ contains a cluster.




Fig. 4. The different cases of extremal points of $H(\cdot, t)$.
 m_i represents the case i) studied above.

Remark 2.1. 1) From the construction of the partition ξ_t , each endpoint of an element of ξ_t satisfy (GVP1) and (GVP2).

2) It follows from D) that an endpoint, which is an accumulation of endpoints from the left, is necessarily a continuous point of F .

3) If F is increasing then there is no successive clusters with positive masses.

3 Extreme points as a function of the position of clusters

For $t > 0, x \in \mathbf{R}$ we set $G(x, z, t) = \int_{\frac{1}{2}}^z [F^{-1}(m) + tu_0(F^{-1}(m))] dm - x(z - \frac{1}{2})$, for $z \in (0, 1)$. The properties refer to G as a function of z with x, t being fixed. G is continuous in z and has, according to the hypothesis on P , the property

$$\lim_{z \rightarrow 1-0} G(z), \quad \lim_{z \rightarrow 0+} G(z) \quad \text{exist,}$$

or equal to $+\infty$. Hence G attains its smallest value for one or several values of z , the smallest and the largest of which are denoted by $M_*(x, t)$ and $M^*(x, t)$, respectively,

$$M_*(x, t) \leq M^*(x, t).$$

Following the same proof as in [6] we have,

Theorem 3.1. *The functions M_* and M^* have the properties*

- (a) $M^*(x, t) \leq M_*(x', t)$ if $x < x'$,
- (b) $M^*(x - 0, t) = M_*(x, t)$, $M_*(x + 0, t) = M^*(x, t)$
- (c) As functions of (x, t) , $M_*(x, t), M^*(x, t)$ are respectively lower and upper-semicontinuous. At a point where $M_*(x, t) = M^*(x, t)$ both functions are continuous.

3.1 The Physical meaning of $M_*(x, t)$, $M^*(x, t)$

Let us consider the convex hull $H(\cdot, t)$ defined in Section 2. The left inverse $\sup\{z : \partial_z H(z, t) < x\}$ of $\partial_z H(\cdot, t)$ coincides with $M_*(x, t)$, and its right inverse

$$\inf\{z : \partial_z H(z, t) > x\}$$

coincides with $M^*(x, t)$. So $M_*(x, t)$ is the mass at time t of the set of clusters located in $(-\infty, x)$, and $M^*(x, t)$ is the mass of the set of clusters located in $(-\infty, x]$. More precisely let (X_t) be the process defined by (3), then

$$M^*(x, t) = \mathbf{E}[H(x - X_t)] , \quad (7)$$

where $H = 1_{[0, +\infty)}$ is the Heaviside function.

The following result shows that collisions occur when $M_* < M^*$.

Proposition 3.1. *If $M_*(x, t) < M^*(x, t)$, then $M_*(x, t), M^*(x, t)$ are two extreme points of $H(\cdot, t)$, and x is a position of some cluster at time t and $M^*(x, t) - M_*(x, t)$ is the mass of this cluster. If $M_*(x, t) = M^*(x, t) := M(x, t)$, then x is not a position of any cluster at time t or $F_t^{-1}(M_*(x, t)) = F_t^{-1}(M^*(x, t)) := \alpha$ is a cluster situated at $x = \alpha + tu_0(\alpha)$ at time t . Moreover the velocity $u(x, t)$ defined by (2) is given by*

$$u(x, t) = \frac{\int_{M_*(x, t)}^{M^*(x, t)} u_0(F^{-1}(m)) dm}{M^*(x, t) - M_*(x, t)} , \quad \text{if } M_*(x, t) < M^*(x, t) ,$$

and $u(x, t) = u_0(F^{-1}(M(x, t)))$ if $M_*(x, t) = M^*(x, t) := M(x, t)$.

3.2 Characteristics

We set $y_*(x_1, t_1) = F_t^{-1}(M_*(x_1, t_1))$, $y^*(x_1, t_1) = F_t^{-1}(M^*(x_1, t_1))$, and we define as in the inviscid Burgers equation [6] the segments

$$S_*(x_1, t_1) = [(y_*(x_1, t_1), 0), (x_1, t_1)] , \quad S^*(x_1, t_1) = [(y^*(x_1, t_1), 0), (x_1, t_1)] .$$

Namely $(x, t) \in S_*(x_1, t_1)$, respectively $(x, t) \in S^*(x_1, t_1)$, if

$$x = x_1 + \frac{x_1 - y_*(x_1, t_1)}{t_1}(t - t_1) , \quad t \in (0, t_1) ,$$

respectively

$$x = x_1 + \frac{x_1 - y^*(x_1, t_1)}{t_1}(t - t_1) , \quad t \in (0, t_1) .$$

Theorem 3.2. *At every point x, t of the segment $S_*(x_1, t_1)$, respectively $S^*(x_1, t_1)$, $M_*(x, t) = M^*(x, t) = M_*(x_1, t_1)$, and $y_*(x, t) = y^*(x, t) = y_*(x_1, t_1)$ respectively $M_*(x, t) = M^*(x, t) = M^*(x_1, t_1)$, and $y_*(x, t) = y^*(x, t) = y^*(x_1, t_1)$ and they are continuous.*

Proof. Let $x = x_1 + \frac{x_1 - y_*(x_1, t_1)}{t_1}(t - t_1)$, for $t \in (0, t_1)$. First we have

$$G(x, z, t) = t \int_{\frac{1}{2}}^z \left[\frac{F^{-1}(m) - x}{t} + u_0(F^{-1}(m)) \right] dm$$

and

$$\frac{F^{-1}(m) - x}{t} = \frac{F^{-1}(m) - x_1}{t_1} + \frac{t_1 - t}{tt_1}(F^{-1}(m) - y_*(x_1, t_1)) .$$

So $t^{-1}\{G(x, z, t) - G(x, M_*(x_1, t_1), t)\} =$

$$\begin{aligned} & \int_{M_*(x_1, t_1)}^z \left[\frac{F^{-1}(m) - x}{t} + u_0(F^{-1}(m)) \right] dm = \\ & t_1^{-1}\{G(x_1, z, t_1) - G(x_1, M_*(x_1, t_1), t_1)\} \\ & + \frac{t_1 - t}{tt_1} \int_{M_*(x_1, t_1)}^z [F^{-1}(m) - y_*(x_1, t_1)] dm . \end{aligned}$$

If $m > M_*(x_1, t_1)$, then $F^{-1}(m) > y_*(x_1, t_1)$, which implies that

$$\int_{M_*(x_1, t_1)}^z [F^{-1}(m) - y_*(x_1, t_1)] dm > 0, \quad \forall z > M_*(x_1, t_1) ,$$

and thus $G(x, z, t) - G(x, M_*(x_1, t_1), t) > 0$ for all $z > M_*(x_1, t_1)$.

If $z < M_*(x_1, t_1)$, then $G(x_1, z, t_1) - G(x_1, M_*(x_1, t_1), t_1) > 0$ and

$$\int_{M_*(x_1, t_1)}^z [F^{-1}(m) - y_*(x_1, t_1)] dm \geq 0, \quad \forall z < M_*(x_1, t_1) .$$

We conclude that $M_*(x_1, t_1)$ is the unique minimum of the map $z \in (0, 1) \rightarrow G(x, z, t)$.

The proof of the case

$$x = x_1 + \frac{x_1 - y^*(x_1, t_1)}{t_1}(t - t_1), \quad t \in (0, t_1)$$

is similar.

4 Pressureless gas equations and scalar conservation law

Let $P_n = \sum_j m_j \delta(x - x_j)$ be a sequence of finite probabilities such that $P_n \rightarrow P$. The particles $\{x_j : j\}$ move following the model of sticky particles. We denote by $x_j(t)$ the position of the particle x_j at time t . Brenier and Grenier [2] have proved that

$$\sum_j m_j \delta(x - x_j(t)) \rightarrow \partial_x M(x, t) ,$$

where M is the unique entropy solution of the scalar conservation (5).

First let us recall the definition of the entropy solution. Let f be a locally Lipschitz continuous function. The equation

$$\partial_t u(x, t) + \partial_x (f(u(x, t))) = 0, \quad u(0, x) = u_0(x) \text{ is given,} \quad (8)$$

is called a scalar conservation law. The entropy solution is a locally integrable function such that, for all positive smooth function ϕ ,

$$\int \int \partial_t \phi(x, t) I(u(x, t)) + \partial_x \phi(x, t) F(u(x, t)) dx dt + \int \phi(0, x) u_0(x) dx \geq 0, \quad (9)$$

where $I(u) = \int_0^u h(x) dx$, $F(u) = \int_0^u h(x) df(x)$ and h is any nondecreasing function.

In this part we show that $M_*(x, t)$ (respectively $M^*(x, t)$) is the left continuous version (respectively the right continuous) of the entropy solution of (5). We define the function M on the set of the points (x, t) such that $M_*(x, t) = M^*(x, t)$. So M is continuous at every point where $M_* = M^*$, and if $M_*(x_1, t_1) < M^*(x_1, t_1)$ then (x_1, t_1) is a discontinuity point of M . We have $\lim_{x \rightarrow x_1 - 0} M(x, t_1) = M_*(x_1, t_1)$ and $\lim_{x \rightarrow x_1 + 0} M(x, t_1) = M^*(x_1, t_1)$.

Now we show that M is the entropy solution of (5). It is known [1] (see also [2]) that the map

$$(x, t) \rightarrow \Psi(x, t) := \int^x m(y, t) dy,$$

is a viscosity solution (in the sense of Crandall Lions) of the Hamilton-Jacobi equation

$$\partial_t \Psi + A(\partial_x \Psi) = 0,$$

if and only if $(x, t) \rightarrow m(x, t)$ is an entropy solution of

$$\partial_t m(x, t) + \partial_x (A(m(x, t))) = 0.$$

Since the initial condition $\Psi(x, 0) := \frac{1}{2} F^{-1}(\frac{1}{2}) + \int_{\frac{1}{2}}^x F(y) dy$ is convex, the second Hopf formula [1] asserts that the unique viscosity solution with $\Psi(\cdot, 0)$ as initial conditions is given by

$$\Psi(x, t) = \sup_{m \in (0, 1)} \{xm - \Psi(\cdot, 0)^*(m) - tA(m)\},$$

where

$$\Psi(\cdot, 0)^*(m) = \sup_{x \in \mathbb{R}} \{xm - \Psi(x, 0)\}$$

is the Legendre-Fenchel transform of $\Psi(\cdot, 0)$. It is known that for each $t \geq 0$ fixed, $\Psi(\cdot, t)^*$ is the convex hull of the map

$$m \in (0, 1) \rightarrow \Psi(\cdot, 0)^*(m) + tA(m)$$

and the inverse of $\partial_m \Psi(\cdot, t)^*$ coincides with $\partial_x \Psi(\cdot, t)$. We can show for

$$A(m) = \int_{\frac{1}{2}}^m u_0(F^{-1}(z)) dz, \quad m \in (0, 1),$$

and

$$\Psi(x, 0) := \frac{1}{2}F^{-1}\left(\frac{1}{2}\right) + \int_{\frac{1}{2}}^x F(y)dy ,$$

that

$$\Psi(\cdot, 0)^*(m) + tA(m) = \int_{\frac{1}{2}}^m F^{-1}(z) + tu_0(F^{-1}(z))dz , \quad \forall m \in (0, 1).$$

In Section 3 we have denoted by $H(\cdot, t)$ the convex hull of the function

$$m \in (0, 1) \rightarrow \int_{\frac{1}{2}}^m F^{-1}(z)dz + t \int_{\frac{1}{2}}^m u_0(F^{-1}(m'))dm' .$$

So, the inverse $M(\cdot, t)$ of the function $\partial_m H(\cdot, t)$ is equal to the function $\partial_x \Psi(\cdot, t)$. We derive that $M(x, t) = \partial_x \Psi(x, t)$ is the entropy solution of (5).

References

- [1] M. Bardi, L. C. Evans, On Hopf's formula for solutions of Hamilton-Jacobi equations, *Nonlinear Anal.*, 8 (1984), pp. 1373-1381.
- [2] Y. Brenier, E. Grenier, Sticky particles and scalar conservation laws. *Siam. J. Numer. Anal.* Vol. 35, No. 6, pp. 2317-2328, December 1998.
- [3] A. Dermoune, Probabilistic interpretation for system of conservation law arising in adhesion particle dynamics. *C. R. Acad. Sci. Paris*, 1998, tome 5.
- [4] A. Dermoune, Probabilistic interpretation of sticky particles model. *The Annals of Probability*, 1999, Vol. 27, No. 3, 1357-1367.
- [5] A. Dermoune, Sticky particles and propagation of chaos. *Nonlinear Analysis* 45 (2001), 529-541.
- [6] E. Hopf, The partial differential equation $u_t + uu_x = \mu u_{xx}$. *Comm. Pure Appl. Math.* 3, 201-230, (1950).
- [7] Weinan E, Yu. G. Rykov, Ya. G. Sinai, Generalized variational principles, global weak solutions and behavior with random initial data for systems of conservation laws arising in adhesion particle dynamics. *Commun. Math. Phys.* 177, 349-380, (1996).