

SÉMINAIRE DE THÉORIE SPECTRALE ET GÉOMÉTRIE

ANTON ZORICH

Asymptotic flag of an orientable measured foliation

Séminaire de Théorie spectrale et géométrie, tome 11 (1992-1993), p. 113-131

http://www.numdam.org/item?id=TSG_1992-1993__11__113_0

© Séminaire de Théorie spectrale et géométrie (Chambéry-Grenoble), 1992-1993, tous droits réservés.

L'accès aux archives de la revue « Séminaire de Théorie spectrale et géométrie » implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>

ASYMPTOTIC FLAG OF AN ORIENTABLE MEASURED FOLIATION

Anton ZORICH

ABSTRACT.

We state several conjectures on asymptotic “spectral properties” of transformation operators involved in Rauzy induction for a generic interval exchange transformation. Modulo these conjectures we get a very precise approximation for dynamics of leaves of a generic orientable measured foliation on a surface. The main object, which we get is a flag of subspaces in the first (co)homology group of the surface of dimensions $1, \dots, g$, where g is a genus of the surface. This flag of subspaces generalizes asymptotic cycle; in particular the smallest subspace is spanned by the asymptotic cycle. Presumably this flag of subspaces provides a new invariant of foliation.

We illustrate the conjectures by treating a specific example, which comes from a model of electron dynamics on a Fermi-surface suggested by I. Dinnikov.

Though we can not provide any strict mathematical proofs of the conjectures proclaimed, authors belief in their validity is strongly supported by numerous computer experiments, which gave affirmative results.

1. INTRODUCTION.

It is well known, that leaves of a generic orientable measured foliation on a surface M_g^2 of genus g wind around the surface along one and the same cycle from the first homology group $H_1(M_g^2, \mathbb{R})$ of the surface, which is called asymptotic cycle [Kerck]. In a sense asymptotic cycle gives the first term of approximation of dynamics of leaves. Here we study other terms of approximation. It turns out¹, that taking the next term of approximation we get a two-dimensional subspace in $H_1(M_g^2, \mathbb{R})$, i.e., with a good precision leaves deviate from asymptotic cycle not arbitrary, but inside one at the same two-dimensional subspace in the first homology. Taking further steps $n = 3, \dots, g$ of approximation we get subspaces of dimension k for the k -th step: collection of the subspaces generates a flag of subspaces in the first homology group. The largest,

¹Here and below the presentation is valid modulo conjectures formulated in section 2.

g -dimensional subspace, gives a Lagrangian subspace in $2g$ -dimensional symplectic space $H_1(M_g^2, \mathbb{R})$, with the intersection form considered as a symplectic form. We stop at level g since in a sense at this level we get the best possible approximation — it looks like the error can be in a sense uniformly bounded.

Having a measured foliation generated by a generic closed 1-form on a surface, one can consider interval exchange transformation induced by the first return map on a closed transversal. This interval exchange transformation would be minimal and uniquely ergodic, provided we started from a generic closed 1-form. Our hypothetical approximation is based on several conjectures on asymptotic “spectral properties” of transformation operators $^{(k)}A$ involved in Rauzy induction corresponding to this interval exchange transformation. The conjectures are stated in section 2.

In section 3 we describe behavior of trajectories modulo conjectures on asymptotic “spectral properties” of Rauzy induction.

In section 4 we list some properties of operators $^{(k)}A$ and suggest some speculations on possible proofs of conjectures.

In section 5 we apply general constructions to some particular case arising from an example suggested by I. Dinnikov. This example came from study of Novikov’s problem on electron trajectories on Fermi-surfaces in a weak homogeneous magnetic field. Here closed 1-form under consideration is obtained as a restriction of a specific 1-form on three-dimensional torus with constant coefficients to a specific surface of genus 3 embedded into the torus. Rauzy process in this case is periodic, which simplifies the picture. Besides, unfolding the torus we can “make visible” our trajectories.

In section 6 we present several illustrations for sections of Dinnikov surface.

We wish to thank I. Dinnikov for communicating his example long before it became accessible even as a written text, and to J. Smillie for numerous discussions, and helpful comments.

2. CONJECTURES ON “SPECTRAL PROPERTIES” OF RAUZY INDUCTION.

Consider a minimal uniquely ergodic interval exchange transformation with probability vector $(\lambda^1, \dots, \lambda^n)$ and nondegenerate permutation $\sigma \in \mathfrak{S}_n$. To settle notations we remind construction of Rauzy induction [Rauzy]. Our notations are almost the same as in [Kerck].

Let us describe one step of Rauzy induction. Denote by $I_{i,j}$ square $n \times n$ -matrix, which has only one nonzero entry, which equals one, at the (i, j) place. By E we denote identity $n \times n$ -matrix. Let

$$^{(1)}A = \begin{cases} E + I_{n, \sigma^{-1}(n)} & \text{if } \lambda_n > \lambda_{\sigma(n)}, \\ E + I_{\sigma^{-1}(n), n} & \text{if } \lambda_n < \lambda_{\sigma(n)} \end{cases}$$

Let

$$^{(1)}\lambda = ^{(1)}A^{-1}\lambda$$

Let $\sigma_{\text{dom}} = (1, 2, \dots, n)$ and $\sigma_{\text{im}} = \sigma$.

If $\lambda_n > \lambda_{\sigma(n)}$ modify σ_{im} by cyclically moving forward one step all those entries occurring after the last entry in σ_{dom} , i.e., after $\sigma_{\text{dom}}(n)$. Denote the permutation obtained by $^{(1)}\sigma_{\text{im}}$, and let $^{(1)}\sigma_{\text{dom}} = \sigma_{\text{dom}}$ unchanged. If $\lambda_n < \lambda_{\sigma(n)}$ modify σ_{dom} by cyclically moving forward one step all those entries occurring after the last entry in σ_{im} , i.e., after $\sigma_{\text{im}}(n)$. Denote the permutation obtained by $^{(1)}\sigma_{\text{dom}}$, and let $^{(1)}\sigma_{\text{im}} = \sigma_{\text{im}}$ unchanged. Let

$$^{(1)}\sigma = ^{(1)}\sigma_{\text{dom}}^{-1} \cdot ^{(1)}\sigma_{\text{im}}$$

Here the product of permutations should be understood as a composition of operators, from right to left.

Vector $^{(1)}\sigma_{\text{dom}}^{-1} \left(^{(1)}\lambda \right)$ and permutation $^{(1)}\sigma$ determine a new interval exchange transformation. This interval exchange transformation is just an induction of original interval exchange transformation to subinterval $[0, 1 - \eta[$, where $\eta = \min(\lambda_n, \lambda_{\sigma^{-1}(n)})$. Note, that vector $^{(1)}\lambda$ has L^1 -norm smaller than λ ; we do not renormalize it.

By $^{(k)}\lambda$, $^{(k)}\sigma$, $^{(k)}\sigma_{\text{im}}$, $^{(k)}\sigma_{\text{dom}}$ we denote the data obtained after k steps of Rauzy induction. By $^{(0)}\lambda = \lambda$, $^{(0)}\sigma = \sigma$, $^{(0)}\sigma_{\text{im}} = \sigma$, $^{(0)}\sigma_{\text{dom}} = (1, 2, \dots, n)$ we denote the initial data. By $^{(k)}A$ we denote a product of k elementary matrices corresponding to first k steps of induction, so that

$$(2.1) \quad ^{(0)}\lambda = ^{(k)}A ^{(k)}\lambda$$

or in coordinates

$$(2.2) \quad ^{(0)}\lambda^i = ^{(k)}A_j^i ^{(k)}\lambda^j$$

Recall, that having an interval exchange transformation one can construct a Riemann surface and a closed (harmonic) 1-form, which defines a measured foliation on Riemann surface (see [Masur] and [Veech]). Initial interval exchange transformation would be generated as a first return map to a specific transversal to the foliation. Denote genus of corresponding Riemann surface by g . Though value of g is determined by combinatorics of permutation σ , we referred to construction of Riemann surface to emphasize topological meaning of g , which is rather essential in this paper.

Let $^{(k)}x_1, \dots, ^{(k)}x_n$ be eigenvalues of $^{(k)}A$ enumerated according to decreasing order of their norms: $|^{(k)}x_1| \geq |^{(k)}x_2| \geq \dots \geq |^{(k)}x_n|$.

We formulate propositions and conjectures below everywhere assuming k is sufficiently large, and initial vector λ is generic. We start with reminding a well-known fact, concerning the greatest eigenvalue.

Proposition 1. *The greatest eigenvalue $^{(k)}x_1$ is real and positive: it tends to infinity as k tends to infinity: it is much greater than norms of other eigenvalues*

$$\lim_{k \rightarrow \infty} ^{(k)}x_1 = +\infty$$

$$\lim_{k \rightarrow \infty} \frac{{}^{(k)}x_i}{{}^{(k)}x_1} = 0 \quad \text{for } i = 2, \dots, n$$

In particular ${}^{(k)}x_1$ has multiplicity one. Corresponding eigenvector ${}^{(k)}\lambda_1$ has positive coefficients. Being normalized in L^1 -norm it tends to ${}^{(0)}\lambda$.

$$\lim_{k \rightarrow \infty} {}^{(k)}\lambda_1 = {}^{(0)}\lambda$$

Conjecture 1. Eigenvalues ${}^{(k)}x_1, \dots, {}^{(k)}x_g$ and ${}^{(k)}x_{n-g+1}, \dots, {}^{(k)}x_n$ are all real provided k is sufficiently large.

Conjecture 2. Eigenvalues ${}^{(k)}x_1, \dots, {}^{(k)}x_g$ tend to infinity; their ratios $\frac{{}^{(k)}x_{i+1}}{{}^{(k)}x_i}$ for $i = 1, \dots, g-1$ tend to zero, i.e., ${}^{(k)}x_1 \gg |{}^{(k)}x_2| \gg \dots \gg |{}^{(k)}x_g| \gg 1$

$$\lim_{k \rightarrow \infty} {}^{(k)}x_i = \pm \infty \quad \text{for } i = 1, \dots, g$$

$$\lim_{k \rightarrow \infty} \frac{{}^{(k)}x_{i+1}}{x_i} = 0 \quad \text{for } i = 1, \dots, g-1$$

Conjecture 3. Eigenvalues x_{n-g+1}, \dots, x_n tend to zero; ratios $\frac{x_{i+1}}{x_i}$ for $i = n-g+1, \dots, n-1$ tend to zero, i.e., $1 \gg |x_{n-g+1}| \gg |x_{n-g+2}| \gg \dots \gg |x_n|$

$$\lim_{k \rightarrow \infty} x_i = 0 \quad \text{for } i = n-g+1, \dots, n$$

$$\lim_{k \rightarrow \infty} \frac{x_{i+1}}{x_i} = 0 \quad \text{for } i = n-g+1, \dots, n-1$$

Conjecture 4. Eigenvalues ${}^{(k)}x_{g+1}, \dots, {}^{(k)}x_{n-g}$ can be complex, but with probability p their absolute values are uniformly bounded by a constant $C(g, p)$, for any $p < 1$. (As a probability measure we consider a natural measure on simplex Δ^{n-1} , parametrizing λ .)

$$|{}^{(k)}x_i| \leq C(g, p) \quad \text{for } i = g+1, \dots, n-g$$

In other words

$${}^{(k)}x_{g+1} \sim \dots \sim {}^{(k)}x_{n-g} \sim 1$$

Conjecture 5. Pairwise products of eigenvalues ${}^{(k)}x_i {}^{(k)}x_{n-i+1}$ for $i = 1, \dots, g$ are close to 1. i.e., ${}^{(k)}x_1 {}^{(k)}x_n \sim 1; \dots; {}^{(k)}x_g {}^{(k)}x_{n-g+1} \sim 1$

Note that $\det {}^{(k)}A = 1$, and hence $\prod_{i=1}^n {}^{(k)}x_i = 1$.

Morally we claim, that operator ${}^{(k)}A$ behaves "similar" to a high power of a self-adjoint symplectic operator (cf. example 1).

Consider a flag of subspaces ${}^{(k)}\mathcal{L}^1 \subset {}^{(k)}\mathcal{L}^2 \subset \dots \subset {}^{(k)}\mathcal{L}^g$, where subspace ${}^{(k)}\mathcal{L}^i$, $1 \leq i \leq g$, is spanned by eigenvectors ${}^{(k)}v_1, \dots, {}^{(k)}v_i$ corresponding to "top" i eigenvalues of operator ${}^{(k)}A$. According to Conjecture 1 above, subspace ${}^{(k)}\mathcal{L}^i$, where $1 \leq i \leq g$, is real and has dimension i . Consider this flag as a point of corresponding flag manifold $F_{1,2,\dots,g}(\mathbb{R}^{2g})$.

Conjecture 6. *Flags ${}^{(k)}\mathcal{L}^1 \subset {}^{(k)}\mathcal{L}^2 \subset \dots \subset {}^{(k)}\mathcal{L}^g$ have a limit as $k \rightarrow \infty$ with respect to natural topology on flag manifold.*

Consider much more general problem. Let $f : M \rightarrow M$ be a transitive Anosov diffeomorphism. Let f^* be induced mapping in cohomology. It is known, that the largest by absolute value eigenvalue x_1 of f^* is real, and that $1/x_1$ is also eigenvalue of f^* ; corresponding eigenvectors are called Ruelle—Sullivan classes of f , they are Poincaré dual one to the other.

Problem 1. Does f^* have any other "spectral properties"? Are there any generalizations of Ruelle—Sullivan classes, say, some invariant subspaces in cohomology?

3. HYPOTHETICAL BEHAVIOR OF LEAVES OF ORIENTABLE MEASURED FOLIATION.

Having an interval exchange transformation one can associate to it a Riemann surface and a holomorphic 1-form (see [Masur] and [Veech]), which determines a measured foliation on the surface. By construction we have a specific transversal to the foliation: first return map to this transversal induces initial interval exchange transformation. We may assume, that we started from orientable measured foliation, and then choosing a transversal got interval exchange transformation; in any case, what we are interested in is homological behavior of leaves of corresponding measured foliation.

Recall, that one can associate to each subinterval under exchange a cycle in the first homology group of a surface. The cycle N_i , corresponding to subinterval X_i is represented by the following closed pass on our surface M_g^2 : we start at the left endpoint of the interval X (i.e., at the left endpoint of our transversal), and go to the right along transversal till we get to some point $x \in X_i$ inside subinterval X_i . Then we follow (in positive direction) leaf of foliation starting at the point x till we hit our transversal for the first time. We hit it at the point $T(x)$, where T is our interval exchange transformation. Then we go to the left along interval X till we come back to its left endpoint.

Choose some basis c_1, \dots, c_m in the first homology group of M_g^2 with real coefficients. In fact we do not care, whether it is a basis in absolute or relative homology, so we do not want to specify dimension m precisely. It would be convenient to organize our cycles in a $n \times m$ -dimensional matrix N as follows: row number i of matrix N is just our cycle N_i represented in components N_i^1, \dots, N_i^m with respect to the basis c_1, \dots, c_m .

Let us trace how Rauzy induction affects the cycles N_i . Denote the cycles obtained after k steps of Rauzy induction by ${}^{(k)}N_i$. (Note, that ordering of the subintervals, and hence of the cycles, is determined by permutation ${}^{(k)}\sigma_{\text{dom}}$.) We use initial basis c_1, \dots, c_m in homology to decompose cycles ${}^{(k)}N_i$ in components. It is easy to see, that

$$(3.1) \quad {}^{(k)}N = {}^{(k)}A^T \cdot {}^{(0)}N$$

or in coordinates

$$(3.2) \quad {}^{(k)}N_j^q = {}^{(0)}N_i^q {}^{(k)}A_j^i$$

where index q enumerates components of cycles, and indices i and j enumerate cycles.

Remark 1. We would like to emphasize, that according to transformation rule (3.2) columns of matrix N are transformed as covariant objects with respect to linear transformation defined by matrix ${}^{(k)}A^T$, while vector ${}^{(k)}\lambda$ of lengths ${}^{(k)}\lambda^i$ of subintervals is transformed as a contravariant object with respect to the same linear transformation (c.f. equations (2.1) and (2.2)). In other words, if we consider equation (3.2) as an action of a linear operator ${}^{(k)}R$ with matrix ${}^{(k)}A^T$ on covariant objects, then equations (2.1) and (2.2) define an action of adjoint operator on contravariant objects.

Probably we had to choose operator ${}^{(k)}R$ with matrix ${}^{(k)}R = {}^{(k)}A^T$ as a starting object in our presentation, otherwise “unexpected transposition” leads to some confusion. On the other hand these would lead to contradiction with existing notations in [Kerck] and other papers.

We want to describe now image ${}^{(k)}L = {}^{(k)}A \cdot {}^{(0)}L$ of a generic covariant object ${}^{(0)}L = L$ under our transformation, assuming k is sufficiently large.

Matrix ${}^{(k)}A^T$ of our transformation has the same collection ${}^{(k)}x_1, \dots, {}^{(k)}x_n$ of eigenvalues as ${}^{(k)}A$. According to Conjecture 2 eigenvalues ${}^{(k)}x_1, \dots, {}^{(k)}x_g$ are all distinct. Denote corresponding eigenvectors by ${}^{(k)}\mathbb{V}_1, \dots, {}^{(k)}\mathbb{V}_g$. We have a natural projection to one-dimensional subspaces spanned by these eigenvectors. Denote the projection of a vector L to the subspace spanned by eigenvector ${}^{(k)}\mathbb{V}_i$ by $L|_{{}^{(k)}\mathbb{V}_i}$. Then

$$(3.3) \quad L = L|_{{}^{(k)}\mathbb{V}_1} + \dots + L|_{{}^{(k)}\mathbb{V}_g} + \dots$$

where the tail of decomposition belongs to the invariant subspace corresponding to eigenvalues ${}^{(k)}x_{g+1}, \dots, {}^{(k)}x_n$.

Consider eigen(co)vector ${}^{(k)}W_i$, where $1 \leq i \leq g$, of adjoint operator (having matrix ${}^{(k)}A^{-1}$) corresponding to eigennumber $\frac{1}{{}^{(k)}x_i}$. Note, that it coincides with eigen(co)vector of inverse to adjoint operator (having matrix ${}^{(k)}A$) corresponding to eigennumber ${}^{(k)}x_i$. Normalize our eigenvectors so that under a natural pairing (of covariant and contravariant objects) we get

$$\langle {}^{(k)}W_i, {}^{(k)}W_i \rangle = 1 \quad \text{for any } i = 1, \dots, g$$

Projection $L|_{{}^{(k)}W_i}$ of vector L to subspace generated by vector ${}^{(k)}W_i$ can be expressed now as

$$(3.4) \quad L|_{{}^{(k)}W_i} = \langle L, {}^{(k)}W_i \rangle \cdot {}^{(k)}W_i \quad \text{where } i = 1, \dots, g$$

where \langle, \rangle is again the natural pairing between covariant and contravariant objects. We can rewrite now (3.3) as follows:

$$(3.5) \quad L = \langle L, {}^{(k)}W_1 \rangle \cdot {}^{(k)}W_1 + \dots + \langle L, {}^{(k)}W_g \rangle \cdot {}^{(k)}W_g + \dots$$

where the tail of decomposition belongs to the invariant subspace corresponding to eigennumbers ${}^{(k)}x_{g+1}, \dots, {}^{(k)}x_n$ as before.

Having decomposition (3.5) we can easily describe action of our operator ${}^{(k)}R$ (represented in our coordinates by matrix ${}^{(k)}A^T$).

$$(3.6) \quad {}^{(k)}A^T L = {}^{(k)}x_1 \langle L, {}^{(k)}W_1 \rangle \cdot {}^{(k)}W_1 + \dots + {}^{(k)}x_g \langle L, {}^{(k)}W_g \rangle \cdot {}^{(k)}W_g + O(1)$$

We remind, that according to Conjectures 3 and 4 the tail in (3.6) is small with respect to the leading terms, since projections to eigenvectors ${}^{(k)}W_{n-g+1}, \dots, {}^{(k)}W_n$ would be multiplied by corresponding eigennumbers ${}^{(k)}x_{n-g+1}, \dots, {}^{(k)}x_n$, which tend to zero, while projections to the “middle” eigenvectors ${}^{(k)}W_{g+1}, \dots, {}^{(k)}W_{n-g}$ would be multiplied by eigennumbers, which presumably remain bounded.

Let us use equation (3.6) to rewrite equations (3.1) and (3.2) for the columns ${}^{(k)}N^q$, $q = 1, \dots, m$ of matrix ${}^{(k)}N$.

$$(3.7) \quad {}^{(k)}N^q = {}^{(k)}x_1 \langle {}^{(0)}N^q, {}^{(k)}W_1 \rangle \cdot {}^{(k)}W_1 + \dots + {}^{(k)}x_g \langle {}^{(0)}N^q, {}^{(k)}W_g \rangle \cdot {}^{(k)}W_g + O(1)$$

Consider the following cycles ${}^{(k)}Z_1, \dots, {}^{(k)}Z_g$ in the first homology group (same where cycles N_i live):

$$(3.8) \quad {}^{(k)}Z_i = \langle {}^{(0)}N^1, {}^{(k)}W_i \rangle c_1 + \dots + \langle {}^{(0)}N^m, {}^{(k)}W_i \rangle c_m$$

We are interested, actually, in the rows of matrix ${}^{(k)}N$, representing cycles in the first homology group of our surface. Combining equation (3.7) with definition (3.8) we obtain

$$(3.9) \quad {}^{(k)}N_i = {}^{(k)}x_1 \langle {}^{(k)}W_1, {}^{(k)}Z_i \rangle + \dots + {}^{(k)}x_g \langle {}^{(k)}W_g, {}^{(k)}Z_i \rangle + O(1)$$

We are going to analyze now equation 3.9, which is a key equation in this section.

Recall now, that according to Proposition 1 we have $^{(k)}x_1 \gg |^{(k)}x_i|$ for $i = 2, \dots, n$. Hence the first term of approximation in (3.9) is defined by cycle $^{(k)}Z_1$. This means, that if we will rescale cycles $^{(k)}N_i$ by $1/^{(k)}x_1$ we get

$$(3.10) \quad ^{(k)}N_i = ^{(k)}\mathbb{V}_1^i \cdot ^{(k)}Z_1 + o(1)$$

i.e., cycle $^{(k)}N_i$ is proportional to $^{(k)}Z_1$ with a coefficient of proportionality $^{(k)}\mathbb{V}_1^i$ up to an error, which tends to zero as $k \rightarrow +\infty$. We would like to note that this result is based only on Proposition 1, it does not depend on conjectures, so it is quite rigorous. Still for this case we get nothing new. According to the same Proposition 1 one has

$$\lim_{k \rightarrow +\infty} ^{(k)}\mathbb{V}_1^i = ^{(0)}\lambda$$

Hence (3.8) leads to

$$\lim_{k \rightarrow +\infty} ^{(k)}Z_1 = ^{(0)}\lambda_1 \cdot ^{(0)}N_1 + \dots + ^{(0)}\lambda_n \cdot ^{(0)}N_n$$

i.e., cycle $^{(k)}Z_1$ tends to asymptotic cycle (see [Schw]).

Recall now, that according to Conjecture 2 we have $^{(k)}x_1 \gg \dots \gg |^{(k)}x_g| \gg 1$. Hence if we take leading r terms in approximation (3.9), $1 \leq r \leq g$, we get

$$^{(k)}N_i \approx ^{(k)}x_1 ^{(k)}\mathbb{V}_1^i ^{(k)}Z_1 + \dots + ^{(k)}x_r ^{(k)}\mathbb{V}_r^i ^{(k)}Z_r$$

In other words up to a relatively small error all the cycles belong to a r -dimensional subspace in the first homology group spanned by cycles $^{(k)}Z_1, \dots, ^{(k)}Z_r$. Compare this r -dimensional subspace with one obtained after some other number k' of steps in Rauzy induction. New cycles $^{(k')}Z_1, \dots, ^{(k')}Z_r$ may change, since they are defined in terms of eigen(co)vectors $^{(k')}\mathbb{V}_1, \dots, ^{(k')}\mathbb{V}_r$, which may change. Still, according to Conjecture 6, the space $^{(k')}\mathcal{L}_r$ generated by eigen(co)vectors $^{(k')}\mathbb{V}_1, \dots, ^{(k')}\mathbb{V}_r$ is close to the space $^{(k)}\mathcal{L}_r$ generated by eigen(co)vectors $^{(k)}\mathbb{V}_1, \dots, ^{(k)}\mathbb{V}_r$ in the sense of natural topology of Grassmann manifold $G_r(\mathbb{R}^n)$. Hence (see definition (3.8)) of cycles Z_i) subspaces generated by cycles $^{(k)}Z_1, \dots, ^{(k)}Z_r$ and $^{(k')}Z_1, \dots, ^{(k')}Z_r$ would be also close.


Denote the subspace of the space of first homology of M_g^2 with real coefficients spanned by cycles $^{(k)}Z_1, \dots, ^{(k)}Z_r$ by $^{(k)}\mathcal{H}^r$. We showed that Conjectures 1, 2, 3, 4, and 6 imply the following statement:

Main Conjecture. *Flags $^{(k)}\mathcal{H}^1 \subset ^{(k)}\mathcal{H}^2 \subset \dots \subset ^{(k)}\mathcal{H}^g$ have a limit as $k \rightarrow \infty$ with respect to natural topology on flag manifold.*

We checked this statement by computer experiments with small genera (up to genus 5) using *Mathematica* package ([W]). We used random initial data, and high precision to be able to take approximately a thousand steps in Rauzy induction and compared relative differences in Plucker coordinates. Typical result for the tail of the sequence is 10^{-10} for small genera.

The other obvious computer experiment is as follows. Chose arbitrary two dimensional vectors N_1, \dots, N_n , playing a role of cycles, which satisfy $\sum \lambda^i N_i = 0$. Consider a "trajectory" for some large number of iterations of interval exchange transformations. According to Main Conjecture our "trajectory" is supposed to follow a straight line with direction Z_2 . This hypothetical straight line becomes already visible (see figure 1) starting with 100000 iterations for small genres; for greater values of g and n one has to take more iterations.

FIGURE 1. Computer simulation of "trajectory" for the case, when "asymptotic cycle" equals zero. Initial permutation $\sigma = (6, 5, 3, 8, 7, 4, 2, 1)$ corresponds to a surface of genus 3. Number of iterations is 100.000.



4. PROPERTIES OF OPERATORS ${}^{(k)}A$ AND SOME SPECULATIONS ON POSSIBLE PROOFS OF CONJECTURES.

Remind some properties of operators ${}^{(k)}A$.

Given an interval exchange transformation T corresponding to a pair (λ, σ) , $\lambda \in \mathbb{R}_+^n$, $\sigma \in \mathfrak{S}_n$, set $\beta_0 = 0$, $\beta_i = \sum_{j=1}^i \lambda_j$, and $X_i = [\beta_{i-1}, \beta_i[$. Define skew-symmetric $n \times n$ -matrix $S(\sigma)$ as follows:

$$(4.1) \quad S(\sigma)_{ij} = \begin{cases} 1 & \text{if } i < j \text{ and } \sigma^{-1}(i) > \sigma^{-1}(j) \\ -1 & \text{if } i > j \text{ and } \sigma^{-1}(i) < \sigma^{-1}(j) \\ 0 & \text{otherwise} \end{cases}$$

Consider a translation vector

$$(4.2) \quad \tau = S(\sigma)\lambda$$

Our interval exchange transformation T is defined as follows:

$$T(x) = x + \tau_i, \quad \text{for } x \in X_i, 1 \leq i \leq n$$

To each permutation $\pi \in S^n$ we assign $n \times n$ -matrix which we will denote by $P(\pi)$:

$$(4.3) \quad P(\pi)_{i,j} = \begin{cases} 1 & \text{if } j = \sigma(i), \\ 0 & \text{otherwise} \end{cases}$$

Our first comment is that operators ${}^{(k)}A$ preserve skew-symmetric scalar product $S(\sigma)$ in the following sense (see [N-R]):

$$(4.4) \quad P^T({}^{(k)}\sigma_{\text{dom}})S({}^{(k)}\sigma)P({}^{(k)}\sigma_{\text{dom}}) = {}^{(k)}A^T \cdot S({}^{(0)}\sigma) \cdot {}^{(k)}A$$

In particular for those values of k , when ${}^{(k)}\sigma = {}^{(0)}\sigma$ and ${}^{(k)}\sigma_{\text{dom}} = {}^{(0)}\sigma_{\text{dom}}$ equation (4.4) simplifies as follows:

$$(4.5) \quad S({}^{(0)}\sigma) = {}^{(k)}A^T \cdot S({}^{(0)}\sigma) \cdot {}^{(k)}A$$

i.e., for those values of k operators ${}^{(k)}A$ preserve “degenerate symplectic form” $S({}^{(0)}\sigma)$.

The other comment concerns kernels of operators $S({}^{(k)}\sigma)$ (see (4.2)). Recall construction of a Riemann surface and a measured foliation on the surface corresponding to a given interval exchange transformation (see [Masur] and [Veech]). Due to this construction our initial interval exchange transformation $({}^{(0)}\sigma, {}^{(0)}\lambda)$ is represented as a first return map to a transversal generated by the measured foliation. Enumerate saddles P_1, P_2, \dots, P_s on our surface. Assign to each endpoint of subintervals X_i , $i = 1, \dots, n$ under exchange corresponding saddle. To each saddle point P assign a vector $K \in \mathbb{R}^n$ as follows:

$$(4.6) \quad K^j = \begin{cases} 1 & \text{if } P \text{ is assigned to the left endpoint of } X_j, \\ -1 & \text{if } P \text{ is assigned to the right endpoint of } X_j, \\ 0 & \text{otherwise} \end{cases}$$

We got s vectors K_1, \dots, K_s corresponding to saddles P_1, \dots, P_s .

Proposition 2. *Vectors K_i , $i = 1, \dots, s$ belong to the kernel of operator $S(\sigma)$, i.e.,*

$$S(\sigma)K_i = 0.$$

Kernel of operator $S(\sigma)$ has dimension $s-1$: it coincides with a linear span of vectors K_1, \dots, K_s .

Since a step of Rauzy induction can be considered as induction to a proper subinterval of the transversal of the first return map, we get a natural identification of saddles corresponding to interval exchange transformations $(^{(k)}\sigma, ^{(k)}\lambda)$. Consider vectors $^{(k)}K_i, i = 1, \dots, s$ corresponding to interval exchange transformation obtained after k steps of Rauzy induction.

Proposition 3. *Operator $^{(k)}A$ maps vector $^{(k)}K_i$ to vector $^{(0)}K_i$:*

$$^{(k)}A(^{(k)}K_i) = ^{(0)}K_i \quad \text{for } i = 1, \dots, s.$$

Construction of a Riemann surface in [Masur] and [Veech] by given interval exchange transformation in fact provides us with a natural basis in the first relative (co)homology of the surface with respect to subset of saddle points. Recall, that a measured foliation in this construction is obtained as a foliation of leaves of a closed 1-form. Note, that values λ_i represent integrals over the basic relative 1-cycles. Note also, that values τ_i of the translation vector (4.2) represent integrals of the 1-form over cycles N_i (see previous section). Consider the following terms of exact sequence of a pair (set of saddle points) \subset (Riemann surface M_g^2):

$$\dots \rightarrow H^0(\text{saddles}; \mathbb{R}) \rightarrow H^1(M_g^2, \{\text{saddles}\}; \mathbb{R}) \rightarrow H^1(M_g^2; \mathbb{R}) \rightarrow H^1(\text{saddles}; \mathbb{R}) = 0$$

Under identification with cohomology suggested above, mapping (4.2) can be considered as a mapping from relative to absolute cohomology from the exact sequence of the pair, while the set of vectors K_i defined by (4.6) represents image of the mapping $H^0(\text{saddles}; \mathbb{R}) \rightarrow H^1(M_g^2, \{\text{saddles}\}; \mathbb{R})$. Moreover, under identification of our space (where vector λ lives) with the first cohomology of the surface, skew symmetric matrix $S(\sigma)$ represents intersection form on (co)homology.

* * *

To complete this section we want to suggest several ideas on possible proofs of conjectures.

We start with discussion of Conjecture 1. We suspect, that one can associate with operator $^{(k)}A$ an automorphism of the Riemann surface, such that induced linear mapping in the first cohomology group would be described by operator $^{(k)}A$. As a possible mapping one can take proper pseudo Anosov map from [Veech] or some relative of it, say mapping \mathcal{S} from ([Veech], (7.19)). We suspect, that corresponding operator in cohomology is self-adjoint with respect to some natural scalar product in cohomology. As a candidate for such scalar product we can suggest the following one.

Note that construction in [Masur] and [Veech] provides us with a complex structure, and hence with a metric on the surface. To define a product of two first cohomology classes consider (uniquely determined) harmonic representatives of these classes. In the presence of metric one has a natural pointwise pairing of vector fields, and differential forms as well. Consider a scalar function on the surface obtained as a pointwise pairing of our harmonic representatives. Integrate this function over the surface with respect to the natural volume element, defined by the metric. Define the result to be a value of the scalar product of initial cohomology classes. The bad thing in this construction is that naive metric $dz d\bar{z}$ is singular in our case.

Conjecture 5 should be related to equation (4.4). In particular exact equality

$${}^{(k)}x_i = 1/{}^{(k)}x_{n-i} \quad \text{for } i = 1, \dots, g$$

generically should be valid for infinite subsequence of values of k , for which one has ${}^{(k)}\sigma = {}^{(0)}\sigma$ and ${}^{(k)}\sigma_{\text{dom}} = {}^{(0)}\sigma_{\text{dom}}$ since for these values of k operators ${}^{(k)}A$ are "symplectic" (see (4.5)).

Conjecture 4 presumably is related to Propositions 2 and 3.

In the next section we illustrate how our conjectures work for the easiest case, when Rauzy process is periodic. We hope, that in general, quasiperiodic case, the whole picture is similar.

5. ELECTRON TRAJECTORIES IN DINNIKOV'S EXAMPLE.

In this section we want to illustrate ideas of section 3 by treating a particular measured foliation. On the one hand the structure of Rauzy induction is very easy for this case. On the other hand this example has some independent interest since it came from the framework of S. Novikov problem on behavior of electron trajectories on a Fermi-surface in the presence of a weak homogeneous magnetic field (see [Nov82], [Nov91], [Zorich], and [Dinn1]).

We remind briefly mathematical formulation of initial problem ([Nov82], [Nov91]). Let $\hat{M}_g^2 \subset \mathbb{R}^3$ be a periodic surface in \mathbb{R}^3 , i.e., a surface invariant under translations of cubic lattice in \mathbb{R}^3 . Consider its intersection lines with a plane $ax + by + cz = \text{const}$. What can one say about behavior of these lines? S. Novikov conjectured, that generically nonclosed curves as defined go along a straight line in the plane "from negative infinity to positive infinity".

It was proved in [Zorich], that for a fixed embedding conjecture is valid for an open dense set of directions of planes (union of neighbourhoods of rational directions). For this set of directions all curves can not deviate too far from the lines along which they go — they all belong to stripes of finite width. Paper [Dinn1] assumes that our surface is a level surface of a periodic function, and proves that for any fixed direction of a plane the same behavior of curves is valid for all but at most one level of the

function. There is an example due to S.Tzarev, when Novikov's conjecture is not valid.

We need to reformulate the problem as follows. Consider a closed surface M_g^2 of genus g ("Fermi-surface") embedded into a three-dimensional torus T^3 . We identify torus T^3 with the space \mathbb{R}^3 factored over a cubic lattice. Having a closed 1-form with constant coefficients $a dx + b dy + c dz$ on T^3 one can confine it to the surface. One gets a closed 1-form on the surface, which generically has nondegenerate singularities. This 1-form determines a measured foliation on M_g^2 . Consider universal covering $\mathbb{R}^3 \rightarrow T^3$ and induced covering $\hat{M}_g^2 \rightarrow M_g^2$. Consider leaves of induced measured foliation on the surface \hat{M}_g^2 . By construction they can be obtained as intersection lines of \hat{M}_g^2 with a plane $ax + by + cz = \text{const}$.

Generically measured foliation on a surface obtained by construction above splits into several minimal components (tori with holes). For a long time it was not known whether one can get in this way a minimal foliation. We can assume, that homological class of a surface is equal to zero in the second homology of torus (the case when it is nonzero is trivial). Hence, due to a remark by J.Smillie, the image of asymptotic cycle of foliation equals zero in the first homology of torus. This means that curves in \mathbb{R}^3 obtained by unfolding of leaves of a minimal uniquely ergodic foliation do not have any natural asymptotic direction. Hence examples of minimal foliations in this problem could lead to quite peculiar behavior of leaves.

A family of examples of minimal measured foliations on a surface of genus 3 as required was recently constructed in [Dinn2]. One of the tools in the construction is a process similar to Rauzy induction. We treat the case, when this process is periodic. Parameters, determining the surface, and the slope of the plane are obtained as components of an eigenvector of the transformation matrix D (which is morally similar to matrix A in Rauzy induction) corresponding to a period of the process.

Remark 2. We want to make a following side remark. The space of interval exchange transformations arising from foliations determined by closed 1-forms on a surface of genus g has dimension $4g - 4$. Dimension of a subspace, which comes from Dinnikov construction is $2g - 1$. It follows from the construction, that there are open sets (in topology of the subspace), for which interval exchange transformation is always nonminimal, which gives an estimate for dimension of stratum of nonminimal interval exchange transformation in the space of all interval exchange transformations.

We chose a transversal on Dinnikov surface and considered interval exchange transformation induced by foliation. In this example we have a surface of genus $g = 3$, the 1-form has $2g - 2 = 4$ saddles, so we have interval exchange transformation of $n = 4g - 3 = 9$ intervals. One can easily evaluate cycles N_1, \dots, N_9 (see construction in section 3). It would be convenient for us to consider images of these cycles in $H_1(T^3; \mathbb{R})$, so we will identify cycles N_i with vectors in \mathbb{R}^3 .

For completeness of presentation we display numerical data for this example: interval exchange transformation has permutation

$$\sigma = (3, 8, 5, 2, 7, 4, 9, 1, 6)$$

and vector

$$\lambda \approx (0.558, 2.871, 1.227, 1.558, 0.700, 0.368, 2.730, 0.558, 0.141).$$

Matrix N of cycles given in natural coordinates in $H_1(T^3; \mathbb{R})$ is as follows:

$$(5.1) \quad N = \begin{pmatrix} -1 & -2 & -6 \\ -1 & 0 & -1 \\ 0 & 1 & 2 \\ 0 & -1 & -2 \\ 0 & 1 & 3 \\ 0 & -1 & -2 \\ 1 & 0 & 1 \\ 1 & 2 & 5 \\ 1 & 0 & 0 \end{pmatrix}$$

Having such data it is easy to get computer pictures for the leaves of our foliation (unfolded in \mathbb{R}^3). Figure 2 illustrates a piece of curve obtained by random choice of initial point.


It is easy to see, that the leaf goes rather close to a straight line. Still one should not think, that our leaf just goes straight in one direction — it walks along the line to and fro many times (see section 6 for more details). We stress once more, that such behavior of the leaf can not be explained by means of asymptotic cycle which is equal to zero in the first homology of the torus.

The “straight line” behavior of leaves immediately follows from our Main Conjecture in the end of section 3. Consider images of the subspaces $\mathcal{H}^1, \mathcal{H}^2, \mathcal{H}^3$ in the first homology $H_1(T^3; \mathbb{R})$ of the torus. We know, that asymptotic cycle, which spans \mathcal{H}^1 maps to zero. Hence the image of \mathcal{H}^2 is a one-dimensional subspace in $H_1(T^3; \mathbb{R})$ (unless it also maps to zero, which is not the case in our example). This one-dimensional subspace gives the direction of the line, which one sees at figure 2. One can also check, that two-dimensional image of \mathcal{H}^3 coincides with the plane $ax + by + cz = 0$.

Fortunately Rauzy process for interval exchange transformation in our example is so simple, that we can prove all conjectures in this particular case.

Example 1. Consider interval exchange transformation corresponding to first return map to a transversal in Dinnikov’s example. Under specific choice of transversal one has the following picture. After 12 steps the procedure starts to go cyclically with a period 162. Here is the list of eigennumbers of the matrix $A_{\text{cycle}} = {}^{(12)}A^{-1} \cdot {}^{(174)}A$ corresponding to a cycle in Rauzy induction: $x_1 \approx 25520$, $x_2 \approx 1260$, $x_3 \approx$

FIGURE 2. A piece of leaf after 100 000 returns to the transversal. Unit of measurement is one unit of our cubic lattice. Starting point is at the origin.



20. $x_4 = x_5 = x_6 = 1$. $x_7 \approx 0.05$. $x_8 \approx 0.0008$, $x_9 \approx 0.00004$. Taking a large power of this matrix one gets a picture as in Conjectures above.

We checked cyclic behavior of Rauzy induction in this example as follows: having initial data from Dinnikov process we got approximate initial data for interval exchange transformation with precision sufficient to be sure in first several hundred of steps. Then using computer we generated Rauzy process for our data, and got information on probable length of cycle (162) and number of starting steps (12) before going cyclically. We calculated corresponding matrices $^{(12)}A$ and $^{(174)}A$; these matrices are integer, so they were calculated precisely. Then we checked that these integer matrices obey some algebraic equation containing matrix D of period of Dinnikov process, which proved that interval exchange transformation obtained from periodic point in Dinnikov process gives periodic point (with period 162) in Rauzy induction. Unfortunately we do not see any mapping or any other direct relations between Dinnikov process and Rauzy induction, though morally they represent on and the same process (it was noticed by J. Smillie). In particular we can not prove in general, that periodic Dinnikov process generates periodic process in Rauzy induction.

Let us give some explanation of the properties of eigennumbers of matrix A_{cycle} .

For simplicity take $^{(12)}\lambda$ and $^{(12)}\sigma$ as initial data. Then Rauzy process would be purely cyclic with period 162, i.e.,

$$(5.2) \quad {}^{(162)}\sigma = {}^{(0)}\sigma$$

$$(5.3) \quad {}^{(162)}\sigma_{\text{dom}} = {}^{(0)}\sigma_{\text{dom}}$$

$$(5.4) \quad {}^{(0)}\lambda = {}^{(162)}A \cdot {}^{(162)}\lambda = {}^{(162)}x_1 \cdot {}^{(162)}\lambda$$

i.e., λ is exactly the eigenvector of $^{(162)}A$ corresponding to largest eigennumber $^{(162)}x_1$.

Consider matrix $S = {}^{(0)}S$ defined by (4.1). Due to (5.2) $^{(162)}S = S$, and due to (5.3) change of coordinates (4.3) determined by $^{(162)}\sigma_{\text{dom}}$ is trivial — it is identity matrix. Hence in our case equation (4.4) simplifies as follows:

$$(5.5) \quad S = ({}^{(162)}A)^T \cdot S \cdot {}^{(162)}A$$

It means that transformation $^{(162)}A$ preserves three-dimensional kernel of operator S (see proposition 2). Moreover, due to (5.2), (5.3), and using proposition 3 we see, that operator $^{(162)}A$ acts on the space $\text{Ker} S$ as identity mapping. This way we get three unity eigenvalues $x_4 = x_5 = x_6 = 1$ (cf. Conjecture 4).

We have a well-defined action of operator $^{(162)}A$ on the quotient space $\mathbb{R}^9/\text{Ker} S$, since we factorize over invariant subspace. On the quotient space we have skew-symmetric bilinear form, which comes from skew-symmetric bilinear form on \mathbb{R}^9 determined by matrix S . On the quotient space our bilinear form is already nondegenerate, and according to (5.5) we get a symplectic operator on this six-dimensional vector space. This explains why $x_1 = 1/x_9$, $x_2 = 1/x_8$, $x_3 = 1/x_7$ (cf. Conjecture 5).

Taking powers of matrix $^{(162)}A$ we will get a picture of distribution of eigennumbers as in Conjectures 2 and 3.

Let us discuss behavior of flags $^{(k)}\mathcal{L}^1, {}^{(k)}\mathcal{L}^2, {}^{(k)}\mathcal{L}^3$. It is easy to see, that for $k_q = 162 \cdot q$ we have

$${}^{(k_1)}\mathcal{L}^i = {}^{(k_2)}\mathcal{L}^i = \dots \stackrel{\text{def}}{=} \mathcal{L}^i \quad \text{for } i = 1, 2, 3$$

Consider some intermediate k , say, $k = 162 \cdot q + r$, where $0 < r < 162$. Then ${}^{(k)}A = {}^{(k_q)}A \cdot {}^{(r)}A$. Note, that ${}^{(r)}A$ is nondegenerate operator. Since we have a finite number of possible values for r , we can get any uniform estimates for action of ${}^{(r)}A$, so morally we can consider this operator as a “small perturbation of identity operator” with respect to “significant” operator ${}^{(k_q)}A$ (assuming k_q is sufficiently large).

More precisely we can express this idea as follows. Suppose we have a linear projection operator $P : X \rightarrow X$ on a finite-dimensional vector space X , which maps the whole space to some invariant subspace $Y \subset X$, i.e., $\text{Im}(P) = Y$, and $P(Y) = Y$. Let Q be an automorphism of the vector space X . Then composition $P \cdot Q$ (first apply Q , then P) is again projection to the subspace Y , i.e. $\text{Im}(P \cdot Q) = Y$, and for almost all automorphisms Q one has $(P \cdot Q)(Y) = Y$.


Morally operator ${}^{(k_q)}A$ acts as a projection P to the subspace \mathcal{L}^i for $i = 1, 2, 3$ depending how many steps (1, 2, or 3) of approximation we want to consider, while operator ${}^{(r)}A$ plays a role of automorphism Q . This idea can be easily formalized in our case, which implies that intermediate subspaces ${}^{(k_q+r)}\mathcal{L}^i$, where $i = 1, 2, 3$ converge to \mathcal{L}^i as q tends to infinity.

6. APPENDIX. SECTIONS OF DINNIKOV SURFACE

This is just to present several illustrations to section 5. Consider a section of Dinnikov surface in \mathbb{R}^3 by a plane $ax + by + cz = \text{const}$, where coefficients a, b, c are as in section 5. Consider a square in the (x, y) plane with a side d . Cut a

FIGURE 3. Slice of Dinnikov surface.

Accessible area is 40x40 units



parallelogram from the plane $ax + by + cz = \text{const}$ which projects to our square under projection along z -axes. A piece of section of Dinnikov surface which got into our parallelogram splits into several connected components. Take one of them. Here we present two pictures of such components for different values of d (we measure d in terms of units of our lattice). It would not be interesting to show the whole picture for large values of d . Since our components are just unions of pieces of trajectories,


we would see just a straight line for large values of d . Figure 4 demonstrates only a small part of the whole picture, as if we use a zoom.

Problem 2. It would be rather interesting to know, how many connected components has a generic section of Dinnikov surface: two, finite number, or countable number?

The picture presented is schematic — it is represented by a plane graph. The actual picture is obtained by replacement of edges of the graph by thin ribbons, and by proper conjugation of the ribbons near the vertices.

FIGURE 4. Slice of Dinnikov surface.

Accessible area is 500x500 units
 Graphic area is 50 units
 Number of vertices: 6869
 Number of branches: 1858
 Starting point: $t = 1.763092$ at interval 3



The second picture illustrates, that our trajectories may “wonder along the line” in a quite complicated way. Lacunas in the graph would be filled up after enlarging the size of the rectangle under consideration. But the picture shows, that trajectories have to go far enough before they come back and fill up the lacunas.

REFERENCES

- [Anos] D.V. Anosov, *How curves on the universal covering plane that cover nonselfintersecting curves on a closed surface can go to infinity*, Proc. of the Steklov Inst. of Math., **191:2** (1989), 35–46.
- [Arn92] V.I. Arnold, *Problems on singularities and dynamical systems*, Progress in Soviet Math., Chapman and Hall, 1992.
- [CFS] I.P. Cornfeld, S.V. Fomin, Ya.G. Sinai, *Ergodic Theory*, Springer-Verlag, 1982.
- [Dinn1] I.A. Dinnikov, *Proof of Novikov's conjecture on semiclassical motion of an electron*.
- [Dinn2] I.A. Dinnikov, to appear.
- [FLP] A. Fathi, F. Laudenbach, V. Poénaru. *Travaux de Thurston sur les surfaces*, Astérisque **66–67** (1979).
- [Fried] D. Fried, *Anosov foliations and cohomology*, Ergod. Th & Dynam. Sys. **6** (1986), 9–16; **8** (1988), 491–492.
- [Katok] A.B. Katok, *Invariant measures of flows on oriented surfaces*, Soviet Math. Dokl. **14** (1973), 1104–1108.
- [Keane] M. Keane *Interval exchange transformations*, Math. Z. **141** (1975), 21–31.
- [Kerck] S.P. Kerckhoff, *Simplicial systems for interval exchange maps and measured foliations*, Ergod. Th. & Dynam. Sys. **5** (1985), 257–271
- [KN] H.B. Keynes, D. Newton, *A “minimal”, non-uniquely ergodic interval exchange transformation*, Math. Z., **148** (1976), 101–105.
- [Lev1] G. Levitt, *Flots topologiquement transitifs sur les surfaces compactes sans bord: contre-exemples à une conjecture de Katok*, Ergod. Th. & Dynam. Sys., **3** (1983), 241–249.
- [Lev2] G. Levitt, *Feuilletages des surfaces*. These de Doctorat D'Etat, Universite Paris VII, 1983.
- [Masur] H. Masur, *Interval exchange transformations and measured foliations*, Ann. of Math., **115:1** (1982), 169–200.
- [N-R] A. Nogueira, D. Rudolph, *Topological weakly mixing of interval exchange maps*
- [Nov82] S.P. Novikov, *The Hamiltonian formalism and a many-valued analogue of Morse theory*, Russian Math. Surveys, **37:5**, (1982), 1–56.
- [Nov91] S.P. Novikov, *Quasiperiodic structures in topology*, Proc. of the Conference on Topological Methods in Math. , dedicated to 60-th anniv. of J. Milnor, Stony Brook, 1991.
- [Rauzy] G. Rauzy. *Echanges d'intervalles et transformations induites*. Acta Arith. **34** (1979), 315–328.
- [R-S] D. Ruelle, D. Sullivan, *Currents, flows and diffeomorphisms*. Topology **14** (1975), 319–328.
- [Schw] S. Schwartzman, *Asymptotic cycles*, Annals of Mathematics **66** (1957), 270–284
- [SK] Ya.G. Sinai. K.M. Khanin. *Mixing of some classes of special flows over rotation of a circle*, Functional Anal. Appl., **26:3** (1992).
- [Veech] W.A. Veech, *Gauss measures for transformations on the space of interval exchange maps*. Annals of Mathematics **115** (1982), 201–242
- [W] S. Wolfram. *Mathematica: a system for doing mathematics by computer*, Wolfram Research, Inc.; Addison-Wesley Pub. Co., second edition, 1991.
- [Zorich] A. Zorich, *The S.P. Novikov Problem on Semiclassical motion of electron in homogeneous Magnetic Field*, Russian Math. Surveys, **39:5** (1984), 287–288.

Anton ZORICH
 SISSA (ISAS)
 Via Beirut 2–4
 34013 TRIESTE
 (Italie)