

SÉMINAIRE DE THÉORIE SPECTRALE ET GÉOMÉTRIE

URSULA HAMENSTÄDT

Geometric and ergodic properties of the stable foliation

Séminaire de Théorie spectrale et géométrie, tome 13 (1994-1995), p. 55-62

http://www.numdam.org/item?id=TSG_1994-1995__13__55_0

© Séminaire de Théorie spectrale et géométrie (Grenoble), 1994-1995, tous droits réservés.

L'accès aux archives de la revue « Séminaire de Théorie spectrale et géométrie » implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>

GEOMETRIC AND ERGODIC PROPERTIES OF THE STABLE FOLIATION

Ursula HAMENSTÄDT

In 1983 appeared an article of **Lucy Garnett** in the Journal of Functional Analysis ([G]) in which she studies ergodic properties of a foliation \mathcal{F} on a compact manifold N . Principal assumption is that for every $x \in N$ the leaf $\mathcal{F}(x)$ of \mathcal{F} through x is a smoothly immersed submanifold of N depending continuously on $x \in N$ in the C^∞ -topology. (In her paper she only considers smooth foliations, but her arguments immediately carry over to foliations \mathcal{F} satisfying the assumptions just mentioned, see [Y]).

Any smooth Riemannian metric g on N restricts to a leafwise smooth Riemannian metric $g_{\mathcal{F}}$ on the tangent bundle $T\mathcal{F}$ of \mathcal{F} . With respect to this metric the leaves of \mathcal{F} are smooth Riemannian manifolds of bounded geometry. In particular each leaf carries a natural Laplacean, and these Laplaceans group together to define a global second order differential operator $\Delta_{\mathcal{F}}$ on N with continuous coefficients which is leafwise elliptic.

For every $x \in N$ the Laplacean of $g_{\mathcal{F}}$ on the leaf $\mathcal{F}(x)$ of \mathcal{F} through x induces a Brownian motion on $\mathcal{F}(x)$, described by the heat kernel $p(x, y, t)$ ($y \in \mathcal{F}(x), t > 0$) and the Lebesgue measure $\lambda_{\mathcal{F}}$ on $\mathcal{F}(x)$ induced by $g_{\mathcal{F}}$. For each $t > 0$ we now obtain a Borel-probability measure ω_t on N whose support equals the closure $\overline{\mathcal{F}(x)}$ of $\mathcal{F}(x)$ in N by defining

$$\omega_t(A) = \frac{1}{t} \int_0^t \left(\int_A p(x, y, s) d\lambda_{\mathcal{F}}(y) \right) ds.$$

This measure is the time- t -average of the diffusions of the Dirac mass at x . Since $\overline{\mathcal{F}(x)} \subset N$ is compact we can find a sequence $\{t_j\}_j$ such that $t_j \rightarrow \infty$ ($j \rightarrow \infty$) and such that the measures ω_{t_j} converge weakly to a Borel-probability measure ω on $\overline{\mathcal{F}(x)} \subset N$. This measure ω is *stationary* for the process obtained by considering simultaneously all Brownian motions on all leaves of \mathcal{F} .

More precisely, if P^x denotes the Wiener measure on paths defined by Brownian

motion on $\mathcal{F}(x)$ with starting point x then the measure P on the space of paths Ω in N which is defined by $P(A) = \int P^x(A) d\omega(x)$ is invariant under the one-parameter group $\{T^t \mid t \geq 0\}$ of *shift transformations* $T^t\xi(s) = \xi(s+t)$ (see [G] and the survey [Y]).

The stationary measure ω is also called a *harmonic measure* for the operator $\Delta_{\mathcal{F}}$ since it is characterized by the property that $\int \Delta_{\mathcal{F}}(f) d\omega = 0$ for every smooth function f on N . It disintegrates locally into a transversal sum of leaf measures, where almost every leaf measure is a positive harmonic function times the Riemannian leaf measure ([G]).

In contrast to the case of the trivial foliation ($\dim \mathcal{F} = \dim N$) a harmonic measure for $\Delta_{\mathcal{F}}$ needs not be unique. If \mathcal{F} has two distinct compact leaves $\mathcal{F}(x_1), \mathcal{F}(x_2)$ then the normalized Lebesgue measures on these leaves are harmonic measures for $\Delta_{\mathcal{F}}$ which are mutually singular.

However there are nontrivial foliations for which a harmonic measure is unique. Once again, the first example of such a foliation was described by Garnett ([G]).

Namely let M be a compact Riemannian manifold of negative sectional curvature. The *geodesic flow* Φ^t is a smooth dynamical system on the unit tangent bundle T^1M of M generated by the *geodesic spray* X .

There are four Φ^t -invariant Hölder continuous foliations on T^1M with smooth leaves which depend continuously in the C^∞ -topology on the points in T^1M ([S]). These foliations can be described as follows: Let d be a distance on T^1M defined by a smooth Riemannian metric. Then for $v \in T^1M$ the leaf $W^{ss}(v)$ through v of the *strong stable foliation* W^{ss} is the set $\{w \in T^1M \mid d(\Phi^t v, \Phi^t w) \rightarrow 0 \text{ (} t \rightarrow \infty \text{)}\}$. Its tangent bundle TW^{ss} is a Hölder continuous subbundle of TT^1M . The tangent bundle TW^s of the *stable foliation* W^s is $TW^s = \mathbf{R}X \oplus TW^{ss}$, and the *strong unstable foliation* W^{su} (resp. the *unstable foliation* W^u) is the image of W^{ss} (resp. W^s) under the *flip* $v \rightarrow -v$ of T^1M .

The canonical projection $P: T^1M \rightarrow M$ maps each leaf of W^s locally diffeomorphically onto M . Thus the Riemannian metric on M lifts to a Riemannian metric g^s on TW^s which gives rise to a *stable Laplacean* Δ^s along the leaves of W^s whose coefficients as a global operator on T^1M are Hölder continuous.

Garnett showed ([G]) that if M is a surface of constant curvature, then Δ^s admits a unique harmonic measure. However her arguments are also valid for an arbitrary compact negatively curved manifold M , a fact which was explicitly pointed out by Yue ([Y]). Ledrappier gave independently a proof using the same arguments ([L2]).

The above considerations indicate that the structure of the convex compact space of harmonic measures for a leafwise Laplacean $\Delta_{\mathcal{F}}$ reflects ergodic properties of the foliation \mathcal{F} , but it might not depend in a sensitive way on the Riemannian metric on N used to define the operator $\Delta_{\mathcal{F}}$. Some additional evidence for this was given by Kaimanovich ([K]). To formulate his result, recall that a *completely invariant* transverse measure for a foliation \mathcal{F} is a measure defined on transversals T for \mathcal{F} and such that the following holds: If T, T' are transversals, if $\varphi: T \rightarrow T'$ is a homeomorphism such that $\varphi(x) \in \mathcal{F}(x)$ for all $x \in T$ (i.e. φ is defined by sliding T along the leaves of \mathcal{F}) then

φ maps the measure on T to the measure on T' . According to Plante ([Pl]), completely invariant measures exist if \mathcal{F} has sub-exponential growth. Any such invariant transverse measure can be combined with the Lebesgue measure on the leaves of \mathcal{F} to define a finite Borel-measure on N which we call a *completely invariant harmonic measure*. Now Kaimanovich showed the following ([K]):

THEOREM 1. — *If \mathcal{F} has sub-exponential growth, then every harmonic measure ν for $\Delta_{\mathcal{F}}$ is a completely invariant harmonic measure, and ν -almost every leaf of \mathcal{F} is Liouville.*

The arguments of Kaimanovich go as follows: First he induces a notion of entropy for the leafwise diffusion, the so called *Kaimanovich entropy* h_K which depends on the choice of a stationary measure ν . He shows that $h_K = 0$ if and only if almost every leaf of \mathcal{F} is Liouville, i.e. does not admit nonconstant bounded harmonic functions.

If \mathcal{F} is of subexponential growth, then necessarily $h_K = 0$ for every harmonic measure for $\Delta_{\mathcal{F}}$. Now let ν be a harmonic measure for $\Delta_{\mathcal{F}}$ and consider reversal of time of the diffusion induced by $\Delta_{\mathcal{F}}$ with respect to ν . Since $h_K = 0$, this reversal of time coincides with the diffusion itself, and hence ν is a *self-adjoint harmonic measure* for $\Delta_{\mathcal{F}}$, i.e.

$$\int f(\Delta_{\mathcal{F}}u) d\nu = \int u(\Delta_{\mathcal{F}}f) d\nu$$

for all smooth functions f, u on T^1M (compare [H1]). But a self adjoint harmonic measure corresponds to constant harmonic functions on the leaves of \mathcal{F} in the description of harmonic measures by Garnett ([G]) and hence is completely invariant (see [H1] for a detailed discussion).

Consider now the strong stable foliation W^{ss} on T^1M as above, equipped with the restriction g^{ss} of the Riemannian metric g^s on TW^s and the induced *strong stable Laplacean* Δ^{ss} . The foliation W^{ss} is of subexponential growth and by a classical result of Bowen and Marcus ([B-M]) it admits a unique transverse invariant measure (defined by conditionals of unstable manifolds of the *Bowen-Margulis measure* on T^1M). Thus the above considerations imply.

COROLLARY. — *The strong stable Laplacean Δ^{ss} admits a unique harmonic measure.*

In the terminology of Knieper ([Kn]) the harmonic measure for Δ^{ss} is just the *horospherical measure* ν given with respect to a local product structure by $d\nu = d\lambda^s \times d\mu^{su}$ where μ^{su} is a family of conditional measures on strong unstable manifolds for the Bowen-Margulis measure and λ^s is the family of Lebesgue measures on stable manifolds induced by the Riemannian metric g^s . Moreover, if we change the metric on the leaves of W^{ss} to obtain a new leafwise Laplacean $\bar{\Delta}$, then $\bar{\Delta}$ admits again a unique harmonic measure which is self-adjoint and contained in the measure class of ν .

To summarize the above considerations, we see that a leafwise Laplacean on a foliation \mathcal{F} on N of subexponential growth with the additional property that every leaf of \mathcal{F} is dense in N induces a leafwise diffusion process whose ergodic properties are easy to describe and from which we can not hope to extract geometric properties of \mathcal{F} or N .

For foliations of exponential growth, however, the situation is much more complicated and interesting. A principal and relatively well understood example is the stable foliation W^s of a compact negatively curved manifold M . In fact, if the leaf $W^s(v)$ through $v \in T^1M$ does not contain a periodic orbit of the geodesic flow (and there are only countably many such leaves), then $(W^s(v), g^s)$ is isometric to the universal covering \widetilde{M} of M . This means that we can study Brownian motion on \widetilde{M} by studying the diffusion induced by Δ^s on the compact space T^1M .

Recall that Δ^s admits a unique harmonic measure ω , and properties of ω reflect geometric properties of M and \widetilde{M} . One of the first observations in this direction is due to Ledrappier ([L1]); it can be combined with deep results of Benoist, Foulon, Labourie ([B-F-L], [F-L]) and Besson, Courtois, Gallot ([B-C-G]) to show:

THEOREM 2. — *The unique harmonic measure ω for Δ^s is invariant under the geodesic flow if and only if M is locally symmetric.*

If ω is Φ^t -invariant, then ω necessarily coincides with the Lebesgue Liouville measure λ on T^1M . An open question is whether M is locally symmetric if only ω is contained in the Lebesgue measure class.

Following Ledrappier ([L1]), the measure ω can explicitly be constructed as follows:

Denote by Δ the Laplace operator on the universal covering \widetilde{M} of M . The operator Δ is *weakly coercive* in the sense of Ancona ([A]), i.e. there is a number $\varepsilon > 0$ such that $\Delta + \varepsilon$ admits a positive superharmonic function (i.e. a positive function f such that $(\Delta + \varepsilon)(f) \geq 0$). By the results of Ancona ([A]) Δ admits a Green's function and the Martin boundary of Δ can naturally be identified with the *ideal boundary* $\partial\widetilde{M}$ of \widetilde{M} . This means that for every $\xi \in \partial\widetilde{M}$ and every $x \in \widetilde{M}$ there is a unique minimal positive Δ -harmonic function $y \rightarrow K(x, y, \xi)$ on \widetilde{M} with *pole* at ξ which is normalized by $K(x, x, \xi) = 1$. The function $K: \widetilde{M} \times \widetilde{M} \times \partial\widetilde{M} \rightarrow (0, \infty)$ is Hölder continuous.

Now recall that the stable foliation on T^1M lifts to a foliation on $T^1\widetilde{M}$ which we denote again by W^s . This foliation defines a natural closed equivalence relation \sim on $T^1\widetilde{M}$ by writing $v \sim w$ if and only if $w \in W^s(v)$. The ideal boundary $\partial\widetilde{M}$ of \widetilde{M} is then naturally homeomorphic to the quotient $T^1\widetilde{M}/\sim$. In other words, there is a natural projection $\pi: T^1\widetilde{M} \rightarrow \partial\widetilde{M}$ such that for every $\zeta \in \partial\widetilde{M}$ the pre-image $\pi^{-1}(\zeta)$ is a leaf of W^s .

For every $v \in T^1\widetilde{M}$ the restriction to $W^s(v)$ of the canonical projection $P: T^1\widetilde{M} \rightarrow \widetilde{M}$ is a diffeomorphism. Thus for every fixed $x \in \widetilde{M}$ and every $\zeta \in \partial\widetilde{M}$ the gradient of the logarithm of the function $y \rightarrow K(x, y, \zeta)$ lifts to a vector field on $\pi^{-1}(\zeta)$ not depending on the base-point x .

These vector fields group together to a Hölder continuous leafwise smooth section \widetilde{Y} of TW^s over $T^1\widetilde{M}$ which is equivariant under the action of the fundamental group $\pi_1(M)$ of M on $T^1\widetilde{M}$ and hence projects to a Hölder continuous leafwise smooth section Y of TW^s over T^1M .

Let \mathcal{M} be the convex compact space of Φ^t -invariant Borel-probability measures

on T^1M equipped with the weak*-topology. For $\eta \in \mathcal{M}$ denote by h_η the *entropy* of η (see [W]). The *pressure* of a Hölder continuous function f on T^1M is defined by $pr(f) = \sup\{h_\eta - \int f d\eta \mid \eta \in \mathcal{M}\}$. There is a unique measure $\nu_f \in \mathcal{M}$, the so called *Gibbs equilibrium state* of f , such that $h_{\nu_f} = \int f d\nu_f = pr(f)$ (see [W]). The measure ν_f admits a family ν_f^{su} of conditional measures on strong unstable manifolds which transform under the geodesic flow via $\frac{d}{dt}\Phi^t \circ \nu^{su} |_{t=0} = f + pr(f)$.

Let again X be the geodesic spray and let Y be the section of TW^s over T^1M as above. Then the pressure of the function $g^s(X, Y)$ is zero ([L1]) and the unique harmonic measure ω for Δ^s is of the form $d\omega = d\lambda^s \times d\nu^{su}$ where ν^{su} is a family of conditional measures of the Gibbs equilibrium state induced by $g^s(X, Y)$.

For $v \in T^1M$ denote now by P^v the Wiener measure on paths on $W^s(v)$ induced by $\Delta^s |_{W^s(v)}$ with starting point v . Let $\tilde{v} \in T^1\tilde{M}$ be a lift of v to $T^1\tilde{M}$ and let P^x be the Wiener measure on paths on \tilde{M} induced by Brownian motion on \tilde{M} with starting point $x = P\tilde{v}$. If A is a family of paths on $W^s(v)$ starting at v , then A lifts to a unique family \tilde{A} of paths on $W^s(\tilde{v})$ starting at \tilde{v} , and we have $P^x\{Pc \mid c \in \tilde{A}\} = P^v(A)$.

By a result of Prat ([P]), for P^x -almost every path c in \tilde{M} the limit $\lim_{t \rightarrow \infty} c(t)$ exists in $\tilde{M} \cup \partial\tilde{M}$ and is contained in $\partial\tilde{M}$. Thus P^x projects to a *hitting measure* ω^x on $\partial\tilde{M}$ defined by $\omega^x(A) = P^x\{c \mid c(\infty) \in A\}$. The measures ω^x, ω^y for $x, y \in \tilde{M}$ are equivalent and do not have atoms. Moreover the above convergence is with *positive speed*, which means that $\liminf_{t \rightarrow \infty} \frac{1}{t} \text{dist}(c(0), c(t)) > 0$ for P^x -almost every path c .

While the result of Prat is valid for every simply connected Riemannian manifold \tilde{M} of bounded negative curvature, more can be said for the universal covering of a compact space using methods from ergodic theory applied to the diffusion on (T^1M, ω) induced by Δ^s . Namely for $w \in T_x^1\tilde{M}$ let Θ_w be the *Busemann function* at $\pi(w)$ normalized by $\Theta_w(P\dot{w}) = 0$. The lift of Θ_w to $(W^s(w), g^s)$ is a function whose gradient is just the negative $-X$ of the geodesic spray X .

For $v \in T^1M$ denote by $trU(v)$ the trace of the second fundamental form of the horosphere $PW^{ss}(v)$ at Pv , normalized to be positive. Then for every $x \in \tilde{M}, w \in T_x^1\tilde{M}$ and P^x -almost every path c in \tilde{M} the limit $\lim_{t \rightarrow \infty} \frac{1}{t} \Theta_w(c(t))$ exists and equals $l = \int (trU) d\omega$ (see [K], [L1]). This means that the asymptotic escape rate for a typical path c does not depend on c , moreover Brownian motion does not have a preferred escape direction. For the diffusion on T^1M induced by Δ^s this shows that a typical paths follows (roughly) an orbit of the geodesic flow with positive speed, but in the negative direction (recall that the gradient of Θ_w on $W^s(w) \subset T^1\tilde{M}$ is $-X$). We call such a diffusion a *diffusion of positive escape* and say also in short that Δ^s is of *positive escape* with respect to its (unique) harmonic measure ω .

Let now f be a minimal positive harmonic function on \tilde{M} with pole at $\zeta \in \partial\tilde{M}$. The diffusion induced by the operator $\Delta + 2\nabla \log f$ on \tilde{M} is a conditional Brownian motion, and a typical path c satisfies $\lim_{t \rightarrow \infty} c(t) = \zeta$ in $\tilde{M} \cup \partial\tilde{M}$.

The collection of all those diffusions given by all possible positive minimal harmonic functions can be described by the diffusion induced by $\Delta^s + 2\tilde{Y}$ on $T^1\tilde{M}$. The

operator $\Delta^s + 2\tilde{Y}$ projects to the operator $\Delta^s + 2Y$ on T^1M (notations as above), and the diffusion induced by $\Delta^s + 2Y$ can again be studied using ergodic theory on the compact space T^1M . Now a typical path of $\Delta^s + 2Y$ follows a flow line of the geodesic flow with positive speed in the positive direction. We say that this diffusion is of *negative escape* and call $\Delta^s + 2Y$ of *negative escape*. Observe here that this qualitative behaviour may depend on the choice of a harmonic measure for $\Delta^s + 2Y$. One particular harmonic measure for $\Delta^s + 2Y$ is just ω , the harmonic measure for Δ^s . In fact we have ([H1]):

LEMMA. — *The reversal of time of the diffusion induced by Δ^s on (T^1M, ω) is the diffusion induced by $\Delta^s + 2Y$ on (T^1M, ω) .*

The above considerations are valid in a larger context. Let now g be any smooth Riemannian metric on T^1M and denote by Δ the leafwise Laplacean along the stable foliation induced by g . Recall that g induces an isomorphism of TW^s with its dual T^*W^s . Let Z be Hölder continuous section of TW^s which is differentiable along the leaves of W^s and such that its restriction to every leaf of W^s is dual with respect to g to a closed one-form along the leaf. Write $L = \Delta + Z$ and call L *weakly coercive* if there is $v \in T^1M$ such that the restriction of L to $W^s(v) \sim \tilde{M}$ is weakly coercive in the sense of Ancona. Observe that $\Delta^s + 2Y$ is an operator of this type which is weakly coercive.

Call an operator L of this form of *positive escape* resp. *negative escape* if a typical path with respect to every harmonic measure for L follows (roughly) an orbit of the geodesic flow with positive speed in the negative direction (resp. the positive direction). As we have seen, Δ^s is of positive escape, and the fact that $\Delta^s + 2Y$ is of negative escape is contained in the following theorem ([H1]):

THEOREM 3.

- 1) If $\text{pr}(g(X, Z)) > 0$ then $L = \Delta + Z$ admits a unique harmonic measure ν . Moreover L is weakly coercive, of positive escape, and the Kaimanovich entropy h_K of the diffusion induced by L on (T^1M, ν) is positive.
- 2) If $\text{pr}(g(X, Z)) = 0$ then L admits a unique self-adjoint harmonic measure ν . Moreover L is not weakly coercive, of zero escape, and the Kaimanovich entropy of the diffusion induced by L on (T^1M, ν) vanishes.
- 3) If $\text{pr}(g(X, Z)) < 0$ then L is weakly coercive, of negative escape with respect to every harmonic measure ν , and the Kaimanovich entropy vanishes.

In the case $\text{pr}(g(X, Z)) < 0$ a harmonic measure for L needs not be unique; we'll describe an example for this in Theorem 5 below.

Operators of the above type are suitable to study eigenfunctions of the Laplacean Δ on \tilde{M} . Namely let $\delta_0 > 0$ be the bottom of the positive spectrum for Δ . Ledrappier related δ_0 to the *topological entropy* h of the geodesic flow on T^1M ; he showed:

THEOREM 4 [L3]. — $\delta_0 \leq \frac{h^2}{4}$, with equality if and only if M is asymptotically harmonic and hence locally symmetric.

For $\varepsilon > 0$ the operator $\Delta_\varepsilon = \Delta + \delta_0 - \varepsilon$ on \tilde{M} is weakly coercive and hence as before its Martin kernel is a Hölder continuous function $K_\varepsilon: \tilde{M} \times \tilde{M} \times \partial\tilde{M} \rightarrow (0, \infty)$ which

gives rise to a Hölder continuous section ξ_ε of TW^s over T^1M as before.

Let $p(x, y, t)$ be the heat kernel of Δ and let f be a positive solution of the equation $\Delta_\varepsilon = 0$. Then the fundamental solution of the parabolic equation $\frac{\partial}{\partial t} - \Delta - 2\nabla \log f = 0$ equals the function $(x, y, t) \in \widetilde{M} \times \widetilde{M} \times (0, \infty) \rightarrow e^{(\delta_0 - \varepsilon)t} p(x, y, t) f(y) / f(x)$.

In other words, we may study the operator $\Delta + 2\nabla \log f$ without zero order term to find properties of Δ_ε .

Recall the definition of the section $\tilde{\xi}_\varepsilon$ of TW^s over $T^1\widetilde{M}$ ($\varepsilon \in (0, \delta_0]$) from above. For every $v \in T^1\widetilde{M}$ the restriction of $\Delta^s + 2\xi_\varepsilon$ to $W^s(v)$ is an operator of the kind just described. But $\tilde{\xi}_\varepsilon|_{W^s(v)}$ is the gradient of the logarithm of a minimal positive Δ_ε -harmonic function on $W^s(v) \sim \widetilde{M}$ and hence a typical path of the diffusion induced by $\Delta^s + 2\tilde{\xi}_\varepsilon|_{W^s(v)}$ converges as $t \rightarrow \infty$ to the distinguished point $\pi(v) \in \partial\widetilde{M}$ (with positive speed). Thus the operator $\Delta^s + 2\xi_\varepsilon$ on T^1M falls into category 3) in Theorem 3 above. In fact it admits many harmonic measures (see [H2]):

THEOREM 5. — *Let $\bar{\eta}$ a Gibbs equilibrium state of a flip invariant Hölder continuous function on T^1M . Let $\bar{\eta}^{su}$ be a family of conditional measures on strong unstable manifolds for $\bar{\eta}$. Then for every $\varepsilon \in (0, \delta_0)$ the operator $\Delta^s + 2\xi_\varepsilon$ admits a harmonic measure in the measure class of η where $d\eta = d\lambda^s \times d\bar{\eta}^{su}$.*

Fix now a point $w \in T^1\widetilde{M}$ and consider the restriction of $\tilde{\xi}_\varepsilon$ to $W^s(w)$. The operators Δ_ε satisfy a uniform infinitesimal Harnack inequality, independent of $\varepsilon \in (0, \delta]$, and hence there is a sequence $\{\varepsilon_i\} \subset (0, \delta]$ such that $\varepsilon_i \rightarrow 0$ ($i \rightarrow \infty$) and that $\tilde{\xi}_{\varepsilon_i}|_{W^s(w)}$ converge uniformly on compact subsets of $W^s(w) \sim \widetilde{M}$ to a vector field $\tilde{\xi}_0$ on $W^s(w) \sim \widetilde{M}$. Then $\tilde{\xi}_0$ is the gradient of the logarithm of a positive $\Delta_0 = \Delta + \delta_0$ -harmonic function on $W^s(w) \sim \widetilde{M}$. The next theorem says that every positive Δ_0 -harmonic function on \widetilde{M} is in fact a combination of function of this kind; it is contained in [H2]:

THEOREM 6. — *The sections $\tilde{\xi}_\varepsilon$ of TW^s over $T^1\widetilde{M}$ converge uniformly to a section $\tilde{\xi}_0$. The restriction of $\tilde{\xi}_0$ to a leaf $W^s(w)$ is the gradient of the logarithm of a minimal positive Δ_0 -harmonic function on $W^s(w) \sim \widetilde{M}$ with pole at $\pi(w)$. Every minimal positive Δ_0 -harmonic function is of this kind.*

The vector fields $\tilde{\xi}_0$ projects to a section ξ_0 of TW^s over T^1M . The operator $\Delta^s + 2\xi_0$ admits a unique self-adjoint harmonic measure.

The above describes the minimal Martin boundary for Δ_0 ; it can be identified with the ideal boundary $\partial\widetilde{M}$ of \widetilde{M} . We do not know however how the full Martin boundary of Δ_0 looks like. We also do not know whether the Martin topology for the minimal Martin boundary $\partial\widetilde{M}$ of Δ_0 induces on $\partial\widetilde{M}$ a Hölder structure compatible with the usual Hölder structure of $\partial\widetilde{M}$ (which is the case for the Martin boundary of the operators Δ_ε for $\varepsilon > 0$).

References

- [A] A. ANCONA. — *Negatively curved manifolds, elliptic operators and the Martin boundary*, Ann. Math. **125** (1987), 495–536.
- [A-S] M. ANDERSON, R. SCHOEN. — *Positive harmonic functions on complete manifolds of negative curvature*, Journal AMS **4** (1992), 33–74.
- [B-F-L] Y. BENOIST, P. FOULON, F. LABOURIE. — *Flots d'Anosov à distributions stable et instable différentiables*, Journal AMS **4** (1992), 33–74.
- [B-C-G] G. BESSON, G. COURTOIS, S. GALLOT. — *Entropies et rigidités des espaces localement symétrique de courbure strictement négative*, preprint, 1994.
- [B-M] R. BOWEN, B. MARCUS. — *Unique ergodicity for horocycle foliations*, Israel J. of Math. **26** (1977), 43–67.
- [F-L] P. FOULON, F. LABOURIE. — *Sur les variétés asymptotiquement harmonique*, Inventiones Math. **109** (1992), 97–111.
- [G] L. GARNETT. — *Foliations, the ergodic theorem and Brownian motion*, J. Funct. Anal. **51** (1983), 285–311.
- [H1] U. HAMENSTÄDT. — *Harmonic measures for compact negatively curved manifolds*, preprint n°399, SFB 256, Feb., 1995.
- [H2] U. HAMENSTÄDT. — *Positive eigenfunctions on the universal covering of a compact negatively curved manifold*, preprint.
- [H3] U. HAMENSTÄDT. — *Harmonic measures for leafwise elliptic operators along foliations*, Proceedings ECM Birkhäuser (1994), 73–95.
- [K] V. KAIMANOVICH. — *Brownian motion on foliations: Entropy, invariant measures, mixing*, J. Funct. Anal. **22** (1989), 326–328.
- [Kn] G. KNIEPER. — *Spherical and means on compact Riemannian manifolds of negative curvature*, Diff. Geom. & its Appl. **4** (1994), 361–390.
- [L1] F. LEDRAPPIER. — *Ergodic properties of Brownian motion on covers of compact negatively curved manifolds*, Bol. Soc. Mat. Bras. **19** (1988), 115–140.
- [L2] F. LEDRAPPIER. — *Ergodic properties of the stable foliation*, Proceedings Güstrow (1990), Springer LNM **1514** (1992), 131–145.
- [L3] F. LEDRAPPIER. — *Harmonic measures and Bowen Margulis measures*, Israel J. Math. **71** (1990), 275–287.
- [Pl] J. PLANTE. — *Foliations with measure preserving holonomy*, Ann. Math. **102** (1975), 327–361.
- [P] J.-J. PRAT. — *Mouvement Brownien sur une variété à courbure négative*, C. R. Acad. Sci. Paris Sér. I Math. **280** (1975), 1539–1542.
- [S] M. SHUB. — *Global stability of dynamical systems*, Springer, Berlin, 1987.
- [W] P. WALTERS. — *An introduction to ergodic theory*, Springer Graduate Text in Math. **79**, New York, 1982.
- [Y] C.-B. YUE. — *Brownian motion on Anosov foliations and manifolds of negative curvature*, J. Diff. Geo. **41** (1995), 159–183.

Ursula HAMENSTÄDT
 Math. Institut der Universität Bonn
 Beringstrasse 1
 53115 BONN (Germany)