

SÉMINAIRE DE THÉORIE SPECTRALE ET GÉOMÉTRIE

TATAO YAMAGUCHI

Collapsing and soul theorem in three-dimension

Séminaire de Théorie spectrale et géométrie, tome 15 (1996-1997), p. 163-166

http://www.numdam.org/item?id=TSG_1996-1997__15__163_0

© Séminaire de Théorie spectrale et géométrie (Grenoble), 1996-1997, tous droits réservés.

L'accès aux archives de la revue « Séminaire de Théorie spectrale et géométrie » implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>

COLLAPSING AND SOUL THEOREM IN THREE-DIMENSION

Tatao YAMAGUCHI

In this article, we present a recent joint work with Takashi Shioya [SY]. We study the topology of three-dimensional Riemannian manifolds with a lower sectional curvature bound collapsing to an Alexandrov space of lower dimension.

For a positive integer n and $D > 0$, we denote by $\mathcal{M}(n, D)$ the isometry classes of closed n -dimensional Riemannian manifolds M with sectional curvature ≥ -1 and diameter $\leq D$. The Gromov precompactness theorem [G] implies that for any infinite sequence $\{M_i\}$ in $\mathcal{M}(n, D)$ there is a convergent subsequence of M_i , with respect to the Gromov-Hausdorff distance, whose limit is a compact Alexandrov space X of dimension $\leq n$ and curvature ≥ -1 . We now assume that M_i itself converges to X and consider the following:

PROBLEM 1. — *Describe the topology of M_i by using the geometry and topology of X .*

In some extremal cases, the following results are known: Perelman [P1, P2] proved that if $\dim X = n$, then M_i is homeomorphic to X . Fukaya and Yamaguchi [FY] proved that if $\dim X = 0$, then $\pi_1(M_i)$ contains a nilpotent subgroup of finite index. Therefore, it suffices to consider only the cases $1 \leq \dim X \leq n - 1$ for which there are no general results known. For the special case when the singularity is small in a sense, we have the fibration theorem; there exists a locally trivial fibration $M_i \rightarrow X$ with almost non-negatively curved fibre ([Y1], [Y2]). It is also announced in [P3] that if X has no serious singularities (precisely called extremal subsets), there is a homotopical fibration $M_i \rightarrow X$. When $1 \leq \dim X \leq n - 1$ and X has serious singularities, any solution to Problem 1 was

not known. In the work [SY], we completely solve the problem in the case when $n = 3$ and $\dim X = 1, 2$. In [FY], it was obtained that if $n = 3$ and $\dim X = 0$, then some finite cover of M_i is homeomorphic to either a homotopy sphere, $S^1 \times S^2$, T^3 , or a nilmanifold.

We first consider the case $\dim X = 2$. It is known [BGP] that X is a topological manifold possibly with boundary.

From now on, we always assume that all the M_i are orientable.

THEOREM 1. — *If $\dim X = 2$ and X has empty boundary, then M_i is homeomorphic to a Seifert fibred space over X , for which any singular fibre over a point $x \in X$ has type (μ, ν) satisfying $\mu \leq 2\pi/L(\Sigma_x)$.*

Here $L(\Sigma_x)$ is the length of the space of directions Σ_x at x .

We describe the situation of Theorem 1 in more detail. It is easy to see that for each $\epsilon > 0$, the set $S_\epsilon(X)$ of singular points of X at which the spaces of directions have lengths less than $2\pi - \epsilon$ is finite. Note that one can apply the fibration theorem on a compact domain X_ϵ of X away from $X - S_\epsilon(X)$ for a small ϵ . Thus a compact domain $M_i(\epsilon)$ of M_i converging to X_ϵ has the structure of S^1 -fibration over X_ϵ . We proved that each component of the closure of $M_i - M_i(\epsilon)$ is homeomorphic to $S^1 \times D^2$ and it has a Seifert structure which is compatible to the S^1 -bundle structure over $M_i(\epsilon)$.

THEOREM 2. — *If $\dim X = 2$ and X has non-empty boundary, then there is a Seifert fibred space $\text{Seif}_i(X)$ over X such that*

1. M_i is homeomorphic to the union $\text{Seif}_i(X) \cup (\partial X \times D^2)$ glued along their boundaries;
2. any singular fibre of $\text{Seif}_i(X)$ over a point $x \in \text{int } X$ has type (μ, ν) satisfying $\mu \leq 2\pi/L(\Sigma_x)$.

Applying Theorem 1 to a compact domain M'_i of M_i converging to a compact domain X' away from the boundary of X , we see that M'_i is homeomorphic to a Seifert fibred space, $\text{Seif}_i(X)$, over X' . An essential part in the proof of Theorem 2 is to prove that $M_i - M'_i$ is homeomorphic to $\partial X \times D^2$.

As a corollary of Theorem 2, we have the following

COROLLARY 1. — *Let M_i, X , and $\text{Seif}_i(X)$ be as in Theorem 2, and let g and k denote the genus of X and the number of components of ∂X respectively. Then we have the following factorization:*

$$M_i \cong S^3 \# \underbrace{S^2 \times S^1 \# \cdots \# S^2 \times S^1}_{f(g,k)} \# L(\mu_1, \nu_1) \# \cdots \# L(\mu_\ell, \nu_\ell),$$

where

$$f(g, k) = \begin{cases} 2g + k - 1 & \text{if } X \text{ is orientable,} \\ g + k - 1 & \text{if } X \text{ is non-orientable,} \end{cases}$$

and (μ_j, ν_j) , $1 \leq j \leq \ell$, denote the orbit types of singular fibres of $\text{Seif}_i(X)$.

In particular, the set of homeomorphism classes in the sequence $\{M_i\}$ is finite.

Note that a regular fibre of $\text{Seif}_i(X)$ represents the trivial element in $\pi_1(M_i)$. This is the essential reason for the factorization above.

Let us next consider the case when $\dim X = 1$, i.e., X is isometric to either a circle or a closed interval. When X is isometric to a circle, we observe from the fibration theorem ([Y1]) that M_i is a fibre bundle over S^1 whose fibre is S^2 or T^2 . The rest in our study is the case when X is isometric to a closed interval.

THEOREM 3. — *If X is isometric to a closed interval, then M_i is homeomorphic to the gluing of U and V along their boundaries, where U and V are respectively either D^3 , the twisted product $P^2 \tilde{\times} I$, $S^1 \times D^2$ or $\text{Mö} \tilde{\times} S^1$, the twisted product of Möbius band and circle.*

More precisely, we have a decomposition $M_i = B_i \cup A_i \cup C_i$, where A_i is a compact domain converging to a subinterval of X away from the endpoints and B_i and C_i are the components of the closure of the complement of A_i . It follows from the fibration theorem that A_i is homeomorphic to $F_i \times I$, where F_i is either S^2 or T^2 . We proved that if $F_i \simeq S^2$ (resp. $F_i \simeq T^2$), then B_i and C_i are homeomorphic to either D^3 or $P^2 \tilde{\times} I$ (resp. $S^1 \times D^2$ or $\text{Mö} \tilde{\times} S^1$).

Theorem 1 is the most basic result; for the proof of Theorem 2, we need Theorem 1 and for the proof of Theorem 3, we need both Theorems 1 and 2. Here we give an idea of the proof of Theorem 1.

Let p be an (essential) singular point of the two-dimensional Alexandrov surface X without boundary, and $p_i \in M_i$ be such that $p_i \rightarrow p$ under the convergence $M_i \rightarrow X$. We have to show that for a fixed small $r > 0$, $B(p_i, r)$ is homeomorphic to $S^1 \times D^2$ for large i . From the fibration theorem applied to the convergence, $\partial B(p_i, r) \rightarrow \partial B(p, r)$, we see that $\partial B(p_i, r)$ is homeomorphic to T^2 . To get the topological information of collapsed balls $B(p_i, r)$, we use a rescaling argument of the metrics with a suitable choice of base point. In general, we need to retake a base point different from p_i but close to p_i . Let \hat{p}_i be a point of $B(p_i, r)$ which attains a maximum of the averaged distance function from the points of $\partial B(p_i, r)$. Let $\delta_i \rightarrow 0$ be the maximum distance from \hat{p}_i to the critical point set of \hat{p}_i . Then we can prove that the limit (Y, y_0) of a subsequence of $(\frac{1}{\delta_i} M_i, \hat{p}_i)$ has dimension three; namely, no collapsing occurs in this situation. Note that

1. Y is three-dimensional complete open Alexandrov space with nonnegative curvature without boundary which is a topological manifold;
2. we can recover the topology of $B(p_i, r)$ from that of Y .

Now we need a geometry on Y to classify such spaces. Let us denote by S the soul of Y .

THEOREM 4. — *Y is homeomorphic to the normal bundle of S .*

This is a natural extension of the soul theorem ([CG]) to Alexandrov spaces in three-dimension. It should also be noted that the above result does not hold when Y is not a topological manifold and S is a point. Actually we have a more general classification of three-dimensional complete open Alexandrov spaces with nonnegative curvature without boundary which are not necessarily topological manifolds.

Finally, together with the boundary information $\partial B(p_i, r) \simeq T^2$, the above soul theorem implies that S has dimension one and that $B(p_i, r)$ is homeomorphic to $S^1 \times D^2$, as required.

References

- [BGP] YU. BURAGO, M. GROMOV and G. PERELMAN. — *A.D. Alexandrov's spaces with curvature bounded below*, *Uspehi Mat. Nauk* 47 1992, pp. 3–51. English Transl. *Russian Math. Survey* 47 1992, pp. 1–58.
- [CG] J. CHEEGER and D. GROMOLL. — *On the structure of complete manifolds of nonnegative curvature*, *Ann. of Math.*, 96 1972, pp. 413–443.
- [FY] K. FUKAYA and T. YAMAGUCHI. — *The fundamental groups of almost nonnegatively curved manifolds*, *Ann. of Math.*, 136 1992, pp. 253–333.
- [G] M. GROMOV. — *Structure métrique pour les variétés riemanniennes*, rédigé par J. Lafontaine et P. Pansu, Cedic/Fernand Nathan, Paris, 1981.
- [P1] G. PERELMAN. — *A. D. Alexandrov's spaces with curvatures bounded from below II*, preprint.
- [P2] G. PERELMAN. — *Elements of Morse theory on Alexandrov spaces*, *St. Petersburg Math. Jour.*, 5 1994, pp. 207–214.
- [P3] G. PERELMAN. — *Collapsing with no proper extremal subsets*, preprint.
- [SY] T. SHIOYA and T. YAMAGUCHI. — *Collapsing three-manifolds under a lower curvature bound*, in preparation.
- [Y1] T. YAMAGUCHI. — *Collapsing and Pinching under a lower curvature bound*, *Ann. of Math.*, 133 1991, pp. 317–357.
- [Y2] T. YAMAGUCHI. — *A convergence theorem in the geometry of Alexandrov spaces* *Actes de la table ronde de géométrie différentielle en l'honneur de Marcel Berger*, ed. Arthur L. Besse, Collection SMF, Séminaires et Congrès, 1 Société Mathématique de France, 1996, pp. 601–642.

Tatao YAMAGUCHI
 Graduate School of Mathematics
 Kyushu University
 FUKUOKA 812-81 (Japon)
 email: takao@math.kyushu-u.ac.jp