

# Short time existence and uniqueness in Hölder spaces for the 2D dynamics of dislocation densities

A. El Hajj

*Université d'Orléans, Laboratoire MAPMO, Route de Chartres, 45000 Orléans cedex 2, France*

Received 2 February 2009; received in revised form 5 July 2009; accepted 5 July 2009

Available online 10 July 2009

---

## Abstract

In this paper, we study the model of Groma and Balogh [I. Groma, P. Balogh, Investigation of dislocation pattern formation in a two-dimensional self-consistent field approximation, *Acta Mater.* 47 (1999) 3647–3654] describing the dynamics of dislocation densities. This is a two-dimensional model where the dislocation densities satisfy a system of two transport equations. The velocity vector field is the shear stress in the material solving the equations of elasticity. This shear stress can be related to Riesz transforms of the dislocation densities. Basing on some commutator estimates type, we show that this model has a unique local-in-time solution corresponding to any initial datum in the space  $C^r(\mathbb{R}^2) \cap L^p(\mathbb{R}^2)$  for  $r > 1$  and  $1 < p < +\infty$ , where  $C^r(\mathbb{R}^2)$  is the Hölder–Zygmund space.

© 2009 Elsevier Masson SAS. All rights reserved.

## Résumé

Dans ce papier, nous étudions le modèle de Groma et Balogh [I. Groma, P. Balogh, Investigation of dislocation pattern formation in a two-dimensional self-consistent field approximation, *Acta Mater.* 47 (1999) 3647–3654] qui décrit la dynamique des densités de dislocations. Il s'agit d'un modèle bidimensionnel où les densités de dislocations satisfont un système de deux équations de transport. Le champ de vitesse dans ce système est la contrainte de cisaillement du matériau, calculée à partir de l'équation de l'élasticité linéaire. Cette contrainte de cisaillement peut être liée aux densités de dislocations par certaines transformations de Riesz. En se basant sur des estimations de type commutateurs, nous montrons que ce modèle admet une unique solution locale pour toutes données initiales dans  $C^r(\mathbb{R}^2) \cap L^p(\mathbb{R}^2)$  pour  $r > 1$  et  $1 < p < +\infty$ , où  $C^r(\mathbb{R}^2)$  est l'espace Hölder–Zygmund.

© 2009 Elsevier Masson SAS. All rights reserved.

MSC: 54C70; 35L45; 35Q72; 74H20; 74H25

Keywords: Cauchy's problem; Non-linear transport equations; Non-local transport equations; System of hyperbolic equations; Riesz transform; Hölder–Zygmund space; Dynamics of dislocation densities

---

---

E-mail address: [elhajj@cermics.enpc.fr](mailto:elhajj@cermics.enpc.fr).

## 1. Introduction

### 1.1. Physical motivation and presentation of the model

Real crystals show certain defects in the organization of their crystalline structure, called dislocations. These defects were introduced in the Thirties by Taylor [20], Orowan [17] and Polanyi [18] as the principal explanation of plastic deformation of materials at the microscopic scale. Dislocations can move under the action of exterior stresses applied to the material.

Groma and Balogh in [11] considered the particular case where these defects are parallel lines in the three-dimensional space, that can be viewed as points in a plane considering their cross-section.

In this model we consider two types of “edge dislocations” in the plane  $(x_1, x_2)$ . Typically, for a given velocity field, those dislocations of type (+) propagate in the direction  $+\vec{e}_1$  where  $\vec{e}_1 = (1, 0)$  is the Burgers vector, while those of type (−) propagate in the direction  $-\vec{e}_1$ . We refer the reader to the book of Hirth and Lothe [13], for a detailed description of the classical notion in physics of edge dislocations and of the Burgers vector associated to these dislocations.

In [11] Groma and Balogh have considered the case of densities of dislocations. More precisely, this 2D system is given by the following coupled non-local and non-linear transport equations (see Cannone et al. [4, Section 2] for more modeling details):

$$\begin{cases} \frac{\partial \rho^+}{\partial t}(x, t) + u \frac{\partial \rho^+}{\partial x_1}(x, t) = 0 & \text{on } \mathbb{R}^2 \times (0, T), \\ \frac{\partial \rho^-}{\partial t}(x, t) - u \frac{\partial \rho^-}{\partial x_1}(x, t) = 0 & \text{on } \mathbb{R}^2 \times (0, T), \\ u = R_1^2 R_2^2 (\rho^+ - \rho^-). \end{cases} \quad (1.1)$$

The unknowns of this system are the two scalar functions  $\rho^+$  and  $\rho^-$  at the time  $t$  and the position  $x = (x_1, x_2)$ , that we denote for simplification by  $\rho^\pm$ . This term correspond to the plastic deformations in a crystal. Its derivative in the  $x_1$ -direction  $\frac{\partial \rho^\pm}{\partial x_1}$  represents the dislocation densities of type  $(\pm)$ . Physically, these quantities are non-negative. The function  $u$  is the velocity vector field which is equal to the shear stress in the material, solving the equations of elasticity. The operators  $R_1$  (resp.  $R_2$ ) are the 2D Riesz transform associated to  $x_1$  (resp.  $x_2$ ). More precisely, the Fourier transform of these 2D Riesz transforms  $R_1$  and  $R_2$  are given by

$$\widehat{R_k f}(\xi) = \frac{\xi_k}{|\xi|} \hat{f}(\xi) \quad \text{for } \xi \in \mathbb{R}^2, \quad k = 1, 2.$$

The goal of this work is to establish local existence and uniqueness result of the solution of (1.1) when the initial datum

$$\rho^\pm(x_1, x_2, t = 0) = \rho_0^\pm(x_1, x_2) = \bar{\rho}_0^\pm(x_1, x_2) + Lx_1, \quad L \in \mathbb{R} \quad (1.2)$$

with  $\bar{\rho}_0^\pm \in C^r(\mathbb{R}^2) \cap L^p(\mathbb{R}^2)$ , for  $r > 1$ ,  $p \in (1, +\infty)$ , where  $C^r(\mathbb{R}^2)$  is the Hölder–Zygmund space defined in Section 2. The choice  $L > 0$  guarantee the possibility to choose  $\bar{\rho}_0^\pm \in L^p(\mathbb{R}^2)$  such that the assumption is compatible with the non-negativity of  $\frac{\partial \rho^\pm}{\partial x_1}$ . In a particular case where the initial datum is increasing, the global existence of a solution was proved by Cannone et al. [4], using especially an entropy inequality satisfies by the dislocation densities. However, in the case where the initial datum is decreasing, the solutions of system (1.1) can create shocks, like the case in the classical Burgers equation. Therefore, the fundamental issue of uniqueness for global solutions in the general case remains open.

In a particular sub-case of model (1.1) where the dislocation densities depend on a single variable  $x_1 + x_2$ , the existence and uniqueness of a Lipschitz solution was proved by El Hajj et al. in [10] in the framework of viscosity solutions. Also the existence and uniqueness of a strong solution in  $W_{loc}^{1,2}(\mathbb{R} \times [0, +\infty))$  was proved by El Hajj [9] in the framework of Sobolev spaces. For a similar model describing moreover boundary layer effects (see Groma, Csikor and Zaiser [12]), we refer the reader to Ibrahim [14] where a result of existence and uniqueness is established, using the framework of viscosity solutions and also entropy solution for non-linear hyperbolic equations.

Our study of the dynamics of dislocation densities in a special geometry is related to the more general dynamics of dislocation lines. We refer the interested reader to the work of Alvarez et al. [1], for a local existence and uniqueness of some non-local Hamilton–Jacobi equation. We also refer to Barles et al. [2] for some long-time existence results.

### 1.2. Main results

We shall show that the system (1.1) possesses a unique local-in-time solution for any initial datum satisfy (1.2) such that  $\bar{\rho}_0^\pm \in C^r(\mathbb{R}^2) \cap L^p(\mathbb{R}^2)$ , for  $r > 1$  and for  $p \in (1, +\infty)$ . This functional setting allows us to control the velocity field  $u$  in terms of  $\rho^+ - \rho^-$  (see the third line of (1.1)). As we wrote it before, the velocity  $u$  is related to  $\rho^+ - \rho^-$  through the two-dimensional Riesz transforms  $R_1, R_2$ . Riesz transforms do not map  $C^r(\mathbb{R}^2)$  into  $C^r(\mathbb{R}^2)$ , but they are bounded on  $C^r(\mathbb{R}^2) \cap L^p(\mathbb{R}^2)$ , for  $r \in [0, +\infty)$  and for  $p \in (1, +\infty)$ , as we will see later.

For notational convenience, we define the space  $Y_{r,p}$ , for  $r \in [0, +\infty)$  and  $p \geq 1$  as follows

$$Y_{r,p} = \{f = (f_1, f_2) \text{ such that } f_k \in C^r(\mathbb{R}^2) \cap L^p(\mathbb{R}^2), \text{ for } k = 1, 2\},$$

where  $C^r(\mathbb{R}^2)$  is the inhomogeneous Hölder–Zygmund space (see Section 2, for more precise definition). This space is a Banach space endowed with the following norm: for  $f = (f_1, f_2)$

$$\|f\|_{r,p} = \max_{k=1,2} (\|f_k\|_{C^r}) + \max_{k=1,2} (\|f_k\|_{L^p}).$$

In order to avoid technical difficulties, we first consider (see Theorem 1.1) the case  $L = 0$ . Then (see Theorem 1.2) we treat the general case  $L \in \mathbb{R}$ .

**Theorem 1.1** (Local existence and uniqueness, case  $L = 0$ ). Consider the initial data

$$\rho_0 = (\rho_0^+, \rho_0^-) \in Y_{r,p}. \tag{1.3}$$

If  $r > 1$  and  $p \in (1, +\infty)$ , then (1.1) has a unique solution  $\rho = (\rho^+, \rho^-) \in L^\infty([0, T]; Y_{r,p})$ , where the time  $T > 0$  depends only on  $\|\rho_0\|_{r,p}$ . Moreover, the solution  $\rho$  satisfies

$$\rho \in Lip([0, T]; Y_{r-1,p}).$$

In order to prove this theorem, we strongly use the fact that the Riesz transforms are continuous on  $C^r(\mathbb{R}^2) \cap L^p(\mathbb{R}^2)$  for  $r \in [0, +\infty)$ ,  $p \in (1, +\infty)$ . This result ensures that the velocity vector field remains bounded on  $C^r(\mathbb{R}^2) \cap L^p(\mathbb{R}^2)$ . Using this property and some commutator estimates, we can prove that there exists some  $T > 0$  such that the solution  $\rho^n$  of an approached system of (1.1) (see system (4.28) in Section 4.2) is uniformly bounded in  $L^\infty([0, T]; Y_{r,p})$  for  $r > 1$ ,  $1 < p < +\infty$ . Finally, we show that the sequence of the approximate solutions  $\rho^n$  is a Cauchy sequence in  $L^\infty([0, T]; Y_{r-1,p})$ , which gives the local existence and uniqueness of the solution of (1.1).

The next theorem treats the general case  $L \in \mathbb{R}$ .

**Theorem 1.2** (Local existence and uniqueness, case  $L \in \mathbb{R}$ ). Consider Eq. (1.1) corresponding to initial data (1.2), where  $L \in \mathbb{R}$  and  $\bar{\rho}_0 = (\bar{\rho}_0^+, \bar{\rho}_0^-) \in Y_{r,p}$ . If  $r > 1$  and  $1 < p < +\infty$ , then (1.1) has a unique solution  $\rho = (\rho^+, \rho^-) \in L^\infty([0, T]; Y_{r,p})$ , where the time  $T > 0$  depends only on  $L$  and  $\|\bar{\rho}_0\|_{r,p}$ . Moreover,

$$\rho^\pm(x_1, x_2, t) = \bar{\rho}^\pm(x_1, x_2, t) + Lx_1,$$

where

$$\bar{\rho} = (\bar{\rho}^+, \bar{\rho}^-) \in Lip([0, T]; Y_{r-1,p}).$$

**Remark 1.3.** If at the initial time we have  $\frac{\partial \rho^\pm}{\partial x_1}(\cdot, \cdot, t = 0) \geq 0$  two positive quantities, then this remains true for  $0 \leq t \leq T$ , i.e.,  $\frac{\partial \rho^\pm}{\partial x_1} \geq 0$  for all  $(x, t) \in \mathbb{R} \times [0, T]$ .

Related to our analysis in the present paper, we get the following theorem as a by-product.

**Theorem 1.4** (Global existence and uniqueness for linear transport equations). Take  $g_0 \in C^r(\mathbb{R}^2) \cap L^p(\mathbb{R}^2)$  and  $v = (v^1, v^2) \in L^\infty([0, T]; Y_{r,p})$  for all  $T > 0, r > 1$  and  $1 < p < +\infty$ . Then, there exists a unique solution

$$g \in L^\infty([0, T]; C^r(\mathbb{R}^2) \cap L^p(\mathbb{R}^2)) \cap Lip([0, T]; C^{r-1}(\mathbb{R}^2) \cap L^p(\mathbb{R}^2))$$

of the linear transport equation

$$\begin{cases} \frac{\partial g}{\partial t} + v \cdot \nabla g = 0 & \text{on } \mathbb{R}^2 \times (0, T), \\ g(x, 0) = g_0(x) & \text{on } \mathbb{R}^2. \end{cases} \tag{1.4}$$

### 1.3. Organization of the paper

This paper is organized as follows. In Section 2, we recall the characterization of Hölder spaces and gather several important estimates. In particular, the boundedness of Riesz transforms on  $C^r(\mathbb{R}^2) \cap L^p(\mathbb{R}^2)$  is established. Section 3 presents two key commutator estimates (Lemma 3.1). Finally in Section 4, we prove Theorem 1.4 and a basic *a priori* estimate. Then, thanks to this *a priori* estimate, we give in Sections 4.2 and 4.3 the proofs of Theorems 1.1 and 1.2 respectively.

## 2. Some results on Hölder–Zygmund spaces

This is a preparatory section in which we recall some results on Hölder–Zygmund spaces, and gather several estimates that will be used in the subsequent sections. A major part of the following results can be found in Meyer [15] and Meyer and Coifman [16].

We start with a dyadic decomposition of  $\mathbb{R}^d$ , where  $d > 0$  is an integer. To this end, we take an arbitrary radial function  $\chi \in C_0^\infty(\mathbb{R}^d)$ , such that

$$\text{supp } \chi \subset \left\{ \xi: |\xi| \leq \frac{4}{3} \right\}, \quad \chi \equiv 1 \text{ for } |\xi| \leq \frac{3}{4}, \quad \|\chi\|_{L^1} = 1.$$

It is a classical result that, for  $\phi(\xi) = \chi(\frac{\xi}{2}) - \chi(\xi)$ , we have  $\phi \in C_0^\infty(\mathbb{R}^d)$  and

$$\begin{aligned} \text{supp } \phi &\subset \left\{ \xi: \frac{3}{4} \leq |\xi| \leq \frac{8}{3} \right\}, \\ \chi(\xi) + \sum_{j \geq 0} \phi(2^{-j}\xi) &= 1, \quad \text{for all } \xi \in \mathbb{R}^d. \end{aligned}$$

For the purpose of isolating different Fourier frequencies, define the operators  $\Delta_i$  for  $i \in \mathbb{Z}$  as follows:

$$\Delta_i f = \begin{cases} 0 & \text{if } i \leq -2, \\ \chi(D)f = \int \check{\chi}(y)f(x-y)dy & \text{if } i = -1, \\ \phi(2^{-i}D)f = 2^{id} \int \check{\phi}(2^i y)f(x-y)dy & \text{if } i \geq 0, \end{cases} \tag{2.5}$$

where  $\check{\chi}$  and  $\check{\phi}$  are the inverse Fourier transforms of  $\chi$  and  $\phi$ , respectively.

For  $i \in \mathbb{Z}$ ,  $S_i$  is the sum of  $\Delta_j$  with  $j \leq i - 1$ , i.e.

$$S_i f = \Delta_{-1} f + \Delta_0 f + \Delta_1 f + \dots + \Delta_{i-1} f = 2^{id} \int \check{\chi}(2^i y)f(x-y)dy.$$

It can be shown for any tempered distribution  $f$  that  $S_i f \rightarrow f$  in the distributional sense, as  $i \rightarrow \infty$ .

For any  $r \in \mathbb{R}$  and  $p, q \in [1, \infty]$ , the inhomogeneous Besov space  $B_{p,q}^r(\mathbb{R}^d)$  consists of all tempered distributions  $f$  such that the sequence  $\{2^{jr} \|\Delta_j f\|_{L^p}\}_{j \in \mathbb{Z}}$  belongs to  $l^q(\mathbb{Z})$ . When both  $p$  and  $q$  are equal to  $\infty$ , the Besov space  $B_{p,q}^r(\mathbb{R}^d)$  reduces to the inhomogeneous Hölder–Zygmund space  $C^r(\mathbb{R}^d)$ , i.e.  $B_{\infty,\infty}^r(\mathbb{R}^d) = C^r(\mathbb{R}^d)$ . More explicitly,  $C^r(\mathbb{R}^d)$  with  $r \in \mathbb{R}$  contains any function  $f$  satisfying

$$\|f\|_{C^r} = \sup_{j \in \mathbb{Z}} 2^{jr} \|\Delta_j f\|_{L^\infty} < \infty. \tag{2.6}$$

It is easy to check that  $C^r(\mathbb{R}^d)$  endowed with the norm defined in (2.6) is a Banach space.

For  $r \geq 0$ ,  $C^r(\mathbb{R}^d)$  is closely related to the classical Hölder space  $\tilde{C}^r(\mathbb{R}^d)$  equipped with the norm

$$\|f\|_{\tilde{C}^r} = \sum_{|\beta| \leq [r]} \|\partial^\beta f\|_{L^\infty} + \sup_{x \neq y} \frac{|\partial^{[r]} f(x) - \partial^{[r]} f(y)|}{|x - y|^{r-[r]}}. \tag{2.7}$$

In fact, if  $r$  is not an integer, then the norms (2.6) and (2.7) are equivalent, and  $C^r(\mathbb{R}^d) = \tilde{C}^r(\mathbb{R}^d)$ . The proof for this equivalence is classical and can be found in Chemin [7]. When  $r$  is an integer, say  $r = k$ ,  $\tilde{C}^k(\mathbb{R}^d)$  is the space of bounded functions with bounded  $j$ -th derivatives for any  $j \leq k$ . In particular,  $\tilde{C}^1(\mathbb{R}^d)$  contains the usual Lipschitz functions and is sometimes denoted by  $Lip(\mathbb{R}^d)$ . As a consequence of Bernstein’s Lemma (stated below),  $\tilde{C}^r(\mathbb{R}^d)$  is a subspace of  $C^r(\mathbb{R}^d)$ . Explicit examples can be constructed to show that such an inclusion is genuine. In addition, according to Proposition 2.2,  $\tilde{C}^r(\mathbb{R}^d)$  includes  $C^{r+\varepsilon}(\mathbb{R}^d)$  for any  $\varepsilon > 0$ . In summary, for any integer  $k \geq 0$  and  $\varepsilon > 0$ ,

$$C^{k+\varepsilon}(\mathbb{R}^d) \subset \tilde{C}^k(\mathbb{R}^d) \subset C^k(\mathbb{R}^d).$$

**Proposition 2.1** (Bernstein’s Lemma). (See Meyer [15].) *Let  $d > 0$  be an integer and  $0 < \alpha_1 < \alpha_2$  be two real numbers.*

(1) *If  $1 \leq p \leq q \leq \infty$  and  $\text{supp } \hat{f} \subset \{\xi \in \mathbb{R}^d : |\xi| \leq \alpha_1 2^j\}$ , then*

$$\max_{|\beta|=k} \|\partial^\beta f\|_{L^q} \leq C 2^{jk+d(\frac{1}{p}-\frac{1}{q})} \|f\|_{L^p},$$

where  $C > 0$  is a constant depending only on  $k$  and  $\alpha_1$ .

(2) *If  $1 \leq p \leq \infty$  and  $\text{supp } \hat{f} \subset \{\xi \in \mathbb{R}^d : \alpha_1 2^j \leq |\xi| \leq \alpha_2 2^j\}$ , then*

$$C^{-1} 2^{jk} \|f\|_{L^p} \leq \max_{|\beta|=k} \|\partial^\beta f\|_{L^p} \leq C 2^{jk} \|f\|_{L^p},$$

where  $C > 0$  is a constant depending only on  $k$ ,  $\alpha_1$  and  $\alpha_2$ .

**Proposition 2.2** (Inequalities in Hölder–Zygmund space). (See Meyer [15].) *Let  $d > 0$  be an integer. Then, we have following two inequalities:*

(1) *There exists a constant  $C = C(d)$  such that for any  $r > 0$  and  $f \in C^r(\mathbb{R}^d)$ , we have*

$$\|f\|_{L^\infty} \leq \frac{C}{r} \|f\|_{C^r}. \tag{2.8}$$

(2) *There exists a constant  $C = C(d)$  such that for any  $r > 1$  and  $f \in C^{r-1}(\mathbb{R}^d)$ , we have*

$$\left\| \frac{\partial f}{\partial x_k} \right\|_{C^{r-1}} \leq C \|f\|_{C^r} \quad \text{for all } k = 1, \dots, d. \tag{2.9}$$

In the system (1.1), the velocity field  $u$  is determined by  $\rho^+ - \rho^-$  through the 2D Riesz transforms. These Riesz transforms do not map a  $C^r(\mathbb{R}^d)$  Hölder–Zygmund space to itself, but their action on  $C^r(\mathbb{R}^d)$  is indeed bounded in  $C^r(\mathbb{R}^d) \cap L^p(\mathbb{R}^d)$  for  $p \in (1, +\infty)$  (see Proposition 2.4). We first recall a general result concerning the boundedness of Fourier multiplier operators on Hölder spaces.

**Proposition 2.3** (Fourier multiplier operators on Hölder spaces). (See Meyer [15].) *Let  $d > 0$  be an integer and  $F$  be an infinitely differentiable function on  $\mathbb{R}^d$ . Assume that for some  $R > 0$  and  $m \in \mathbb{R}$ , we have*

$$F(\lambda \xi) = \lambda^m F(\xi)$$

for any  $\xi \in \mathbb{R}^d$  with  $|\xi| > R$  and  $\lambda \geq 1$ . Then the Fourier multiplier operator  $F(D)$  maps continuously  $C^r(\mathbb{R}^d)$  into  $C^{r-m}(\mathbb{R}^d)$  for any  $r \in \mathbb{R}$ .

**Proposition 2.4** (Boundedness of Riesz transforms on  $C^r(\mathbb{R}^d) \cap L^p(\mathbb{R}^d)$ ). *Let  $r \in \mathbb{R}$  and  $p \in (1, +\infty)$ . Then there exists a positive constant  $C$  depending only on  $r$  and  $p$  such that*

$$\|R_k f\|_{C^r \cap L^p} \leq C \|f\|_{C^r \cap L^p} \quad \text{for } k = 1, 2,$$

where  $\|\cdot\|_{C^r \cap L^p} = \|\cdot\|_{C^r} + \|\cdot\|_{L^p}$ .

**Proof.** Using the operator  $\Delta_{-1}$  defined in (2.5), we divide  $R_k f$  into two parts,

$$R_k f = \Delta_{-1} R_k f + (1 - \Delta_{-1}) R_k f. \tag{2.10}$$

Since  $\text{supp } \chi(\xi) \cap \text{supp } \phi(2^{-j}\xi) = \emptyset$  for  $j \geq 1$ , the operator  $\Delta_j \Delta_{-1} = 0$  when  $j \geq 1$ . Thus, according to (2.6),

$$\begin{aligned} \|\Delta_{-1} R_k f\|_{C^r} &= \sup_{j \in \mathbb{Z}} 2^{jr} \|\Delta_j \Delta_{-1} R_k f\|_{L^\infty} \\ &= \max[2^{-r} \|\Delta_{-1} \Delta_{-1} R_k f\|_{L^\infty}, \|\Delta_0 \Delta_{-1} R_k f\|_{L^\infty}] \\ &\leq \max[1, 2^{-r}] \|\Delta_{-1} R_k f\|_{L^\infty}. \end{aligned}$$

Let  $q$  be the conjugate of  $p$ , namely  $\frac{1}{p} + \frac{1}{q} = 1$ . It then follow, since Riesz transforms are bounded on  $L^p(\mathbb{R}^d)$ , that for all  $p \in (1, +\infty)$ :

$$\begin{aligned} \|\Delta_{-1} R_k f\|_{C^r} &\leq \max[1, 2^{-r}] \|\check{\chi} * R_k f\|_{L^\infty} \\ &\leq \max[1, 2^{-r}] \|\check{\chi}\|_{L^q} \|R_k f\|_{L^p} \\ &= C \|f\|_{L^p}, \end{aligned}$$

where  $C$  is a constant depending only on  $r$  and  $p$ . To estimate the second part in (2.10), we apply Proposition 2.3 with  $F(\xi) = (1 - \chi(\xi)) \frac{\xi_k}{|\xi|}$  and  $m = 0$ , and hence we conclude that it maps  $C^r(\mathbb{R}^d)$  into  $C^r(\mathbb{R}^d)$ . This gives that Riesz transforms are continuous from  $C^r(\mathbb{R}^d) \cap L^p(\mathbb{R}^d)$  into  $C^r(\mathbb{R}^d)$ , for  $r \in \mathbb{R}$  and  $p \in (1, +\infty)$ . Moreover, using the fact that the Riesz transforms are bounded on  $L^p(\mathbb{R}^d)$ , for all  $p \in (1, +\infty)$ , we terminate the proof of Proposition 2.4.  $\square$

Finally, we recall the notion of Bony’s paraproduct (see Bony [3]). The usual product  $uv$  of two functions  $u$  and  $v$  can be decomposed into three parts. More precisely, using  $v = \sum_{j \in \mathbb{Z}} \Delta_j v$ ,  $u = \sum_{j \in \mathbb{Z}} \Delta_j u$  and

$$\Delta_j \Delta_k v = 0 \quad \text{if } |j - k| \geq 1, \quad \Delta_j (S_{k-1} v \Delta_k v) = 0 \quad \text{if } |j - k| \geq 5,$$

we can write

$$uv = T_u v + T_v u + R(u, v),$$

where

$$T_u v = \sum_{j \geq 1} S_{j-1}(u) \Delta_j v, \quad R(u, v) = \sum_{|i-j| \leq 1} \Delta_i u \Delta_j v.$$

We remark that the previous decomposition allows one to distinguish different types of terms in the product of  $uv$ . The Fourier frequencies of  $u$  and  $v$  in  $T_u v$  and  $T_v u$  are separated from each other while those of the terms in  $R(u, v)$  are close to each other. Using this decomposition, one can show that

$$\|uv\|_{C^s} \leq \|u\|_{C^s} \|v\|_{L^\infty} + \|u\|_{L^\infty} \|v\|_{C^s} \quad \text{for } s > 0. \tag{2.11}$$

For the proof of (2.11) see Chen et al. [8, Proposition 5.1].

### 3. Two commutator estimates

Two major commutator estimates are stated and proved in this section. We remark that this commutator estimates was often used to resolve the Navier–Stokes equations (see for instance Cannone et al. [5,6]). Here, we apply these techniques on our system (1.1).

**Lemma 3.1** ( *$L^\infty$  commutator estimates*). *Let  $j \geq -1$  be an integer and  $r > 0$ . Then, for some absolute constant  $C$ , we have*

$$\begin{aligned}
 (1) \quad & \left\| \left[ u \frac{\partial}{\partial x_\alpha}, \Delta_j \right] f \right\|_{L^\infty} \leq C 2^{-jr} \left( \left\| \frac{\partial f}{\partial x_\alpha} \right\|_{L^\infty} \|u\|_{C^r} + \|\nabla u\|_{L^\infty} \|f\|_{C^r} \right), \quad \text{for } \alpha = 1, 2, \\
 (2) \quad & \left\| \left[ u \frac{\partial}{\partial x_\alpha}, \Delta_j \right] f \right\|_{L^\infty} \leq C 2^{-jr} (\|f\|_{L^\infty} \|u\|_{C^{r+1}} + \|\nabla u\|_{L^\infty} \|f\|_{C^r}), \quad \text{for } \alpha = 1, 2,
 \end{aligned}$$

where the bracket  $[ \cdot, \cdot ]$  represents the commutator, namely

$$\left[ u \frac{\partial}{\partial x_\alpha}, \Delta_j \right] f = u \frac{\partial}{\partial x_\alpha} (\Delta_j f) - \Delta_j \left( u \frac{\partial f}{\partial x_\alpha} \right), \quad \text{for } \alpha = 1, 2. \tag{3.12}$$

**Proof.** (1) Using the paraproduct notations  $T$  and  $R$ , we decompose  $[u \frac{\partial}{\partial x_\alpha}, \Delta_j] f$ , for  $\alpha = 1, 2$ , into five parts,

$$\left[ u \frac{\partial}{\partial x_\alpha}, \Delta_j \right] \rho = I_1 + I_2 + I_3 + I_4 + I_5,$$

where

$$\begin{aligned}
 I_1 &= \left[ T_u \frac{\partial}{\partial x_\alpha}, \Delta_j \right] f = T_u \left( \frac{\partial}{\partial x_\alpha} (\Delta_j f) \right) - \Delta_j \left( T_u \frac{\partial f}{\partial x_\alpha} \right), \\
 I_2 &= -\Delta_j T_{\frac{\partial f}{\partial x_\alpha}}(u), \\
 I_3 &= T_{\frac{\partial(\Delta_j f)}{\partial x_\alpha}}(u), \\
 I_4 &= R \left( u, \frac{\partial(\Delta_j f)}{\partial x_\alpha} \right), \\
 I_5 &= -\Delta_j R \left( u, \frac{\partial f}{\partial x_\alpha} \right).
 \end{aligned}$$

Back to the definition of  $T$ , we can write

$$\begin{aligned}
 I_1 &= \sum_{k \geq 1} S_{k-1}(u) \Delta_k \left( \frac{\partial(\Delta_j f)}{\partial x_\alpha} \right) - \Delta_j \left( \sum_{k \geq 1} S_{k-1}(u) \Delta_k \frac{\partial f}{\partial x_\alpha} \right) \\
 &= \sum_{k \geq 1} \left[ S_{k-1}(u) \Delta_j \left( \frac{\partial(\Delta_k f)}{\partial x_\alpha} \right) - \Delta_j \left( S_{k-1}(u) \frac{\partial(\Delta_k f)}{\partial x_\alpha} \right) \right].
 \end{aligned} \tag{3.13}$$

Since  $\Delta_j \Delta_k = 0$  for  $|j - k| > 1$  and

$$\text{supp} \left[ S_{k-1}(u) \frac{\partial(\Delta_k f)}{\partial x_\alpha} \right] \subset \left\{ \xi : \frac{1}{3} 2^{k-2} \leq |\xi| \leq \frac{5}{3} 2^{k+1} \right\},$$

the sum in (3.13) only involves those terms with  $k$  satisfying  $|j - k| \leq 4$ . We only take  $j \geq 0$  since the case  $j = -1$  can be handled similarly. Applying the definition of  $\Delta_j$  in (2.5), we obtain

$$\begin{aligned}
 I_1 &= \sum_{|j-k| \leq 4} 2^{jd} \int \check{\phi}(2^j(x-y)) [S_{k-1}(u(x)) - S_{k-1}(u(y))] \frac{\partial(\Delta_k f)}{\partial x_\alpha}(y) dy \\
 &= \sum_{|j-k| \leq 4} \int \check{\phi}(y) [S_{k-1}(u(x)) - S_{k-1}(u(x-2^{-j}y))] \frac{\partial(\Delta_k f)}{\partial x_\alpha}(x-2^{-j}y) dy.
 \end{aligned}$$

Using the fact that  $\check{\phi} \in S(\mathbb{R}^d)$  and  $S_j$  are continuous from  $L^\infty$  onto itself, we get for  $r \in \mathbb{R}$  and an absolute constant  $C$ :

$$\begin{aligned} \|I_1\|_{L^\infty} &\leq C2^{-j}\|\nabla u\|_{L^\infty}\left\|\frac{\partial(\Delta_j f)}{\partial x_\alpha}\right\|_{L^\infty} \\ &\leq C\|\nabla u\|_{L^\infty}\|\Delta_j f\|_{L^\infty} \\ &\leq C2^{-jr}\|\nabla u\|_{L^\infty}\|f\|_{C^r}, \end{aligned} \tag{3.14}$$

where we have used Proposition 2.1 in the second inequality. To estimate  $I_2$  and  $I_3$ , we first write them as

$$I_2 = - \sum_{|j-k|\leq 4} \Delta_j \left( S_{k-1} \left( \frac{\partial f}{\partial x_\alpha} \right) \Delta_k u \right), \quad I_3 = \sum_{|j-k|\leq 4} S_{k-1} \left( \frac{\partial(\Delta_j f)}{\partial x_\alpha} \right) \Delta_k u.$$

Similarly, only terms with  $k$  satisfying  $|j - k| \leq 4$  are considered in the above sums. Thus, since  $\Delta_j$  and  $S_j$  are continuous from  $L^\infty$  onto itself, we have for  $r \in \mathbb{R}$ :

$$\begin{aligned} \|I_2\|_{L^\infty} &\leq C \sum_{k_1=-4}^4 \left\| \Delta_j \left( S_{j+k_1-1} \left( \frac{\partial f}{\partial x_\alpha} \right) \Delta_{j+k_1} u \right) \right\|_{L^\infty} \\ &\leq C \sum_{k_1=-4}^4 \left\| S_{j+k_1-1} \left( \frac{\partial f}{\partial x_\alpha} \right) \Delta_{j+k_1} u \right\|_{L^\infty} \\ &\leq C \left\| \frac{\partial f}{\partial x_\alpha} \right\|_{L^\infty} \|u\|_{C^r} \sum_{k_1=-4}^4 2^{-(j+k_1)r} \leq C2^{-jr} \left\| \frac{\partial f}{\partial x_\alpha} \right\|_{L^\infty} \|u\|_{C^r}, \end{aligned} \tag{3.15}$$

$$\begin{aligned} \|I_3\|_{L^\infty} &\leq C \sum_{k_1=-4}^4 \left\| S_{j+k_1-1} \left( \frac{\partial(\Delta_j f)}{\partial x_\alpha} \right) \Delta_{j+k_1} u \right\|_{L^\infty} \\ &\leq C \left\| \frac{\partial f}{\partial x_\alpha} \right\|_{L^\infty} \|u\|_{C^r} \sum_{k_1=-4}^4 2^{-(j+k_1)r} \leq C2^{-jr} \left\| \frac{\partial f}{\partial x_\alpha} \right\|_{L^\infty} \|u\|_{C^r}, \end{aligned} \tag{3.16}$$

where the  $C$ 's in the above inequalities are absolute constants. From the definition of  $R$ , we have

$$I_4 = \sum_{|k_1-k_2|\leq 1, |j-k_2|\leq 1} \left( \Delta_{k_1}(u) \Delta_{k_2} \left( \frac{\partial(\Delta_j f)}{\partial x_\alpha} \right) \right).$$

Obviously, only a finite number of terms involved in the above sums are non-zeros. Then,

$$\|I_4\|_{L^\infty} \leq C2^{-jr} \sup_{j \in \mathbb{Z}} (2^{jr} \|\Delta_j u\|_{L^\infty}) \left\| \Delta_j \frac{\partial f}{\partial x_\alpha} \right\|_{L^\infty} \leq C2^{-jr} \|u\|_{C^r} \left\| \frac{\partial f}{\partial x_\alpha} \right\|_{L^\infty}. \tag{3.17}$$

Note that from the definition of  $R$  and  $\Delta_j$ ,  $j \geq -1$ , we can write  $I_5$  as

$$I_5 = - \sum_{k \geq j-3} \sum_{k_1=k-1}^{k+1} \Delta_j \left( \Delta_k(u) \Delta_{k_1} \left( \frac{\partial f}{\partial x_\alpha} \right) \right).$$

Therefore, for an absolute constant  $C$ , we have

$$\begin{aligned} \|I_5\|_{L^\infty} &\leq C \sum_{k \geq j-3} \sum_{k_1=k-1}^{k+1} \|\Delta_k u\|_{L^\infty} \left\| \Delta_{k_1} \frac{\partial f}{\partial x_\alpha} \right\|_{L^\infty} \\ &\leq C \left\| \frac{\partial f}{\partial x_\alpha} \right\|_{L^\infty} \|u\|_{C^r} \sum_{k \geq j-3} 2^{-kr} \\ &\leq C2^{-jr} \left\| \frac{\partial f}{\partial x_\alpha} \right\|_{L^\infty} \|u\|_{C^r}. \end{aligned} \tag{3.18}$$



Gathering the estimates in (3.14)–(3.18), we establish the desired inequality in (1).

(2) As in the proof of (1), we decompose  $[u \frac{\partial}{\partial x_\alpha}, \Delta_j]f$  as the sum of  $I_1, I_2, I_3, I_4$  and  $I_5$ . The estimate on  $I_1$  remains untouched, while different bounds are needed for  $I_2, I_3, I_4$  and  $I_5$ . Indeed, for  $j \geq 1$ :

$$\begin{aligned} \|I_2\|_{L^\infty} &\leq C \|\Delta_j u\|_{L^\infty} \left\| \frac{\partial S_{j-1} f}{\partial x_\alpha} \right\|_{L^\infty} \\ &\leq C 2^j \|\Delta_j u\|_{L^\infty} \|S_{j-1} f\|_{L^\infty} \\ &\leq C 2^{-jr} \|u\|_{C^{r+1}} \|f\|_{L^\infty}, \end{aligned} \tag{3.19}$$

where we have used Proposition 2.1 in the second inequality.  $I_3$  and  $I_4$  can be similarly estimated as  $I_2$ :

$$\begin{aligned} \|I_3\|_{L^\infty} &\leq C \left\| \frac{\partial \Delta_j f}{\partial x_\alpha} \right\|_{L^\infty} \|\Delta_j u\|_{L^\infty} \leq C 2^j \|\Delta_j u\|_{L^\infty} \|\Delta_j f\|_{L^\infty} \\ &\leq C 2^{-jr} \|u\|_{C^{r+1}} \|f\|_{L^\infty}, \end{aligned} \tag{3.20}$$

$$\begin{aligned} \|I_4\|_{L^\infty} &\leq C \|\Delta_j u\|_{L^\infty} \left\| \frac{\partial \Delta_j f}{\partial x_\alpha} \right\|_{L^\infty} \leq C 2^j \|\Delta_j u\|_{L^\infty} \|\Delta_j f\|_{L^\infty} \\ &\leq C 2^{-jr} \|u\|_{C^{r+1}} \|f\|_{L^\infty}. \end{aligned} \tag{3.21}$$

Finally, we have

$$\begin{aligned} \|I_5\|_{L^\infty} &\leq C \sum_{k \geq j-3} \sum_{k_1=k-1}^{k+1} \|\Delta_{k_1} u\|_{L^\infty} \left\| \frac{\partial \Delta_{k_1} f}{\partial x_\alpha} \right\|_{L^\infty} \\ &\leq C \sum_{k \geq j-3} \sum_{k_1=k-1}^{k+1} 2^{k_1} \|\Delta_{k_1} u\|_{L^\infty} \|\Delta_{k_1} f\|_{L^\infty} \leq C \|f\|_{L^\infty} \sum_{k \geq j-3} 2^k \|\Delta_k u\|_{L^\infty} \\ &\leq C \|f\|_{L^\infty} \|u\|_{C^{r+1}} \sum_{k \geq j-3} 2^{-kr} \leq C 2^{-jr} \|f\|_{L^\infty} \|u\|_{C^{r+1}}. \end{aligned} \tag{3.22}$$

Combining (3.19)–(3.22) yields (2).  $\square$

#### 4. Local existence and uniqueness results

This section is devoted to the proofs of Theorems 1.1 and 1.2. For the sake of a clear presentation, we divide it into three subsections. In the first subsection, we show a basic *a priori* estimate and we prove Theorem 1.4. With the aid of this estimate, we prove Theorems 1.1 and 1.2 in the next subsections.

##### 4.1. An a priori estimate

**Proposition 4.1** (*A priori estimate*). *Let  $r > 1$  and  $p > 1$ . For all  $T > 0$ ,  $\rho_0 = (\rho_0^+, \rho_0^-) \in Y_{r,q}$  and  $u \in L^\infty([0, T]; C^r(\mathbb{R}^2) \cap L^p(\mathbb{R}^2))$ , there exists a unique solution  $\rho = (\rho^+, \rho^-) \in L^\infty([0, T]; Y_{r,p})$  of the following system of linear transport equations*

$$\frac{\partial \rho^\pm}{\partial t} \pm u \frac{\partial \rho^\pm}{\partial x_1} = 0. \tag{4.23}$$

Moreover, for all  $t \in [0, T]$ , we have

$$\|\rho(\cdot, t)\|_{r,p} \leq \|\rho_0\|_{r,p} \exp\left(C \int_0^t \|u(\cdot, \tau)\|_{C^r \cap L^p} d\tau\right),$$

where  $C > 0$  is a constant depending only on  $r$  and  $p$ .

**Proof.** From the fact that  $u(\cdot, t) \in C^r(\mathbb{R}^2) \cap L^p(\mathbb{R}^2)$ , for  $t \in [0, T]$ , we can define the flow map  $X^\pm(\cdot, t)$  satisfying

$$\begin{cases} \frac{\partial X^\pm(x, t)}{\partial t} = \pm \bar{u}(X(x, t), t), & \text{where } \bar{u} = (u, 0), \\ X^\pm(x, 0) = x. \end{cases} \quad (4.24)$$

By the characteristics method, we know that, if  $(X^\pm)^{-1}$  is the inverse function of  $X^\pm$  with respect to  $x$ , then  $\rho^\pm(x, t) = \rho_0^\pm((X^\pm)^{-1}(x, t))$  is the unique solution of system (4.23) (see Serre [19] for more details).

Let  $j \geq -1$ . Applying the operator  $\Delta_j$  to both sides of the system (1.1) yields

$$\frac{\partial \Delta_j \rho^\pm}{\partial t} \pm u \frac{\partial \Delta_j \rho^\pm}{\partial x_1} = \pm \left[ u \frac{\partial}{\partial x_1}, \Delta_j \right] \rho^\pm,$$

where  $[u \frac{\partial}{\partial x_1}, \Delta_j] \rho^\pm$  is defined in (3.12). This equation can be rewritten in the following form

$$\Delta_j \rho^\pm(x, t) = \Delta_j \rho_0^\pm((X^\pm)^{-1}(x, t)) \pm \int_0^t \left[ u \frac{\partial}{\partial x_1}, \Delta_j \right] \rho^\pm(X^\pm((X^\pm)^{-1}(x, t), s), s) ds.$$

Taking the  $L^\infty$ -norm of both sides of this equality, we get

$$\|\Delta_j \rho^\pm(\cdot, t)\|_{L^\infty} \leq \|\Delta_j \rho_0^\pm\|_{L^\infty} + \int_0^t \left\| \left[ u \frac{\partial}{\partial x_1}, \Delta_j \right] \rho^\pm(\cdot, s) \right\|_{L^\infty} ds.$$

Applying Lemma 3.1(1), we obtain

$$\|\rho^\pm(\cdot, t)\|_{C^r} \leq \|\rho_0^\pm\|_{C^r} + C \int_0^t \left( \left\| \frac{\partial \rho^\pm}{\partial x_1}(\cdot, s) \right\|_{L^\infty} \|u(\cdot, s)\|_{C^r} + \|\nabla u(\cdot, s)\|_{L^\infty} \|\rho^\pm(\cdot, s)\|_{C^r} \right) ds.$$

According to (2.8)–(2.9), we know that for  $r > 1$  and a constant  $C = C(r) > 0$ , we have

$$\left\| \frac{\partial \rho^\pm}{\partial x_1} \right\|_{L^\infty} \leq C \|\rho^\pm\|_{C^r}.$$

In a similar way, we can obtain  $\|\nabla u\|_{L^\infty} \leq C \|u\|_{C^r}$ . Therefore, for  $C = C(r) > 0$ ,

$$\begin{aligned} \|\rho^\pm(\cdot, t)\|_{C^r} &\leq \|\rho_0^\pm\|_{C^r} + C \int_0^t \|u(\cdot, s)\|_{C^r} \|\rho^\pm(\cdot, s)\|_{C^r} ds \\ &\leq \max_{\pm} (\|\rho_0^\pm\|_{C^r}) + C \int_0^t \|u(\cdot, s)\|_{C^r} \|\rho(\cdot, s)\|_{r,p} ds, \end{aligned}$$

where  $\rho = (\rho^+, \rho^-)$ . Moreover, integrating in time the system (4.23), we get the following  $L^p$  estimate:

$$\begin{aligned} \|\rho^\pm(\cdot, t)\|_{L^p} &\leq \|\rho_0^\pm\|_{L^p} + \int_0^t \|u(\cdot, s)\|_{L^p} \left\| \frac{\partial \rho^\pm}{\partial x_1}(\cdot, s) \right\|_{L^\infty} ds \\ &\leq \|\rho_0^\pm\|_{L^p} + C \int_0^t \|u(\cdot, s)\|_{L^p} \|\rho(\cdot, s)\|_{r,p} ds, \end{aligned}$$

where we have used Hölder inequality in the first line and (2.8)–(2.9) in the second line. Now, adding the two previous inequalities, we obtain

$$\|\rho(\cdot, t)\|_{r,p} \leq \|\rho_0\|_{r,p} + C \int_0^t \|u(\cdot, s)\|_{C^r \cap L^p} \|\rho(\cdot, s)\|_{r,p} ds.$$

By Gronwall’s Lemma, we obtain

$$\|\rho(\cdot, t)\|_{r,p} \leq \|\rho_0\|_{r,p} \exp\left(C \int_0^t \|u(\cdot, s)\|_{C^r \cap L^p} ds\right). \tag{4.25}$$

Which completes the proof of Proposition 4.1.  $\square$

**Proof of Theorem 1.4.** The proof of Theorem 1.4 is a consequence of the proof of Proposition 4.1. Indeed, just consider the characteristic equation

$$\begin{cases} \frac{\partial X(x, t)}{\partial t} = v(X(x, t), t), \\ X(x, 0) = x. \end{cases} \tag{4.26}$$

Then, as in the proof of Proposition 4.1, we use the commutator estimates proved in Lemma 3.1(1), to show the following estimate

$$\|g(\cdot, t)\|_{C^r \cap L^p} \leq \|g_0\|_{C^r \cap L^p} \exp\left(C \int_0^t \|v(\cdot, s)\|_{r,p} ds\right), \tag{4.27}$$

which proves the result.  $\square$

#### 4.2. Proof of Theorem 1.1

The proof starts with the construction of a successive approximation sequence  $\{\rho^n = (\rho^{+,n}, \rho^{-,n})\}_{n \geq 1}$  satisfying

$$\begin{cases} \rho^1 = (\rho_0^+, \rho_0^-) = \rho_0, \\ \frac{\partial \rho^{\pm, n+1}}{\partial t} \pm u^n \frac{\partial \rho^{\pm, n+1}}{\partial x_1} = 0, \quad \text{on } \mathbb{R}^2 \times (0, T), \\ u^n = R_1^2 R_2^2 (\rho^{+,n} - \rho^{-,n}), \\ \rho^{\pm, n+1}(x, 0) = \rho_0^\pm. \end{cases} \tag{4.28}$$

First of all, according to Proposition 2.4,  $\rho^1 \in Y_{r,p}$  implies that  $u^1 \in C^r(\mathbb{R}^2) \cap L^p(\mathbb{R}^2)$ . Thus, applying Proposition 4.1, we can prove that, for all  $T > 0$ , there exists a unique solution  $\rho^2 \in L^\infty([0, T]; Y_{r,p})$  for (4.28) with  $n = 2$ . Arguing in a similar manner we can show that this approached problem (4.28) has a unique solution  $\rho^n$  for all  $n \geq 1$ .

The rest of the proof can be divided into two major steps. The first step establishes the existence of  $T_1 > 0$  such that  $\{\rho^n = (\rho^{+,n}, \rho^{-,n})\}_{n \geq 1}$  is uniformly bounded in  $Y_{r,p}$  for any  $t \in [0, T_1]$ . The second step shows that for some  $T_2 \in [0, T_1]$ , we have  $\{\rho^n = (\rho^{+,n}, \rho^{-,n})\}_{n \geq 1}$  is a Cauchy sequence in  $C([0, T_2], Y_{r-1,p})$ .

*Step 1 (A uniform bound):* Using similar arguments as in the proof of Proposition 4.1, estimate (4.25) yields, by Proposition 2.4, the following bound on  $\{\rho^n = (\rho^{+,n}, \rho^{-,n})\}_{n \geq 1}$ :

$$\begin{aligned} \|\rho^{n+1}(\cdot, t)\|_{r,p} &\leq \|\rho_0\|_{r,p} \exp\left(C_0 \int_0^t \|u^n(\cdot, s)\|_{C^r \cap L^p} ds\right), \\ &\|\rho_0\|_{r,p} \exp\left(C_0 \int_0^t \|\rho^n(\cdot, s)\|_{r,p} ds\right), \end{aligned}$$

where  $r > 1, p \in (1, +\infty)$  and  $C_0 = C_0(r, p)$ . Choose  $T_1$  and  $M$  satisfying

$$M = 2\|\rho_0\|_{r,p} \quad \text{and} \quad \left( \exp(C_0 T_1 M) \leq 2 \text{ or } T_1 = \frac{\ln(2)}{2C_0\|\rho_0\|_{r,p}} \right).$$

Then  $\|\rho^n(\cdot, t)\|_{r,p} \leq M$  for all  $n \geq 1$  and  $t \in [0, T_1]$ . Since,

$$\|\rho^1\|_{r,p} \leq \|\rho_0\|_{r,p} < M$$

and  $\|\rho^k(\cdot, t)\|_{r,p} < M$ , we obtain

$$\|\rho^{n+1}(\cdot, t)\|_{r,p} \leq \|\rho_0\|_{r,p} \exp(C_0 T_1 M) \leq M. \quad (4.29)$$

Furthermore, since  $r > 1$ , we use (2.11) and Proposition 2.4 to get

$$\begin{aligned} \left\| \frac{\partial \rho^{\pm, n}}{\partial t} \right\|_{C^{r-1}} &\leq \left\| u^n \frac{\partial \rho^{\pm, n+1}}{\partial x_1} \right\|_{C^{r-1}} \\ &\leq \|u^n\|_{C^{r-1}} \left\| \frac{\partial \rho^{\pm, n+1}}{\partial x_1} \right\|_{L^\infty} + \|u^n\|_{L^\infty} \left\| \frac{\partial \rho^{\pm, n+1}}{\partial x_1} \right\|_{C^{r-1}} \\ &\leq C \|u^n\|_{C^{r-1}} \|\rho^{\pm, n+1}\|_{C^r} \\ &\leq CM^2, \end{aligned}$$

where we have used (2.8)–(2.9) in the third line. We can also check that the following  $L^p$  estimate on  $\rho^{\pm, n}$  is valid:

$$\begin{aligned} \left\| \frac{\partial \rho^{\pm, n}}{\partial t} \right\|_{L^p} &\leq \left\| u^n \frac{\partial \rho^{\pm, n+1}}{\partial x_1} \right\|_{L^p} \leq \|u^n\|_{L^p} \left\| \frac{\partial \rho^{\pm, n+1}}{\partial x_1} \right\|_{L^\infty} \\ &\leq C \|u^n\|_{L^p} \|\rho^{\pm, n+1}\|_{C^r} \leq CM^2, \end{aligned}$$

where we have used Hölder inequality in the first line, then (2.8)–(2.9) and Proposition 2.4 in the second line. Adding the two previous inequalities, we deduce that

$$\max_{\pm} \left( \left\| \frac{\partial \rho^{\pm, n}}{\partial t} \right\|_{C^{r-1}} \right) + \max_{\pm} \left( \left\| \frac{\partial \rho^{\pm, n}}{\partial t} \right\|_{L^p} \right) \leq CM^2, \quad (4.30)$$

where  $C = C(r)$ . Thus, by (4.29)–(4.30), we obtain that

$$\rho^n \in L^\infty([0, T_1]; Y_{r,p}) \cap Lip([0, T_1]; Y_{r-1,p})$$

is uniformly bounded.

*Step 2 (Cauchy sequence):* To show that  $\{\rho^n = (\rho^{+,n}, \rho^{-,n})\}_{n \geq 1}$  is a Cauchy sequence in  $Y_{r-1,q}$ , we consider the difference  $\eta^{\pm, n} = \rho^{\pm, n} - \rho^{\pm, n-1}$ . Rigorously speaking, we should consider the more general difference  $\eta^{\pm, m, n} = \rho^{\pm, m} - \rho^{\pm, n}$ , but the analysis for  $\eta^{m, n} = (\eta^{+, m, n}, \eta^{-, m, n})$  is parallel to what we shall present for  $\eta^n = (\eta^{+, n}, \eta^{-, n})$  and thus we consider  $\eta^n$  for the sake of a concise presentation. It follows from (4.28) that  $\{\eta^n = (\eta^{+, n}, \eta^{-, n})\}_{n \geq 1}$  satisfies

$$\begin{cases} \eta^{\pm, 1} = \rho_0^{\pm}, \\ \frac{\partial \eta^{\pm, n+1}}{\partial t} \pm u^n \frac{\partial \eta^{\pm, n+1}}{\partial x_1} = \mp w^n \frac{\partial \rho^{\pm, n}}{\partial x_1}, \\ w^n = R_1^2 R_2^2 (\eta^{+, n} - \eta^{-, n}), \\ \eta^{\pm, n+1}(x, 0) = \eta_0^{\pm, n+1}(x) = 0. \end{cases} \quad (4.31)$$

Proceeding as in the proof of Proposition 4.1, we obtain for any integer  $j \geq -1$ ,

$$\|\Delta_j \eta^{\pm, n+1}(\cdot, t)\|_{L^\infty} \leq \underbrace{\int_0^t \left\| \left[ u^n \frac{\partial}{\partial x_1}, \Delta_j \right] \eta^{\pm, n+1}(\cdot, s) \right\|_{L^\infty} ds}_{K_1} + \underbrace{\int_0^t \left\| \Delta_j \left( w^n \frac{\partial \rho^{\pm, n}}{\partial x_1}(\cdot, s) \right) \right\|_{L^\infty} ds}_{K_2}.$$

Estimating  $K_1$  by Lemma 3.1(2), and  $K_2$  by (2.11), we get

$$\begin{aligned} \|\eta^{\pm,n+1}(\cdot, t)\|_{C^{r-1}} &\leq C \int_0^t (\|\nabla u^n(\cdot, s)\|_{L^\infty} \|\eta^{\pm,n+1}(\cdot, s)\|_{C^{r-1}} + \|u^n(\cdot, s)\|_{C^r} \|\eta^{\pm,n+1}(\cdot, s)\|_{L^\infty}) ds \\ &\quad + C \int_0^t \left( \|w^n(\cdot, s)\|_{L^\infty} \left\| \frac{\partial \rho^{\pm,n}}{\partial x_1}(\cdot, s) \right\|_{C^{r-1}} + \|w^n(\cdot, s)\|_{C^{r-1}} \left\| \frac{\partial \rho^{\pm,n}}{\partial x_1}(\cdot, s) \right\|_{L^\infty} \right) ds. \end{aligned}$$

Since  $r > 1$ , Proposition 2.2 implies,

$$\begin{aligned} \|\nabla u^n\|_{L^\infty} &\leq C \|u^n\|_{C^r}, \quad \|\eta^{\pm,n+1}\|_{L^\infty} \leq C \|\eta^{\pm,n+1}\|_{C^{r-1}}, \\ \left\| \frac{\partial \rho^{\pm,n}}{\partial x_1} \right\|_{L^\infty} &\leq C \|\rho^{\pm,n}\|_{C^r} \quad \text{and} \quad \|w^n\|_{L^\infty} \leq C \|w^n\|_{C^{r-1}}. \end{aligned}$$

Therefore, for a constant  $C$  depending only on  $r$ ,

$$\begin{aligned} \|\eta^{\pm,n+1}(\cdot, t)\|_{C^{r-1}} &\leq C \int_0^t \|u^n(\cdot, s)\|_{C^r} \|\eta^{\pm,n+1}(\cdot, s)\|_{C^{r-1}} ds \\ &\quad + C \int_0^t \|w^n(\cdot, s)\|_{C^{r-1}} \|\rho^{\pm,n}(\cdot, s)\|_{C^r} ds. \end{aligned} \tag{4.32}$$

However, it follows from a basic  $L^p$  estimate that

$$\|\eta^{\pm,n+1}(\cdot, t)\|_{L^p} \leq C \int_0^t \|\nabla u^n(\cdot, s)\|_{L^\infty} \|\eta^{\pm,n+1}(\cdot, s)\|_{L^p} ds + C \int_0^t \|w^n(\cdot, s)\|_{L^p} \left\| \frac{\partial \rho^{\pm,n}}{\partial x_1}(\cdot, s) \right\|_{L^\infty} ds.$$

Since  $r > 1$ , using Proposition 2.2, we deduce that

$$\|\eta^{\pm,n+1}(\cdot, t)\|_{L^p} \leq C \int_0^t \|u^n(\cdot, s)\|_{C^r} \|\eta^{\pm,n+1}(\cdot, s)\|_{L^p} ds + C \int_0^t \|w^n(\cdot, s)\|_{L^p} \|\rho^{\pm,n}(\cdot, s)\|_{C^r} ds. \tag{4.33}$$

Adding the two inequalities (4.32) and (4.33), yields

$$\begin{aligned} \|\eta^{n+1}(\cdot, t)\|_{r-1,p} &\leq C \int_0^t \|u^n(\cdot, s)\|_{C^r \cap L^p} \|\eta^{n+1}(\cdot, s)\|_{r-1,p} ds \\ &\quad + C \int_0^t \|w^n(\cdot, s)\|_{C^{r-1} \cap L^p} \|\rho^n(\cdot, s)\|_{r,p} ds. \end{aligned}$$

The components of  $w^n$  are the Riesz transforms of  $\eta^n$  and thus, according to Proposition 2.4:

$$\|w^n\|_{C^{r-1} \cap L^p} \leq C \|\eta^n\|_{r-1,p}.$$

We thus have reached an iterative relationship between  $\|\eta^n\|_{r-1,p}$  and  $\|\eta^{n+1}\|_{r-1,p}$ :

$$\begin{aligned} \|\eta^{n+1}(\cdot, t)\|_{r-1,p} &\leq C_1 \int_0^t \|\rho^n(\cdot, s)\|_{r,p} \|\eta^{n+1}(\cdot, s)\|_{r-1,p} ds \\ &\quad + C_1 \int_0^t \|\eta^n(\cdot, s)\|_{r-1,p} \|\rho^n(\cdot, s)\|_{r,p} ds, \end{aligned} \tag{4.34}$$

where the constants are labeled as  $C_1$  for the purpose of defining  $T_2$ . It has been shown in Step 1 that for  $t \leq T_1$ ,

$$\|\rho^n\|_{r,p} \leq M.$$

Now, choose  $T_2 > 0$  satisfying

$$T_2 \leq T_1, \quad C_1MT_2 \leq \frac{1}{4}.$$

In the following, we will prove that the sequence  $\{\rho^n(\cdot, t)\}_{n \geq 1}$  is a Cauchy sequence in  $Y_{r-1,p}$  for  $t \leq T_2$ . Indeed, for any given  $\varepsilon > 0$  small enough, if we assume that  $\|\eta^n\|_{r-1,p} \leq \varepsilon$  for  $t \leq T_2$ , then (4.34) implies that

$$\|\eta^{n+1}\|_{r-1,p} \leq C_1\varepsilon MT_2 + C_1M \int_0^t \|\eta^{n+1}(\cdot, s)\|_{r-1,p} ds,$$

is valid for any  $t \leq T_2$ . It then follows from Gronwall’s inequality that

$$\|\eta^{n+1}\|_{r-1,p} \leq \varepsilon,$$

which implies that  $\{\rho^n(\cdot, t)\}_{n \geq 1}$  is a Cauchy sequence in  $Y_{r-1,p}$  for  $t \leq T_2$  and then completes the proof of Step 2.

We conclude from Steps 1 and 2 that there exists  $\rho = (\rho^+, \rho^-)$  satisfying

$$\rho \in L^\infty([0, T_2]; Y_{r,p}) \cap Lip([0, T_2]; Y_{r-1,p})$$

such that  $\rho^n$  converges to  $\rho$  in  $C([0, T_2]; Y_{r-1,p})$ .

The proof of uniqueness follows directly from Step 2. This completes the proof of Theorem 1.1.

### 4.3. Proof of Theorem 1.2

It is worth mentioning that the ideas of the proof of Theorem 1.2 are already contained in the proof of Theorem 1.1.

First of all, we note that for all  $L \in \mathbb{R}$ , if  $\rho^\pm$  are solutions of (1.1) then

$$\bar{\rho}^\pm(x_1, x_2, t) = \rho^\pm(x_1, x_2, t) - Lx_1$$

solves the following system:

$$\begin{cases} \frac{\partial \bar{\rho}^\pm}{\partial t}(x, t) \pm u \frac{\partial \bar{\rho}^\pm}{\partial x_1}(x, t) = \mp Lu & \text{on } \mathbb{R}^2 \times (0, T), \\ u = R_1^2 R_2^2 (\bar{\rho}^+ - \bar{\rho}^-), \end{cases} \tag{4.35}$$

and respects the following initial data:

$$\bar{\rho}_0^\pm(x_1, x_2, t) = \rho_0^\pm(x_1, x_2) - Lx_1.$$

Now, to prove Theorem 1.2, it suffices to show that, for all initial data  $\bar{\rho}_0^\pm \in Y_{r,p}$ , the system (4.35) has a unique local solution  $\bar{\rho}^\pm \in L^\infty([0, T]; Y_{r,p})$  for  $r > 1$  and  $p \in (1, +\infty)$ .

In order to do this, we proceed as in the proof of Theorem 1.1. We consider the following approached system:

$$\begin{cases} \bar{\rho}^1 = (\bar{\rho}_0^+, \bar{\rho}_0^-) = \bar{\rho}_0, \\ \frac{\partial \bar{\rho}^{\pm, n+1}}{\partial t} \pm u^n \frac{\partial \bar{\rho}^{\pm, n+1}}{\partial x_1} = \mp Lu^n, & \text{on } \mathbb{R}^2 \times (0, T), \\ u^n = R_1^2 R_2^2 (\bar{\rho}^{+, n} - \bar{\rho}^{-, n}), \\ \bar{\rho}^{\pm, n+1}(x, 0) = \bar{\rho}_0^\pm. \end{cases} \tag{4.36}$$

We remark that, the only change that appears here, compared to the approached system (4.28) is the right-hand side  $Lu^n$  of the second equation of (4.36). However, by Proposition 2.4, we know that this term remains bounded in  $L^\infty([0, T]; C^r(\mathbb{R}^2) \cap L^p(\mathbb{R}^2))$  for  $r > 1$  and  $p \in (1, +\infty)$ . Which permits us to easily follow the same steps of the proof of Theorem 1.1. This finally proves that, for some small  $T > 0$ , we have on the one hand: the sequence  $\bar{\rho}^n = (\bar{\rho}^{+, n}, \bar{\rho}^{-, n})$  is uniformly bounded in  $L^\infty([0, T]; Y_{r,p})$ , and on the other hand, this sequence is a Cauchy sequence in  $L^\infty([0, T]; Y_{r-1,p})$ . This terminate the proof.

## Acknowledgements

The author would like to thank M. Cannone and R. Monneau for fruitful remarks that helped in the preparation of the paper, and H. Ibrahim for carefully reading it. This work was partially supported by the program “PPF, programme pluri-formations mathématiques financières et EDP” (2006–2010), Université Paris-Est.

## References

- [1] O. Alvarez, P. Hoch, Y. Le Bouar, R. Monneau, Dislocation dynamics: Short-time existence and uniqueness of the solution, *Arch. Ration. Mech. Anal.* 181 (2006) 449–504.
- [2] G. Barles, P. Cardaliaguet, O. Ley, R. Monneau, Global existence results and uniqueness for dislocation equations, *SIAM J. Math. Anal.* 40 (2008) 44–69.
- [3] J.-M. Bony, Calcul symbolique et propagation des singularités pour les équations aux dérivées partielles non linéaires, *Ann. Sci. École Norm. Sup. (4)* 14 (1981) 209–246.
- [4] M. Cannone, A. El Hajj, R. Monneau, R. Ribaud, Global existence for a system of non-linear and non-local transport equations describing the dynamics of dislocation densities, *Arch. Ration. Mech. Anal.*, in press.
- [5] M. Cannone, Y. Meyer, Littlewood–Paley decomposition and Navier–Stokes equations, *Methods Appl. Anal.* 2 (1995) 307–319.
- [6] M. Cannone, F. Planchon, On the regularity of the bilinear term for solutions to the incompressible Navier–Stokes equations, *Rev. Mat. Iberoamericana* 16 (2000) 1–16.
- [7] J.-Y. Chemin, *Perfect Incompressible Fluids*, Oxford Lecture Ser. Math. Appl., vol. 14, The Clarendon Press/Oxford University Press, New York, 1998, translated from the 1995 French original by Isabelle Gallagher and Dragos Ifitimie.
- [8] Q. Chen, C. Miao, Z. Zhang, A new Bernstein’s inequality and the 2D dissipative quasi-geostrophic equation, *Comm. Math. Phys.* 271 (2007) 821–838.
- [9] A. El Hajj, Well-posedness theory for a nonconservative Burgers-type system arising in dislocation dynamics, *SIAM J. Math. Anal.* 39 (2007) 965–986.
- [10] A. El Hajj, N. Forcadel, A convergent scheme for a non-local coupled system modelling dislocations densities dynamics, *Math. Comp.* 77 (2008) 789–812.
- [11] I. Groma, P. Balogh, Investigation of dislocation pattern formation in a two-dimensional self-consistent field approximation, *Acta Mater.* 47 (1999) 3647–3654.
- [12] I. Groma, F. Csikor, M. Zaiser, Spatial correlations and higher-order gradient terms in a continuum description of dislocation dynamics, *Acta Mater.* 51 (2003) 1271–1281.
- [13] J. Hirth, J. Lothe, *Theory of Dislocations*, second ed., Krieger Publishing Company, Florida 32950, 1982.
- [14] H. Ibrahim, Existence and uniqueness for a non-linear parabolic/Hamilton–Jacobi system describing the dynamics of dislocation densities, *Ann. Inst. H. Poincaré Anal. Non Linéaire* (2009) 415–435.
- [15] Y. Meyer, Ondelettes et opérateurs. I, II, *Actualités Mathématiques (Current mathematical topics)*, in: *Ondelettes (Wavelets)*, Hermann, Paris, 1990.
- [16] Y. Meyer, R.R. Coifman, Ondelettes et opérateurs. III, *Actualités Mathématiques (Current mathematical topics)*, in: *Opérateurs multilinéaires (Multilinear Operators)*, Hermann, Paris, 1991.
- [17] E. Orowan, Zur kristallplastizität i–iii, *Z. Phys.* 89 (1934) 605–634.
- [18] M. Polanyi, Über eine art gitterstörung, die einem kristall plastisch machen konnte, *Z. Phys.* 89 (1934) 660–664.
- [19] D. Serre, Systems of conservation laws. I, II, in: *Geometric Structures, Oscillations, and Initial–Boundary Value Problems*, Cambridge University Press, Cambridge, 1999/2000, translated from the 1996 French original by I.N. Sneddon.
- [20] G.I. Taylor, The mechanism of plastic deformation of crystals, *Proc. R. Soc. Lond. Ser. A* 145 (1934) 362–387.