

Invertibility of Sobolev mappings under minimal hypotheses

Leonid V. Kovalev^{a,1}, Jani Onninen^{a,2}, Kai Rajala^{b,*,3}

^a Department of Mathematics, Syracuse University, Syracuse, NY 13244, USA

^b Department of Mathematics and Statistics, University of Jyväskylä, PO Box 35 (MaD), FI-40014, University of Jyväskylä, Finland

Received 15 May 2009; received in revised form 14 September 2009; accepted 14 September 2009

Available online 1 October 2009

Abstract

We prove a version of the Inverse Function Theorem for continuous weakly differentiable mappings. Namely, a nonconstant $W^{1,n}$ mapping is a local homeomorphism if it has integrable inner distortion function and satisfies a certain differential inclusion. The integrability assumption is shown to be optimal.

© 2009 Elsevier Masson SAS. All rights reserved.

MSC: primary 30C65; secondary 26B10, 26B25

Keywords: Local homeomorphism; Differential inclusion; Finite distortion

1. Introduction

Throughout this paper Ω is a bounded domain in \mathbb{R}^n . The classical Inverse Function Theorem states that if $f: \Omega \rightarrow \mathbb{R}^n$ is continuously differentiable and the differential matrix $Df(x)$ is invertible at some point x , then f is a homeomorphism in a neighborhood of x . We are interested in a version of the Inverse Function Theorem for continuous weakly differentiable mappings. In this context the invertibility of the differential matrix is not sufficient. As an example, consider the winding mapping $f: \mathbb{R}^3 \rightarrow \mathbb{R}^3$ written in cylindrical coordinates as $f(r, \theta, z) = (r, 2\theta, z)$. Although f is Lipschitz and its Jacobian determinant $J(x, f)$ equals 2 for a.e. $x \in \mathbb{R}^n$, this mapping is not a local homeomorphism.

Let us introduce the following subset of $n \times n$ matrices.

$$\mathcal{M}(\delta) = \{A \in \mathbb{R}^{n \times n}: \langle A\xi, \xi \rangle \geq \delta |A\xi| |\xi| \text{ for all } \xi \in \mathbb{R}^n\},$$

where $-1 \leq \delta \leq 1$. Note that $\delta = -1$ imposes no condition on the matrix. When $-1 < \delta < 0$, the set $\mathcal{M}(\delta)$ is not convex and the differential inclusion

* Corresponding author.

E-mail addresses: lvkvale@syr.edu (L.V. Kovalev), jkonnine@syr.edu (J. Onninen), kirajala@maths.jyu.fi (K. Rajala).

¹ Supported by the NSF grant DMS-0913474.

² Supported by the NSF grant DMS-0701059.

³ Supported by the Academy of Finland.

$$Df(x) \in \mathcal{M}(\delta) \quad \text{for a.e. } x \in \Omega, \quad (1.1)$$

cannot be integrated to yield a pointwise inequality for f .

The winding mapping does not satisfy (1.1) for any $\delta > -1$. Even so, this differential inclusion does not by itself guarantee that f is locally invertible, e.g., $f(x_1, x_2) = (x_1, 0)$. There are also such examples with strictly positive Jacobian [14, Example 18]. To quantify the invertibility of a matrix $A \in \mathbb{R}^{n \times n}$, we introduce the inner distortion $K_I(A) \in [1, \infty]$.

$$K_I(A) = \begin{cases} \frac{\|A^\sharp\|^n}{(\det A)^{n-1}}, & \det A > 0, \\ 1, & A = 0, \\ \infty, & \text{otherwise.} \end{cases} \quad (1.2)$$

Here A^\sharp stands for the cofactor matrix of A and $\|\cdot\|$ is the operator norm. To shorten the notation we write $K_I(x, f) = K_I(Df(x))$ and

$$\mathcal{K}_\Omega[f] := \frac{1}{|\Omega|} \int_\Omega K_I(x, f) \, dx,$$

where $|\Omega|$ is the Lebesgue measure of Ω . If $f \in W^{1,n}(\Omega, \mathbb{R}^n)$ and $K_I(x, f) < \infty$ a.e., then f has a logarithmic modulus of continuity [4,9]; that is,

$$|f(a) - f(b)|^n \leq \frac{C(n) \int_{2B} \|Df\|^n}{\log(e + \frac{2 \operatorname{diam} B}{|a-b|})}, \quad a, b \in B, \quad 2B \Subset \Omega.$$

In this paper we always take f to be its continuous representative.

If moreover $\mathcal{K}_\Omega[f] < \infty$ and f is invertible, then the inverse $h := f^{-1}$ is a $W^{1,n}$ -mapping and

$$\int_\Omega K_I(x, f) \, dx = \int_{f(\Omega)} \|Dh\|^n,$$

see [1, Theorem 9.1]. Thus $\mathcal{K}_\Omega[f]$ controls the modulus of continuity of f^{-1} , should it exist. Our main result addresses its existence.

Theorem 1.1. *Suppose that $f \in W_{\text{loc}}^{1,n}(\Omega, \mathbb{R}^n)$ is a nonconstant mapping such that $\mathcal{K}_\Omega[f] < \infty$. If there exists $\delta > -1$ such that $Df(x) \in \mathcal{M}(\delta)$ for almost every $x \in \Omega$, then f is a local homeomorphism.*

This theorem is already known in the planar case $n = 2$ [14, Theorem 4]. The assumption $\mathcal{K}_\Omega[f] < \infty$ cannot be replaced by $\int_\Omega K_I^q(x, f) \, dx < \infty$ for any $q < 1$, see [14, Example 18] or [2, Example 1].

Our proof of Theorem 1.1 is based on two results of independent interest. The first step toward proving that a mapping is a local homeomorphism is to show that it is discrete and open; that is, preimages of points are discrete sets and images of open sets are open.

Theorem 1.2. *Let $f : \Omega \rightarrow \mathbb{R}^n$ be a mapping in $W_{\text{loc}}^{1,n}(\Omega, \mathbb{R}^n)$ such that $J(x, f) > 0$ a.e. If $(Df)^{-1} \in L^\infty(\Omega, \mathbb{R}^{n \times n})$, then f is discrete and open.*

The challenging Iwaniec–Šverák conjecture asserts even more: a nonconstant mapping $f \in W_{\text{loc}}^{1,n}(\Omega, \mathbb{R}^n)$ with $\mathcal{K}_\Omega[f] < \infty$ is discrete and open. So far this conjecture was proved only for $n = 2$ in [10]. Partial results in this direction were recently obtained in [6–8,15,19,20].

Another crucial ingredient of our proof of Theorem 1.1 is an estimate for the multiplicity $N(y, f, A) := \#(f^{-1}(y) \cap A)$ of a local homeomorphism f in terms of the integral of $K_I(\cdot, f)$ in dimensions $n \geq 3$. This result (Theorem 5.1) continues the line of development that began in 1967 with the celebrated Global Homeomorphism Theorem of Zorich [24].

The proof of Theorem 1.1 proceeds as follows. The differential inclusion (1.1) allows us to approximate f by mappings $f^\lambda(x) := f(x) + \lambda x$ to which Theorem 1.2 can be applied. The results of [14] yield that f^λ is a local

homeomorphism. By virtue of Theorem 5.1 the mappings f^λ have uniformly bounded multiplicity, which leads to a bound for the essential multiplicity of f . This additional information suffices to show that f is discrete and open, see Proposition 2.2 below. Since f is a limit of local homeomorphisms f^λ , the conclusion follows.

Different approaches to the invertibility of Sobolev mappings were pursued in [2,3,5,16,18,22], see also references therein.

2. Background

In this section we collect necessary notation and preliminaries. An open ball with center a and radius r is denoted by $B(a, r) := \{x \in \mathbb{R}^n : |x - a| < r\}$. Its boundary is the sphere $S(a, r)$. If $\lambda > 0$ and $B = B(a, r)$, then $\lambda B = B(a, \lambda r)$ and $\lambda S = S(a, \lambda r)$. In addition, $\mathbb{B} = B(0, 1)$, $\mathbb{B}_r = B(0, r)$, $\mathbb{S} = S(0, 1)$ and $\mathbb{S}_r = S(0, r)$.

Let \mathcal{H}^d stand for the d -dimensional Hausdorff measure which agrees with the Lebesgue measure when d coincides with the space dimension. The Hausdorff distance $d_{\mathcal{H}}(E, F)$ between nonempty bounded sets E and F is defined as the infimum of numbers $\epsilon > 0$ such that the ϵ -neighborhood of E contains F and vice versa.

Given a continuous mapping $f : \Omega \rightarrow \mathbb{R}^n$ and a set $E \subset \Omega$, we denote by $N(y, f, E)$ the cardinality (possibly infinite) of the set $f^{-1}(y) \cap E$. If $y \in \mathbb{R}^n \setminus f(\partial\Omega)$, the local degree of f at y with respect to a domain $G \subset \Omega$ is denoted by $\text{deg}(y, f, G)$. We write $f : A \xrightarrow{\text{hom}} B$ to indicate that f is a homeomorphism from A onto B .

Let Γ be a family of paths (parametrized curves) in \mathbb{R}^n , $n \geq 2$. The image of $\gamma \in \Gamma$ is denoted by $|\gamma|$. We let \mathcal{Y}_Γ be the set of all Borel functions $\rho : \mathbb{R}^n \rightarrow [0, \infty]$ such that

$$\int_\gamma \rho \, ds \geq 1$$

for every locally rectifiable path $\gamma \in \Gamma$. The functions in \mathcal{Y}_Γ are called admissible for Γ . For a given weight $\omega : \mathbb{R}^n \rightarrow [0, \infty]$ we define

$$M_\omega \Gamma = \inf_{\rho \in \mathcal{Y}_\Gamma} \int \rho(x)^n \omega(x) \, dx,$$

and call $M_\omega \Gamma$ the weighted conformal modulus of Γ . Here it suffices to have ω defined on a Borel set containing $\bigcup_{\gamma \in \Gamma} |\gamma|$. When $\omega \equiv 1$ we obtain the conformal modulus $M\Gamma$. We will also use the spherical modulus with respect to a sphere S ,

$$M^S \Gamma = \inf_{\rho \in \mathcal{Y}_\Gamma} \int_S \rho(y)^n \, d\mathcal{H}^{n-1}(y).$$

The reader may wish to consult the monographs [21,23] for basic properties of moduli of path families. The following generalization of the Poletsky inequality relates the moduli of Γ and of its image under f , denoted by $f\Gamma$.

Proposition 2.1. (See [12].) *Suppose that $f \in W^{1,n}(\Omega, \mathbb{R}^n)$ is a discrete and open mapping with $\mathcal{K}_\Omega[f] < \infty$. If Γ is a family of paths contained in Ω , then*

$$Mf\Gamma \leq M_{K_I(\cdot, f)}\Gamma. \tag{2.1}$$

We will use the following result, which establishes the Iwaniec–Šverák conjecture under an additional assumption on the multiplicity of f .

Proposition 2.2. *Suppose that $f \in W_{\text{loc}}^{1,n}(\Omega, \mathbb{R}^n)$ is a nonconstant mapping with $\mathcal{K}_\Omega[f] < \infty$. Let B be a ball such that $2B \Subset \Omega$. If*

$$\text{ess lim sup}_{r \rightarrow 0} r^{1-n} \int_{S(a,r)} N(y, f, B) \, d\mathcal{H}^{n-1}(y) < \infty \tag{2.2}$$

for every $a \in \mathbb{R}^n$, then f is discrete and open in B .

This proposition is a consequence of [20, Theorem 2.2]. Although [20, Theorem 2.2] requires that

$$\operatorname{ess\,sup}_{0 < t < 1} \int_{\partial(tB)} \frac{\|D^\sharp f(x)\|}{|f(x) - a|^{n-1}} d\mathcal{H}^{n-1}(x) < \infty,$$

this condition is only used to obtain (2.2).

3. Preliminary results

For the sake of brevity, the connected component of a set A that contains a point $x \in A$ will be called the x -component of A .

Proposition 3.1. *Suppose that $f \in W_{\text{loc}}^{1,n}(\Omega, \mathbb{R}^n)$ is a mapping such that $\mathcal{K}_\Omega[f] < \infty$. Let $x \in \Omega$ and $y = f(x)$. If the x -component of $f^{-1}(y)$ is $\{x\}$, then f is discrete and open in some neighborhood of x .*

Proof. Pick $r > 0$ such that $B(x, r) \Subset \Omega$ and let U_j be the x -component of $(f^{-1}B(y, 1/j)) \cap B(x, r)$, $j = 1, 2, \dots$. Since the sets $\bar{U}_j \subset \mathbb{R}^n$ are nested, compact, and connected, their intersection E is also connected. On the other hand, $x \in E \subset f^{-1}(y)$, hence $E = \{x\}$. It follows that $\operatorname{diam}(U_j) \rightarrow 0$ as $j \rightarrow \infty$. Let us fix j such that $U_j \Subset \Omega$. Note that U_j coincides with the x -component of $f^{-1}B(y, 1/j)$.

We claim that f is quasilight in U_j ; that is, the connected components of $f^{-1}(w) \cap U_j$ are compact for all $w \in \mathbb{R}^n$. If not, then there exists $z \in U_j$ such that the z -component of $f^{-1}(f(z))$ intersects ∂U_j at some point b . Since $f(b) = f(z) \in B(y, 1/j)$, there exists $t > 0$ such that $fB(b, t) \subset B(y, 1/j)$. This contradicts the definition of U_j . Therefore, f is quasilight in U_j . By [19, Theorem 1.1] f is discrete and open in U_j . \square

For the convenience of the reader we state two preliminary results from [21, III.3].

Lemma 3.2. (See [21, III.3.1].) *Let $f: \Omega \rightarrow \mathbb{R}^n$ be a local homeomorphism and let Q be a simply connected and locally pathwise connected set in \mathbb{R}^n . Suppose P is a component of $f^{-1}Q$ such that $\bar{P} \subset \Omega$. Then $f: P \xrightarrow{\text{hom}} Q$. If in addition Q is relatively locally connected, then $f: \bar{P} \xrightarrow{\text{hom}} \bar{Q}$.*

Lemma 3.3. (See [21, III.3.3].) *Let $f: \Omega \rightarrow \mathbb{R}^n$ be a local homeomorphism and let $A, B \subset \Omega$ be two sets such that f is homeomorphic in A and in B . If $A \cap B \neq \emptyset$ and if $fA \cap fB$ is connected, then f is homeomorphic in $A \cup B$.*

Given a sphere $S = S(a, r) \subset \mathbb{R}^n$, and a point $p \in S$, let $C_S(p, \phi)$ be the open spherical cap of S with center p and opening angle $\phi \in (0, \pi]$,

$$C_S(p, \phi) = \{y \in S: \langle y - a, p - a \rangle > r^2 \cos \phi\}.$$

For instance $C_S(p, \pi/2)$ is a hemisphere and $C_S(p, \pi)$ is a punctured sphere. For any $\phi \in (0, \pi]$ the cap $C_S(p, \phi)$ contains the point p .

The following topological lemma forms the main step of the proof of Zorich Global Homeomorphism Theorem, see [21, III.3].

Lemma 3.4. *Let $f: \Omega \rightarrow \mathbb{R}^n$ be a local homeomorphism, $\Omega \subset \mathbb{R}^n$, $n \geq 3$. Suppose we have the following:*

- (i) $G \Subset \Omega$ such that $f: G \xrightarrow{\text{hom}} G'$ where G' is convex;
- (ii) $G \subset D \Subset \Omega$ and there is $a \in \partial G \cap \partial D$;
- (iii) a ball $\mathcal{B} \subset \mathbb{R}^n$ that contains $a' = f(a)$ and such that $S = \partial \mathcal{B}$ meets G' at some point b' .

Let $b = f^{-1}(b') \cap G$ and denote by $C_S^*(b', \phi)$ the component of $f^{-1}C_S(b', \phi)$ containing b . Then there exists $0 < \phi_0 < \pi$ such that $C_S^*(b', \phi_0) \subset D$ and the closure of $C_S^*(b', \phi_0)$ meets ∂D .

Proof. Let ϕ_0 be the supremum of all ϕ such that $C_S^*(b', \phi) \subset D$. First we observe that $\phi_0 > 0$. Indeed, since f is a local homeomorphism, there exists a neighborhood $V \subset D$ of b such that $f : V \xrightarrow{\text{hom}} f(V)$. If ϕ is sufficiently small, then $C_S(b', \phi) \Subset f(V)$, hence $C_S^*(b', \phi) \Subset V \subset D$. It remains to show that $\phi_0 < \pi$.

Suppose to the contrary that $\phi_0 = \pi$. Since $C_S^*(b', \pi) \subset D$, it follows from Lemma 3.2 that $f : \overline{C_S^*(b', \pi)} \xrightarrow{\text{hom}} \overline{C_S(b', \pi)} = S$ (here the assumption $n \geq 3$ is used). Since $S^* := \overline{C_S^*(b', \pi)}$ is homeomorphic to S , it separates \mathbb{R}^n into two components. Let U be the bounded component of $\mathbb{R}^n \setminus S^*$. Then the boundary of $f(U)$ is contained in S which implies $f(U) = \emptyset$. Moreover, $f : \overline{U} \xrightarrow{\text{hom}} \overline{\emptyset}$ by Lemma 3.2. Since $b \in \overline{U} \cap \overline{G}$ and since $f(\overline{U}) \cap f(\overline{G}) = \overline{\emptyset} \cap \overline{G}'$ is convex (hence connected), Lemma 3.3 yields that f is homeomorphic in $\overline{U} \cup \overline{G}$.

This leads to a contradiction. Since $\overline{U} \cup \overline{G} \subset \overline{D}$ it follows that a lies on the boundary of $\overline{U} \cup \overline{G}$. On the other hand, $f(a) = a' \in f(U)$ is an interior point of $f(\overline{U} \cup \overline{G})$. \square

We shall use a geometric lemma which is essentially contained in [13].

Lemma 3.5. *Suppose we are given a ball $B(y_0, r) \subset \mathbb{R}^n$, a point $y_1 \in S(y_0, r)$ and a connected set E that contains y_0 and some point $y_2 \in S(y_0, r)$. Then there exist $q \in B(y_0, r)$ and $0 < \sigma < 2r$ such that for every $\sigma < t < 4\sigma/3$,*

- (i) $y_1 \in B(q, t)$;
- (ii) $S(q, t) \cap E \neq \emptyset$;
- (iii) $S(q, t) \subset B(y_0, 2r) \setminus B(y_0, r/10)$.

Proof. Let α be the angle at the point $(y_0 + y_1)/2$ formed by the line segments from y_0 to $(y_0 + y_1)/2$ and from $(y_0 + y_1)/2$ to y_2 . There are two cases possible.

Case 1. $0 \leq \alpha < \pi/2$, or, equivalently, $|y_1 - y_2| > r$. In this case we choose $q = (y_0 + y_1)/2$ and $\sigma = 3r/5$. For $\sigma < t < 4\sigma/3$ we have $B(y_0, r/10) \subset B(q, t)$ and $y_1 \in B(q, t)$. At the same time, $y_2 \notin \overline{B}(q, t)$ because

$$|y_2 - q| > \frac{\sqrt{3}}{2}r = \frac{5}{2\sqrt{3}}\sigma > \frac{4}{3}\sigma.$$

Thus, all conditions (i)–(iii) are satisfied.

Case 2. $\pi/2 \leq \alpha \leq \pi$, or, equivalently, $|y_1 - y_2| \leq r$. This time we choose $q = (y_1 + y_2)/2$ and $\sigma = |y_1 - y_2|/2$. Since $|y_0 - q| \geq (\sqrt{3}/2)r$, it follows that $\overline{B}(q, t) \cap B(y_0, r/10) = \emptyset$ provided that

$$t < \left(\frac{\sqrt{3}}{2} - \frac{1}{10}\right)r.$$

This is indeed the case, because

$$\frac{4}{3}\sigma \leq \frac{2}{3}r < \left(\frac{\sqrt{3}}{2} - \frac{1}{10}\right)r.$$

All conditions (i)–(iii) are met. \square

4. Proof of Theorem 1.2

Let $\|(Df)^{-1}\|_\infty = L$. First we observe that the inner distortion of f is locally integrable because

$$K_I(x, f) = \|(Df(x))^{-1}\|^n J(x, f) \leq L^n \|Df\|^n \quad \text{for a.e. } x \in \Omega. \tag{4.1}$$

We may assume that $\mathbb{B}_4 = B(0, 4) \Subset \Omega$. It suffices to show that f is discrete and open in \mathbb{B} . We will do this by proving that (2.2) holds. Without loss of generality, a in (2.2) equals 0. Fix $1 < t < 2$ and $3 < T < 4$ so that $\mathcal{H}^{n-1}(fS_t) < \infty$ and $\mathcal{H}^{n-1}(fS_T) < \infty$. By the area formula we have

$$\int_{\mathbb{R}^n} N(y, f, \mathbb{B}_T) dy = \int_{\mathbb{B}_T} J(x, f) dx < \infty.$$

Therefore, for almost every $0 < R < \infty$ we have

$$\int_{\mathbb{S}_R} N(y, f, \mathbb{B}_T) d\mathcal{H}^{n-1}(y) < \infty \quad \text{and} \quad \mathcal{H}^{n-1}(f(\mathbb{S}_T) \cap \mathbb{S}_R) = 0. \quad (4.2)$$

We fix such $R < 1/(2L)$ so that (4.2) holds, and let

$$M := R^{1-n} \int_{\mathbb{S}_R} N(y, f, \mathbb{B}_T) d\mathcal{H}^{n-1}(y).$$

Our goal is to prove that

$$r^{1-n} \int_{\mathbb{S}_r} N(y, f, \mathbb{B}_T) d\mathcal{H}^{n-1}(y) \leq M \quad \text{for a.e. } 0 < r < R. \quad (4.3)$$

Let $r < R$ be such that $\mathcal{H}^{n-1}(f(\mathbb{S}_r) \cap \mathbb{S}_r) = 0$, and denote by $E \subset \mathbb{S}$ the set of unit vectors v for which

$$\deg(Rv, f, \mathbb{B}_T) < \deg(rv, f, \mathbb{B}_T). \quad (4.4)$$

Let $I_v : [r, R] \rightarrow \mathbb{R}^n$ be the parametrized line segment $I_v(s) = sv$. By Proposition 3.1, either $f^{-1}(sv)$ has a nontrivial component for some $r \leq s \leq R$, or f is discrete and open in a neighborhood of $f^{-1}(I_v[r, R])$, denoted by U_v . By using the co-area formula as in [20, Lemma 2.4], we see that the former possibility only occurs for $v \in F_1$ where $\mathcal{H}^{n-1}(F_1) = 0$. The mapping f is discrete and open in the open set $U := \bigcup \{U_v : v \in E \setminus F_1\}$. It follows from (4.4) and basic properties of path lifting [21, Section II.3] that for each $v \in E \setminus F_1$ the segment I_v has a maximal f -lifting I_v^* starting at \mathbb{B}_r and leaving \mathbb{B}_T .

Denote

$$\ell_f(x) := \liminf_{z \rightarrow x} \frac{|f(z) - f(x)|}{|z - x|}.$$

By our assumption on $(Df)^{-1}$ there exists a Borel null set $F \subset \Omega$ such that $\ell_f(x) \geq 1/L$ for $x \in \Omega \setminus F$. Let F_2 be the set of $v \in E \setminus F_1$ such that either I_v^* is unrectifiable or $\mathcal{H}^1(I_v^* \cap F) > 0$. Since the measure of F is zero, it follows that the family of curves $\Gamma_F := \{I_v^* : v \in F_2\}$ has zero weighted modulus for any locally integrable weight. In particular, $M_{K_I} \Gamma_F = 0$. Since $\Gamma_F \subset U$ we can apply (2.1) and obtain $M\{I_v^* : v \in F_2\} = 0$, which implies $\mathcal{H}^{n-1}(F_2) = 0$.

For $v \in E \setminus (F_1 \cup F_2)$ we have

$$\mathcal{H}^1(I_v^*) \leq L\mathcal{H}^1(I_v) < LR < \frac{1}{2}, \quad (4.5)$$

which contradicts the fact that I_v^* begins at \mathbb{B}_r and leaves \mathbb{B}_T . Thus $E \subset F_1 \cup F_2$. As a consequence, $\mathcal{H}^{n-1}(E) = 0$, which means $\deg(rv, f, \mathbb{B}_T) \leq \deg(Rv, f, \mathbb{B}_T)$ for \mathcal{H}^{n-1} -a.e. $v \in \mathbb{S}$. Since $\deg(y, f, \mathbb{B}_T) = N(y, f, \mathbb{B}_T)$ for a.e. $y \in \mathbb{R}^n$ [8, Proposition 2], inequality (4.3) follows. This completes the proof of Theorem 1.2 via Proposition 2.2.

5. Multiplicity of local homeomorphisms

In 1967 Zorich [24] proved that a local homeomorphism $f : \mathbb{R}^n \rightarrow \mathbb{R}^n$, $n \geq 3$, with $K_I(\cdot, f) \in L^\infty(\mathbb{R}^n)$ must be a global homeomorphism. Martio, Rickman and Väisälä [16] gave a local version of this result. Namely, if $f : 2B \rightarrow \mathbb{R}^n$, $n \geq 3$, is a local homeomorphism with bounded distortion K_I , then its radius of injectivity in B is bounded from below by a constant depending only on n and $\text{ess sup } K_I$. As a consequence, the multiplicity $N(y, f, B)$ is bounded by $C(n, \text{ess sup } K_I)$ for all $y \in \mathbb{R}^n$.

The boundedness of K_I can be replaced by the condition

$$\exp(\lambda K_I^{1/(n-1)}) \in L^1(2B),$$

but this cannot be relaxed any further [13,17]. Surprisingly, the multiplicity bound remains true under a much weaker condition, namely $K_I \in L^1$. Example 7.2 below shows that $K_I^q \in L^1$ with $q < 1$ does not suffice. The mappings $f_j(z) = e^{jz}$ show that all results discussed here fail when $n = 2$.

Theorem 5.1. *Let $f \in W_{loc}^{1,n}(\Omega, \mathbb{R}^n)$, $n \geq 3$, be a local homeomorphism such that $\mathcal{K}_\Omega[f] < \infty$. If B is a ball such that $4B \Subset \Omega$, then $N(y, f, B) \leq C(n, \mathcal{K}_{4B}[f])$ for all $y \in \mathbb{R}^n$.*

Proof. We may assume that B is the unit ball \mathbb{B} . Let $x_1, \dots, x_m \in f^{-1}(y) \cap \mathbb{B}$. Moreover, let r_j be the largest radius r so that the x_j -component $U(x_j, r)$ of $f^{-1}B(y, r)$ satisfies $U(x_j, r) \subset \mathbb{B}_3$. By Lemma 3.2 f is a homeomorphism from $U(x_j, r_j)$ onto $B(y, r_j)$. We denote by s_j the largest radius s such that $\overline{B}(x_j, s) \subset \overline{U}(x_j, r_j)$. Then $f\overline{B}(x_j, s_j)$ intersects both y and $S(y, r_j)$. We notice that since $x_j \in \mathbb{B}$ and since the balls $B(x_j, s_j)$ are pairwise disjoint, there exist at most $N(n)$ indices j for which $s_j \geq 1$. Thus we may assume that $B(x_j, s_j) \subset \mathbb{B}_2$ for every $1 \leq j \leq m$.

We now fix $1 \leq j \leq m$ and a point $a_j \in \overline{U}(x_j, r_j) \cap \mathbb{S}_3$. We apply Lemma 3.5 with $B(y_0, r) = B(y, r_j)$, $y_1 = f(a_j)$ and $E = f(\overline{B}(x_j, s_j))$, obtaining a point q_j and a number $\sigma_j > 0$. For $\sigma_j < t < 4\sigma_j/3$ choose $w_t \in \overline{B}(x_j, s_j)$ such that $f(w_t) \in S(q_j, t)$. We apply Lemma 3.4 with $G = U(x_j, r_j)$, $D = \mathbb{B}_3$, $a = a_j$, $\mathcal{B} = B(q_j, t)$ and $b' = f(w_t)$. As a result we obtain $0 < \phi_t < \pi$ such that the spherical cap $\mathcal{C}_t := C_{S(q_j, t)}(f(w_t), \phi_t)$ satisfies $\mathcal{C}_t^* \subset \mathbb{B}_3$ and $\overline{\mathcal{C}_t^*} \cap \mathbb{S}_3$ contains some point c_t . Consequently, for every path γ joining $f(w_t)$ and $f(c_t)$ in \mathcal{C}_t , the maximal f -lifting γ^* of γ starting at w_t starts from \mathbb{B}_2 and leaves \mathbb{B}_3 . Following [23, 10.2], we will choose a particular family Γ_t of such paths.

Let us say that a circular arc is *short* if it is contained in a half-circle. The family Γ_t will consist of all short circular arcs that connect $f(w_t)$ to $f(c_t)$ within \mathcal{C}_t . More precisely, let h be a Möbius transformation that maps $f(w_t)$ to infinity and $S(q_j, t) \setminus \{f(w_t)\}$ to \mathbb{R}^{n-1} . Observe that $h(\mathcal{C}_t)$ is the complement of a ball in \mathbb{R}^{n-1} . The convexity of $\mathbb{R}^{n-1} \setminus h(\mathcal{C}_t)$ implies that there exists an $(n - 2)$ -hemisphere V such that $h(f(c_t)) + sv \in h(\mathcal{C}_t)$ for every $s > 0$ and $v \in V$.

Introduce a family of curves $I_v : [0, \infty) \rightarrow \mathcal{C}_t$, defined by

$$I_v(s) = h^{-1}(h(f(c_t)) + s^{-1}v),$$

and denote by I_v^* the maximal f -lifting of I_v starting at w_t . Now let $0 < \ell(v) < \infty$ be the smallest number such that $I_v^*(\ell(v)) \in \mathbb{S}_3$. Let

$$\Gamma_t = \{I_v^*|_{[0, \ell(v)]} : v \in V_t\}.$$

We write $f\Gamma_t$ for the image of Γ_t under f .

There is a lower bound for the spherical modulus of $f\Gamma_t$, namely [23, Theorem 10.2]

$$M^S(f\Gamma_t) \geq \frac{C(n)}{t}. \tag{5.1}$$

Let

$$\Gamma'_j = \{\gamma : \gamma \in f\Gamma_t \text{ for some } \sigma_j < t < 4\sigma_j/3\},$$

and let Γ_j^* be the family of the corresponding lifts γ^* starting at w_t . Then integrating (5.1) we obtain

$$M\Gamma'_j \geq \int_{\sigma_j}^{4\sigma_j/3} \frac{C(n)}{t} dt \geq C(n). \tag{5.2}$$

As observed earlier, every $\gamma \in \Gamma_j^*$ starts at \mathbb{B}_2 and leaves \mathbb{B}_3 . We denote by E_j the smallest closed subset of $\overline{\mathbb{B}_3} \setminus \mathbb{B}_2$ that contains $|\gamma| \cap (\overline{\mathbb{B}_3} \setminus \mathbb{B}_2)$ for all $\gamma \in \Gamma_j^*$. Note that

$$E_j \subset f^{-1}(\overline{B}(y, 2r_j) \setminus B(y, r_j/10)) \tag{5.3}$$

by part (iii) of Lemma 3.5. Since the characteristic function χ_{E_j} is an admissible function for Γ_j^* , we have

$$M_{K_I} \Gamma_j^* \leq \int_{E_j} K_I(x, f) dx. \tag{5.4}$$

The generalized Poletsky inequality $M \Gamma_j' \leq M_{K_I} \Gamma_j^*$ [14, Theorem 4.1], together with (5.2) and (5.4) yield

$$mC(n) \leq \sum_{j=1}^m M \Gamma_j' \leq \sum_{j=1}^m \int_{E_j} K_I(x) dx \leq \left(\sup_{x \in \mathbb{B}_3 \setminus \mathbb{B}_2} \sum_{j=1}^m \chi_{E_j}(x) \right) \times \int_{3B} K_I(x, f) dx. \tag{5.5}$$

Claim 1. *There exists $M = M(n, \mathcal{K}_{4B}[f])$ such that*

$$\sum_{j=1}^m \chi_{E_j}(x) \leq M \quad \text{for every } x \in \mathbb{B}_3 \setminus \mathbb{B}_2. \tag{5.6}$$

By virtue of (5.5), Theorem 5.1 follows from Claim 1. In the rest of this section we prove (5.6).

Let $x \in \mathbb{B}_3 \setminus \mathbb{B}_2$ be a point covered by M of the sets E_j . After relabeling we have $x \in E_j$ for $1 \leq j \leq M$, and $r_1 \leq r_2 \leq \dots \leq r_M$. Since disjoint sets have disjoint preimages, (5.3) implies $r_M \leq 20r_1$.

Choose $\tau > 0$ such that $B(x, \tau) \subset \mathbb{B}_3$ and f is injective in $\bar{B}(x, \tau)$. For $1 \leq j \leq M$ there exists $\gamma_j^* \in \Gamma_j^*$ which meets $B(x, \tau)$. Let w_j be the starting point of γ_j^* , and let γ_j be the subcurve of γ_j^* that begins at w_j and ends once it meets $\bar{B}(x, \tau)$.

Claim 2. *For $1 \leq j \leq M$ there is a curve τ_j that joins y to $f(w_j)$ within $\bar{B}(y, r_j)$ in such a way that the union of $|\tau_j|$ and $|f \circ \gamma_j|$ can be mapped onto a line segment by an L -biLipschitz mapping $g: \mathbb{R}^n \rightarrow \mathbb{R}^n$. Here L is a universal constant.*

Proof. Note that the image $f \circ \gamma_j$ is a short circular arc contained in the sphere $S(q, t)$ of Lemma 3.5. Part (iii) of Lemma 3.5 implies

$$\text{dist}(y, |f \circ \gamma_j|) \geq \text{dist}(y, S(q, t)) \geq \frac{1}{10} r_j \geq \frac{1}{40} \text{diam } |f \circ \gamma_j|. \tag{5.7}$$

There are two cases. If $y \in B(q, t)$, then τ_j is the line segment connecting y to $f(w_j)$. By virtue of (5.7), the distance from y to $S(q, t)$ is comparable to t . Therefore, the angle between τ_j and the sphere $S(q, t)$ is bounded from below by a universal constant, and the claim follows.

Suppose that $y \notin B(q, t)$. Let $\rho_j := |f(w_j) - y|$. Note that $r_j/10 \leq \rho_j \leq r_j$. Let p be the point of the sphere $S(y, \rho_j)$ that is farthest from q , namely

$$p = y - \rho_j \frac{q - y}{|q - y|}.$$

We choose τ_j as the union of the line segment connecting y to p and the geodesic arc on $S(y, \rho_j)$ from p to $f(w_j)$. Once again, the angle between τ_j and the sphere $S(q, t)$ is bounded from below by a universal constant. \square

Let η_j , $1 \leq j \leq M$, be the curve obtained by concatenating $-(f \circ \gamma_j)$ with $-\tau_j$, where $-$ indicates the reversal of orientation. Note that η_j begins in $f\bar{B}(x, \tau)$, proceeds along a circular arc to $f(w_j)$, and ends at y . Its f -lifting η_j^* starting in $\bar{B}(x, \tau)$ is contained in $\bar{\mathbb{B}}_3$ and ends at x_j .

Claim 3. *There exists $\epsilon = \epsilon(n, M)$ such that $\epsilon \rightarrow 0$ as $M \rightarrow \infty$, and*

$$\min_{1 \leq i < j \leq M} d_{\mathcal{H}}(|\eta_i|, |\eta_j|) \leq \epsilon r_1 / L. \tag{5.8}$$

Proof. We begin our proof of Claim 3 by observing that $|\eta_j| \subset B(y, 2r_M) \subset B(y, 40r_1)$. For $\epsilon > 0$ let $Z = \{z_1, \dots, z_N\}$ be an $(\epsilon r_1 / L)$ -net in $B(y, 40r_1)$, where $N = N(\epsilon, n)$. The set of all nonempty subsets of Z is an $(\epsilon r_1 / L)$ -net in the set of all nonempty closed subsets of $B(y, 40r_1)$ equipped with the Hausdorff metric. If $M > 2^N$, then by

the pigeonhole principle there exist $i < j$ such that $|\eta_i|$ and $|\eta_j|$ are within the distance $(\epsilon r_1/L)$ from the same subset of Z . Claim 3 follows. \square

Fix i, j , and ϵ as in Claim 3, and let $g : \mathbb{R}^n \rightarrow \mathbb{R}^n$ be the L -biLipschitz mapping from Claim 2. By replacing f with $g \circ f$, which has a comparable distortion function K_I , we may assume that $|\eta_j|$ is a line segment. For $\delta > 0$ we denote by $W(\delta)$ the open δ -neighborhood of $|\eta_j|$. Let $W^*(\delta)$ be the x_j -component of $f^{-1}W(\delta)$.

Claim 4. *If $\delta > \epsilon r_1$, then $W^*(\delta) \cap \mathbb{S}_4 \neq \emptyset$.*

Proof. Since $\delta > \epsilon r_1$, we have $|\eta_i| \subset W(\delta)$. Suppose to the contrary that $W^*(\delta) \subset \mathbb{B}_4$. Then $W^*(\delta) \Subset \Omega$, which by Lemma 3.2 implies that $f : W^*(\delta) \rightarrow W(\delta)$ is a homeomorphism. This contradicts the fact that the f -liftings of η_i and η_j starting in $\bar{B}(x, \tau)$ end at different points, namely x_i and x_j . \square

Let δ_0 be the supremum of all numbers δ such that $W^*(\delta) \subset \mathbb{B}_4$. Since f is a local homeomorphism, $\delta_0 > 0$. By Lemma 3.2, $f : W^*(\delta) \xrightarrow{\text{hom}} W(\delta)$ for every $0 < \delta < \delta_0$. By Claim 4 we have $\delta_0 \leq \epsilon r_1$.

Choose a point $a \in \partial W^*(\delta_0) \cap \mathbb{S}_4$. Let $a' = f(a)$. Since $a' \in \partial W(\delta_0)$, there exists $p \in |\eta_j|$ such that $|a' - p| = \delta_0$. For $\delta_0 < t < \frac{1}{2} \text{diam}|\eta_j|$ choose $b'_t \in |\eta_j| \cap S(p, t)$. We apply Lemma 3.4 with $G = W^*(\delta_0)$, $D = \mathbb{B}_4$, $a = a$, $\mathcal{B} = B(p, t)$ and $b' = b'_t$. As a result we obtain $0 < \phi_t < \pi$ such that the spherical cap $\mathcal{C}_t := C_{S(p,t)}(b'_t, \phi_t)$ satisfies $\mathcal{C}_t^* \subset \mathbb{B}_4$ and $\bar{\mathcal{C}}_t^* \cap \mathbb{S}_4$ contains some point c_t . Consequently, for every path γ joining b'_t and $f(c_t)$ in \mathcal{C}_t , the maximal f -lifting γ^* of γ starting at $f^{-1}(b'_t) \cap |\eta_j^*|$ starts from \mathbb{B}_3 and leaves \mathbb{B}_4 . Let Γ be the family of all such paths γ and let Γ^* be the family of the lifts γ^* . From [23, Theorem 10.2] we have

$$M\Gamma \geq C(n) \int_{\epsilon r_1}^{\text{diam}(\eta_j)/2} \frac{dt}{t} \geq C(n) \log \frac{\text{diam}(\eta_j)}{2\epsilon r_1}.$$

By (5.7) we have $\text{diam}|\eta_j| \geq cr_1$ with a universal constant $c > 0$. Therefore,

$$M\Gamma \geq C(n) \log \frac{1}{\epsilon}. \tag{5.9}$$

On the other hand, since the characteristic function $\chi_{\mathbb{B}_4 \setminus \mathbb{B}_3}$ is an admissible function for Γ_j , we obtain

$$M_{K_I} \Gamma^* \leq \int_{\mathbb{B}_4 \setminus \mathbb{B}_3} K_I(x, f) dx.$$

Combining this with (5.9) and using the Poletsky inequality again, we have $\epsilon \geq C(n, \mathcal{K}_{4B}[f])$, hence $M \leq C(n, \mathcal{K}_{4B}[f])$. This gives (5.6). The proof of Theorem 5.1 is complete. \square

6. Proof of Theorem 1.1

Denote $f^\lambda(x) = f(x) + \lambda x$, $\lambda > 0$. Then $f^\lambda \in W_{\text{loc}}^{1,n}(\Omega, \mathbb{R}^n)$. Moreover, by [14, Lemma 10],

$$K_I(x, f^\lambda) \leq C(\delta, n) K_I(x, f) \quad \text{and} \quad \|(Df^\lambda)^{-1}(x)\| \leq C(\delta, \lambda) \tag{6.1}$$

for almost every $x \in \Omega$. Thus f^λ is discrete and open for every $\lambda > 0$ by Theorem 1.2. Furthermore, by [14, Lemma 13] f^λ is a local homeomorphism. (Although [14, Lemma 13] imposes a stronger condition on the distortion of f , this condition is only used to ensure that f is discrete and open.) Since $f^\lambda \rightarrow f$ locally uniformly, the following proposition implies that f is a local homeomorphism, completing the proof of Theorem 1.1.

Proposition 6.1. *Suppose that a mapping $f \in W_{\text{loc}}^{1,n}(\Omega, \mathbb{R}^n)$ with $\mathcal{K}_\Omega[f] < \infty$ can be uniformly approximated by local homeomorphisms $f_j \in W_{\text{loc}}^{1,n}(\Omega, \mathbb{R}^n)$ such that $\sup_j \mathcal{K}_\Omega[f_j] < \infty$. Then f is a local homeomorphism.*

Proof. By [14, Proposition 7] it suffices to show that f is discrete and open. If $n = 2$, this is due to Iwaniec and Šverák [10]. Thus we assume that $n \geq 3$. Let $B = B(x_0, R)$ be a ball such that $8B \Subset \Omega$. We will show that

$$N(y, f, B) \leq C \quad \text{for a.e. } y \in \mathbb{R}^n, \quad (6.2)$$

where $C < \infty$ does not depend on y . Proposition 2.2 will then imply that f is discrete and open in B .

Applying Theorem 5.1 to f_j , we obtain

$$N(y, f_j, 2B) \leq C \quad \text{for every } y \in \mathbb{R}^n,$$

where C depends only on $\sup_j \mathcal{K}_\Omega[f_j]$ and n .

We fix $R < t < 2R$ so that $\mathcal{H}^{n-1}(fS(x_0, t)) < \infty$, and a point $y \in fB \setminus fS(x_0, t)$. Let $d = \text{dist}(y, fS(x_0, t))$. Since $f_j \rightarrow f$ locally uniformly, there exists j_0 such that $|f_j(x) - f(x)| < d/2$ for all $j \geq j_0$ and all $x \in S(x_0, t)$. Consequently, the restrictions of f_j and f to $S(x_0, t)$ are homotopic via the straight-line homotopy that takes values in $\mathbb{R}^n \setminus \{y\}$. It follows that

$$\deg(y, f, B(x_0, t)) = \deg(y, f_j, B(x_0, t)) \leq N(y, f_j, 2B) \leq C$$

for all $j \geq j_0$. Since $N(y, f, B) \leq N(y, f, B(x_0, t)) = \deg(y, f, B(x_0, t))$ for almost every $y \in \mathbb{R}^n$, we conclude that (6.2) indeed holds. The proof is complete. \square

7. Concluding remarks

Corollary 7.1. *Suppose that $f \in W_{\text{loc}}^{1,n}(\mathbb{R}^n, \mathbb{R}^n)$ is a nonconstant mapping such that $K_I(\cdot, f) \in L_{\text{loc}}^1(\mathbb{R}^n)$. If there exists $\delta > -1$ such that $Df(x) \in \mathcal{M}(\delta)$ for almost every $x \in \mathbb{R}^n$, then f is a homeomorphism.*

Proof. As in the proof of Theorem 1.1 we have that $f^\lambda(x) = f(x) + \lambda x$ is a local homeomorphism for all $\lambda > 0$. Since $(Df^\lambda)^{-1} \in L^\infty(\mathbb{R}^n)$, it follows from [14, Lemma 12] that

$$\liminf_{x \rightarrow a} \frac{|f^\lambda(x) - f^\lambda(a)|}{|x - a|} \geq \frac{\lambda}{2} > 0 \quad (7.1)$$

for all $a \in \mathbb{R}^n$. By a theorem of John [11, p. 87], f^λ is a homeomorphism. Since f is discrete and open by Theorem 1.1, we can apply [14, Proposition 7] and conclude that f is a homeomorphism. \square

Sharpness of Theorem 5.1 is demonstrated by the following example which combines the ideas from [2] and [13].

Example 7.2. For any $q < 1$ there exists a sequence of mappings $f_j \in W^{1,3}(\mathbb{B}, \mathbb{R}^3)$ such that

$$\sup_j \int_{\mathbb{B}} K_I^q(x, f_j) dx < \infty \quad \text{and} \quad N(0, f_j, B(0, 1/4)) \rightarrow \infty.$$

Proof. By a version of Zorich's construction (see [9,21]) there exists a mapping $\phi \in W^{1,3}(\mathbb{R}^3, \mathbb{R}^3)$ such that $K_I(\cdot, \phi) \in L^\infty(\mathbb{R}^3)$, ϕ is a local homeomorphism outside of $\mathbb{R} \times (2\mathbb{Z} + 1)^2$, and ϕ is 4-periodic in the last two variables. Therefore, it suffices for us to construct biLipschitz homeomorphisms $f_j : \mathbb{B} \rightarrow \mathbb{R}^3$ such that

- (i) $\sup_j \int_{\mathbb{B}} K_I^q(x, f_j) dx < \infty$;
- (ii) $f_j(\mathbb{B}) \subset \mathbb{D} \times \mathbb{R}$ (here $\mathbb{D} \subset \mathbb{R}^2$ is the unit disc);
- (iii) $f_j(\mathbb{B}_{1/4})$ contains a line segment $\{0\} \times [-L, L] \subset \mathbb{R}^2 \times \mathbb{R}$ where $L \rightarrow \infty$ as $j \rightarrow \infty$.

The compositions $\phi \circ f_j$ will be mappings with large multiplicity.

For $y \in \mathbb{R}^3$ let $s(y) = \sqrt{y_1^2 + y_2^2}$. For $\alpha > 2$ we define a mapping $x = g(y)$ by

$$x_i = s(y)^{\alpha-1} y_i, \quad i = 1, 2;$$

$$x_3 = s(y) y_3.$$

Since $s(x) = s(y)^\alpha$, the inverse mapping $y = f(x)$ outside the set $\{s(x) = 0\}$ is given by

$$\begin{aligned} y_i &= s(x)^{1/\alpha-1} x_i, \quad i = 1, 2; \\ y_3 &= s(x)^{-1/\alpha} x_3, \quad s(x) \neq 0. \end{aligned}$$

Let $\Omega = \{x \in \mathbb{R}^3: s(x) < 1, |x_3| < 1\}$ and $\Omega' = f(\Omega)$. We restrict our attention to $y \in \Omega'$, where in particular $s(y) < 1$. Elementary computations show that

$$\begin{aligned} \|Dg(y)\| &\leq C \max(s(y), |y_3|) \quad \text{and} \\ J(y, g) &\geq C s(y)^{2(\alpha-1)+1}. \end{aligned}$$

Therefore,

$$\frac{\|Dg(y)\|^3}{J(y, g)} \leq C s(y)^{2(1-\alpha)-1} \max(s(y)^3, |y_3|^3). \tag{7.2}$$

Since

$$\frac{\|Dg(y)\|^3}{J(y, g)} = K_I(x, f),$$

inequality (7.2) can be used to estimate $K_I(x, f)$ as follows.

$$K_I(x, f) \leq C s(x)^{(2(1-\alpha)-1)/\alpha} \max(s(x)^{3/\alpha}, s(x)^{-3/\alpha} |x_3|^3) \leq C s(x)^{-(2\alpha+2)/\alpha},$$

where at the last step we used $|x_3| < 1$. We achieve $\int_\Omega K_I(x, f)^q dx < \infty$ by choosing α large enough so that

$$\frac{2\alpha + 2}{\alpha} q < 2.$$

The mapping f constructed thus far is not in $W^{1,3}$, and is not even continuous. However, this can be corrected by replacing $s(y)$ with $s_j(y) = \sqrt{y_1^2 + y_2^2 + 1/j^2}$. The mapping $x = g_j(y)$ given by

$$\begin{aligned} x_i &= s_j(y)^{\alpha-1} y_i, \quad i = 1, 2; \\ x_3 &= s_j(y) y_3, \end{aligned}$$

is biLipschitz; we denote the inverse by f_j . The computation of $\|Dg_j\|$ and $J(\cdot, g_j)$ goes through exactly as before and shows that the integral of $K_I^q(\cdot, f_j)$ is bounded independently of ϵ_j . Since $g_j(0, 0, y_3) = (0, 0, y_3/j)$, we have $f_j(0, 0, x_3) = (0, 0, jx_3)$. Thus, this mapping f_j fulfills the requirements (i)–(iii). \square

Acknowledgements

The research was partially carried out during Rajala’s visit to Syracuse University. He wishes to thank the Department of Mathematics for its hospitality. Part of this research was done when Kovalev and Onninen were visiting Princeton University. Also, the authors are thankful to the referee for careful reading of the manuscript and several valuable comments.

References

[1] K. Astala, T. Iwaniec, G.J. Martin, J. Onninen, Extremal mappings of finite distortion, Proc. London Math. Soc. (3) 91 (3) (2005) 655–702.
 [2] J.M. Ball, Global invertibility of Sobolev functions and the interpenetration of matter, Proc. Roy. Soc. Edinburgh Sect. A 88 (3–4) (1981) 315–328.
 [3] V.M. Gol’dshstein, The behavior of mappings with bounded distortion when the distortion coefficient is close to one, Sibirsk. Mat. Zh. 12 (1971) 1250–1258.
 [4] V.M. Gol’dshstein, S.K. Vodopyanov, Quasiconformal mappings, and spaces of functions with first generalized derivatives, Sibirsk. Mat. Zh. 17 (3) (1976) 515–531.
 [5] J. Heinonen, T. Kilpeläinen, BLD-mappings in $W^{2,2}$ are locally invertible, Math. Ann. 318 (2) (2000) 391–396.
 [6] J. Heinonen, P. Koskela, Sobolev mappings with integrable dilatations, Arch. Ration. Mech. Anal. 125 (1) (1993) 81–97.

- [7] S. Hencl, P. Koskela, Mappings of finite distortion: Discreteness and openness for quasi-light mappings, *Ann. Inst. H. Poincaré Anal. Non Linéaire* 22 (3) (2005) 331–342.
- [8] S. Hencl, J. Malý, Mappings of finite distortion: Hausdorff measure of zero sets, *Math. Ann.* 324 (3) (2002) 451–464.
- [9] T. Iwaniec, G. Martin, *Geometric Function Theory and Non-Linear Analysis*, Oxford Univ. Press, New York, 2001.
- [10] T. Iwaniec, V. Šverák, On mappings with integrable dilatation, *Proc. Amer. Math. Soc.* 118 (1) (1993) 181–188.
- [11] F. John, On quasi-isometric mappings. I, *Comm. Pure Appl. Math.* 21 (1968) 77–110.
- [12] P. Koskela, J. Onninen, Mappings of finite distortion: Capacity and modulus inequalities, *J. Reine Angew. Math.* 599 (2006) 1–26.
- [13] P. Koskela, J. Onninen, K. Rajala, Mappings of finite distortion: Injectivity radius of a local homeomorphism, in: *Future Trends in Geometric Function Theory*, in: *Rep. Univ. Jyväskylä Dep. Math. Stat.*, vol. 92, Univ. Jyväskylä, Jyväskylä, 2003, pp. 169–174.
- [14] L.V. Kovalev, J. Onninen, On invertibility of Sobolev mappings, preprint, 2008, arXiv:0812.2350.
- [15] J.J. Manfredi, E. Villamor, An extension of Reshetnyak’s theorem, *Indiana Univ. Math. J.* 47 (3) (1998) 1131–1145.
- [16] O. Martio, S. Rickman, J. Väisälä, Topological and metric properties of quasiregular mappings, *Ann. Acad. Sci. Fenn. Ser. A I* 488 (1971).
- [17] J. Onninen, Mappings of finite distortion: Minors of the differential matrix, *Calc. Var. Partial Differential Equations* 21 (4) (2004) 335–348.
- [18] K. Rajala, The local homeomorphism property of spatial quasiregular mappings with distortion close to one, *Geom. Funct. Anal.* 15 (5) (2005) 1100–1127.
- [19] K. Rajala, Reshetnyak’s theorem and the inner distortion, *Pure Appl. Math. Q.*, in press, University of Jyväskylä preprint, No. 336, 2007.
- [20] K. Rajala, Remarks on the Iwaniec–Šverák conjecture, University of Jyväskylä preprint, No. 377, 2009.
- [21] S. Rickman, *Quasiregular Mappings*, Springer-Verlag, Berlin, 1993.
- [22] Q. Tang, Almost-everywhere injectivity in nonlinear elasticity, *Proc. Roy. Soc. Edinburgh Sect. A* 109 (1–2) (1988) 79–95.
- [23] J. Väisälä, *Lectures on n -Dimensional Quasiconformal Mappings*, *Lecture Notes in Math.*, vol. 229, Springer-Verlag, Berlin, 1971.
- [24] V.A. Zorich, M.A. Lavrentyev’s theorem on quasiconformal space maps, *Mat. Sb. (N.S.)* 74 (116) (1967) 417–433.