



Partial Differential Equations

# Uniform asymptotic formulae for Green's kernels in regularly and singularly perturbed domains

Vladimir Maz'ya<sup>a,b,c</sup>, Alexander Movchan<sup>a</sup>

<sup>a</sup> Department of Mathematical Sciences, University of Liverpool, Liverpool L69 3BX, UK

<sup>b</sup> Department of Mathematics, Ohio State University, 231 W 18th Avenue, Columbus, OH 43210, USA

<sup>c</sup> Department of Mathematics, Linköping University, SE-581 83 Linköping, Sweden

Received 13 December 2005; accepted after revision 15 May 2006

Presented by Évariste Sanchez Palencia

---

## Abstract

Asymptotic formulae for Green's kernels  $G_\varepsilon(\mathbf{x}, \mathbf{y})$  of various boundary value problems for the Laplace operator are obtained in regularly perturbed domains and certain domains with small singular perturbations of the boundary, as  $\varepsilon \rightarrow 0$ . The main new feature of these asymptotic formulae is their uniformity with respect to the independent variables  $\mathbf{x}$  and  $\mathbf{y}$ . **To cite this article:** V. Maz'ya, A. Movchan, C. R. Acad. Sci. Paris, Ser. I 343 (2006).

© 2006 Académie des sciences. Published by Elsevier SAS. All rights reserved.

## Résumé

**Formules asymptotiques uniformes pour des noyaux de Green dans des domaines avec perturbations régulières et singulières.** Des formules asymptotiques sont obtenues pour des noyaux de Green  $G_\varepsilon(\mathbf{x}, \mathbf{y})$  de divers problèmes aux limites pour l'opérateur de Laplace dans des domaines régulièrement perturbés et certains domaines avec des petites perturbations singulières du bord, quand  $\varepsilon \rightarrow 0$ . Le caractère novateur de ces formules asymptotiques réside dans leur uniformité par rapport aux variables indépendantes  $\mathbf{x}$  et  $\mathbf{y}$ . **Pour citer cet article :** V. Maz'ya, A. Movchan, C. R. Acad. Sci. Paris, Ser. I 343 (2006).

© 2006 Académie des sciences. Published by Elsevier SAS. All rights reserved.

---

## Version française abrégée

Cette Note s'intéresse à la question des approximations asymptotiques uniformes des fonctions de Green pour des domaines à la fois régulièrement et singulièrement perturbés. Tout d'abord, nous affinons l'une des formules classiques d'Hadamard dans le cas d'une perturbation régulière. Puis, nous considérons plusieurs types de perturbations singulières, notamment dans des domaines avec un nombre fini de trous, de tiges minces et un cône tronqué. Pour ces géométries et différents types de conditions limites, nous déduisons les représentations asymptotiques uniformes des noyaux de Green pour l'opérateur Laplacien. Les techniques utilisées dans les preuves des théorèmes du texte principal de la Note se fondent sur la méthode des développements asymptotiques combinés, qui est discutée en détail dans la monographie en deux tomes [4]. Les formules asymptotiques uniformes pour les noyaux de Green peuvent

---

E-mail addresses: [vlmaz@mai.liu.se](mailto:vlmaz@mai.liu.se) (V. Maz'ya), [abm@liv.ac.uk](mailto:abm@liv.ac.uk) (A. Movchan).

être avantageusement utilisées dans des algorithmes numériques pour résoudre des problèmes avec conditions limites, dans des domaines avec des frontières perturbées singulièrement.

## 1. Introduction

In 1907, Hadamard was awarded the Prix Vaillant by the Académie des Sciences de Paris for his work [2]. Among much else, this memoir includes investigation of Green's kernels of some boundary value problems, and, in particular, their dependence on a small regular perturbation of the domain. Hadamard's asymptotic formulae played a significant role at the dawn of functional analysis (see [3]). The list of subsequent applications includes extremal problems in the complex function theory, a biharmonic maximum principle for hyperbolic surfaces, shape sensitivity and optimisation analysis, free boundary problems, Brownian motion on hypersurfaces, theory of reproducing kernels. Analogues of Hadamard's formulae were obtained for general elliptic boundary value problems [1] and for the heat equation [5].

The issue of asymptotic formulae for Green's kernels in singularly perturbed domains attracted less attention. In this respect, we mention the papers [6,7], where certain estimates for Green's functions in domains with small holes were obtained.

An important drawback of the estimates just mentioned, which is also inherent in the classical Hadamard formulae, is the non-uniformity with respect to the independent variables. The present article settles the question of uniformity left open in earlier work, both for regularly and singularly perturbed domains. First, we sharpen one of Hadamard's classical formulae in the case of a regular perturbation. Next, we consider several types of singularly perturbed domains including domains with a finite number of holes, thin rods and a truncated cone. For these geometries and different types of boundary conditions, we derive uniform asymptotic representations of Green's kernels for the Laplacian. The asymptotic formulae presented here for singularly perturbed problems involve Green's functions and other auxiliary solutions of certain model boundary value problems independent of  $\varepsilon$ .

The techniques used in the proofs of theorems of the main text of the paper are based on the method of compound asymptotic expansions discussed in detail in the two-volume monograph [4]. The uniform asymptotic formulae for Green's kernels can be efficiently used in numerical algorithms for solving boundary value problems in domains with singularly perturbed boundaries.

## 2. Uniform Hadamard's type formula

Let  $\Omega$  be a planar domain with compact closure  $\bar{\Omega}$  and smooth boundary  $\partial\Omega$ . Also, let another domain  $\Omega(\varepsilon)$ , depending on a small positive parameter  $\varepsilon$ , lie inside  $\Omega$ . By  $\delta_z$  we denote a smooth positive function defined on  $\partial\Omega$ , and assume that  $\varepsilon\delta_z$  is the distance between a point  $\mathbf{z} \in \partial\Omega$  and  $\partial\Omega_\varepsilon$ . One of the results in Hadamard's paper [2] is the following formula, which relates Green's functions  $G$  and  $G_\varepsilon$  for the Dirichlet boundary value problem for the Laplacian in  $\Omega$  and  $\Omega_\varepsilon$ :

$$G_\varepsilon(\mathbf{x}, \mathbf{y}) - G(\mathbf{x}, \mathbf{y}) + \varepsilon \int_{\partial\Omega} \frac{\partial G}{\partial \nu_z}(\mathbf{x}, \mathbf{z}) \frac{\partial G}{\partial \nu_z}(\mathbf{z}, \mathbf{y}) \delta_z \, ds_z = O(\varepsilon^2). \quad (1)$$

This asymptotic relation holds, in particular, when either  $\mathbf{x}$  or  $\mathbf{y}$  is placed at a positive distance from  $\partial\Omega$ , independent of  $\varepsilon$ . However, one can see from the simplest example of two concentric disks  $\Omega$  and  $\Omega_\varepsilon$  that (1) may fail when  $\mathbf{x}$  and  $\mathbf{y}$  approach the same point at  $\partial\Omega_\varepsilon$ . In the next theorem, we improve Hadamard's formula by obtaining a uniform asymptotic representation for  $G_\varepsilon - G$ .

**Theorem 1.** *Let  $\mathbf{x}$  and  $\mathbf{y}$  be points of  $\bar{\Omega}_\varepsilon$  situated in a sufficiently thin neighbourhood of  $\partial\Omega$ . The right-hand side in (1) can be replaced by*

$$\begin{aligned} & \frac{1}{4\pi} \log \frac{|z(\mathbf{x}) - z(\mathbf{y})|^2 + (\rho_x - \varepsilon\delta_{z(x)} + \rho_y - \varepsilon\delta_{z(y)})^2}{|z(\mathbf{x}) - z(\mathbf{y})|^2 + (\rho_x + \rho_y)^2} \\ & + \frac{1}{2\pi} \frac{\varepsilon(\delta_{z(x)} + \delta_{z(y)})(\rho_x + \rho_y)}{|z(\mathbf{x}) - z(\mathbf{y})|^2 + (\rho_x + \rho_y)^2} + O\left(\frac{\varepsilon^2(\rho_x + \rho_y)}{|z(\mathbf{x}) - z(\mathbf{y})|^2 + (\rho_x + \rho_y)^2}\right), \end{aligned} \quad (2)$$

where  $\mathbf{z}(\mathbf{x})$  is the point of  $\partial\Omega$  nearest to  $\mathbf{x}$ ,  $\rho_x = |\mathbf{x} - \mathbf{z}(\mathbf{x})|$ , and the notations  $\mathbf{z}(\mathbf{y})$ ,  $\rho_y$  have the same meaning for the point  $\mathbf{y}$ .

We note that the principal term in (2) plays the role of a boundary layer, and when  $|\mathbf{z}(\mathbf{x}) - \mathbf{z}(\mathbf{y})| < \text{Const } \varepsilon$  and  $\rho_x + \rho_y < \text{Const } \varepsilon$ , it has the order  $O(1)$ , whereas the remainder term has the order  $O(\varepsilon)$ .

### 3. Dirichlet problem for a domain with a small inclusion

Let  $\Omega$  and  $\omega$  be bounded domains in  $\mathbf{R}^n$ . We assume that  $\Omega$  and  $\omega$  contain the origin  $O$  and introduce the domain  $\omega_\varepsilon = \{\mathbf{x}: \varepsilon^{-1}\mathbf{x} \in \omega\}$ . Without loss of generality, it is assumed that the minimum distance between the origin and the points of  $\partial\Omega$  as well as the maximum distance between the origin and the points of  $\partial\omega$  are equal to 1. Let  $G_\varepsilon$  be Green's function of the Dirichlet problem for the Laplace operator in  $\Omega_\varepsilon = \Omega \setminus \bar{\omega}_\varepsilon$ . We use the notation  $|S^{n-1}|$  for the  $(n - 1)$ -dimensional measure of the unit sphere.

**Theorem 2.** *Let  $n > 2$ . By  $G$  and  $\mathcal{G}$  we denote Green's functions of the Dirichlet problems in  $\Omega$  and  $\mathbf{R}^n \setminus \bar{\omega}$ , respectively.*

*Let  $H$  be the regular part of  $G$ , that is  $H(\mathbf{x}, \mathbf{y}) = ((n - 2)|S^{n-1}|)^{-1}|\mathbf{x} - \mathbf{y}|^{2-n} - G(\mathbf{x}, \mathbf{y})$ , and let  $P$  stand for the harmonic capacity potential of  $\bar{\omega}$ . Then*

$$\begin{aligned} G_\varepsilon(\mathbf{x}, \mathbf{y}) &= G(\mathbf{x}, \mathbf{y}) + \varepsilon^{2-n}\mathcal{G}(\varepsilon^{-1}\mathbf{x}, \varepsilon^{-1}\mathbf{y}) - ((n - 2)|S^{n-1}|)^{-1}|\mathbf{x} - \mathbf{y}|^{2-n} \\ &\quad + H(0, \mathbf{y})P(\varepsilon^{-1}\mathbf{x}) + H(\mathbf{x}, 0)P(\varepsilon^{-1}\mathbf{y}) - H(0, 0)P(\varepsilon^{-1}\mathbf{x})P(\varepsilon^{-1}\mathbf{y}) \\ &\quad - \varepsilon^{n-2} \text{cap } \bar{\omega} H(\mathbf{x}, 0)H(0, \mathbf{y}) + O\left(\frac{\varepsilon^{n-1}}{(\min\{|\mathbf{x}|, |\mathbf{y}|\})^{n-2}}\right) \end{aligned} \tag{3}$$

*uniformly with respect to  $\mathbf{x}$  and  $\mathbf{y}$  in  $\Omega_\varepsilon$ . (Note that the remainder term in (3) is  $O(\varepsilon)$  on  $\partial\omega_\varepsilon$  and  $O(\varepsilon^{n-1})$  on  $\partial\Omega$ .)*

The next theorem contains a result of the same nature for  $n = 2$ . As before,  $G$  and  $\mathcal{G}$  are Green's functions for  $\Omega$  and  $\mathbf{R}^2 \setminus \bar{\omega}$ , respectively, whereas  $H$  is the regular part of  $G$ .

**Theorem 3.** *Let*

$$\zeta(\eta) = \lim_{|\xi| \rightarrow \infty} \mathcal{G}(\xi, \eta) \quad \text{and} \quad \zeta_\infty = \lim_{|\eta| \rightarrow \infty} \{\zeta(\eta) - (2\pi)^{-1} \log |\eta|\}.$$

*Then the asymptotic representation, uniform with respect to  $\mathbf{x}, \mathbf{y} \in \Omega_\varepsilon$ , holds*

$$\begin{aligned} G_\varepsilon(\mathbf{x}, \mathbf{y}) &= G(\mathbf{x}, \mathbf{y}) + \mathcal{G}(\varepsilon^{-1}\mathbf{x}, \varepsilon^{-1}\mathbf{y}) + (2\pi)^{-1} \log(\varepsilon^{-1}|\mathbf{x} - \mathbf{y}|) \\ &\quad + \frac{((2\pi)^{-1} \log \varepsilon + \zeta(\frac{\mathbf{x}}{\varepsilon}) - \zeta_\infty + H(\mathbf{x}, 0))((2\pi)^{-1} \log \varepsilon + \zeta(\frac{\mathbf{y}}{\varepsilon}) - \zeta_\infty + H(0, \mathbf{y}))}{(2\pi)^{-1} \log \varepsilon + H(0, 0) - \zeta_\infty} \\ &\quad - \zeta(\varepsilon^{-1}\mathbf{x}) - \zeta(\varepsilon^{-1}\mathbf{y}) + \zeta_\infty + O(\varepsilon). \end{aligned} \tag{4}$$

The next assertion is a direct consequence of (4). It shows that asymptotic representation of  $G_\varepsilon(\mathbf{x}, \mathbf{y})$  is simplified if  $\mathbf{x}$  and  $\mathbf{y}$  are subject to additional constraints. We use the same notations as in Theorem 3.

**Corollary 1.** *The following assertions hold:*

(a) *Let  $\mathbf{x}$  and  $\mathbf{y}$  be points of  $\Omega_\varepsilon \subset \mathbf{R}^n$  such that  $\min\{|\mathbf{x}|, |\mathbf{y}|\} > 2\varepsilon$ . Then for  $n = 2$ ,*

$$G_\varepsilon(\mathbf{x}, \mathbf{y}) - G(\mathbf{x}, \mathbf{y}) - \frac{G(\mathbf{x}, 0)G(0, \mathbf{y})}{(2\pi)^{-1} \log \varepsilon + H(0, 0) - \zeta_\infty} = O\left(\frac{\varepsilon}{\min\{|\mathbf{x}|, |\mathbf{y}|\}}\right),$$

*and for  $n > 2$ ,*

$$G_\varepsilon(\mathbf{x}, \mathbf{y}) - G(\mathbf{x}, \mathbf{y}) + \varepsilon^{n-2} \text{cap } \bar{\omega} G(\mathbf{x}, 0)G(0, \mathbf{y}) = O\left(\frac{\varepsilon^{n-1}}{(|\mathbf{x}||\mathbf{y}|)^{n-1} \min\{|\mathbf{x}|, |\mathbf{y}|\}}\right).$$

(b) If  $\max\{|\mathbf{x}|, |\mathbf{y}|\} < 1/2$ , then for  $n = 2$ ,

$$G_\varepsilon(\mathbf{x}, \mathbf{y}) - \mathcal{G}\left(\frac{\mathbf{x}}{\varepsilon}, \frac{\mathbf{y}}{\varepsilon}\right) - \frac{\zeta(\varepsilon^{-1}\mathbf{x})\zeta(\varepsilon^{-1}\mathbf{y})}{(2\pi)^{-1}\log\varepsilon + H(0, 0) - \zeta_\infty} = O(\max\{|\mathbf{x}|, |\mathbf{y}|\}),$$

and for  $n > 2$ ,

$$G_\varepsilon(\mathbf{x}, \mathbf{y}) - \varepsilon^{2-n}\mathcal{G}\left(\frac{\mathbf{x}}{\varepsilon}, \frac{\mathbf{y}}{\varepsilon}\right) + H(0, 0)(P(\varepsilon^{-1}\mathbf{x}) - 1)(P(\varepsilon^{-1}\mathbf{y}) - 1) = O(\max\{|\mathbf{x}|, |\mathbf{y}|\}).$$

One can say that these formulae are closer, in their spirit, to Hadamard's original formula (1) than Theorems 2 and 3 themselves. Analogous simplified formulae can be deduced easily from all other theorems of the present paper, but they are not included here.

#### 4. Dirichlet–Neumann problems in a planar domain with a small hole

The set  $\Omega_\varepsilon$  is assumed to be the same as in Theorem 3, with the additional constraint that  $\partial\omega$  is smooth. First, let  $G_\varepsilon$  denote the kernel of the inverse operator of the mixed boundary value problem in  $\Omega_\varepsilon$  for the operator  $-\Delta$ , with the Dirichlet data on  $\partial\Omega$  and the Neumann data on  $\partial\omega_\varepsilon$ . The notations  $G, \mathcal{G}$  and  $H$  have the same meaning as in Theorem 3, and  $\mathcal{N}$  is the Neumann function in  $\mathbf{R}^2 \setminus \bar{\omega}$ . Let  $\mathcal{D}$  be a vector function, harmonic in  $\mathbf{R}^2 \setminus \bar{\omega}$ , vanishing at infinity and such that  $\partial\mathcal{D}/\partial\nu = \mathbf{v}$  on  $\partial\omega$ . This vector function appears in the asymptotic representation

$$\mathcal{N}(\boldsymbol{\xi}, \boldsymbol{\eta}) \sim (2\pi)^{-1}\log|\boldsymbol{\xi}|^{-1} + (\mathcal{D}(\boldsymbol{\eta}) - \boldsymbol{\eta}) \cdot \nabla((2\pi)^{-1}\log|\boldsymbol{\xi}|^{-1}), \quad \text{as } |\boldsymbol{\xi}| \rightarrow \infty.$$

**Theorem 4.** *The kernel  $G_\varepsilon(\mathbf{x}, \mathbf{y})$  of the inverse operator of the mixed boundary value problem in  $\Omega_\varepsilon$  for the operator  $-\Delta$ , with the Dirichlet data on  $\partial\Omega$  and the Neumann data on  $\partial\omega_\varepsilon$ , has the following asymptotic representation, which is uniform with respect to  $\mathbf{x}, \mathbf{y} \in \Omega_\varepsilon$ ,*

$$\begin{aligned} G_\varepsilon(\mathbf{x}, \mathbf{y}) &= G(\mathbf{x}, \mathbf{y}) + \mathcal{N}(\varepsilon^{-1}\mathbf{x}, \varepsilon^{-1}\mathbf{y}) + (2\pi)^{-1}\log(\varepsilon^{-1}|\mathbf{x} - \mathbf{y}|) \\ &\quad + \varepsilon\mathcal{D}(\varepsilon^{-1}\mathbf{x}) \cdot \nabla_x H(0, \mathbf{y}) + \varepsilon\mathcal{D}(\varepsilon^{-1}\mathbf{y}) \cdot \nabla_y H(\mathbf{x}, 0) \\ &\quad - \varepsilon^2\mathcal{D}(\varepsilon^{-1}\mathbf{y}) \cdot ((\nabla_x \otimes \nabla_y)H(0, 0))\mathcal{D}(\varepsilon^{-1}\mathbf{x}) + O(\varepsilon^2). \end{aligned} \tag{5}$$

Next, consider the mixed boundary value problem in  $\Omega_\varepsilon$  for the Laplace operator, with the Neumann data on the smooth boundary  $\partial\Omega$  and the Dirichlet data on  $\partial\omega_\varepsilon$ , where  $\omega_\varepsilon = \{\mathbf{x}: \varepsilon^{-1}\mathbf{x} \in \omega \subset \mathbf{R}^2\}$  with  $\omega$  being an arbitrary bounded domain. Let  $N(\mathbf{x}, \mathbf{y})$  be the Neumann function in  $\Omega$ , i.e.

$$\begin{aligned} \Delta N(\mathbf{x}, \mathbf{y}) + \delta(\mathbf{x} - \mathbf{y}) &= 0, \quad \mathbf{x}, \mathbf{y} \in \Omega, \\ \frac{\partial}{\partial\nu_x}(N(\mathbf{x}, \mathbf{y}) + (2\pi)^{-1}\log|\mathbf{x}|) &= 0, \quad \mathbf{x} \in \partial\Omega, \mathbf{y} \in \Omega, \end{aligned}$$

and

$$\int_{\partial\Omega} N(\mathbf{x}, \mathbf{y}) \frac{\partial}{\partial\nu_x} \log|\mathbf{x}| \, ds_x = 0,$$

with the last condition implying the symmetry of  $N(\mathbf{x}, \mathbf{y})$ . The regular part of the Neumann function is defined by

$$R(\mathbf{x}, \mathbf{y}) = (2\pi)^{-1}\log|\mathbf{x} - \mathbf{y}|^{-1} - N(\mathbf{x}, \mathbf{y}).$$

Note that

$$R(0, \mathbf{y}) = -(2\pi)^{-2} \int_{\partial\Omega} \log|\mathbf{x}| \frac{\partial}{\partial\nu} \log|\mathbf{x}| \, ds_x.$$

Let  $\mathcal{D}$  be a vector function, harmonic in  $\mathbf{R}^2 \setminus \bar{\omega}$ , bounded at infinity and subject to the Dirichlet condition  $\mathcal{D}(\boldsymbol{\xi}) = \boldsymbol{\xi}$ ,  $\boldsymbol{\xi} \in \partial\omega$ . This vector function appears in the asymptotic representation of Green's function in  $\mathbf{R}^2 \setminus \bar{\omega}$

$$\mathcal{G}(\boldsymbol{\xi}, \boldsymbol{\eta}) \sim \mathcal{G}(\infty, \boldsymbol{\eta}) + (\mathcal{D}(\boldsymbol{\eta}) - \boldsymbol{\eta}) \cdot \nabla((2\pi)^{-1}\log|\boldsymbol{\xi}|^{-1}), \quad \text{as } |\boldsymbol{\xi}| \rightarrow \infty.$$

**Theorem 5.** *The function  $G_\varepsilon(\mathbf{x}, \mathbf{y})$  has the asymptotic behaviour, uniform with respect to  $\mathbf{x}, \mathbf{y} \in \Omega_\varepsilon$ :*

$$\begin{aligned}
 G_\varepsilon(\mathbf{x}, \mathbf{y}) &= \mathcal{G}(\varepsilon^{-1}\mathbf{x}, \varepsilon^{-1}\mathbf{y}) + N(\mathbf{x}, \mathbf{y}) - (2\pi)^{-1} \log |\mathbf{x} - \mathbf{y}|^{-1} + R(0, 0) \\
 &\quad + \varepsilon \mathbf{D}(\varepsilon^{-1}\mathbf{y}) \cdot \nabla_{\mathbf{y}} R(\mathbf{x}, 0) + \varepsilon \mathbf{D}(\varepsilon^{-1}\mathbf{x}) \cdot \nabla_{\mathbf{x}} R(0, \mathbf{y}) \\
 &\quad - \varepsilon^2 \mathbf{D}(\varepsilon^{-1}\mathbf{y}) \cdot ((\nabla_{\mathbf{x}} \otimes \nabla_{\mathbf{y}}) R)(0, 0) \mathbf{D}(\varepsilon^{-1}\mathbf{x}) + O(\varepsilon^2).
 \end{aligned} \tag{6}$$

**5. The Dirichlet–Neumann problem in a thin rod**

Let  $C$  be the infinite cylinder  $\{(\mathbf{x}', x_n): \mathbf{x}' \in \omega, x_n \in \mathbf{R}\}$ , where  $\omega$  is a bounded domain in  $\mathbf{R}^{n-1}$  with smooth boundary. Also let  $C^\pm$  denote Lipschitz subdomains of  $C$  separated from  $\pm\infty$  by surfaces  $\gamma^\pm$ , respectively. We introduce a positive number  $a$ , the vector  $\mathbf{a} = (\mathbf{O}', a)$ , where  $\mathbf{O}'$  is the origin of  $\mathbf{R}^{n-1}$ , and the small parameter  $\varepsilon > 0$ .

By  $G_\varepsilon$  we denote the fundamental solution for  $-\Delta$  in the thin domain  $C_\varepsilon = \{\mathbf{x}: \varepsilon^{-1}(\mathbf{x} - \mathbf{a}) \in C^+, \varepsilon^{-1}(\mathbf{x} + \mathbf{a}) \in C^-\}$  subject to zero Neumann condition on the cylindrical part of  $C$  and zero Dirichlet condition on the remaining part of  $\partial C_\varepsilon$ . Similarly,  $G^+$  and  $G^-$  stand for the fundamental solutions for  $-\Delta$  in the domains  $C^\pm$ , and satisfy zero Dirichlet condition on  $\gamma^\pm$ , zero Neumann condition on  $\partial C^\pm \setminus \gamma^\pm$ , and are bounded as  $x_n \rightarrow \mp\infty$ . Let  $\zeta^\pm$  be harmonic functions in  $C^\pm$  subject to the same boundary conditions as  $G^\pm$  and asymptotically equivalent to  $\mp|\omega|^{-1}x_n + \zeta_\infty^\pm$  as  $x_n \rightarrow \mp\infty$ , where  $\zeta_\infty^\pm$  are constants. We note that  $\zeta^\pm(\mathbf{y}) = \lim_{\mathbf{x} \rightarrow \infty} G^\pm(\mathbf{x}, \mathbf{y})$ . Finally, by  $G_\infty(\mathbf{x}, \mathbf{y})$  we denote Green's function of the Neumann problem in  $C$  such that

$$G_\infty(\mathbf{x}, \mathbf{y}) = -(2|\omega|)^{-1} |x_n - y_n| + O(\exp(-\alpha|x_n - y_n|)) \quad \text{as } |x_n| \rightarrow \infty,$$

where  $\alpha$  is a positive constant, and  $|\omega|$  is the  $(n - 1)$ -dimensional measure of  $\omega$ .

**Theorem 6.** *The following asymptotic formula for  $G_\varepsilon(\mathbf{x}, \mathbf{y})$ , uniform with respect to  $\mathbf{x}, \mathbf{y} \in \Omega_\varepsilon$ , holds*

$$\begin{aligned}
 G_\varepsilon(\mathbf{x}, \mathbf{y}) &= \varepsilon^{2-n} \left\{ G^+(\varepsilon^{-1}(\mathbf{x} - \mathbf{a}), \varepsilon^{-1}(\mathbf{y} - \mathbf{a})) + G^-(\varepsilon^{-1}(\mathbf{x} + \mathbf{a}), \varepsilon^{-1}(\mathbf{y} + \mathbf{a})) - G^\infty(\varepsilon^{-1}\mathbf{x}, \varepsilon^{-1}\mathbf{y}) \right. \\
 &\quad - \varepsilon \frac{(\frac{x_n}{\varepsilon|\omega|} - \frac{1}{2}(\zeta_\infty^- - \zeta_\infty^+) + \zeta^+(\frac{\mathbf{x}-\mathbf{a}}{\varepsilon}) - \zeta^-(\frac{\mathbf{x}+\mathbf{a}}{\varepsilon}))(\frac{y_n}{\varepsilon|\omega|} - \frac{1}{2}(\zeta_\infty^- - \zeta_\infty^+) + \zeta^+(\frac{\mathbf{y}-\mathbf{a}}{\varepsilon}) - \zeta^-(\frac{\mathbf{y}+\mathbf{a}}{\varepsilon}))}{2|\omega|^{-1}a + \varepsilon(\zeta_\infty^+ + \zeta_\infty^-)} \\
 &\quad \left. + \frac{1}{4} \left( (\varepsilon|\omega|)^{-1} 2a + \zeta_\infty^- + \zeta_\infty^+ - 2 \sum_{\pm} (\zeta^\pm(\varepsilon^{-1}(\mathbf{x} \mp \mathbf{a})) + \zeta^\pm(\varepsilon^{-1}(\mathbf{y} \mp \mathbf{a}))) \right) \right\} + O(\exp(-\beta/\varepsilon)),
 \end{aligned}$$

where  $\beta$  is a positive constant independent of  $\varepsilon$ .

**6. The Dirichlet problem in a truncated cone**

Let  $K$  be an infinite cone  $\{\mathbf{x}: |\mathbf{x}| > 0, |\mathbf{x}|^{-1}\mathbf{x} \in \omega\}$ , where  $\omega$  is a subdomain of the unit sphere  $S^{n-1}$  such that  $S^{n-1} \setminus \omega$  has a positive  $(n - 1)$ -dimensional harmonic capacity. Also let  $K_0$  and  $K_\infty$  denote subdomains of  $K$  separated from the vertex of  $K$  and from  $\infty$  by surfaces  $\gamma$  and  $\Gamma$ , respectively.

By  $G_\varepsilon$  and  $G_{\text{cone}}$  we denote Green's function of the Dirichlet problem for  $-\Delta$  in the domains  $K_\varepsilon = \{\mathbf{x} \in K_0: \varepsilon^{-1}\mathbf{x} \in K_\infty\}$  and  $K$ , respectively. Similarly,  $G_0$  and  $G_\infty$  stand for Green's functions of the Dirichlet problem for  $-\Delta$  in the domains  $K_0$  and  $K_\infty$  satisfying zero Dirichlet boundary conditions on  $\partial K_0 \setminus \{O\}$  and  $\partial K_\infty$ , with the asymptotic representations

$$Z_0(\mathbf{x}) = |\mathbf{x}|^{2-n-\lambda} \Psi(|\mathbf{x}|^{-1}\mathbf{x})(1 + o(1)) \quad \text{as } |\mathbf{x}| \rightarrow 0,$$

and

$$Z_\infty(\mathbf{x}) = |\mathbf{x}|^\lambda \Psi(|\mathbf{x}|^{-1}\mathbf{x})(1 + o(1)) \quad \text{as } |\mathbf{x}| \rightarrow \infty,$$

where  $\lambda$  is a positive number such that  $\lambda(\lambda + n - 2)$  is the first eigenvalue of the Dirichlet spectral problem in  $\omega$  for the Beltrami operator on  $S^{n-1}$ , and  $\Psi$  is the corresponding eigenfunction. Similarly, we introduce  $\lambda_2 > 0$  such that  $\lambda_2(\lambda_2 + n - 2)$  is the second eigenvalue of the same spectral problem.

**Theorem 7.** Let  $\xi = \varepsilon^{-1}\mathbf{x}$  and  $\eta = \varepsilon^{-1}\mathbf{y}$ . Green's function  $G_\varepsilon(\mathbf{x}, \mathbf{y})$  has the asymptotic behaviour

$$G_\varepsilon(\mathbf{x}, \mathbf{y}) = G_0(\mathbf{x}, \mathbf{y}) + \varepsilon^{2-n}G_\infty(\xi, \eta) - G_{\text{cone}}(\mathbf{x}, \mathbf{y}) + \frac{\varepsilon^\lambda}{2\lambda + n - 2} \left\{ \left( |\xi|^\lambda \Psi\left(\frac{\xi}{|\xi|}\right) - Z_\infty(\xi) \right) \left( |\mathbf{y}|^{2-n-\lambda} \Psi\left(\frac{\mathbf{y}}{|\mathbf{y}|}\right) - Z_0(\mathbf{y}) \right) + \left( |\eta|^\lambda \Psi\left(\frac{\eta}{|\eta|}\right) - Z_\infty(\eta) \right) \left( |\mathbf{x}|^{2-n-\lambda} \Psi\left(\frac{\mathbf{x}}{|\mathbf{x}|}\right) - Z_0(\mathbf{x}) \right) \right\} + O(\varepsilon^{\min(2\lambda, \lambda_2)}). \tag{7}$$

This representation is uniform with respect to  $\mathbf{x}$  and  $\mathbf{y}$  in the truncated cone  $K_\varepsilon$ .

**7. Dirichlet problem in a domain containing several inclusions**

It is straightforward to generalise the results of sections dealing with a domain containing a small inclusion/void to the case of a body containing a finite number of inclusions. As an example, we formulate a generalisation of Theorem 2. Let  $\Omega \subset \mathbf{R}^n$ ,  $n > 2$ , be a bounded domain, and let  $\mathbf{O}^{(1)}, \mathbf{O}^{(2)}, \dots, \mathbf{O}^{(N)}$  be interior points in  $\Omega$ . Small sets  $\omega_\varepsilon^{(j)}$ ,  $j = 1, \dots, N$ , are defined by  $\omega_\varepsilon^{(j)} = \{\mathbf{x}: \varepsilon^{-1}(\mathbf{x} - \mathbf{O}^{(j)}) \in \omega^{(j)} \subset \mathbf{R}^n\}$ , where  $\omega^{(j)}$ ,  $j = 1, \dots, N$ , are bounded domains in  $\mathbf{R}^n$ , and they contain the origin  $\mathbf{O}$ . Similar to Section 2, it is assumed that the minimum distance between  $\mathbf{O}^{(j)}$ ,  $j = 1, \dots, N$ , and the points of  $\partial\Omega$  is equal to 1. Also, it is supposed that the distance between  $\mathbf{O}$  and the points of  $\partial\omega^{(j)}$ ,  $j = 1, \dots, N$ , does not exceed 1. By  $G_\varepsilon$  we denote Green's function for the Laplacian in the domain  $\Omega_\varepsilon = \Omega \setminus \bigcup_j \overline{\omega_\varepsilon^{(j)}}$ .

**Theorem 8.** Let  $G$  and  $\mathcal{G}^{(j)}$  stand for Green's functions of the Dirichlet problems in  $\Omega$  and  $\mathbf{R}^n \setminus \overline{\omega^{(j)}}$ , respectively. Also let  $H$  be the regular part of  $G$ , and  $P^{(j)}$  denote the harmonic capacity potentials of the sets  $\overline{\omega^{(j)}}$ . Then

$$G_\varepsilon(\mathbf{x}, \mathbf{y}) = G(\mathbf{x}, \mathbf{y}) + \varepsilon^{2-n} \sum_{j=1}^N \mathcal{G}^{(j)}\left(\frac{\mathbf{x} - \mathbf{O}^{(j)}}{\varepsilon}, \frac{\mathbf{y} - \mathbf{O}^{(j)}}{\varepsilon}\right) - \frac{N}{(n-2)|S^{n-1}||\mathbf{x} - \mathbf{y}|^{n-2}} + \sum_{j=1}^N \left\{ H(\mathbf{O}^{(j)}, \mathbf{y}) P^{(j)}\left(\frac{\mathbf{x} - \mathbf{O}^{(j)}}{\varepsilon}\right) + H(\mathbf{x}, \mathbf{O}^{(j)}) P^{(j)}\left(\frac{\mathbf{y} - \mathbf{O}^{(j)}}{\varepsilon}\right) - H(\mathbf{O}^{(j)}, \mathbf{O}^{(j)}) P^{(j)}\left(\frac{\mathbf{x} - \mathbf{O}^{(j)}}{\varepsilon}\right) P^{(j)}\left(\frac{\mathbf{y} - \mathbf{O}^{(j)}}{\varepsilon}\right) - \varepsilon^{n-2} \text{cap} \overline{\omega^{(j)}} H(\mathbf{x}, \mathbf{O}^{(j)}) H(\mathbf{O}^{(j)}, \mathbf{y}) \right\} + \sum_{j=1}^N \sum_{1 \leq i \leq N, i \neq j} G(\mathbf{O}^{(j)}, \mathbf{O}^{(i)}) P^{(j)}\left(\frac{\mathbf{x} - \mathbf{O}^{(j)}}{\varepsilon}\right) P^{(i)}\left(\frac{\mathbf{y} - \mathbf{O}^{(i)}}{\varepsilon}\right) + O\left(\sum_{j=1}^N \frac{\varepsilon^{n-1}}{(\min\{|\mathbf{x} - \mathbf{O}^{(j)}|, |\mathbf{y} - \mathbf{O}^{(j)}|\})^{n-2}}\right) \tag{8}$$

uniformly with respect to  $\mathbf{x}$  and  $\mathbf{y}$  in  $\Omega_\varepsilon$ . (Note that the remainder term in (8) is  $O(\varepsilon)$  on  $\partial(\bigcup_j \omega_\varepsilon^{(j)})$  and  $O(\varepsilon^{n-1})$  on  $\partial\Omega$ .)

**References**

[1] D. Fujiwara, S. Ozawa, The Hadamard variational formula for the Green functions of some normal elliptic boundary value problems, Proc. Japan Acad. Ser. A Math. Sci. 54 (8) (1978) 215–220.  
 [2] J. Hadamard, Sur le problème d'analyse relatif à l'équilibre des plaques élastiques encastrées, Mémoire couronné en 1907 par l'Académie des Sciences 33 (4) 515–629.  
 [3] P. Levy, Leçons d'analyse fonctionnelle, Gauthier-Villars, Paris, 1922.  
 [4] V. Maz'ya, S. Nazarov, B. Plamenevskij, Asymptotic Theory of Elliptic Boundary Value Problems in Singularly Perturbed Domains, vols. 1–2, Birkhäuser, 2000.  
 [5] S. Ozawa, Perturbation of domains and Green kernels of heat equations, Proc. Japan Acad. Ser. A Math. Sci. 54 (1978) 322–325; Part II, 55 (1979) 172–175; Part III, 55 (1979) 227–230.  
 [6] S. Ozawa, Spectra of the Laplacian with small Robin conditional boundary, Proc. Japan Acad. Ser. A Math. Sci. 72 (3) (1996) 53–54.  
 [7] S. Ozawa, S. Roppongi, Heat kernel and singular variation of domains, Osaka J. Math. 32 (4) (1995) 941–957.