



ELSEVIER

Contents lists available at ScienceDirect

C. R. Acad. Sci. Paris, Ser. I

www.sciencedirect.com



Numerical analysis

Error estimates for stabilized finite element methods applied to ill-posed problems



Estimations d'erreurs pour des méthodes d'éléments finis stabilisées appliquées à des problèmes mal posés

Erik Burman

Department of Mathematics, University College London, Gower Street, London, WC1E 6BT, United Kingdom

ARTICLE INFO

Article history:

Received 10 March 2014

Accepted after revision 17 June 2014

Available online 21 July 2014

Presented by the Editorial Board

ABSTRACT

We propose an analysis for the stabilized finite element methods proposed in Burman (2013) [2] valid in the case of ill-posed problems for which only weak continuous dependence can be assumed. A priori and a posteriori error estimates are obtained without assuming coercivity or inf-sup stability of the continuous problem.

© 2014 Académie des sciences. Published by Elsevier Masson SAS. All rights reserved.

R É S U M É

Dans cette note, nous proposons une nouvelle analyse pour les méthodes d'éléments finis stabilisées introduites dans Burman (2013) [2], appliquées à des problèmes mal posés avec des propriétés de dépendance continue faibles. Nous obtenons des estimations a priori et a posteriori sans supposer ni coercitivité ni stabilité inf-sup de la forme bilinéaire du problème continu.

© 2014 Académie des sciences. Published by Elsevier Masson SAS. All rights reserved.

1. Introduction

We are interested in the numerical approximation of ill-posed problems. The abstract theory will be illustrated by the following linear elliptic Cauchy problem. Let Ω be a convex polygonal (polyhedral) domain in \mathbb{R}^d , $d = 2, 3$, and consider the equation

$$\begin{cases} -\Delta u = f, & \text{in } \Omega \\ u = 0 \text{ and } \nabla u \cdot n = \psi & \text{on } \Gamma \end{cases} \quad (1)$$

where $\Gamma \subset \partial\Omega$ denotes a simply connected part of the boundary and $f \in L^2(\Omega)$, $\psi \in H^{\frac{1}{2}}(\Gamma)$. Introducing the spaces $V := \{v \in H^1(\Omega) : v|_{\Gamma} = 0\}$ and $W := \{v \in H^1(\Omega) : v|_{\Gamma'} = 0\}$, where $\Gamma' := \partial\Omega \setminus \Gamma$ and the forms $a(u, w) := \int_{\Omega} \nabla u \cdot \nabla w \, dx$, and $l(w) := \int_{\Omega} f w \, dx + \int_{\Gamma} \psi w \, ds$ Eq. (1) may be cast in the abstract weak formulation, find $u \in V$ such that

E-mail address: E.Burman@ucl.co.uk.

$$a(u, w) = l(w) \quad \forall w \in W. \tag{2}$$

It is well known that the Cauchy problem (1) is not well-posed in the sense of Hadamard. If $l(w)$ is such that a sufficiently smooth, exact solution exists, conditional continuous dependence estimates can nevertheless be obtained [1].

The objective of the present paper is to study numerical methods for ill-posed problems of the form (2), where $a : V \times W \mapsto \mathbb{R}$ and $l : W \mapsto \mathbb{R}$ are a bilinear and a linear form. Assume that the linear form $l(w)$ is such that the problem (2) admits a unique solution $u \in V$. Define the following dual norm on l , $\|l\|_{W'} := \sup_{\|w\|_W=1} |l(w)|$. Observe that we do not assume that (2) admits a unique solution for all $l(w)$ such that $\|l\|_{W'} < \infty$. The stability property we assume to be satisfied by (2) is the following continuous dependence.

Assumption: continuous dependence on data. Consider the functional $j : V \mapsto \mathbb{R}$. Let $\mathcal{E} : \mathbb{R}^+ \mapsto \mathbb{R}^+$ be a continuous, monotone increasing function with $\lim_{x \rightarrow 0^+} \mathcal{E}(x) = 0$. Let $\epsilon > 0$.

Assume that there holds $\|l\|_{W'} \leq \epsilon$ in (2) then, for ϵ sufficiently small, $|j(u)| \leq \mathcal{E}(\epsilon)$.

(3)

For the example of the Cauchy problem (1), it is known [1, Theorems 1.7 and 1.9] that if (1) admits a unique solution $u \in H^1(\Omega)$, a continuous dependence of the form (3), with $0 < \epsilon < 1$, holds for

$$j(u) := \|u\|_{L^2(\omega)}, \omega \subset \Omega : \text{dist}(\omega, \partial\Omega) =: d_{\omega, \partial\Omega} > 0 \quad \text{with } \mathcal{E}(x) := C_{u\zeta} x^\zeta, C_{u\zeta} > 0, \zeta := \zeta(d_{\omega, \partial\Omega}) \in (0, 1) \tag{4}$$

and for

$$j(u) := \|u\|_{L^2(\Omega)} \quad \text{with } \mathcal{E}(x) := C_u (|\log(x)| + C)^{-\zeta} \quad \text{with } C_u, C > 0, \zeta \in (0, 1). \tag{5}$$

Note that to derive these results, $l(\cdot)$ is first associated with its Riesz representant in W (cf. [1, Eq. (1.31)] and discussion). The constant $C_{u\zeta}$ in (4) grows monotonically in $\|u\|_{L^2(\Omega)}$ and C_u in (5) grows monotonically in $\|u\|_{H^1(\Omega)}$.

2. Finite element discretization

Let \mathcal{K}_h be a shape regular, conforming, subdivision of Ω into non-overlapping triangles κ . The family of meshes $\{\mathcal{K}_h\}_h$ is indexed by the mesh parameter $h := \max(\text{diam}(\kappa))$. Let \mathcal{F}_I be the set of interior faces in \mathcal{K}_h and $\mathcal{F}_\Gamma, \mathcal{F}_{\Gamma'}$ the set of element faces of \mathcal{K}_h whose interior intersects Γ and Γ' respectively. We assume that the mesh matches the boundary of Γ so that $\mathcal{F}_\Gamma \cap \mathcal{F}_{\Gamma'} = \emptyset$. Let X_h^1 denote the standard finite element space of continuous, affine functions. Define $V_h := V \cap X_h^1$ and $W_h := W \cap X_h^1$. We may then write the finite element method: find $(u_h, z_h) \in V_h \times W_h$ such that,

$$\left. \begin{aligned} a(u_h, w_h) - s_W(z_h, w_h) &= l(w_h) \\ a(v_h, z_h) + s_V(u_h, v_h) &= s_V(u, v_h) \end{aligned} \right\} \quad \text{for all } (v_h, w_h) \in V_h \times W_h. \tag{6}$$

A possible choice of stabilization operators for the problem (1) are

$$s_V(u_h, v_h) := \sum_{F \in \mathcal{F}_I \cup \mathcal{F}_\Gamma} \int_F h_F [\partial_n u_h][\partial_n v_h] \, ds, \quad \text{with } h_F := \text{diam}(F) \tag{7}$$

and

$$s_W(z_h, w_h) := a(z_h, w_h) \quad \text{or} \quad s_W(z_h, w_h) := \sum_{F \in \mathcal{F}_I \cup \mathcal{F}_{\Gamma'}} \int_F h_F [\partial_n z_h][\partial_n w_h] \, ds \tag{8}$$

where $[\partial_n u_h]$ denotes the jump of $\nabla u_h \cdot n_F$ for $F \in \mathcal{F}_I$ and when $F \in \mathcal{F}_\Gamma$ or $F \in \mathcal{F}_{\Gamma'}$ define $[\partial_n u_h]|_F := \nabla u_h \cdot n_{\partial\Omega}$. Unique existence of (u_h, z_h) solution to (6)–(8) follows using the arguments of [2, Proposition 3.3]. By inspection we have that the system (6) is consistent with (2) for $z_h = 0$. Taking the difference of (6) and the relation (2), with $w = w_h$, we obtain the Galerkin orthogonality,

$$a(u_h - u, w_h) - s_W(z_h, w_h) + a(v_h, z_h) + s_V(u_h - u, v_h) = 0 \quad \text{for all } (v_h, w_h) \in V_h \times W_h. \tag{9}$$

3. Hypotheses on forms and interpolants

Consider the general, positive semi-definite, symmetric stabilization operators, $s_V : V_h \times V_h \mapsto \mathbb{R}$, $s_W : W_h \times W_h \mapsto \mathbb{R}$. We assume that $s_V(u, v_h)$, with u the solution of (2), is explicitly known; it may depend on data from $l(w)$ or measurements of u . Assume that both s_V and s_W define semi-norms on $H^s(\Omega) + V_h$ and $H^s(\Omega) + W_h$ respectively, for some $s \geq 1$,

$$|v + v_h|_{s_Z} := s_Z(v + v_h, v + v_h)^{\frac{1}{2}}, \quad \forall v \in H^s(\Omega), v_h \in Z_h, \quad \text{with } Z = V, W. \tag{10}$$

Then assume that there exist interpolation operators $i_V : V \mapsto V_h$ and $i_W : W \mapsto W_h$ and norms $\|\cdot\|_{*,V}$ and $\|\cdot\|_{*,W}$ defined on V and W respectively, such that the form $a(u, v)$ satisfies the continuities

$$a(v - i_V v, w_h) \leq \|v - i_V v\|_{*,V} |w_h|_{s_W}, \quad \forall v \in V, w_h \in W_h \tag{11}$$

and for u solution of (2),

$$a(u - u_h, w - i_W w) \leq \delta_l(h) \|w\|_W + \|w - i_W w\|_{*,W} |u - u_h|_{s_V}, \quad \forall w \in W. \tag{12}$$

In practice, $\delta_l(h)$ only depends on the properties of the interpolant i_W and the data of the problem (and satisfies $\lim_{h \rightarrow 0} \delta_l(h) = 0$ provided the data are unperturbed). We also assume that the interpolants have the following approximation and stability properties. For sufficiently smooth $v \in V$ there holds, for $t > 0$:

$$|v - i_V v|_{s_V} + \|v - i_V v\|_{*,V} \leq C_V(v) h^t. \tag{13}$$

The factor $C_V(v) > 0$ will typically depend on some Sobolev norm of v . For i_W we assume that for some $C_W > 0$ there holds

$$\|w - i_W w\|_{*,W} + |i_W w|_{s_W} \leq C_W \|w\|_W, \quad \forall w \in W. \tag{14}$$

3.1. Satisfaction of hypothesis for the formulation (6)–(8)

Let i_V and i_W be defined by Scott–Zhang interpolation operators preserving the Dirichlet boundary conditions. The consistency of $s_V(\cdot, \cdot)$ holds for solutions $u \in H^2(\Omega)$. Consider first the form of $s_W(\cdot, \cdot)$ in the left definition of (8). Define $\|v\|_{*,V} := \|\nabla v\|_{L^2(\Omega)}$ and $\|w\|_{*,W} := \|h^{-1} w\|_{L^2(\Omega)} + (\sum_{F \in \mathcal{F}_I \cup \mathcal{F}_T} h^{-1} \|w\|_{L^2(F)}^2)^{1/2}$. Using local trace inequalities and the stability and approximation properties of the Scott–Zhang interpolant, we deduce that the inequalities (13)–(14) hold with $t = 1$ and $C_V(v) := C \|v\|_{H^2(\Omega)}$. The inequality (11) follows by the Cauchy–Schwarz inequality. To prove (12), with $\delta(h) = C_W h \|f\|_{L^2(\Omega)}$, integrate by parts in $a(u - u_h, w - i_W w)$, and use Eq. (1), to obtain:

$$a(u - u_h, w - i_W w) = (f, w - i_W w)_{L^2(\Omega)} + \sum_{F \in \mathcal{F}_I \cup \mathcal{F}_T} ([\partial_n(u - u_h)], w - i_W w)_{L^2(F)}.$$

The bound (12) then follows by the Cauchy–Schwarz inequality, the definitions of $s_V(\cdot, \cdot)$ and $\|\cdot\|_{*,W}$ and the approximation (14). For the variant where $s_W(w_h, z_h) := \sum_{F \in \mathcal{F}_I \cup \mathcal{F}_{T'}} \int_F h [\partial_n z_h] [\partial_n w_h] ds$ let $\|w\|_{*,V} := \|h^{-1} w\|_{L^2(\Omega)} + (\sum_{F \in \mathcal{F}_I \cup \mathcal{F}_{T'}} h^{-1} \|w\|_{L^2(F)}^2)^{1/2}$ and prove inequality (11) similarly as (12) above, but integrating by parts the other way. This latter method has enhanced adjoint consistency.

4. Error analysis

We will now prove an error analysis using only the continuous dependence (3). First we prove that assuming smoothness of the exact solution the error converges with the rate h^t in the stabilization semi-norms defined in Eq. (10). Then we show that the computational error satisfies a perturbation equation of the form (2), and that the right-hand side of the perturbation equation can be upper bounded by the stabilization semi-norm. Our error bounds are then a consequence of the assumption (3).

Lemma 4.1. *Let u be the solution of (2) and (u_h, z_h) the solution of the formulation (6) for which (10), (11) and (13) hold. Then*

$$|u - u_h|_{s_V} + |z_h|_{s_W} \leq (1 + \sqrt{2}) C_V(u) h^t.$$

Proof. Let $\xi_h := i_V u - u_h$ and write $|\xi_h|_{s_V}^2 + |z_h|_{s_W}^2 = s_V(\xi_h, \xi_h) + a(\xi_h, z_h) - a(\xi_h, z_h) + s_W(z_h, z_h)$. Using Eq. (9) we then have $|\xi_h|_{s_V}^2 + |z_h|_{s_W}^2 = s_V(i_V u - u, \xi_h) + a(i_V u - u, z_h)$. Applying the Cauchy–Schwarz inequality in the first term of the right-hand side and the continuity (11) in the second, followed by (13), we may deduce:

$$|\xi_h|_{s_V}^2 + |z_h|_{s_W}^2 \leq |i_V u - u|_{s_V} |\xi_h|_{s_V} + \|i_V u - u\|_{*,V} |z_h|_{s_W} \leq C_V(u) h^t (|\xi_h|_{s_V}^2 + |z_h|_{s_W}^2)^{1/2}.$$

The claim follows by the triangle inequality $|u - u_h|_{s_V} \leq |u - i_V u|_{s_V} + |\xi_h|_{s_V}$. \square

Theorem 4.2. *Let u be the solution of (2) and (u_h, z_h) the solution of the formulation (6) for which (10)–(13) hold. Assume that the problem (2) has the stability property (3). Then*

$$|j(u - u_h)| \leq \mathcal{E}(\eta(u_h, z_h)) \tag{15}$$

where the a posteriori quantity $\eta(u_h, z_h)$ is defined by $\eta(u_h, z_h) := \delta_l(h) + C_W (|u - u_h|_{s_V} + |z_h|_{s_W})$. For sufficiently smooth u there holds:

$$\eta(u_h, z_h) \leq \delta_l(h) + (1 + \sqrt{2}) C_W C_V(u) h^t. \tag{16}$$




Fig. 1. Left: convergence of global L^2 -errors (dashed) and stabilization semi-norms (full). Middle: convergence of local L^2 -errors. Right: study of error under variation of the parameter $\gamma_V = \gamma_W$. (P_1 approximation marked with squares, P_2 with circles.)

Proof. Let $e := u - u_h \in V$. By the Galerkin orthogonality, there holds for all $w \in W$:

$$a(e, w) = a(e, w - i_W w) - s_W(z_h, i_W w) = l(w - i_W w) - a(u_h, w - i_W w) - s_W(z_h, i_W w)$$

and we identify $r \in W'$ such that $\forall w \in W$,

$$(r, w)_{W', W} = l(w - i_W w) - a(u_h, w - i_W w) - s_W(z_h, i_W w). \tag{17}$$

We have shown that e satisfies equation (2) with right-hand side $(r, w)_{W', W}$. Now apply the continuity (12), Cauchy-Schwarz inequality and the stability (14) in the right-hand side of (17), leading to:

$$|(r, w)_{W', W}| = |a(e, w - i_W w) - s_W(z_h, i_W w)| \leq (\delta_l(h) + C_W |u - u_h|_{S_V} + C_W |z_h|_{S_W}) \|w\|_W.$$

We conclude that $\|r\|_{W'} \leq \delta_l(h) + C_W (|u - u_h|_{S_V} + |z_h|_{S_W})$ and the claim (15) follows by assumption (3). The upper bound of (16) is a consequence of Lemma 4.1. \square

Corollary 4.3. Let $u \in H^2(\Omega)$ be the solution of (1) and u_h, z_h the solution of (6)–(8). Then the conclusions of Lemma 4.1 and Theorem 4.2 hold for $u - u_h, z_h$ with $t = 1$ and $j(\cdot), \mathcal{E}(\cdot)$ given by (4) or (5). Moreover C_{u_ζ} and C_u of (4) and (5) are independent of h .

Proof. In Section 3.1 above we showed that the formulation (6)–(8) satisfies (10)–(13), and we conclude that Lemma 4.1 and Theorem 4.2 hold. For C_{u_ζ} and C_u of (4) and (5) to be bounded uniformly in h , $\|u - u_h\|_{H^1(\Omega)}$ must be bounded by some constant independent of h . To this end, one may prove a discrete Poincaré inequality $\|\nabla u_h\|_{L^2(\Omega)} \leq C_P h^{-1} |u_h|_{S_V}$. Using this result together with Lemma 4.1, we deduce that $\|\nabla u_h\|_{L^2(\Omega)} \leq C \|u\|_{H^2(\Omega)}$, which proves the claim. \square

5. Numerical example

To illustrate the theory we recall a numerical example from [2]. We solve the Cauchy problem (1) on the unit square $\Omega \in (0, 1) \times (0, 1)$ with exact solution $u(x, y) = 30x(1 - x)y(1 - y)$, $\psi = \nabla u \cdot n_{\partial\Omega}$ and $\Gamma := \{x \in (0, 1), y = 0\} \cup \{x = 1, y \in (0, 1)\}$. We compute piecewise affine approximations on a sequence of unstructured meshes using the method (6) and the stabilizations (7) and (8)₂ ($\gamma_V = \gamma_W = 0.01$). We also make a similar series of computations using piecewise quadratic elements and an added penalty term on the jump of the elementwise Laplacian following [2] ($\gamma_V = \gamma_W = 0.001$). The results are reported in Fig. 1. The convergence of the global L^2 -error and the stabilization semi-norm is given in the left plot, compared with theoretically motivated logarithmic bounds. The local $L^2(\omega)$ errors in the subdomain $\omega = (0.5, 1) \times (0, 0.5)$ are presented in the middle plot and we observe that they have $O(h^k)$ convergence where k denotes the polynomial order. Finally we report a study of the error on a fixed mesh with 64×64 elements under variation of the penalty parameter in the right plot.

6. Conclusion and further perspectives

Herein we have proposed a framework for the analysis of the stabilized methods introduced in [2] when applied to ill-posed problems. The upshot is that error estimates can be obtained using only continuous dependence properties, without relying on a well-posedness theory of the continuous problem. For numerical examples, we refer the reader to [2]. Important extensions of the results presented herein are the inclusion of perturbed data and the exploration of the consequences

of adjoint consistency. The latter may allow for improved estimates, when the error is measured by linear functionals that are in the range of the adjoint problem.

References

- [1] G. Alessandrini, L. Rondi, E. Rosset, S. Vessella, The stability for the Cauchy problem for elliptic equations, *Inverse Probl.* 25 (12) (2009) 123004, 47 p.
- [2] E. Burman, Stabilized finite element methods for nonsymmetric, noncoercive, and ill-posed problems. Part I: Elliptic equations, *SIAM J. Sci. Comput.* 35 (6) (2013) A2752–A2780.