



ELSEVIER

Contents lists available at ScienceDirect

C. R. Acad. Sci. Paris, Ser. I

www.sciencedirect.com



Group theory/Topology

Unbounded asymmetry of stretch factors



Asymétrie non bornée des facteurs d'étirement

Spencer Dowdall, Ilya Kapovich, Christopher J. Leininger

Department of Mathematics, University of Illinois at Urbana-Champaign, 1409 West Green Street, Urbana, IL 61801, United States

ARTICLE INFO

Article history:

Received 18 May 2014

Accepted 11 September 2014

Available online 1 October 2014

Presented by the Editorial Board

ABSTRACT

A result of Handel–Mosher guarantees that the ratio of logarithms of stretch factors of any fully irreducible automorphism of the free group F_N and its inverse is bounded by a constant C_N . In this short note, we show that this constant C_N cannot be chosen independent of the rank N .

© 2014 Académie des sciences. Published by Elsevier Masson SAS. All rights reserved.

R É S U M É

Un énoncé de Handel–Mosher certifie que le rapport des logarithmes des facteurs d'étirement d'un automorphisme irréductible du groupe libre F_N et de son inverse est borné par une constante C_N . Nous montrons dans la présente courte note que cette constante C_N ne peut pas être choisie indépendante de N .

© 2014 Académie des sciences. Published by Elsevier Masson SAS. All rights reserved.

Let F_N be the free group of rank $N \geq 2$. An outer automorphism $\varphi \in \text{Out}(F_N)$ is said to be *fully irreducible* if no power of φ preserves the conjugacy class of any proper free factor of F_N . In this case φ has a well defined *stretch factor* $\lambda(\varphi)$, which, for any non- φ -periodic conjugacy class α in F_N and a free basis X of F_N , is given by

$$\lambda(\varphi) = \lim_{n \rightarrow \infty} \sqrt[n]{\|\varphi^n(\alpha)\|_X},$$

where $\|\cdot\|_X$ denotes cyclically reduced word length with respect to X . As was observed in [2] (see also [7]), there exist fully irreducible elements $\varphi \in \text{Out}(F_N)$ with the property that φ and φ^{-1} have *different* stretch factors:

$$\lambda(\varphi) \neq \lambda(\varphi^{-1}).$$

However, the following result from [6] describes the extent to which they can differ. To state their result precisely, let $N \geq 2$ be an integer and set

$$C_N = \sup_{\varphi} \frac{\log(\lambda(\varphi))}{\log(\lambda(\varphi^{-1}))},$$

where φ ranges over all fully irreducible elements of $\text{Out}(F_N)$.

E-mail addresses: dowdall@illinois.edu (S. Dowdall), kapovich@math.uiuc.edu (I. Kapovich), clein@math.uiuc.edu (C.J. Leininger).

URLs: <http://www.math.uiuc.edu/~dowdall/> (S. Dowdall), <http://www.math.uiuc.edu/~kapovich/> (I. Kapovich), <http://www.math.uiuc.edu/~clein/> (C.J. Leininger).

<http://dx.doi.org/10.1016/j.crma.2014.09.007>

1631-073X/© 2014 Académie des sciences. Published by Elsevier Masson SAS. All rights reserved.

Theorem 1. (Handel–Mosher [6].) For every integer $N \geq 2$, we have $C_N < \infty$.

An alternate proof of this result was more recently given by Algom–Kfir and Bestvina [1]. While the proofs of this theorem appeal to the fact that N is fixed, it is not clear that this dependence is necessary. In this note, we prove that in fact it is.

Theorem 2. With $\{C_N\}_{N \geq 2}$ defined as above, $\limsup_{N \rightarrow \infty} C_N = \infty$.

Proof. The proof will appeal to a construction and analysis carried out in [4] and [5]. To that end, let $F_3 = \langle a, b, c \rangle$ and consider the element $\varphi \in \text{Aut}(F_3)$ defined by

$$\varphi(a) = b, \quad \varphi(b) = b^{-1}a^{-1}bac, \quad \varphi(c) = a.$$

It was shown in [4, Example 2.19] that φ is fully irreducible. Next, let

$$G = F_3 \rtimes_{\varphi} \mathbb{Z} = \langle a, b, c, r \mid r^{-1}xr = \varphi(x) \text{ for all } x \in F_3 \rangle$$

be the free-by-cyclic group determined by φ , and let $u_0: G \rightarrow \mathbb{Z}$ in $\text{Hom}(G; \mathbb{R}) = H^1(G; \mathbb{R})$ be the associated homomorphism obtained by sending r to 1 and all other generators to 0.

In [4], we construct a cone $\mathcal{A} \subset H^1(G; \mathbb{R})$ containing u_0 with the property that every other primitive integral element $u \in \mathcal{A}$ has kernel $\ker(u)$ a finitely generated free group.

The action of $u(G) = \mathbb{Z}$ on $\ker(u)$ is generated by a *monodromy automorphism* $\varphi_u \in \text{Aut}(\ker(u))$ determining an expression of G as a semidirect product $G \cong \ker(u) \rtimes_{\varphi_u} \mathbb{Z}$ with associated homomorphism u . One of the main results of [4] is that all such φ_u are fully irreducible.

In [5], we construct a strictly larger open, convex cone $\mathcal{A} \subsetneq \mathcal{S} \subset H^1(G; \mathbb{R})$ and a function $\mathfrak{h}: \mathcal{S} \rightarrow \mathbb{R}$ that is convex, real analytic, and homogeneous of degree -1 (i.e., $\mathfrak{h}(tu) = \frac{1}{t}\mathfrak{h}(u)$) such that

$$\log(\lambda(\varphi_u)) = \mathfrak{h}(u)$$

for any primitive integral class $u \in \mathcal{A}$. In fact, this holds for all primitive integral $u \in \mathcal{S}$ with the appropriate interpretation of $\lambda(\varphi_u)$. We also show that \mathcal{S} is the cone on the component of the BNS-invariant $\Sigma(G)$ [3] containing u_0 [5, Theorem 1] and that \mathcal{A} lies over the symmetrized BNS-invariant (that is, both \mathcal{A} and $-\mathcal{A}$ project into $\Sigma(G)$) [5, Corollary 13.7]. In fact, a key result of Bieri–Neumann–Strebel is that an integral class $u \in \text{Hom}(G; \mathbb{Z})$ has $\ker(u)$ finitely generated if and only if both u and $-u$ lie in the $\Sigma(G)$ [3].

The homomorphism $-u_0$ has $\ker(-u_0) = \ker(u_0) = F_N$ and associated monodromy φ^{-1} , thus expressing G as $F_N \rtimes_{\varphi^{-1}} \mathbb{Z}$. Since φ^{-1} is also fully irreducible, the main result of [5] provides another open, convex cone $\mathcal{S}_- \subset H^1(G; \mathbb{R})$ containing $-u_0$ and a corresponding convex, real analytic, homogeneous of degree -1 function $\mathfrak{h}_-: \mathcal{S}_- \rightarrow \mathbb{R}$. Since $-\mathcal{A}$ projects into $\Sigma(G)$ and \mathcal{S}_- is the cone on the component of $\Sigma(G)$ containing $-u_0$, we see that $-\mathcal{A} \subset \mathcal{S}_-$. Thus \mathfrak{h}_- calculates the inverse stretch factors

$$\mathfrak{h}_-(-u) = \log(\lambda(\varphi_u^{-1}))$$

for all primitive integral $u \in \mathcal{A}$.

Example 8.3 of [5] exhibits a primitive integral class $u_1 \in \mathcal{S}$ which lies on the boundary of \mathcal{A} (see [5, Fig. 8]) for which $\ker(u_1)$ is *not* finitely generated. It follows that $-u_1$ is *not* in the BNS-invariant. The key observation is that $-u_1$ then necessarily lies on the boundary of \mathcal{S}_- (since $-u_1 \in \overline{-\mathcal{A}} \subset \overline{\mathcal{S}_-}$ but $-u_1 \notin \mathcal{S}_-$).

Let $\{u_n\}_{n=2}^{\infty} \subset \mathcal{A}$ be primitive integral classes protectively converging to u_1 . That is, there exists $\{t_n\}_{n=2}^{\infty} \subset \mathbb{R}$ so that $\lim_{n \rightarrow \infty} t_n u_n = u_1$. Since this convergence occurs inside \mathcal{S} , it follows that

$$\lim_{n \rightarrow \infty} \mathfrak{h}(t_n u_n) = \mathfrak{h}(u_1) < \infty.$$

On the other hand, since $\lim_{n \rightarrow \infty} -t_n u_n = -u_1 \in \partial \mathcal{S}_-$, it follows from [5, Theorem F] that

$$\lim_{n \rightarrow \infty} \mathfrak{h}_-(-t_n u_n) = \infty.$$

Therefore, appealing to the homogeneity of \mathfrak{h} and \mathfrak{h}_- , we have:

$$\lim_{n \rightarrow \infty} \frac{\log(\lambda(\varphi_{u_n}^{-1}))}{\log(\lambda(\varphi_{u_n}))} = \lim_{n \rightarrow \infty} \frac{\mathfrak{h}_-(-u_n)}{\mathfrak{h}(u_n)} = \lim_{n \rightarrow \infty} \frac{\mathfrak{h}_-(-t_n u_n)}{\mathfrak{h}(t_n u_n)} = \infty. \quad \square$$

Acknowledgements

The first author was partially supported by the NSF postdoctoral fellowship, NSF MSPRF No. 1204814. The second author was partially supported by the NSF grant DMS-1405146 and by the Simons Foundation Collaboration grant No. 279836. The third author was partially supported by the NSF grant DMS-1207183. The third author acknowledges support from U.S. National Science Foundation grants DMS 1107452, 1107263, 1107367 “GEAR Network”.

References

- [1] Y. Algom-Kfir, M. Bestvina, Asymmetry of outer space, *Geom. Dedicata* 156 (2012) 81–92.
- [2] M. Bestvina, M. Handel, Train tracks and automorphisms of free groups, *Ann. of Math. (2)* 135 (1) (1992) 1–51.
- [3] R. Bieri, W.D. Neumann, R. Strebel, A geometric invariant of discrete groups, *Invent. Math.* 90 (3) (1987) 451–477.
- [4] S. Dowdall, I. Kapovich, C.J. Leininger, Dynamics on free-by-cyclic-groups, preprint, arXiv:1301.7739v6, 2013.
- [5] S. Dowdall, I. Kapovich, C.J. Leininger, McMullen polynomials and Lipschitz flows for free-by-cyclic groups, preprint, arXiv:1310.7481v3, 2013.
- [6] M. Handel, L. Mosher, The expansion factors of an outer automorphism and its inverse, *Trans. Amer. Math. Soc.* 359 (7) (2007) 3185–3208 (electronic).
- [7] M. Handel, L. Mosher, Parageometric outer automorphisms of free groups, *Trans. Amer. Math. Soc.* 359 (7) (2007) 3153–3183 (electronic).