



ELSEVIER

Contents lists available at ScienceDirect

C. R. Acad. Sci. Paris, Ser. I

www.sciencedirect.com



Numerical analysis

A perturbation-method-based *a posteriori* estimator for the planewave discretization of nonlinear Schrödinger equations



Un estimateur d'erreur a posteriori basé sur une méthode de perturbation pour la discrétisation des équations de Schrödinger non linéaires dans une base d'ondes planes

Éric Cancès^a, Geneviève Dusson^{b,c}, Yvon Maday^{b,d,e}, Benjamin Stamm^b, Martin Vohralík^f

^a Université Paris-Est, CERMICS, École des ponts and INRIA, 6 & 8, av. Blaise Pascal, 77455 Marne-la-Vallée cedex 2, France

^b Sorbonne Universités, UPMC–Université Paris-6 and CNRS, UMR 7598, Laboratoire Jacques-Louis-Lions, 75005 Paris, France

^c Sorbonne Université, UPMC–Université Paris-6, Institut du calcul et de la simulation, 75005 Paris, France

^d Institut universitaire de France, 75005 Paris, France

^e Division of Applied Mathematics, Brown University, 182 George St, Providence, RI 02912, USA

^f INRIA Paris-Rocquencourt, domaine de Voluceau-Rocquencourt, BP 105, 78153 Le Chesnay, France

ARTICLE INFO

Article history:

Received 21 May 2014

Accepted 18 September 2014

Available online 27 September 2014

Presented by Olivier Pironneau

ABSTRACT

In this Note, we propose a new method, based on perturbation theory, to post-process the planewave approximation of the eigenmodes of periodic Schrödinger operators. We then use this post-processing to construct an accurate *a posteriori* estimator for the approximations of the (nonlinear) Gross–Pitaevskii equation, valid at each step of a self-consistent procedure. This allows us to design an adaptive algorithm for solving the Gross–Pitaevskii equation, which automatically refines the discretization along the convergence of the iterative process, by means of adaptive stopping criteria.

© 2014 Académie des sciences. Published by Elsevier Masson SAS. All rights reserved.

R É S U M É

Dans cette Note, nous proposons une nouvelle méthode, basée sur la théorie des perturbations, pour post-traiter l'approximation dans une base d'ondes planes des modes propres d'opérateurs de Schrödinger périodiques. Nous utilisons ensuite ce post-traitement pour construire un estimateur d'erreur *a posteriori* pour les approximations de l'équation de Gross–Pitaevskii (non linéaire), valide à chaque étape de la procédure auto-cohérente. Ceci nous permet de proposer un algorithme adaptatif pour résoudre cette équation, qui

E-mail addresses: cances@cermics.enpc.fr (É. Cancès), dusson@ann.jussieu.fr (G. Dusson), maday@ann.jussieu.fr (Y. Maday), stamm@ann.jussieu.fr (B. Stamm), martin.vohralik@inria.fr (M. Vohralík).

<http://dx.doi.org/10.1016/j.crma.2014.09.014>

1631-073X/© 2014 Académie des sciences. Published by Elsevier Masson SAS. All rights reserved.

raffine automatiquement la discrétisation au cours du processus itératif, par le biais de critères d'arrêt adaptatifs.

© 2014 Académie des sciences. Published by Elsevier Masson SAS. All rights reserved.

1. Introduction

The time-independent Gross–Pitaevskii and Kohn–Sham models are widely used in quantum physics to describe, respectively, steady states of Bose–Einstein condensates and electronic ground states of molecular systems. Although these models dramatically differ from the physical viewpoint (the former describes bosons, the latter describes fermions), their mathematical structures share some similarities: they are constrained optimization problems whose Euler–Lagrange equations are nonlinear elliptic eigenvalue problems. As a consequence, the numerical methods used to simulate these models have common features. In both cases, numerical solutions can be obtained either by minimization of the energy functional on the admissible set, or by solving the Euler–Lagrange equations by a self-consistent field (SCF) procedure. Algorithms mixing the two approaches have also been proposed.

Among the various discretization methods available, the planewave (Fourier) method is one of the most popular ones. It is in particular the state-of-the-art method for Kohn–Sham simulations in solid-state physics and materials science. This method consists in imposing periodic conditions at the boundary of the simulation domain Ω (a cubic box, or more generally the unit cell of a periodic lattice $\mathcal{R} \subset \mathbb{R}^d$), and in expanding the wave function (or more precisely the occupied orbitals in the Kohn–Sham setting) in the discretization space

$$X_N := \text{Span}(e_k, |k| \leq N),$$

where $e_k(x) := |\Omega|^{-1} e^{ik \cdot x}$ is the Fourier mode with wavevector $k \in \mathcal{R}^*$, \mathcal{R}^* denoting the dual lattice of \mathcal{R} . For example, we have for a cubic lattice, $\Omega = (0, L)^d$, $\mathcal{R} = L\mathbb{Z}^d$, and $\mathcal{R}^* = \frac{2\pi}{L}\mathbb{Z}^d$. Let the periodic Lebesgue space $L^2_{\#}(\Omega)$ and the periodic Sobolev spaces $H^r_{\#}(\Omega)$ ($r \in \mathbb{R}$) be defined by

$$L^2_{\#}(\Omega) := \left\{ v \in L^2_{\text{loc}}(\mathbb{R}^d) \mid v \text{ } \mathcal{R}\text{-periodic} \right\} = \left\{ v = \sum_{k \in \mathcal{R}^*} \widehat{v}_k e_k \mid \|v\|_{L^2_{\#}}^2 := \sum_{k \in \mathcal{R}^*} |\widehat{v}_k|^2 < \infty \right\},$$

$$H^r_{\#}(\Omega) := \left\{ v = \sum_{k \in \mathcal{R}^*} \widehat{v}_k e_k \mid \|v\|_{H^r_{\#}}^2 := \sum_{k \in \mathcal{R}^*} (1 + |k|^2)^r |\widehat{v}_k|^2 < \infty \right\}.$$

We denote by Π_N the operator on the space of \mathcal{R} -periodic tempered distributions defined by

$$\Pi_N \left(\sum_{k \in \mathcal{R}^*} \widehat{v}_k e_k \right) = \sum_{|k| \leq N} \widehat{v}_k e_k$$

($\Pi_N|_{H^s_{\#}(\Omega)}$) is in fact the orthogonal projector from $H^s_{\#}(\Omega)$ to X_N for any $s \in \mathbb{R}$), and by $\Pi_N^{\perp} = (1 - \Pi_N)$ the orthogonal projector on the orthogonal space X_N^{\perp} of X_N in $H^s_{\#}(\Omega)$ (for any $s \in \mathbb{R}$). The periodic Gross–Pitaevskii model reads

$$\inf \{ E_{\text{GP}}(v), v \in H^1_{\#}(\Omega), \|v\|_{L^2_{\#}} = 1 \}, \tag{1}$$

where the Gross–Pitaevskii functional is given, in absence of an external magnetic field, by

$$E_{\text{GP}}(v) := \int_{\Omega} |\nabla v|^2 + \int_{\Omega} V |v|^2 + \frac{\mu}{2} \int_{\Omega} |v|^4,$$

where V is a real-valued periodic potential belonging to $H^s_{\#}(\Omega)$ for some $s > d/2$, and $\mu \in \mathbb{R}$. The physical setting corresponds to $d = 3$, but our analysis is valid for any dimension $d \leq 3$. On the other hand, we will assume for simplicity that the real parameter μ , which models the magnitude of the interactions between the condensed particles is non-negative (repulsive interactions). It is well-known that (1) has a unique minimizer u up to a global phase, and that (still up to a global phase) u is real-valued, belongs to $H^2_{\#}(\Omega)$, and is everywhere positive. In addition, the function u satisfies the Euler–Lagrange equation:

$$-\Delta u + Vu + \mu u^3 = \lambda u, \quad \|u\|_{L^2_{\#}} = 1,$$

where λ , the Lagrange multiplier of the constraint $\|v\|_{L^2_{\#}} = 1$, is in fact the lowest eigenvalue of the mean-field operator $H_u := -\Delta + V + \mu u^2$. Note that H_u is a bounded-below self-adjoint operator on $L^2_{\#}(\Omega)$ with domain $H^2_{\#}(\Omega)$ and compact resolvent. The variational approximation of (1) in the discretization space X_N gives rise to the finite-dimensional constrained

optimization problem:

$$\inf\{E_{\text{GP}}(v_N), v_N \in X_N, \|v_N\|_{L^2_{\#}} = 1\}. \tag{2}$$

As Π_N and the Laplace operator commute, the minimizer λ_N of (2) satisfies

$$-\Delta u_N + \Pi_N((V + |u_N|^2)u_N) = \lambda_N u_N, \quad \|u_N\|_{L^2_{\#}} = 1, \tag{3}$$

where λ_N is the Lagrange multiplier associated with the constraint $\|v_N\|_{L^2_{\#}} = 1$. It is proved in [2] that, for N large enough, the discretized problem (2) has a unique real-valued minimizer u_N such that $(u, u_N)_{L^2_{\#}} \geq 0$ (the other minimizers being equal to u_N up to a global phase), and that the following optimal *a priori* error estimates hold

$$\begin{aligned} c\|u - u_N\|_{H^1_{\#}}^2 &\leq E_{\text{GP}}(u_N) - E_{\text{GP}}(u) \leq C\|u - u_N\|_{H^1_{\#}}^2, \\ |\lambda - \lambda_N| &\leq CN^{-2(s+1)}, \quad \|u - u_N\|_{H^r_{\#}} \leq CN^{-(s+2-r)}, \end{aligned} \tag{4}$$

for all $-s \leq r < s + 2$, and where the constants c and C depend on $\|V\|_{H^s_{\#}}$ and μ , but are independent of N .

As mentioned above, the discretized problem (3) can be solved either by constrained minimization or by SCF algorithms. Our approach can be applied to a variety of iteration schemes, but for brevity, we only discuss here the simple case of the Roothaan algorithm. We are aware that the Roothaan algorithm is far from being the most efficient one to solve the Gross–Pitaevskii equation, but we nevertheless chose it for two reasons: first, it allows us to illustrate our approach on a simple example where technicalities are reduced to a minimum, and second this algorithm is somewhat similar to the SCF algorithms actually used in Kohn–Sham simulations.

The purpose of this Note is twofold. First, we propose a method to post-process the plane-wave approximation of the eigenmodes of (linear) periodic Schrödinger operators and obtain a much better accuracy for a limited extra computational cost. This post-processing, although based on a simple application of the perturbation method, seems to be new. Note that this approach can be extended to the case of the (nonlinear) Gross–Pitaevskii equation [3]. Second, we use this post-processing to construct an accurate *a posteriori* error estimator for the Gross–Pitaevskii equation. This estimator allows us to design an adaptive algorithm for solving this equation, which automatically refines the discretization along the convergence of the SCF iterations, by means of adaptive stopping criteria. The first steps of the SCF procedure are carried out in a coarse discretization space (it is indeed a loss of time to use a fine discretization space far from the minimizer), and the cut-off N is gradually increased to balance the two components of the total error: the discretization error on the one hand, and the error due to the fact that the SCF iterations has not converged yet on the other hand. The extension of this approach to more complex nonlinear Schrödinger equations and to the Kohn–Sham model is work in progress.

2. Post-processing of the plane-wave approximation of periodic Schrödinger operators

We first consider a generic linear periodic Schrödinger operator $H = -\Delta + W$ on $L^2_{\#}(\Omega)$, with $W \in H^s_{\#}(\Omega)$ for some $s \geq 0$. In regard of the previous section, H can denote the linearized Gross–Pitaevskii Hamiltonian $H_{u^{j-1}} = -\Delta + V + \mu(u^{j-1})^2$ at step j around an approximation u^{j-1} of u , for example. For simplicity, we focus on the ground-state eigenvalue ν of H , but our argument straightforwardly extends to any non-degenerate eigenvalues by using the minmax principle. The case of degenerate eigenvalue is dealt with in [3]. Denoting by $w \in H^2_{\#}(\Omega)$ the unique positive ground state of H , it is well-known that the optimization problem

$$\nu_N = \inf\{E_S(v_N), v_N \in X_N, \|v_N\|_{L^2_{\#}} = 1\} \quad \text{with } E_S(v) := \int_{\Omega} |\nabla v|^2 + \int_{\Omega} W|v|^2 = (v, Hv)_{L^2_{\#}}$$

has a unique real-valued minimizer w_N such that $(w_N, 1)_{L^2_{\#}} \geq 0$, and that (focusing on the H^1 -norm for brevity)

$$c\|w - w_N\|_{H^1_{\#}}^2 \leq \nu_N - \nu \leq C\|w - w_N\|_{H^1_{\#}}^2 \quad \text{and} \quad \|w - w_N\|_{H^1_{\#}} \leq CN^{-(s+1)},$$

with $0 < c \leq C < \infty$ constants independent of N . The observation our approach is based upon is that (ν_N, w_N) satisfies

$$H_N w_N = \nu_N w_N, \quad \|w_N\|_{L^2_{\#}} = 1 \quad \text{where } H_N = -\Delta + \Pi_N W \Pi_N,$$

while (ν, w) satisfies

$$(H_N + \mathcal{W}_N)w = \nu w, \quad \|w\|_{L^2_{\#}} = 1 \quad \text{with } \mathcal{W}_N = \Pi_N^{\perp} W \Pi_N + \Pi_N W \Pi_N^{\perp} + \Pi_N^{\perp} W \Pi_N^{\perp}.$$

In addition, it can be seen that as soon as $N^2 > \nu_N$, w_N is the ground state of H_N . It therefore makes sense to apply Kato’s perturbation theory using (ν_N, w_N) as the reference solution and (ν, w) as the perturbed solution, in order to build

improved approximations $(\tilde{v}_N, \tilde{w}_N)$ based upon perturbation arguments. A simple calculation shows that the first-order correction to the eigenfunction w_N is given by

$$w_N^{(1)} = -(-\Delta|_{X_N^\perp} - \nu_N)^{-1}r_N \quad \text{where } r_N = Hw_N - \nu_N w_N = \Pi_N^\perp(Ww_N) \in X_N^\perp$$

is the residual of the eigenvalue problem. The first-order correction to the eigenvalue vanishes and the second order correction is (using Dirac’s bracket notation):

$$\nu_N^{(2)} = (w_N^{(1)}, r_N)_{L_\#^2} = -\langle r_N | (-\Delta|_{X_N^\perp} - \nu_N)^{-1} | r_N \rangle.$$

Note that $w_N^{(1)}$ and $\nu_N^{(2)}$ are well-defined (provided that $N^2 > \nu_N$) and computable by a single Fast Fourier Transform on a finer grid. An extension to the result below to more general operators and discretization spaces, higher-order perturbations, and degenerate eigenvalues, is proved in [3].

Lemma 2.1. (See [3].) *Let $W \in H_\#^s(\Omega)$ for some $s > d/2$. For all $N \in \mathbb{N}$ such that $N^2 > \nu_N$, the quantities $\tilde{w}_N = w_N + w_N^{(1)}$ and $\tilde{\nu}_N = \nu_N + \nu_N^{(2)}$ are well-defined and there exists a positive real constant C independent of N such that*

$$|\nu - \tilde{\nu}_N| \leq CN^{-2}|\nu - \nu_N| \quad \text{and} \quad \|w - \tilde{w}_N\|_{H_\#^1} \leq CN^{-2}\|w - w_N\|_{H_\#^1}.$$

3. A posteriori estimators and adaptive algorithms for the Gross–Pitaevskii equation

We now use the results established in the previous section to design an adaptive algorithm based on the Roothaan procedure for the solution to (3), whose iterations are defined by:

$$-\Delta u^j + \Pi_{N_j}(Vu^j + \mu|u^{j-1}|^2u^j) = \lambda^j u^j, \quad u^j \in X_{N_j}, \quad \|u^j\|_{L_\#^2} = 1, \quad (u^j, 1)_{L_\#^2} \geq 0, \tag{5}$$

where $u^0 \in X_{N_0}$ is a given initial guess, and where $(N_j)_{j \in \mathbb{N}}$ is a non-decreasing sequence of integers. Note that u^j is in some sense a shortcut notation for $u_{N_j}^j$, since this function belongs to the approximation space X_{N_j} . Usually, N_j is kept constant (equal to N) along the iterations. In our approach, the sequence $(N_j)_{j \in \mathbb{N}}$ is not specified *a priori*, but automatically constructed by the algorithm. This is achieved by using an *a posteriori* estimator of the error between u^j and u , which can be decomposed into two non-negative contributions:

- i) a contribution η_{scf}^j measuring the error at step j due to the fact that the SCF procedure has not converged;
- ii) a contribution η_{dis}^j measuring the error at step j due to the fact that u^j is a finite-dimensional approximation of u in X_{N_j} .

Following the ideas of [1,4,5], we propose to refine the discretization space whenever $\eta_{\text{scf}}^j \leq \zeta \eta_{\text{dis}}^j$, where ζ is some user-given parameter, typically of order 0.1. The new discretization parameter N_{j+1} can then be chosen by linear regression based on the *a priori* convergence rate in order that the predicted new discretization error is smaller than $\zeta' \eta_{\text{scf}}^j$ for some user-given parameter ζ' .

Let us now provide explicit expressions for η_{scf}^j and η_{dis}^j . As usual in *a posteriori* error estimations, we begin with discussing the choice of the measure of the error. A common choice for nonlinear problems is the energy difference $E_{\text{GP}}(u^j) - E_{\text{GP}}(u)$. In our setting, it appears that we can better estimate the augmented energy difference $J_u(u^j)$ where

$$J_u(v) := E_{\text{GP}}(v) - E_{\text{GP}}(u) + \frac{\mu}{2} \int_{\Omega} (|u|^2 - |v|^2)^2.$$

Note that for all $v \in H_\#^1(\Omega)$ such that $\|v\|_{L_\#^2} = 1$ and $(v, 1)_{L_\#^2} \geq 0$, $J(v) \geq 0$ and $J(v) = 0$ if and only if $v = u$. We also know from [2] that there exists $0 < c \leq C < \infty$ such that $c\|v - u\|_{H_\#^1}^2 \leq J_u(v) \leq C\|v - u\|_{H_\#^1}^2$ for all $v \in H_\#^1(\Omega)$ such that $\|v\|_{L_\#^2} = 1$ and $(v, 1)_{L_\#^2} \geq 0$.

Theorem 3.1. (See [3].) *Let u be the exact solution to (1) satisfying $u > 0$ in Ω and $u^j \in X_{N_j}$ the numerical approximation of u generated by the iterative procedure (5) at step j . Then*

$$J_u(u^j) \leq \eta_{\text{tot}}^j := \left(\lambda^j - \lambda_{1b}^{j+1} + \mu \int_{\Omega} [(u^j)^2 - (u^{j-1})^2](u^j)^2 \right),$$




Fig. 1. (Color online.) Left: convergence rate with respect to N with and without post-processing the eigenvector as well as the quotient $q_N = \|w - \tilde{w}_N\|_{H^1_\#} / \|w - w_N\|_{H^1_\#}$. Right: convergence of the error in the energy and the augmented energy difference $J_u(u^j)$, and *a posteriori* estimators with respect to the Roothaan iterations for $N = 50$. Gauche : taux de convergence par rapport à N avec et sans post-traitement du vecteur propre et quotient $q_N = \|w - \tilde{w}_N\|_{H^1_\#} / \|w - w_N\|_{H^1_\#}$. Droite : convergence de l'erreur sur l'énergie et la différence d'énergie augmentée $J_u(u^j)$, et estimateurs *a posteriori* en fonction des itérations de Roothaan pour $N = 50$.

where λ_{lb}^{j+1} is a lower bound of the exact ground state eigenvalue λ_∞^{j+1} of H_{u^j} . In addition, the error bound η_{tot}^j can be decomposed as $\eta_{tot}^j = \eta_{scf}^j + \eta_{dis}^j$ where

$$\eta_{dis}^j := (\lambda_{N_j}^{j+1} - \lambda_{lb}^{j+1}) \geq 0, \quad \eta_{scf}^j := \left(\mu \int_{\Omega} [(u^j)^2 - (u^{j-1})^2] (u^j)^2 + \lambda^j - \lambda_{N_j}^{j+1} \right) \geq 0,$$

where $\lambda_{N_j}^{j+1}$ is the variational approximation of λ_∞^{j+1} in the discretization space X_{N_j} .

In the numerical results reported in the next section, we first compute $\lambda_{N_j}^{j+1}$ as the lowest eigenvalue of $-\Delta + \Pi_{N_j} W^j \Pi_{N_j}$ with $W^j = V + \mu|u^j|^2$ (which amounts to make one more iteration of the Roothaan algorithm in the discretization space X_{N_j}), and we estimate λ_∞^{j+1} by post-processing $\lambda_{N_j}^{j+1}$ thanks to the perturbation argument provided in Section 2. Although the so-obtained post-processed eigenvalue is not strictly speaking a lower bound of λ_∞^{j+1} , it is an excellent approximation of it and can therefore be used instead of λ_{lb}^{j+1} to compute the error estimators in Theorem 3.1. Note that it is also possible to use perturbation theory to estimate both $\lambda_{N_j}^{j+1}$ and λ_{lb}^{j+1} starting from λ^j (see [3]).

4. Numerical results

We first consider the linear one-dimensional periodic Schrödinger operator with potential $W(x) = |\sin(2\pi x)| + 6$ and $\Omega = (0, 1)$. We observe in Fig. 1 (left) very good agreements with the theoretical results of Lemma 2.1. We then report in Fig. 1 (right) simulations of the one-dimensional Gross–Pitaevskii equation with $V(x) = |\sin(2\pi x)| + 6$ and $\mu = 1$, showing the decay of different quantities related to Theorem 3.1 for a fixed $N = 50$ during the Roothaan iterations, and illustrating the sharpness of the estimator and the splitting of the error components. More extensive numerical results in higher dimensions will be presented in [3].

Acknowledgements

This work was partially undertaken in the framework of CALSIMLAB, supported by the public grant ANR-11-LABX-0037-01 overseen by the French National Research Agency (ANR) as part of the “Investissements d’avenir” program (reference: ANR-11-IDEX-0004-02). Financial support by the ANR grants Manif and Becasim is also acknowledged.

References

[1] R. Becker, C. Johnson, R. Rannacher, Adaptive error control for multigrid finite element methods, *Computing* 55 (1995) 271–288.
 [2] E. Cancès, R. Chakir, Y. Maday, Numerical analysis of nonlinear eigenvalue problems, *J. Sci. Comput.* 45 (2010) 90–117.

- [3] E. Cancès, G. Dusson, Y. Maday, B. Stamm, M. Vohralík, On the use of the perturbation method in planewave numerical simulations of linear and nonlinear Schrödinger equations, in preparation.
- [4] A.L. Chaillou, M. Suri, A posteriori estimation of the linearization error for strongly monotone nonlinear operators, *J. Comput. Appl. Math.* 205 (2007) 72–87.
- [5] A. Ern, M. Vohralík, Adaptive inexact Newton methods with a posteriori stopping criteria for nonlinear diffusion PDEs, *SIAM J. Sci. Comput.* 35 (2013) A1761–A1791.