

MEAN-FIELD STOCHASTIC LINEAR QUADRATIC OPTIMAL CONTROL PROBLEMS: OPEN-LOOP SOLVABILITIES *

JINGRUI SUN¹

Abstract. This paper is concerned with a mean-field linear quadratic (LQ, for short) optimal control problem with deterministic coefficients. It is shown that convexity of the cost functional is necessary for the finiteness of the mean-field LQ problem, whereas uniform convexity of the cost functional is sufficient for the open-loop solvability of the problem. By considering a family of uniformly convex cost functionals, a characterization of the finiteness of the problem is derived and a minimizing sequence, whose convergence is equivalent to the open-loop solvability of the problem, is constructed. Then, it is proved that the uniform convexity of the cost functional is equivalent to the solvability of two coupled differential Riccati equations and the unique open-loop optimal control admits a state feedback representation in the case that the cost functional is uniformly convex. Finally, some examples are presented to illustrate the theory developed.

Mathematics Subject Classification. 49N10, 49N35, 93E20.

Received September 15, 2015. Accepted April 21, 2016.

1. INTRODUCTION

Let $(\Omega, \mathcal{F}, \mathbb{F}, \mathbb{P})$ be a complete filtered probability space on which a standard one-dimensional Brownian motion $W = \{W(t); 0 \leq t < \infty\}$ is defined, where $\mathbb{F} = \{\mathcal{F}_t\}_{t \geq 0}$ is the natural filtration of W augmented by all the \mathbb{P} -null sets in \mathcal{F} . Consider the following controlled linear stochastic differential equation (SDE, for short) on a finite horizon $[t, T]$:

$$\begin{cases} dX(s) = \left\{ A(s)X(s) + \bar{A}(s)\mathbb{E}[X(s)] + B(s)u(s) + \bar{B}(s)\mathbb{E}[u(s)] + b(s) \right\} ds \\ \quad + \left\{ C(s)X(s) + \bar{C}(s)\mathbb{E}[X(s)] + D(s)u(s) + \bar{D}(s)\mathbb{E}[u(s)] + \sigma(s) \right\} dW(s), & s \in [t, T], \\ X(t) = \xi, \end{cases} \quad (1.1)$$

where $A(\cdot)$, $\bar{A}(\cdot)$, $B(\cdot)$, $\bar{B}(\cdot)$, $C(\cdot)$, $\bar{C}(\cdot)$, $D(\cdot)$, $\bar{D}(\cdot)$ are given deterministic matrix-valued functions; $b(\cdot)$, $\sigma(\cdot)$ are vector-valued \mathbb{F} -progressively measurable processes and ξ is an \mathcal{F}_t -measurable random vector. In the above,

Keywords and phrases. Mean-field stochastic differential equation, linear quadratic optimal control, Riccati equation, finiteness, open-loop solvability, feedback representation.

* This work is supported in part by the National Natural Science Foundation of China (11401556) and the Fundamental Research Funds for the Central Universities (WK 2040000012).

¹ Department of Applied Mathematics, The Hong Kong Polytechnic University, Hong Kong, P.R. China. sjr@mail.ustc.edu.cn

$u(\cdot)$ is the *control process* and $X(\cdot)$ is the corresponding *state process* with *initial pair* (t, ξ) . For any $t \in [0, T)$, we define

$$\mathcal{U}[t, T] = \left\{ u : [t, T] \times \Omega \rightarrow \mathbb{R}^m \mid u(\cdot) \text{ is } \mathbb{F}\text{-progressively measurable, } \mathbb{E} \int_t^T |u(s)|^2 ds < \infty \right\}.$$

Any $u(\cdot) \in \mathcal{U}[t, T]$ is called an *admissible control* (on $[t, T]$). Under some mild conditions, for any initial pair (t, ξ) with ξ being square-integrable and any admissible control $u(\cdot) \in \mathcal{U}[t, T]$, (1.1) admits a unique square-integrable solution $X(\cdot) \equiv X(\cdot; t, \xi, u(\cdot))$. Now we introduce the following cost functional:

$$\begin{aligned} J(t, \xi; u(\cdot)) \triangleq & \mathbb{E} \left\{ \langle GX(T), X(T) \rangle + 2\langle g, X(T) \rangle + \langle \bar{G}\mathbb{E}[X(T)], \mathbb{E}[X(T)] \rangle + 2\langle \bar{g}, \mathbb{E}[X(T)] \rangle \right. \\ & + \int_t^T \left[\left\langle \begin{pmatrix} Q(s) & S(s)^\top \\ S(s) & R(s) \end{pmatrix} \begin{pmatrix} X(s) \\ u(s) \end{pmatrix}, \begin{pmatrix} X(s) \\ u(s) \end{pmatrix} \right\rangle + 2 \left\langle \begin{pmatrix} q(s) \\ \rho(s) \end{pmatrix}, \begin{pmatrix} X(s) \\ u(s) \end{pmatrix} \right\rangle \right] ds \\ & \left. + \int_t^T \left[\left\langle \begin{pmatrix} \bar{Q}(s) & \bar{S}(s)^\top \\ \bar{S}(s) & \bar{R}(s) \end{pmatrix} \begin{pmatrix} \mathbb{E}[X(s)] \\ \mathbb{E}[u(s)] \end{pmatrix}, \begin{pmatrix} \mathbb{E}[X(s)] \\ \mathbb{E}[u(s)] \end{pmatrix} \right\rangle + 2 \left\langle \begin{pmatrix} \bar{q}(s) \\ \bar{\rho}(s) \end{pmatrix}, \begin{pmatrix} \mathbb{E}[X(s)] \\ \mathbb{E}[u(s)] \end{pmatrix} \right\rangle \right] ds \right\}, \end{aligned} \quad (1.2)$$

where G, \bar{G} are symmetric matrices and $Q(\cdot), \bar{Q}(\cdot), S(\cdot), \bar{S}(\cdot), R(\cdot), \bar{R}(\cdot)$ are deterministic matrix-valued functions with $Q(\cdot)^\top = Q(\cdot)$, $\bar{Q}(\cdot)^\top = \bar{Q}(\cdot)$, $R(\cdot)^\top = R(\cdot)$, $\bar{R}(\cdot)^\top = \bar{R}(\cdot)$; g is an \mathcal{F}_T -measurable random vector and \bar{g} is a (deterministic) vector; $q(\cdot), \rho(\cdot)$ are vector-valued \mathbb{F} -progressively measurable processes and $\bar{q}(\cdot), \bar{\rho}(\cdot)$ are deterministic vector-valued functions. Our mean-field stochastic LQ optimal control problem can be stated as follows:

Problem (MF-LQ). For any given initial pair $(t, \xi) \in [0, T) \times L^2_{\mathcal{F}_t}(\Omega; \mathbb{R}^n)$, find a $u^*(\cdot) \in \mathcal{U}[t, T]$ such that

$$J(t, \xi; u^*(\cdot)) = \inf_{u(\cdot) \in \mathcal{U}[t, T]} J(t, \xi; u(\cdot)) \triangleq V(t, \xi). \quad (1.3)$$

In the above, $L^2_{\mathcal{F}_t}(\Omega; \mathbb{R}^n)$ is the space of all \mathcal{F}_t -measurable, \mathbb{R}^n -valued random vectors ξ with $\mathbb{E}|\xi|^2 < \infty$. Any $u^*(\cdot) \in \mathcal{U}[t, T]$ satisfying (1.3) is called an (*open-loop*) *optimal control* of Problem (MF-LQ) for the initial pair (t, ξ) , and the corresponding $X^*(\cdot) \equiv X(\cdot; t, \xi, u^*(\cdot))$ is called an *optimal state process*. The function $V(\cdot, \cdot)$ is called the *value function* of Problem (MF-LQ). In the special case where $b(\cdot), \sigma(\cdot), g, \bar{g}, q(\cdot), \bar{q}(\cdot), \rho(\cdot)$, and $\bar{\rho}(\cdot)$ vanish, we denote the corresponding mean-field LQ problem, cost functional, and value function by Problem (MF-LQ)⁰, $J^0(t, \xi; u(\cdot))$, and $V^0(t, \xi)$, respectively.

Comparing with the classical stochastic LQ optimal control problem, a new feature of Problem (MF-LQ) is that both the state equation and the cost functional involve the states and the controls as well as their expectations. In this case, we call (1.1) a controlled mean-field (forward) SDE (MF-SDE, for short). The history of MF-SDEs can be traced back to the work of Kac [18] in 1956 and McKean [21] in 1966. Since then, many researchers have made contributions to such kind of equations and applications; see, for example, [2, 9, 10, 12–15, 24]. For recent development of MF-SDEs, readers may refer to [5, 7, 8, 11, 17, 19, 20, 27] and the references cited therein. Control problems of MF-SDEs were studied by Ahmed *et al.* [3], Ahmed [1], Park *et al.* [23], Buckdahn *et al.* [6], Andersson and Djehiche [4], Meyer *et al.* [22], and so on. More recently, Yong [28] investigated an LQ problem for MF-SDEs in finite horizons and gave some interesting motivation for the control problem with $\mathbb{E}[X(\cdot)]$ and $\mathbb{E}[u(\cdot)]$ being included in the cost functional. Later, Huang *et al.* [16] generalized the results in [28] to the case with an infinite time horizon.

In [28], two coupled differential Riccati equations are derived by decoupling the optimality system. It is shown that under certain conditions, the two Riccati equations are uniquely solvable and Problem (MF-LQ) admits

a unique optimal control which has a state feedback representation. To be precise, if

$$\begin{cases} G, G + \bar{G} \geq 0, & Q(s), Q(s) + \bar{Q}(s) \geq 0, \\ S(s) = \bar{S}(s) = 0, & R(s), R(s) + \bar{R}(s) \geq \delta I, \end{cases} \quad \text{a.e. } s \in [0, T], \quad (1.4)$$

for some $\delta > 0$, then the unique solvability of the two Riccati equations can be obtained from the classical result ([29], Thm. 7.2). However, examples show that the two Riccati equations might still be solvable even if both $R(\cdot)$ and $\bar{R}(\cdot)$ are negative semi-definite (see Example 6.1). On the other hand, it may happen that Problem (MF-LQ) is open-loop solvable, while the optimal control cannot be obtained by solving the corresponding Riccati equations due to the possible singularities of the terms $R + D^\top P D$ and $R + \bar{R} + (D + \bar{D})^\top P (D + \bar{D})$ (see Example 6.2). Thus, some questions arise naturally: (a) what is the relationship between Problem (MF-LQ) and the solvability of the two Riccati equations? (b) How can we characterize the open-loop solvability of Problem (MF-LQ)? (c) How can we find an optimal control in general? The purpose of this paper is to study Problem (MF-LQ) from an open-loop point of view and to address the above issues. Closed-loop mean-field LQ problems will be investigated in a forthcoming paper.

Our main idea and results of this paper can be informally described as follows. By a representation of the cost functional, we first show that for the open-loop solvability of Problem (MF-LQ), a necessary condition is the convexity of the cost functional and a sufficient condition is the uniform convexity of the cost functional. Under the convexity condition, by adding $\varepsilon \mathbb{E} \int_t^T |u(s)|^2 ds$ ($\varepsilon > 0$) to the original cost functional, we get a family of uniformly convex functionals. The corresponding mean-field LQ problems admit unique optimal controls $u_\varepsilon^*(\cdot)$, $\varepsilon > 0$, which form a minimizing sequence of Problem (MF-LQ). Then the open-loop solvability of Problem (MF-LQ) is characterized by the convergence of the sequence, whose limit is an optimal control of Problem (MF-LQ). To construct $u_\varepsilon^*(\cdot)$ explicitly, we further investigate Problem (MF-LQ) with uniformly convex cost functionals. Since the uniform convexity condition is much weaker than (1.4), the result in [28] fails to apply to this case. To overcome this difficulty, we reduce Problem (MF-LQ) to a classical stochastic LQ problem and a deterministic LQ problem. By making use of a result found in [26], we establish the equivalence between the uniform convexity of the cost functional and the solvability of the two Riccati equations. Then by the completion of squares technique, we obtain a state feedback representation of the optimal control *via* the solutions of the two Riccati equations.

The rest of the paper is organized as follows. Section 2 gives some preliminaries. In Section 3, we study Problem (MF-LQ) from a Hilbert space viewpoint and derive necessary and sufficient conditions for the finiteness and open-loop solvability of the problem by considering a family of uniformly convex cost functionals. Section 4 shows that the solvability of two coupled Riccati equations is necessary for the uniform convexity of the cost functional. In Section 5, we further prove that the solvability of the two coupled Riccati equations is also sufficient for the uniform convexity of the cost functional. Moreover, a state feedback representation is obtained for the optimal control. Some illustrative examples are presented in Section 6.

2. PRELIMINARIES

Throughout this paper, we denote by $\mathbb{R}^{n \times m}$ the Euclidean space of all $n \times m$ real matrices, and by \mathbb{S}^n the space of all symmetric $n \times n$ real matrices. Recall that the inner product $\langle \cdot, \cdot \rangle$ on $\mathbb{R}^{n \times m}$ is given by $\langle M, N \rangle \mapsto \text{tr}(M^\top N)$, where the superscript \top denotes the transpose of vectors or matrices, and the induced norm is given by $|M| = \sqrt{\text{tr}(M^\top M)}$. When there is no confusion, we shall use $\langle \cdot, \cdot \rangle$ for inner products in possibly different Hilbert spaces, and denote by $|\cdot|$ the norm induced by $\langle \cdot, \cdot \rangle$. For a matrix $M \in \mathbb{R}^{n \times m}$, we denote by $\mathcal{R}(M)$ the range of M , and if $M \in \mathbb{S}^n$, we use the notation $M > 0$ (≥ 0) to indicate that M is positive (semi-) definite. For a bounded linear operator \mathcal{A} from a Banach \mathcal{X} space into another Banach \mathcal{Y} space, we denote by \mathcal{A}^* the adjoint operator of \mathcal{A} . Let $T > 0$ be a fixed time horizon. For any $t \in [0, T]$ and Euclidean space \mathbb{H} , we let $L^p(t, T; \mathbb{H})$ ($1 \leq p \leq \infty$) be the space of all \mathbb{H} -valued functions that are L^p -integrable on $[t, T]$, and let $C([t, T]; \mathbb{H})$ be the space of all \mathbb{H} -valued continuous functions on $[t, T]$. Next, we introduce the following

spaces:

$$\begin{aligned}
 L^2_{\mathcal{F}_t}(\Omega; \mathbb{H}) &= \left\{ \xi : \Omega \rightarrow \mathbb{H} \mid \xi \text{ is } \mathcal{F}_t\text{-measurable, } \mathbb{E}|\xi|^2 < \infty \right\}, \\
 L^2_{\mathbb{F}}(t, T; \mathbb{H}) &= \left\{ \varphi : [t, T] \times \Omega \rightarrow \mathbb{H} \mid \varphi(\cdot) \text{ is } \mathbb{F}\text{-progressively measurable, } \mathbb{E} \int_t^T |\varphi(s)|^2 ds < \infty \right\}, \\
 L^2_{\mathbb{F}}(\Omega; C([t, T]; \mathbb{H})) &= \left\{ \varphi : [t, T] \times \Omega \rightarrow \mathbb{H} \mid \varphi(\cdot) \text{ is } \mathbb{F}\text{-adapted, continuous, } \mathbb{E} \left(\sup_{s \in [t, T]} |\varphi(s)|^2 \right) < \infty \right\}, \\
 L^2_{\mathbb{F}}(\Omega; L^1(t, T; \mathbb{H})) &= \left\{ \varphi : [t, T] \times \Omega \rightarrow \mathbb{H} \mid \varphi(\cdot) \text{ is } \mathbb{F}\text{-progressively measurable, } \mathbb{E} \left(\int_t^T |\varphi(s)| ds \right)^2 < \infty \right\}.
 \end{aligned}$$

Further, we introduce the following notation: For any \mathbb{S}^n -valued measurable function F on $[t, T]$,

$$\begin{cases}
 F \geq 0 & \iff F(s) \geq 0, & \text{a.e. } s \in [t, T], \\
 F > 0 & \iff F(s) > 0, & \text{a.e. } s \in [t, T], \\
 F \gg 0 & \iff F(s) \geq \delta I, & \text{a.e. } s \in [t, T], \text{ for some } \delta > 0.
 \end{cases}$$

The following assumptions will be in force throughout this paper.

(H1) The coefficients of the state equation satisfy the following:

$$\begin{cases}
 A(\cdot), \bar{A}(\cdot) \in L^1(0, T; \mathbb{R}^{n \times n}), \quad B(\cdot), \bar{B}(\cdot) \in L^2(0, T; \mathbb{R}^{n \times m}), \quad b(\cdot) \in L^2_{\mathbb{F}}(\Omega; L^1(0, T; \mathbb{R}^n)), \\
 C(\cdot), \bar{C}(\cdot) \in L^2(0, T; \mathbb{R}^{n \times n}), \quad D(\cdot), \bar{D}(\cdot) \in L^\infty(0, T; \mathbb{R}^{n \times m}), \quad \sigma(\cdot) \in L^2_{\mathbb{F}}(0, T; \mathbb{R}^n).
 \end{cases}$$

(H2) The weighting coefficients in the cost functional satisfy the following:

$$\begin{cases}
 Q(\cdot), \bar{Q}(\cdot) \in L^1(0, T; \mathbb{S}^n), \quad S(\cdot), \bar{S}(\cdot) \in L^2(0, T; \mathbb{R}^{m \times n}), \quad R(\cdot), \bar{R}(\cdot) \in L^\infty(0, T; \mathbb{S}^m), \\
 g \in L^2_{\mathcal{F}_T}(\Omega; \mathbb{R}^n), \quad q(\cdot) \in L^2_{\mathbb{F}}(\Omega; L^1(0, T; \mathbb{R}^n)), \quad \rho(\cdot) \in L^2_{\mathbb{F}}(0, T; \mathbb{R}^m), \\
 G, \bar{G} \in \mathbb{S}^n, \quad \bar{g} \in \mathbb{R}^n, \quad \bar{q}(\cdot) \in L^1(0, T; \mathbb{R}^n), \quad \bar{\rho}(\cdot) \in L^2(0, T; \mathbb{R}^m).
 \end{cases}$$

By a standard argument using contraction mapping theorem, one can show that under (H1), for any $(t, \xi) \in [0, T] \times L^2_{\mathcal{F}_t}(\Omega; \mathbb{R}^n)$ and any $u(\cdot) \in \mathcal{U}[t, T]$, (1.1) admits a unique solution $X(\cdot) \equiv X(\cdot; t, \xi, u(\cdot)) \in L^2_{\mathbb{F}}(\Omega; C([t, T]; \mathbb{R}^n))$. Hence, under (H1)–(H2), the cost functional (1.2) is well-defined, and Problem (MF-LQ) makes sense. Now we introduce the following definition.

Definition 2.1.

(i) Problem (MF-LQ) is said to be *finite at initial pair* $(t, \xi) \in [0, T] \times L^2_{\mathcal{F}_t}(\Omega; \mathbb{R}^n)$ if

$$V(t, \xi) > -\infty. \tag{2.1}$$

Problem (MF-LQ) is said to be *finite at* $t \in [0, T]$ if (2.1) holds for all $\xi \in L^2_{\mathcal{F}_t}(\Omega; \mathbb{R}^n)$, and Problem (MF-LQ) is said to be *finite* if it is finite at all $t \in [0, T]$.

(ii) Problem (MF-LQ) is said to be *(uniquely) open-loop solvable at initial pair* $(t, \xi) \in [0, T] \times L^2_{\mathcal{F}_t}(\Omega; \mathbb{R}^n)$ if there exists a (unique) $u^*(\cdot) \in \mathcal{U}[t, T]$ satisfying (1.3). Problem (MF-LQ) is said to be *(uniquely) open-loop solvable at* t if for any $\xi \in L^2_{\mathcal{F}_t}(\Omega; \mathbb{R}^n)$, there exists a (unique) $u^*(\cdot) \in \mathcal{U}[t, T]$ satisfying (1.3), and Problem (MF-LQ) is said to be *(uniquely) open-loop solvable (on $[0, T])$* if it is (uniquely) open-loop solvable at all $t \in [0, T]$.

Next, we introduce the following mean-field backward SDE (MF-BSDE, for short) associated with the state process $X(\cdot) \equiv X(\cdot; t, \xi, u(\cdot))$:

$$\begin{cases} dY(s) = -\left\{A^\top Y + \bar{A}^\top \mathbb{E}[Y] + C^\top Z + \bar{C}^\top \mathbb{E}[Z] + QX + \bar{Q}\mathbb{E}[X] \right. \\ \quad \left. + S^\top u + \bar{S}^\top \mathbb{E}[u] + q + \bar{q}\right\} ds + Z dW(s), & s \in [t, T], \\ Y(T) = GX(T) + \bar{G}\mathbb{E}[X(T)] + g + \bar{g}. \end{cases} \quad (2.2)$$

The following result is concerned with the differentiability of the map $u(\cdot) \mapsto J(t, \xi; u(\cdot))$.

Proposition 2.2. *Let (H1)–(H2) hold and $t \in [0, T]$ be given. For any $\xi \in L^2_{\mathcal{F}_t}(\Omega; \mathbb{R}^n)$, $\lambda \in \mathbb{R}$ and $u(\cdot), v(\cdot) \in \mathcal{U}[t, T]$, the following holds:*

$$\begin{aligned} J(t, \xi; u(\cdot) + \lambda v(\cdot)) - J(t, \xi; u(\cdot)) &= \lambda^2 J^0(t, 0; v(\cdot)) + 2\lambda \mathbb{E} \int_t^T \langle B^\top Y + \bar{B}^\top \mathbb{E}[Y] + D^\top Z + \bar{D}^\top \mathbb{E}[Z] \\ &\quad + SX + \bar{S}\mathbb{E}[X] + Ru + \bar{R}\mathbb{E}[u] + \rho + \bar{\rho}, v \rangle ds, \end{aligned} \quad (2.3)$$

where $X(\cdot) = X(\cdot; t, \xi, u(\cdot))$ and $(Y(\cdot), Z(\cdot))$ is the adapted solution to the MF-BSDE (2.2) associated with $X(\cdot)$. Consequently, the map $u(\cdot) \mapsto J(t, \xi; u(\cdot))$ is Fréchet differentiable with the Fréchet derivative given by

$$\begin{aligned} \mathcal{D}J(t, \xi; u(\cdot))(s) &= 2\left\{B(s)^\top Y(s) + \bar{B}(s)^\top \mathbb{E}[Y(s)] + D(s)^\top Z(s) + \bar{D}(s)^\top \mathbb{E}[Z(s)] + S(s)X(s) \right. \\ &\quad \left. + \bar{S}(s)\mathbb{E}[X(s)] + R(s)u(s) + \bar{R}(s)\mathbb{E}[u(s)] + \rho(s) + \bar{\rho}(s)\right\}, \quad s \in [t, T]. \end{aligned} \quad (2.4)$$

Proof. Let $\hat{X}(\cdot) = X(\cdot; t, \xi, u(\cdot) + \lambda v(\cdot))$ and $X_0(\cdot)$ be the solution to the following MF-SDE:

$$\begin{cases} dX_0(s) = \left\{A(s)X_0(s) + \bar{A}(s)\mathbb{E}[X_0(s)] + B(s)v(s) + \bar{B}(s)\mathbb{E}[v(s)]\right\} ds \\ \quad + \left\{C(s)X_0(s) + \bar{C}(s)\mathbb{E}[X_0(s)] + D(s)v(s) + \bar{D}(s)\mathbb{E}[v(s)]\right\} dW(s), & s \in [t, T], \\ X_0(t) = 0, \end{cases}$$

By the linearity of the state equation, $\hat{X}(\cdot) = X(\cdot) + \lambda X_0(\cdot)$. Hence,

$$\begin{aligned} &J(t, \xi; u(\cdot) + \lambda v(\cdot)) - J(t, \xi; u(\cdot)) \\ &= \lambda \mathbb{E} \left\{ \langle G[2X(T) + \lambda X_0(T)], X_0(T) \rangle + 2\langle g, X_0(T) \rangle \right. \\ &\quad \left. + \int_t^T \left[\left\langle \begin{pmatrix} Q & S^\top \\ \bar{S} & \bar{R} \end{pmatrix} \begin{pmatrix} 2X + \lambda X_0 \\ 2u + \lambda v \end{pmatrix}, \begin{pmatrix} X_0 \\ v \end{pmatrix} \right\rangle + 2 \left\langle \begin{pmatrix} q \\ \rho \end{pmatrix}, \begin{pmatrix} X_0 \\ v \end{pmatrix} \right\rangle \right] ds \right\} \\ &\quad + \lambda \left\{ \langle \bar{G}(2\mathbb{E}[X(T)] + \lambda \mathbb{E}[X_0(T)]), \mathbb{E}[X_0(T)] \rangle + 2\langle \bar{g}, \mathbb{E}[X_0(T)] \rangle \right. \\ &\quad \left. + \int_t^T \left[\left\langle \begin{pmatrix} \bar{Q} & \bar{S}^\top \\ \bar{S} & \bar{R} \end{pmatrix} \begin{pmatrix} 2\mathbb{E}[X] + \lambda \mathbb{E}[X_0] \\ 2\mathbb{E}[u] + \lambda \mathbb{E}[v] \end{pmatrix}, \begin{pmatrix} \mathbb{E}[X_0] \\ \mathbb{E}[v] \end{pmatrix} \right\rangle + 2 \left\langle \begin{pmatrix} \bar{q} \\ \bar{\rho} \end{pmatrix}, \begin{pmatrix} \mathbb{E}[X_0] \\ \mathbb{E}[v] \end{pmatrix} \right\rangle \right] ds \right\} \\ &= 2\lambda \mathbb{E} \left\{ \langle GX(T) + g, X_0(T) \rangle + \int_t^T \left[\langle QX + S^\top u + q, X_0 \rangle + \langle SX + Ru + \rho, v \rangle \right] ds \right\} \end{aligned}$$

$$\begin{aligned}
 & + \lambda^2 \mathbb{E} \left\{ \langle GX_0(T), X_0(T) \rangle + \int_t^T \left\langle \begin{pmatrix} Q & S^\top \\ \bar{S} & R \end{pmatrix} \begin{pmatrix} X_0 \\ v \end{pmatrix}, \begin{pmatrix} X_0 \\ v \end{pmatrix} \right\rangle ds \right\} \\
 & + 2\lambda \left\{ \langle \bar{G}\mathbb{E}[X(T)] + \bar{g}, \mathbb{E}[X_0(T)] \rangle + \int_t^T \left[\langle \bar{Q}\mathbb{E}[X] + \bar{S}^\top \mathbb{E}[u] + \bar{q}, \mathbb{E}[X_0] \rangle + \langle \bar{S}\mathbb{E}[X] + \bar{R}\mathbb{E}[u] + \bar{\rho}, \mathbb{E}[v] \rangle \right] ds \right\} \\
 & + \lambda^2 \left\{ \langle \bar{G}\mathbb{E}[X_0(T)], \mathbb{E}[X_0(T)] \rangle + \int_t^T \left\langle \begin{pmatrix} \bar{Q} & \bar{S}^\top \\ \bar{S} & \bar{R} \end{pmatrix} \begin{pmatrix} \mathbb{E}[X_0] \\ \mathbb{E}[v] \end{pmatrix}, \begin{pmatrix} \mathbb{E}[X_0] \\ \mathbb{E}[v] \end{pmatrix} \right\rangle ds \right\} \\
 & = 2\lambda \mathbb{E} \left\{ \langle GX(T) + \bar{G}\mathbb{E}[X(T)] + g + \bar{g}, X_0(T) \rangle \right. \\
 & \quad \left. + \int_t^T \left[\langle QX + \bar{Q}\mathbb{E}[X] + S^\top u + \bar{S}^\top \mathbb{E}[u] + q + \bar{q}, X_0 \rangle + \langle SX + \bar{S}\mathbb{E}[X] + Ru + \bar{R}\mathbb{E}[u] + \rho + \bar{\rho}, v \rangle \right] ds \right\} \\
 & \quad + \lambda^2 J^0(t, 0; v(\cdot)).
 \end{aligned}$$

Now applying Itô's formula to $s \mapsto \langle Y(s), X_0(s) \rangle$, we have

$$\begin{aligned}
 & \mathbb{E} \langle GX(T) + \bar{G}\mathbb{E}[X(T)] + g + \bar{g}, X_0(T) \rangle \\
 & = \mathbb{E} \int_t^T \left\{ - \langle A^\top Y + \bar{A}^\top \mathbb{E}[Y] + C^\top Z + \bar{C}^\top \mathbb{E}[Z] + QX + \bar{Q}\mathbb{E}[X] + S^\top u + \bar{S}^\top \mathbb{E}[u] + q + \bar{q}, X_0 \rangle \right. \\
 & \quad \left. + \langle AX_0 + \bar{A}\mathbb{E}[X_0] + Bv + \bar{B}\mathbb{E}[v], Y \rangle + \langle CX_0 + \bar{C}\mathbb{E}[X_0] + Dv + \bar{D}\mathbb{E}[v], Z \rangle \right\} ds \\
 & = \mathbb{E} \int_t^T \left\{ \langle B^\top Y + \bar{B}^\top \mathbb{E}[Y] + D^\top Z + \bar{D}^\top \mathbb{E}[Z], v \rangle - \langle QX + \bar{Q}\mathbb{E}[X] + S^\top u + \bar{S}^\top \mathbb{E}[u] + q + \bar{q}, X_0 \rangle \right\} ds.
 \end{aligned}$$

Combining the above equalities, we obtain (2.3). □

From the above, we have the following result, which gives a characterization for the optimal controls of Problem (MF-LQ).

Theorem 2.3. *Let (H1)–(H2) hold and $(t, \xi) \in [0, T] \times L^2_{\mathcal{F}_t}(\Omega; \mathbb{R}^n)$ be given. Let $u^*(\cdot) \in \mathcal{U}[t, T]$ and $(X^*(\cdot), Y^*(\cdot), Z^*(\cdot))$ be the adapted solution to the following (decoupled) mean-field forward-backward stochastic differential equation (MF-FBSDE, for short):*

$$\begin{cases}
 dX^*(s) = \left\{ AX^* + \bar{A}\mathbb{E}[X^*] + Bu^* + \bar{B}\mathbb{E}[u^*] + b \right\} ds \\
 \quad + \left\{ CX^* + \bar{C}\mathbb{E}[X^*] + Du^* + \bar{D}\mathbb{E}[u^*] + \sigma \right\} dW(s), & s \in [t, T], \\
 dY^*(s) = - \left\{ A^\top Y^* + \bar{A}^\top \mathbb{E}[Y^*] + C^\top Z^* + \bar{C}^\top \mathbb{E}[Z^*] + QX^* + \bar{Q}\mathbb{E}[X^*] \right. \\
 \quad \left. + S^\top u^* + \bar{S}^\top \mathbb{E}[u^*] + q + \bar{q} \right\} ds + Z^* dW(s), & s \in [t, T], \\
 X^*(t) = \xi, & Y^*(T) = GX^*(T) + \bar{G}\mathbb{E}[X^*(T)] + g + \bar{g}.
 \end{cases} \tag{2.5}$$

Then $u^*(\cdot)$ is an optimal control of Problem (MF-LQ) for the initial pair (t, ξ) if and only if

$$J^0(t, 0; u(\cdot)) \geq 0, \quad \forall u(\cdot) \in \mathcal{U}[t, T], \tag{2.6}$$

and the following stationarity condition holds:

$$\begin{aligned}
 DJ(t, \xi; u^*(\cdot)) & = 2 \left\{ B^\top Y^* + D^\top Z^* + SX^* + Ru^* + \rho \right. \\
 & \quad \left. + \bar{B}^\top \mathbb{E}[Y^*] + \bar{D}^\top \mathbb{E}[Z^*] + \bar{S}\mathbb{E}[X^*] + \bar{R}\mathbb{E}[u^*] + \bar{\rho} \right\} = 0, \quad \text{a.e. a.s.}
 \end{aligned} \tag{2.7}$$

Proof. By (2.3), we see that $u^*(\cdot)$ is an optimal control of Problem (MF-LQ) for the initial pair (t, ξ) if and only if

$$\begin{aligned} & \lambda^2 J^0(t, 0; u(\cdot)) + \lambda \mathbb{E} \int_t^T \langle \mathcal{D}J(t, \xi; u^*(\cdot))(s), u(s) \rangle ds \\ & = J(t, \xi; u^*(\cdot) + \lambda u(\cdot)) - J(t, \xi; u^*(\cdot)) \geq 0, \quad \forall \lambda \in \mathbb{R}, \forall u(\cdot) \in \mathcal{U}[t, T], \end{aligned}$$

which is equivalent to (2.6) and the following:

$$\mathbb{E} \int_t^T \langle \mathcal{D}J(t, \xi; u^*(\cdot))(s), u(s) \rangle ds \leq 0, \quad \forall u(\cdot) \in \mathcal{U}[t, T].$$

Note that the above inequality holds for all $u(\cdot) \in \mathcal{U}[t, T]$ if and only if $\mathcal{D}J(t, \xi; u^*(\cdot))(\cdot) = 0$. The result therefore follows. \square

3. FINITENESS AND OPEN-LOOP SOLVABILITY OF PROBLEM (MF-LQ)

We begin with a representation of the cost functional. For any $u(\cdot) \in \mathcal{U}[t, T]$, let $X_0^u(\cdot)$ be the solution of

$$\begin{cases} dX_0^u(s) = \left\{ A(s)X_0^u(s) + \bar{A}(s)\mathbb{E}[X_0^u(s)] + B(s)u(s) + \bar{B}(s)\mathbb{E}[u(s)] \right\} ds \\ \quad + \left\{ C(s)X_0^u(s) + \bar{C}(s)\mathbb{E}[X_0^u(s)] + D(s)u(s) + \bar{D}(s)\mathbb{E}[u(s)] \right\} dW(s), & s \in [t, T], \\ X_0^u(t) = 0. \end{cases} \quad (3.1)$$

By the linearity of (3.1), we can define bounded linear operators $\mathcal{L}_t : \mathcal{U}[t, T] \rightarrow L_{\mathbb{F}}^2(t, T; \mathbb{R}^n)$ and $\widehat{\mathcal{L}}_t : \mathcal{U}[t, T] \rightarrow L_{\mathcal{F}_T}^2(\Omega; \mathbb{R}^n)$ by $u(\cdot) \mapsto X_0^u(\cdot)$ and $u(\cdot) \mapsto X_0^u(T)$, respectively, via the MF-SDE (3.1). Then

$$\begin{aligned} J^0(t, 0; u(\cdot)) &= \mathbb{E} \left\{ \langle GX_0^u(T), X_0^u(T) \rangle + \langle \bar{G}\mathbb{E}[X_0^u(T)], \mathbb{E}[X_0^u(T)] \rangle \right. \\ & \quad + \int_t^T \left\langle \begin{pmatrix} Q(s) & S(s)^\top \\ S(s) & R(s) \end{pmatrix} \begin{pmatrix} X_0^u(s) \\ u(s) \end{pmatrix}, \begin{pmatrix} X_0^u(s) \\ u(s) \end{pmatrix} \right\rangle ds \\ & \quad \left. + \int_t^T \left\langle \begin{pmatrix} \bar{Q}(s) & \bar{S}(s)^\top \\ \bar{S}(s) & \bar{R}(s) \end{pmatrix} \begin{pmatrix} \mathbb{E}[X_0^u(s)] \\ \mathbb{E}[u(s)] \end{pmatrix}, \begin{pmatrix} \mathbb{E}[X_0^u(s)] \\ \mathbb{E}[u(s)] \end{pmatrix} \right\rangle ds \right\} \\ &= \langle G\widehat{\mathcal{L}}_t u, \widehat{\mathcal{L}}_t u \rangle + \langle \bar{G}\mathbb{E}[\widehat{\mathcal{L}}_t u], \mathbb{E}[\widehat{\mathcal{L}}_t u] \rangle + \langle Q\mathcal{L}_t u, \mathcal{L}_t u \rangle + 2\langle S\mathcal{L}_t u, u \rangle + \langle Ru, u \rangle \\ & \quad + \langle \bar{Q}\mathbb{E}[\mathcal{L}_t u], \mathbb{E}[\mathcal{L}_t u] \rangle + 2\langle \bar{S}\mathbb{E}[\mathcal{L}_t u], \mathbb{E}[u] \rangle + \langle \bar{R}\mathbb{E}[u], \mathbb{E}[u] \rangle \\ &= \langle [\widehat{\mathcal{L}}_t^*(G + \mathbb{E}^*\bar{G}\mathbb{E})\widehat{\mathcal{L}}_t + \mathcal{L}_t^*(Q + \mathbb{E}^*\bar{Q}\mathbb{E})\mathcal{L}_t + (S + \mathbb{E}^*\bar{S}\mathbb{E})\mathcal{L}_t + \mathcal{L}_t^*(S^\top + \mathbb{E}^*\bar{S}^\top\mathbb{E}) + (R + \mathbb{E}^*\bar{R}\mathbb{E})]u, u \rangle. \end{aligned}$$

Denote

$$\mathcal{M}_t \triangleq \widehat{\mathcal{L}}_t^*(G + \mathbb{E}^*\bar{G}\mathbb{E})\widehat{\mathcal{L}}_t + \mathcal{L}_t^*(Q + \mathbb{E}^*\bar{Q}\mathbb{E})\mathcal{L}_t + (S + \mathbb{E}^*\bar{S}\mathbb{E})\mathcal{L}_t + \mathcal{L}_t^*(S^\top + \mathbb{E}^*\bar{S}^\top\mathbb{E}) + (R + \mathbb{E}^*\bar{R}\mathbb{E}), \quad (3.2)$$

which is a bounded self-adjoint linear operator on $\mathcal{U}[t, T]$. Then by Proposition 2.2, the cost functional $J(t, \xi; u(\cdot))$ can be written as

$$\begin{aligned} J(t, \xi; u(\cdot)) &= \langle \mathcal{M}_t u, u \rangle + \langle \mathcal{D}J(t, \xi; 0), u \rangle + J(t, \xi; 0), \\ & \quad \forall (t, \xi) \in [0, T] \times L_{\mathcal{F}_t}^2(\Omega; \mathbb{R}^n), \forall u(\cdot) \in \mathcal{U}[t, T]. \end{aligned} \quad (3.3)$$

Now let us introduce the following conditions.

(H3) The following holds:

$$J^0(t, 0; u(\cdot)) \geq 0, \quad \forall u(\cdot) \in \mathcal{U}[t, T]. \quad (3.4)$$

(H4) There exists a constant $\delta > 0$ such that

$$J^0(t, 0; u(\cdot)) \geq \delta \mathbb{E} \int_t^T |u(s)|^2 ds, \quad \forall u(\cdot) \in \mathcal{U}[t, T]. \tag{3.5}$$

From (3.3), we see that the map $u(\cdot) \mapsto J(t, \xi; u(\cdot))$ is convex if and only if

$$\mathcal{M}_t \geq 0, \tag{3.6}$$

which is also equivalent to (H3), and $u(\cdot) \mapsto J(t, \xi; u(\cdot))$ is uniformly convex if and only if

$$\mathcal{M}_t \geq \delta I, \quad \text{for some } \delta > 0, \tag{3.7}$$

which is also equivalent to (H4). The following result tells us that (H3) is necessary for the finiteness (and open-loop solvability) of Problem (MF-LQ) at t , and (H4) is sufficient for the open-loop solvability of Problem (MF-LQ) at t .

Proposition 3.1. *Let (H1)–(H2) hold and $t \in [0, T]$ be given. We have the following:*

- (i) *If Problem (MF-LQ) is finite at t , then (H3) must hold.*
- (ii) *Suppose (H4) holds. Then Problem (MF-LQ) is uniquely open-loop solvable at t , and the unique optimal control for the initial pair (t, ξ) is given by*

$$u^*(\cdot) = -\frac{1}{2} \mathcal{M}_t^{-1} \mathcal{D}J(t, \xi; 0)(\cdot). \tag{3.8}$$

Moreover,

$$V(t, \xi) = J(t, \xi; 0) - \frac{1}{4} \left| \mathcal{M}_t^{-\frac{1}{2}} \mathcal{D}J(t, \xi; 0) \right|^2. \tag{3.9}$$

Proof.

- (i) We prove the result by contradiction. Suppose that $J^0(t, 0; u(\cdot)) < 0$ for some $u(\cdot) \in \mathcal{U}[t, T]$. By Proposition 2.2, we have

$$J(t, \xi; \lambda u(\cdot)) = J(t, \xi; 0) + \lambda^2 J^0(t, 0; u(\cdot)) + \lambda \mathbb{E} \int_t^T \langle \mathcal{D}J(t, \xi; 0)(s), u(s) \rangle ds, \quad \forall \lambda \in \mathbb{R}.$$

Letting $\lambda \rightarrow \infty$, we obtain that

$$V(t, \xi) \leq \lim_{\lambda \rightarrow \infty} J(t, \xi; \lambda u(\cdot)) = -\infty,$$

which is a contradiction.

- (ii) Suppose (H4) holds. Then the operator \mathcal{M}_t is invertible, and

$$\begin{aligned} J(t, \xi; u(\cdot)) &= \left| \mathcal{M}_t^{\frac{1}{2}} u + \frac{1}{2} \mathcal{M}_t^{-\frac{1}{2}} \mathcal{D}J(t, \xi; 0) \right|^2 + J(t, \xi; 0) - \frac{1}{4} \left| \mathcal{M}_t^{-\frac{1}{2}} \mathcal{D}J(t, \xi; 0) \right|^2, \\ &\geq J(t, \xi; 0) - \frac{1}{4} \left| \mathcal{M}_t^{-\frac{1}{2}} \mathcal{D}J(t, \xi; 0) \right|^2, \quad \forall \xi \in L^2_{\mathcal{F}_t}(\Omega; \mathbb{R}^n), \forall u(\cdot) \in \mathcal{U}[t, T]. \end{aligned}$$

Note that the equality in the above holds if and only if

$$u = -\frac{1}{2} \mathcal{M}_t^{-1} \mathcal{D}J(t, \xi; 0).$$

The result therefore follows. □

Due to the necessity of (H3) for the finiteness of Problem (MF-LQ), we will assume (H3) holds in the rest of this paper. Now for any $\varepsilon > 0$, consider state equation (1.1) and the following cost functional:

$$\begin{aligned} J_\varepsilon(t, \xi; u(\cdot)) &\triangleq J(t, \xi; u(\cdot)) + \varepsilon \mathbb{E} \int_t^T |u(s)|^2 ds \\ &= \langle (\mathcal{M}_t + \varepsilon I)u, u \rangle + \langle \mathcal{D}J(t, \xi; 0), u \rangle + J(t, \xi; 0). \end{aligned} \quad (3.10)$$

Denote the corresponding optimal control problem and value function by Problem (MF-LQ) $_\varepsilon$ and $V_\varepsilon(\cdot, \cdot)$, respectively. By Proposition 3.1, part (ii), for any $\xi \in L^2_{\mathcal{F}_t}(\Omega; \mathbb{R}^n)$, Problem (MF-LQ) $_\varepsilon$ admits a unique optimal control

$$u_\varepsilon^*(\cdot) = -\frac{1}{2}(\mathcal{M}_t + \varepsilon I)^{-1} \mathcal{D}J(t, \xi; 0)(\cdot), \quad (3.11)$$

and the value function is given by

$$V_\varepsilon(t, \xi) = J(t, \xi; 0) - \frac{1}{4} \left| (\mathcal{M}_t + \varepsilon I)^{-\frac{1}{2}} \mathcal{D}J(t, \xi; 0) \right|^2. \quad (3.12)$$

Now, we are ready to state the main result of this section.

Theorem 3.2. *Let (H1)–(H3) hold and $\xi \in L^2_{\mathcal{F}_t}(\Omega; \mathbb{R}^n)$. We have the following:*

- (i) $\lim_{\varepsilon \rightarrow 0} V_\varepsilon(t, \xi) = V(t, \xi)$. In particular, Problem (MF-LQ) is finite at (t, ξ) if and only if $\{V_\varepsilon(t, \xi)\}_{\varepsilon > 0}$ is bounded from below.
- (ii) The sequence $\{u_\varepsilon^*(\cdot)\}_{\varepsilon > 0}$ defined by (3.11) is a minimizing sequence of $u(\cdot) \mapsto J(t, \xi; u(\cdot))$:

$$\lim_{\varepsilon \rightarrow 0} J(t, \xi; u_\varepsilon^*(\cdot)) = \inf_{u(\cdot) \in \mathcal{U}[t, T]} J(t, \xi; u(\cdot)) = V(t, \xi). \quad (3.13)$$

(iii) The following statements are equivalent:

- (a) Problem (MF-LQ) is open-loop solvable at (t, ξ) ;
- (b) The sequence $\{u_\varepsilon^*(\cdot)\}_{\varepsilon > 0}$ is bounded in $\mathcal{U}[t, T]$;
- (c) The sequence $\{u_\varepsilon^*(\cdot)\}_{\varepsilon > 0}$ admits a weakly convergent subsequence;
- (d) The sequence $\{u_\varepsilon^*(\cdot)\}_{\varepsilon > 0}$ admits a strongly convergent subsequence.

In this case, the weak (strong) limit of any weakly (strongly) convergent subsequence of $\{u_\varepsilon^*(\cdot)\}_{\varepsilon > 0}$ is an optimal control of Problem (MF-LQ) at (t, ξ) .

To prove Theorem 3.2, we need the following lemma.

Lemma 3.3. *Let \mathcal{H} be a Hilbert space with norm $|\cdot|$ and $\theta, \theta_n \in \mathcal{H}$, $n = 1, 2, \dots$*

- (i) If $\theta_n \rightarrow \theta$ weakly, then $|\theta| \leq \liminf_{n \rightarrow \infty} |\theta_n|$.
- (ii) $\theta_n \rightarrow \theta$ strongly if and only if

$$|\theta_n| \rightarrow |\theta| \quad \text{and} \quad \theta_n \rightarrow \theta \quad \text{weakly.}$$

Proof of Theorem 3.2.

(i) For any $\varepsilon_2 > \varepsilon_1 > 0$, we have

$$J_{\varepsilon_2}(t, \xi; u(\cdot)) \geq J_{\varepsilon_1}(t, \xi; u(\cdot)) \geq J(t, \xi; u(\cdot)), \quad \forall u(\cdot) \in \mathcal{U}[t, T],$$

which implies that

$$V_{\varepsilon_2}(t, \xi) \geq V_{\varepsilon_1}(t, \xi) \geq V(t, \xi), \quad \forall \varepsilon_2 > \varepsilon_1 > 0.$$

Thus, the limit $\lim_{\varepsilon \rightarrow 0} V_\varepsilon(t, \xi)$ exists and

$$\bar{V}(t, \xi) \equiv \lim_{\varepsilon \rightarrow 0} V_\varepsilon(t, \xi) \geq V(t, \xi). \tag{3.14}$$

On the other hand, for any $K, \delta > 0$, we can find a $u^\delta(\cdot) \in \mathcal{U}[t, T]$, such that

$$V_\varepsilon(t, \xi) \leq J(t, \xi; u^\delta(\cdot)) + \varepsilon \mathbb{E} \int_t^T |u^\delta(s)|^2 ds \leq \max\{V(t, \xi), -K\} + \delta + \varepsilon \mathbb{E} \int_t^T |u^\delta(s)|^2 ds.$$

Letting $\varepsilon \rightarrow 0$, we obtain

$$\bar{V}(t, \xi) \leq \max\{V(t, \xi), -K\} + \delta, \quad \forall K, \delta > 0,$$

from which we see

$$\bar{V}(t, \xi) \leq V(t, \xi). \tag{3.15}$$

Combining (3.14)–(3.15), we obtain the desired result.

(ii) If $V(t, \xi) > -\infty$, then by (i), we have

$$\begin{aligned} \varepsilon \mathbb{E} \int_t^T |u_\varepsilon^*(s)|^2 ds &= J_\varepsilon(t, \xi; u_\varepsilon^*(\cdot)) - J(t, \xi; u_\varepsilon^*(\cdot)) = V_\varepsilon(t, \xi) - J(t, \xi; u_\varepsilon^*(\cdot)) \\ &\leq V_\varepsilon(t, \xi) - V(t, \xi) \rightarrow 0 \quad \text{as } \varepsilon \rightarrow 0. \end{aligned}$$

Hence,

$$\lim_{\varepsilon \rightarrow 0} J(t, \xi; u_\varepsilon^*(\cdot)) = \lim_{\varepsilon \rightarrow 0} \left[V_\varepsilon(t, \xi) - \varepsilon \mathbb{E} \int_t^T |u_\varepsilon^*(s)|^2 ds \right] = V(t, \xi).$$

If $V(t, \xi) = -\infty$, then by (i), we have

$$J(t, \xi; u_\varepsilon^*(\cdot)) \leq J_\varepsilon(t, \xi; u_\varepsilon^*(\cdot)) = V_\varepsilon(t, \xi) \rightarrow -\infty \quad \text{as } \varepsilon \rightarrow 0,$$

and (3.13) still holds.

(iii) (b) \Rightarrow (c) and (d) \Rightarrow (c) are obvious. We next prove (c) \Rightarrow (a). Let $\{u_{\varepsilon_k}^*(\cdot)\}_{k \geq 1}$ be a weakly convergent subsequence of $\{u_\varepsilon^*(\cdot)\}_{\varepsilon > 0}$ with weak limit $u^*(\cdot)$. Then $\{u_{\varepsilon_k}^*(\cdot)\}_{k \geq 1}$ is bounded in $\mathcal{U}[t, T]$. For any $u(\cdot) \in \mathcal{U}[t, T]$, we have

$$J(t, \xi; u_{\varepsilon_k}^*(\cdot)) + \varepsilon_k \mathbb{E} \int_t^T |u_{\varepsilon_k}^*(s)|^2 ds = V_{\varepsilon_k}(t, \xi) \leq J(t, \xi; u(\cdot)) + \varepsilon_k \mathbb{E} \int_t^T |u(s)|^2 ds. \tag{3.16}$$

Note that $u(\cdot) \mapsto J(t, \xi; u(\cdot))$ is sequentially weakly lower semi-continuous. Letting $k \rightarrow \infty$ in (3.16), we obtain

$$J(t, \xi; u^*(\cdot)) \leq \liminf_{k \rightarrow \infty} J(t, \xi; u_{\varepsilon_k}^*(\cdot)) \leq J(t, \xi; u(\cdot)), \quad \forall u(\cdot) \in \mathcal{U}[t, T].$$

Hence, $u^*(\cdot)$ is an optimal control of Problem (MF-LQ) at (t, ξ) . Now it remains to show (a) \Rightarrow (b) and (a) \Rightarrow (d). Suppose $v^*(\cdot)$ is an optimal control of Problem (MF-LQ) at (t, ξ) . Then for any $\varepsilon > 0$, we have

$$\begin{cases} V_\varepsilon(t, \xi) = J_\varepsilon(t, \xi; u_\varepsilon^*(\cdot)) \geq V(t, \xi) + \varepsilon \mathbb{E} \int_t^T |u_\varepsilon^*(s)|^2 ds, \\ V_\varepsilon(t, \xi) \leq J_\varepsilon(t, \xi; v^*(\cdot)) = V(t, \xi) + \varepsilon \mathbb{E} \int_t^T |v^*(s)|^2 ds, \end{cases}$$

from which we see that

$$\mathbb{E} \int_t^T |u_\varepsilon^*(s)|^2 ds \leq \frac{V_\varepsilon(t, \xi) - V(t, \xi)}{\varepsilon} \leq \mathbb{E} \int_t^T |v^*(s)|^2 ds, \quad \forall \varepsilon > 0. \tag{3.17}$$

Thus, $\{u_\varepsilon^*(\cdot)\}_{\varepsilon>0}$ is bounded in the Hilbert space $\mathcal{U}[t, T]$ and hence admits a weakly convergent subsequence $\{u_{\varepsilon_k}^*(\cdot)\}_{k\geq 1}$. Let $u^*(\cdot)$ be the weak limit of $\{u_{\varepsilon_k}^*(\cdot)\}_{k\geq 1}$. By the proof of (c) \Rightarrow (a), we see that $u^*(\cdot)$ is also an optimal control of Problem (MF-LQ) at (t, ξ) . Replacing $v^*(\cdot)$ with $u^*(\cdot)$ in (3.17), we have

$$\mathbb{E} \int_t^T |u_\varepsilon^*(s)|^2 ds \leq \mathbb{E} \int_t^T |u^*(s)|^2 ds, \quad \forall \varepsilon > 0. \quad (3.18)$$

Also, by Lemma 3.3, part (i),

$$\mathbb{E} \int_t^T |u^*(s)|^2 ds \leq \liminf_{k \rightarrow \infty} \mathbb{E} \int_t^T |u_{\varepsilon_k}^*(s)|^2 ds. \quad (3.19)$$

Combining (3.18)–(3.19), we have

$$\mathbb{E} \int_t^T |u^*(s)|^2 ds = \lim_{k \rightarrow \infty} \mathbb{E} \int_t^T |u_{\varepsilon_k}^*(s)|^2 ds.$$

Then it follows from Lemma 3.3, part (ii), that $\{u_{\varepsilon_k}^*(\cdot)\}_{k\geq 1}$ converges to $u^*(\cdot)$ strongly. \square

4. NECESSARY CONDITIONS FOR THE UNIFORM CONVEXITY AND RICCATI EQUATIONS

Theorem 3.2 tells us that in order to solve Problem (MF-LQ), we need only solve mean-field LQ problems with uniformly convex cost functionals and then pass to the limit. By Proposition 3.1, under the uniform convexity condition (H4), the unique optimal control $u^*(\cdot)$ for the initial pair (t, ξ) is determined by (3.8). However, such a representation is not easy to compute, since \mathcal{M}_t^{-1} is in an abstract form and very complicated. Thus, we would like to find some more explicit form of the optimal control. In this section we shall investigate uniform convexity of the cost functional and show the necessity of solvability of two Riccati equations for the uniform convexity of the cost functional.

First, we present the following result concerning the value function of Problem (MF-LQ)⁰.

Proposition 4.1. *Let (H1), (H2) and (H4) hold. Then there exists a constant $\alpha \in \mathbb{R}$ such that*

$$V^0(s, \xi) \geq \alpha \mathbb{E}[|\xi|^2], \quad \forall (s, \xi) \in [t, T] \times L_{\mathcal{F}_s}^2(\Omega; \mathbb{R}^n) \text{ with } \mathbb{E}[\xi] = 0. \quad (4.1)$$

Proof. For any $s \in [t, T]$ and any $u(\cdot) \in \mathcal{U}[s, T]$, we define the zero-extension of $u(\cdot)$ as follows:

$$[0I_{[t,s]} \oplus u(\cdot)](r) = \begin{cases} 0, & r \in [t, s), \\ u(r), & r \in [s, T]. \end{cases} \quad (4.2)$$

Then $v(\cdot) \equiv 0I_{[t,s]} \oplus u(\cdot) \in \mathcal{U}[t, T]$, and due to the initial state being 0, the solution $X_0^v(\cdot)$ of

$$\begin{cases} dX_0^v(r) = \{A(r)X_0^v(r) + \bar{A}(r)\mathbb{E}[X_0^v(r)] + B(r)v(r) + \bar{B}(r)\mathbb{E}[v(r)]\} dr \\ \quad + \{C(r)X_0^v(r) + \bar{C}(r)\mathbb{E}[X_0^v(r)] + D(r)v(r) + \bar{D}(r)\mathbb{E}[v(r)]\} dW(r), & r \in [t, T], \\ X_0^v(t) = 0, \end{cases}$$

satisfies $X_0^v(r) = 0$, $r \in [t, s]$. Hence,

$$J^0(s, 0; u(\cdot)) = J^0(t, 0; 0I_{[t,s]} \oplus u(\cdot)) \geq \delta \mathbb{E} \int_t^T |[0I_{[t,s]} \oplus u(\cdot)](r)|^2 dr = \delta \mathbb{E} \int_s^T |u(r)|^2 dr. \quad (4.3)$$

Now, let $(X(\cdot), Y(\cdot), Z(\cdot))$ be the solution of the following (decoupled) MF-FBSDE:

$$\begin{cases} dX(r) = \{AX + \bar{A}\mathbb{E}[X]\} dr + \{CX + \bar{C}\mathbb{E}[X]\} dW(r), & r \in [s, T], \\ dY(r) = -\{A^\top Y + \bar{A}^\top \mathbb{E}[Y] + C^\top Z + \bar{C}^\top \mathbb{E}[Z] + QX + \bar{Q}\mathbb{E}[X]\} dr + ZdW(r), & r \in [s, T], \\ X(s) = \xi, \quad Y(T) = GX(T) + \bar{G}\mathbb{E}[X(T)]. \end{cases}$$

By Proposition 2.2 and (4.3), we have

$$\begin{aligned}
 & J^0(s, \xi; u(\cdot)) - J^0(s, \xi; 0) \\
 &= J^0(s, 0; u(\cdot)) + 2\mathbb{E} \int_s^T \langle B^\top Y + \bar{B}^\top \mathbb{E}[Y] + D^\top Z + \bar{D}^\top \mathbb{E}[Z] + SX + \bar{S}\mathbb{E}[X], u \rangle dr \\
 &\geq J^0(s, 0; u(\cdot)) - \delta \mathbb{E} \int_s^T |u(r)|^2 dr - \frac{1}{\delta} \mathbb{E} \int_s^T |B^\top Y + \bar{B}^\top \mathbb{E}[Y] + D^\top Z + \bar{D}^\top \mathbb{E}[Z] + SX + \bar{S}\mathbb{E}[X]|^2 dr \\
 &\geq -\frac{1}{\delta} \mathbb{E} \int_s^T |B^\top Y + \bar{B}^\top \mathbb{E}[Y] + D^\top Z + \bar{D}^\top \mathbb{E}[Z] + SX + \bar{S}\mathbb{E}[X]|^2 dr.
 \end{aligned} \tag{4.4}$$

If $\mathbb{E}[\xi] = 0$, then $\mathbb{E}[X(\cdot)] \equiv 0$, and one can verify that

$$X(r) = \mathbb{X}(r)\mathbb{X}(s)^{-1}\xi, \quad Y(r) = \mathbb{Y}(r)\mathbb{X}(s)^{-1}\xi, \quad Z(r) = \mathbb{Z}(r)\mathbb{X}(s)^{-1}\xi, \quad r \in [s, T],$$

where $\mathbb{X}(\cdot)$ is the solution to the following $\mathbb{R}^{n \times n}$ -valued SDE:

$$\begin{cases} d\mathbb{X}(r) = A(r)\mathbb{X}(r)dr + C(r)\mathbb{X}(r)dW(r), & r \in [0, T], \\ \mathbb{X}(0) = I, \end{cases} \tag{4.5}$$

and $(\mathbb{Y}(\cdot), \mathbb{Z}(\cdot))$ is the adapted solution to the following $\mathbb{R}^{n \times n}$ -valued backward SDE (BSDE, for short):

$$\begin{cases} d\mathbb{Y}(r) = -[A(r)^\top \mathbb{Y}(r) + C(r)^\top \mathbb{Z}(r) + Q(r)\mathbb{X}(r)]dr + \mathbb{Z}(r)dW(r), & r \in [0, T], \\ \mathbb{Y}(T) = G\mathbb{X}(T). \end{cases} \tag{4.6}$$

Note that $\mathbb{X}(r)\mathbb{X}(s)^{-1}$, $\mathbb{Y}(r)\mathbb{X}(s)^{-1}$ and $\mathbb{Z}(r)\mathbb{X}(s)^{-1}$ are independent of \mathcal{F}_s . Thus, $\mathbb{E}[X(\cdot)] = \mathbb{E}[Y(\cdot)] = \mathbb{E}[Z(\cdot)] = 0$ and (noting (4.4))

$$\begin{aligned}
 & J^0(s, \xi; u(\cdot)) \geq J^0(s, \xi; 0) - \frac{1}{\delta} \mathbb{E} \int_s^T |B(r)^\top Y(r) + D(r)^\top Z(r) + S(r)X(r)|^2 dr \\
 &= \mathbb{E} \left\{ \langle GX(T), X(T) \rangle + \int_s^T \langle Q(r)X(r), X(r) \rangle dr \right\} - \frac{1}{\delta} \mathbb{E} \int_s^T |B(r)^\top Y(r) + D(r)^\top Z(r) + S(r)X(r)|^2 dr \\
 &= \mathbb{E} \left\{ \xi^\top \left([\mathbb{X}(s)^{-1}]^\top \mathbb{X}(T)^\top G\mathbb{X}(T)\mathbb{X}(s)^{-1} + \int_s^T [\mathbb{X}(s)^{-1}]^\top \mathbb{X}(r)^\top Q(r)\mathbb{X}(r)\mathbb{X}(s)^{-1} dr \right) \xi \right\} \\
 &\quad - \frac{1}{\delta} \mathbb{E} \int_s^T \xi^\top [\mathbb{X}(s)^{-1}]^\top [B(r)^\top \mathbb{Y}(r) + D(r)^\top \mathbb{Z}(r) + S(r)\mathbb{X}(r)]^\top \\
 &\quad \times [B(r)^\top \mathbb{Y}(r) + D(r)^\top \mathbb{Z}(r) + S(r)\mathbb{X}(r)] \mathbb{X}(s)^{-1} \xi dr \\
 &= \mathbb{E} \left\{ \xi^\top \mathbb{E} \left([\mathbb{X}(s)^{-1}]^\top \mathbb{X}(T)^\top G\mathbb{X}(T)\mathbb{X}(s)^{-1} + \int_s^T [\mathbb{X}(s)^{-1}]^\top \mathbb{X}(r)^\top Q(r)\mathbb{X}(r)\mathbb{X}(s)^{-1} dr \right. \right. \\
 &\quad \left. \left. - \frac{1}{\delta} \int_s^T [\mathbb{X}(s)^{-1}]^\top [B(r)^\top \mathbb{Y}(r) + D(r)^\top \mathbb{Z}(r) + S(r)\mathbb{X}(r)]^\top \right. \right. \\
 &\quad \left. \left. \times [B(r)^\top \mathbb{Y}(r) + D(r)^\top \mathbb{Z}(r) + S(r)\mathbb{X}(r)] \mathbb{X}(s)^{-1} dr \right) \xi \right\} \equiv \mathbb{E}[\xi^\top \mathbb{M}(s)\xi].
 \end{aligned}$$

Note that $\mathbb{M}(\cdot) : [t, T] \rightarrow \mathbb{S}^n$ is continuous. The result therefore follows. □

Now, let us introduce the following Riccati equation:

$$\begin{cases} \dot{P} + PA + A^\top P + C^\top PC + Q \\ - (PB + C^\top PD + S^\top) (R + D^\top PD)^{-1} (B^\top P + D^\top PC + S) = 0, & \text{a.e. } s \in [t, T], \\ P(T) = G. \end{cases} \tag{4.7}$$

A solution $P(\cdot)$ of (4.7) is said to be *strongly regular* if

$$R(s) + D(s)^\top P(s)D(s) \geq \delta I, \quad \text{a.e. } s \in [t, T], \quad (4.8)$$

for some $\delta > 0$. The Riccati equation (4.7) is said to be *strongly regularly solvable*, if it admits a strongly regular solution. By a standard argument using Gronwall's inequality, one can show that if the regular solution of (4.7) exists, it must be unique. Compared with the strongly regular solution, the notion of *regular solution*, which is closely related to the closed-loop strategy, was introduced in [25]. The interested reader is referred to [26] for further information.

The following result shows that the strongly regular solvability of the Riccati equation (4.7) is necessary for the uniform convexity of the cost functional.

Theorem 4.2. *Let (H1)–(H2) and (H4) hold. Then the Riccati equation (4.7) is strongly regularly solvable.*

To prove the above result, we need the following lemma, whose proof can be found in [26].

Lemma 4.3. *Let (H1)–(H2) hold. For any $\Theta(\cdot) \in L^2(t, T; \mathbb{R}^{m \times n})$, let $P_\Theta(\cdot) \in C([t, T]; \mathbb{S}^n)$ be the solution to the following Lyapunov equation:*

$$\begin{cases} \dot{P}_\Theta + P_\Theta(A + B\Theta) + (A + B\Theta)^\top P_\Theta + (C + D\Theta)^\top P_\Theta(C + D\Theta) \\ \quad + \Theta^\top R\Theta + S^\top \Theta + \Theta^\top S + Q = 0, & \text{a.e. } s \in [t, T], \\ P_\Theta(T) = G. \end{cases} \quad (4.9)$$

If there exists a constant $\beta > 0$ such that for all $\Theta(\cdot) \in L^2(t, T; \mathbb{R}^{m \times n})$,

$$P_\Theta(s), \quad R(s) + D(s)^\top P_\Theta(s)D(s) \geq \beta I, \quad \text{a.e. } s \in [t, T], \quad (4.10)$$

then the Riccati equation (4.7) is strongly regularly solvable.

Proof of Theorem 4.2. We only need to show that the condition stated in Lemma 4.3 holds. To this end, let $\Theta(\cdot) \in L^2(t, T; \mathbb{R}^{m \times n})$ and $P(\cdot) \equiv P_\Theta(\cdot)$ be the corresponding solution of (4.9). For any deterministic $u(\cdot) \in L^2(t, T; \mathbb{R}^m)$, let $X^u(\cdot)$ be the solution of

$$\begin{cases} dX^u(s) = [(A + B\Theta)X^u + BuW]ds + [(C + D\Theta)X^u + DuW]dW, & s \in [t, T], \\ X^u(t) = 0, \end{cases}$$

and set

$$v(\cdot) \triangleq \Theta(\cdot)X^u(\cdot) + u(\cdot)W(\cdot) \in \mathcal{U}[t, T].$$

Clearly,

$$\mathbb{E}[X^u(s)] = 0, \quad \mathbb{E}[v(s)] = 0, \quad s \in [t, T]. \quad (4.11)$$

By the uniqueness of solutions, $X^u(\cdot)$ also solves

$$\begin{cases} dX^u(s) = \{AX^u + \bar{A}\mathbb{E}[X^u] + Bv + \bar{B}\mathbb{E}[v]\}ds + \{CX^u + \bar{C}\mathbb{E}[X^u] + Dv + \bar{D}\mathbb{E}[v]\}dW, & s \in [t, T], \\ X^u(t) = 0. \end{cases}$$

Thus, by applying Itô's formula to $s \rightarrow \langle P(s)X^u(s), X^u(s) \rangle$, we have (noting (H4) and (4.11))

$$\begin{aligned} & \delta \mathbb{E} \int_t^T |\Theta(s)X^u(s) + u(s)W(s)|^2 ds = \delta \mathbb{E} \int_t^T |v(s)|^2 ds \leq J^0(t, 0; v(\cdot)) \\ & = \mathbb{E} \left\{ \langle GX^u(T), X^u(T) \rangle + \int_t^T [\langle QX^u, X^u \rangle + 2\langle SX^u, v \rangle + \langle Rv, v \rangle] ds \right\} \\ & = \mathbb{E} \int_t^T \left\{ \langle \dot{P}X^u, X^u \rangle + \langle P[(A + B\Theta)X^u + BuW], X^u \rangle + \langle PX^u, (A + B\Theta)X^u + BuW \rangle \right. \\ & \quad + \langle P[(C + D\Theta)X^u + DuW], (C + D\Theta)X^u + DuW \rangle \\ & \quad \left. + \langle QX^u, X^u \rangle + 2\langle SX^u, \Theta X^u + uW \rangle + \langle R(\Theta X^u + uW), \Theta X^u + uW \rangle \right\} ds \\ & = \mathbb{E} \int_t^T \left\{ 2\langle [B^\top P + D^\top PC + S + (R + D^\top PD)\Theta]X^u, uW \rangle + \langle (R + D^\top PD)uW, uW \rangle \right\} ds. \end{aligned}$$

Hence, for any $u(\cdot) \in L^2(t, T; \mathbb{R}^m)$, the following holds:

$$\begin{aligned} & \mathbb{E} \int_t^T \left\{ 2\langle [B^\top P + D^\top PC + S + (R + D^\top PD - \delta I)\Theta]WX^u, u \rangle \right. \\ & \quad \left. + W^2\langle (R + D^\top PD - \delta I)u, u \rangle \right\} ds = \delta \mathbb{E} \int_t^T |\Theta(s)X^u(s)|^2 ds \geq 0. \end{aligned} \tag{4.12}$$

Now, applying Itô's formula again, we have

$$\begin{cases} d\mathbb{E}[W(s)X^u(s)] = \{ [A(s) + B(s)\Theta(s)]\mathbb{E}[W(s)X^u(s)] + sB(s)u(s) \} ds, & s \in [t, T], \\ \mathbb{E}[W(t)X^u(t)] = 0. \end{cases}$$

Fix any $u_0 \in \mathbb{R}^m$ and take $u(s) = u_0 \mathbf{1}_{[t', t'+h]}(s)$, with $t \leq t' < t' + h \leq T$. Then

$$\mathbb{E}[W(s)X^u(s)] = \begin{cases} 0, & s \in [t, t'], \\ \Phi(s) \int_t^{s \wedge (t'+h)} \Phi(r)^{-1} B(r) r u_0 dr, & s \in [t', T], \end{cases}$$

where $\Phi(\cdot)$ is the solution of the following $\mathbb{R}^{n \times n}$ -valued ordinary differential equation (ODE, for short):

$$\begin{cases} \dot{\Phi}(s) = [A(s) + B(s)\Theta(s)]\Phi(s), & s \in [0, T], \\ \Phi(0) = I. \end{cases}$$

Consequently, (4.12) becomes

$$\begin{aligned} & \int_{t'}^{t'+h} \left\{ 2\langle [B^\top P + D^\top PC + S + (R + D^\top PD - \delta I)\Theta]\Phi(s) \int_t^s \Phi(r)^{-1} B(r) r u_0 dr, u_0 \rangle \right. \\ & \quad \left. + s\langle (R + D^\top PD - \delta I)u_0, u_0 \rangle \right\} ds \geq 0. \end{aligned}$$

Dividing both sides by h and letting $h \rightarrow 0$, we obtain

$$t' \langle [R(t') + D(t')^\top P(t')D(t') - \delta I]u_0, u_0 \rangle \geq 0, \quad \forall u_0 \in \mathbb{R}^m, \quad \text{a.e. } t' \in [t, T],$$

which implies that

$$R(s) + D(s)^\top P(s)D(s) \geq \delta I, \quad \text{a.e. } s \in [t, T].$$

Next, for any $(s, x) \in [t, T] \times \mathbb{R}^n$, let $X(\cdot)$ be the solution of

$$\begin{cases} dX(r) = [A(r) + B(r)\Theta(r)]X(r)dr + [C(r) + D(r)\Theta(r)]X(r)dW(r), & r \in [s, T], \\ X(s) = W(s)x, \end{cases}$$

and set

$$w(\cdot) \triangleq \Theta(\cdot)X(\cdot) \in \mathcal{U}[s, T].$$

Similar to the previous argument, by applying Itô's formula to $r \rightarrow \langle P(r)X(r), X(r) \rangle$, we can derive that

$$J^0(s, W(s)x; w(\cdot)) = \mathbb{E}\langle P(s)W(s)x, W(s)x \rangle = s \langle P(s)x, x \rangle.$$

By Proposition 4.1, we have

$$s \langle P(s)x, x \rangle = J^0(s, W(s)x; w(\cdot)) \geq \alpha \mathbb{E}[|W(s)x|^2] = s\alpha|x|^2, \quad \forall (s, x) \in [t, T] \times \mathbb{R}^n,$$

which implies that $P(s) \geq \alpha I, \forall s \in [t, T]$. The proof is completed. □

From Theorem 4.2, we see that the Riccati equation (4.7) is strongly regularly solvable under the uniform convexity condition (H4). With the strongly regular solution $P(\cdot)$ of (4.7), we may further introduce the following deterministic LQ optimal control problem.

Consider the state equation

$$\begin{cases} \dot{y}(s) = [A(s) + \bar{A}(s)]y(s) + [B(s) + \bar{B}(s)]v(s), & s \in [t, T], \\ y(t) = x, \end{cases} \tag{4.13}$$

and the cost functional

$$\bar{J}(t, x; v(\cdot)) \triangleq \langle (G + \bar{G})y(T), y(T) \rangle + \int_t^T [\langle \Upsilon y, y \rangle + 2 \langle \Gamma y, v \rangle + \langle \Sigma v, v \rangle] ds, \tag{4.14}$$

where

$$\begin{cases} \Upsilon = Q + \bar{Q} + (C + \bar{C})^\top P(C + \bar{C}), \\ \Gamma = (D + \bar{D})^\top P(C + \bar{C}) + S + \bar{S}, \\ \Sigma = R + \bar{R} + (D + \bar{D})^\top P(D + \bar{D}). \end{cases}$$

We pose the following deterministic LQ problem.

Problem (DLQ). For any given $(t, x) \in [0, T] \times \mathbb{R}^n$, find a $v^*(\cdot) \in L^2(t, T; \mathbb{R}^m)$, such that

$$\bar{J}(t, x; v^*(\cdot)) = \inf_{v(\cdot) \in L^2(t, T; \mathbb{R}^m)} \bar{J}(t, x; v(\cdot)). \tag{4.15}$$

Note that the Riccati equation associated with Problem (DLQ) is

$$\begin{cases} \dot{\Pi} + \Pi(A + \bar{A}) + (A + \bar{A})^\top \Pi + Q + \bar{Q} + (C + \bar{C})^\top P(C + \bar{C}) \\ \quad - [\Pi(B + \bar{B}) + (C + \bar{C})^\top P(D + \bar{D}) + (S + \bar{S})^\top] [R + \bar{R} + (D + \bar{D})^\top P(D + \bar{D})]^{-1} \\ \quad \times [(B + \bar{B})^\top \Pi + (D + \bar{D})^\top P(C + \bar{C}) + (S + \bar{S})] = 0, & \text{a.e. } s \in [t, T], \\ \Pi(T) = G + \bar{G}. \end{cases} \tag{4.16}$$

We have the following result.

Theorem 4.4. *Let (H1)–(H2) and (H4) hold. Then the map $v(\cdot) \mapsto \bar{J}(t, 0; v(\cdot))$ is uniformly convex, i.e., there exists a $\lambda > 0$ such that*

$$\bar{J}(t, 0; v(\cdot)) \geq \lambda \int_t^T |v(s)|^2 ds, \quad \forall v(\cdot) \in L^2(t, T; \mathbb{R}^m). \tag{4.17}$$

Consequently, the strongly regular solution $P(\cdot)$ of the Riccati equation (4.7) satisfies

$$\Sigma = R + \bar{R} + (D + \bar{D})^\top P(D + \bar{D}) \gg 0, \tag{4.18}$$

and the Riccati equation (4.16) admits a unique solution $\Pi(\cdot) \in C([t, T]; \mathbb{S}^n)$.

Proof. Let $P(\cdot)$ be the strongly regular solution of the Riccati equation (4.7) and set

$$\Theta = -(R + D^\top P D)^{-1}(B^\top P + D^\top P C + S) \in L^2(t, T; \mathbb{R}^{m \times n}).$$

We claim that

$$J^0(t, 0; \Theta(\cdot)X(\cdot) + v(\cdot)) = \bar{J}(t, 0; \Theta(\cdot)y(\cdot) + v(\cdot)), \quad \forall v(\cdot) \in L^2(t, T; \mathbb{R}^m). \tag{4.19}$$

To prove (4.19), take any $v(\cdot) \in L^2(t, T; \mathbb{R}^m)$, let $y(\cdot)$ be the solution of

$$\begin{cases} \dot{y}(s) = [A(s) + \bar{A}(s)]y(s) + [B(s) + \bar{B}(s)][\Theta(s)y(s) + v(s)], & s \in [t, T], \\ y(t) = 0, \end{cases} \tag{4.20}$$

and $X(\cdot)$ be the solution of

$$\begin{cases} dX(s) = \left\{ AX + \bar{A}\mathbb{E}[X] + B(\Theta X + v) + \bar{B}\mathbb{E}[\Theta X + v] \right\} ds \\ \quad + \left\{ CX + \bar{C}\mathbb{E}[X] + D(\Theta X + v) + \bar{D}\mathbb{E}[\Theta X + v] \right\} dW(s), & s \in [t, T], \\ X(t) = 0. \end{cases} \tag{4.21}$$

Note that $v(\cdot)$ is deterministic. Then

$$\begin{cases} d\mathbb{E}[X(s)] = \left\{ (A + \bar{A})\mathbb{E}[X] + (B + \bar{B})(\Theta\mathbb{E}[X] + v) \right\} ds, & s \in [t, T], \\ \mathbb{E}[X(t)] = 0. \end{cases}$$

By the uniqueness of solutions, we see that

$$\mathbb{E}[X(s)] = y(s), \quad s \in [t, T].$$

Now let $z(\cdot) = X(\cdot) - \mathbb{E}[X(\cdot)]$. Then

$$\begin{cases} dz(s) = (A + B\Theta)z ds + \left\{ (C + D\Theta)z + (C + \bar{C})y + (D + \bar{D})(\Theta y + v) \right\} dW(s), & s \in [t, T], \\ z(t) = 0. \end{cases}$$

Keep in mind that $v(\cdot)$ is deterministic and note that

$$\begin{aligned} 0 &= \dot{P} + P(A + B\Theta) + (A + B\Theta)^\top P + (C + D\Theta)^\top P(C + D\Theta) \\ &\quad + \Theta^\top R\Theta + S^\top \Theta + \Theta^\top S + Q. \end{aligned}$$

By applying Itô's formula to $s \mapsto \langle P(s)z(s), z(s) \rangle$, we have (also, noting $\mathbb{E}[z] = 0$)

$$\begin{aligned}
 & J^0(t, 0; \Theta(\cdot)X(\cdot) + v(\cdot)) \\
 &= \mathbb{E} \left\{ \langle GX(T), X(T) \rangle + \langle \bar{G}\mathbb{E}[X(T)], \mathbb{E}[X(T)] \rangle \right. \\
 &\quad \left. + \int_t^T \left[\langle QX, X \rangle + \langle \bar{Q}\mathbb{E}[X], \mathbb{E}[X] \rangle + 2 \langle SX, \Theta X + v \rangle + 2 \langle \bar{S}\mathbb{E}[X], \mathbb{E}[\Theta X + v] \rangle \right. \right. \\
 &\quad \left. \left. + \langle R(\Theta X + v), (\Theta X + v) \rangle + \langle \bar{R}\mathbb{E}[\Theta X + v], \mathbb{E}[\Theta X + v] \rangle \right] ds \right\} \\
 &= \mathbb{E} \left\{ \langle Gz(T), z(T) \rangle + \int_t^T \left[\langle Qz, z \rangle + 2 \langle Sz, \Theta z \rangle + \langle R\Theta z, \Theta z \rangle \right] ds \right\} \\
 &\quad + \langle (G + \bar{G})y(T), y(T) \rangle + \int_t^T \left[\langle (Q + \bar{Q})y, y \rangle + 2 \langle (S + \bar{S})y, \Theta y + v \rangle \right. \\
 &\quad \left. + \langle (R + \bar{R})(\Theta y + v), \Theta y + v \rangle \right] ds \\
 &= \mathbb{E} \int_t^T \left\{ \langle \dot{P}z, z \rangle + \langle P(A + B\Theta)z, z \rangle + \langle Pz, (A + B\Theta)z \rangle \right. \\
 &\quad \left. + \langle P[(C + D\Theta)z + (C + \bar{C})y + (D + \bar{D})(\Theta y + v)], \right. \\
 &\quad \quad \left. (C + D\Theta)z + (C + \bar{C})y + (D + \bar{D})(\Theta y + v) \rangle \right. \\
 &\quad \left. + \langle (Q + S^\top \Theta + \Theta^\top S + \Theta^\top R\Theta)z, z \rangle \right\} ds \\
 &\quad + \langle (G + \bar{G})y(T), y(T) \rangle + \int_t^T \left[\langle (Q + \bar{Q})y, y \rangle + 2 \langle (S + \bar{S})y, \Theta y + v \rangle \right. \\
 &\quad \left. + \langle (R + \bar{R})(\Theta y + v), \Theta y + v \rangle \right] ds \\
 &= \int_t^T \langle P[(C + \bar{C})y + (D + \bar{D})(\Theta y + v)], (C + \bar{C})y + (D + \bar{D})(\Theta y + v) \rangle ds \\
 &\quad + \langle (G + \bar{G})y(T), y(T) \rangle + \int_t^T \left[\langle (Q + \bar{Q})y, y \rangle + 2 \langle (S + \bar{S})y, \Theta y + v \rangle \right. \\
 &\quad \left. + \langle (R + \bar{R})(\Theta y + v), \Theta y + v \rangle \right] ds \\
 &= \langle (G + \bar{G})y(T), y(T) \rangle + \int_t^T \left\{ \langle [Q + \bar{Q} + (C + \bar{C})^\top P(C + \bar{C})]y, y \rangle \right. \\
 &\quad \left. + 2 \langle [(D + \bar{D})^\top P(C + \bar{C}) + S + \bar{S}]y, \Theta y + v \rangle \right. \\
 &\quad \left. + \langle [R + \bar{R} + (D + \bar{D})^\top P(D + \bar{D})](\Theta y + v), \Theta y + v \rangle \right\} ds \\
 &= \bar{J}(t, 0; \Theta(\cdot)y(\cdot) + v(\cdot)).
 \end{aligned}$$

Thus, (4.19) holds. Consequently, by (H4), we have

$$\begin{aligned}
 \bar{J}(t, 0; \Theta(\cdot)y(\cdot) + v(\cdot)) &= J^0(t, 0; \Theta(\cdot)X(\cdot) + v(\cdot)) \geq \delta \mathbb{E} \int_t^T |\Theta(s)X(s) + v(s)|^2 ds \geq \delta \int_t^T |\mathbb{E}[\Theta(s)X(s) + v(s)]|^2 ds \\
 &= \delta \int_t^T |\Theta(s)y(s) + v(s)|^2 ds, \quad \forall v(\cdot) \in L^2(t, T; \mathbb{R}^m),
 \end{aligned}$$

which implies the uniform convexity of $v(\cdot) \mapsto \bar{J}(t, 0; v(\cdot))$. The rest of the theorem follows now immediately from ([26], Thm. 4.6). \square

5. SUFFICIENCY OF THE RICCATI EQUATIONS

In the previous section, we proved that the solvability of the Riccati equations (4.7) and (4.16) is necessary for the uniform convexity of the cost functional. In this section, we shall show that it is also sufficient. Moreover, under the uniform convexity condition, the optimal control can be represented explicitly as a state feedback form *via* the solutions of the Riccati equations.

First we need the following lemma.

Lemma 5.1. *Let (H1)–(H2) hold. For any $u(\cdot) \in \mathcal{U}[t, T]$, let $X_0^u(\cdot)$ be the solution of*

$$\begin{cases} dX_0^u(s) = \left\{ A(s)X_0^u(s) + \bar{A}(s)\mathbb{E}[X_0^u(s)] + B(s)u(s) + \bar{B}(s)\mathbb{E}[u(s)] \right\} ds \\ \quad + \left\{ C(s)X_0^u(s) + \bar{C}(s)\mathbb{E}[X_0^u(s)] + D(s)u(s) + \bar{D}(s)\mathbb{E}[u(s)] \right\} dW(s), & s \in [t, T], \\ X_0^u(t) = 0. \end{cases} \tag{5.1}$$

Then for any $\Theta(\cdot), \bar{\Theta}(\cdot) \in L^2(t, T; \mathbb{R}^{m \times n})$, there exists a constant $\gamma > 0$ such that

$$\begin{aligned} \mathbb{E} \int_t^T |u(s) - \Theta(s)(X_0^u(s) - \mathbb{E}[X_0^u(s)])|^2 ds &\geq \gamma \mathbb{E} \int_t^T |u(s)|^2 ds, & \forall u(\cdot) \in \mathcal{U}[t, T], \\ \int_t^T |\mathbb{E}[u(s)] - \bar{\Theta}(s)\mathbb{E}[X_0^u(s)]|^2 ds &\geq \gamma \int_t^T |\mathbb{E}[u(s)]|^2 ds, & \forall u(\cdot) \in \mathcal{U}[t, T]. \end{aligned} \tag{5.2}$$

Proof. Let $\Theta(\cdot) \in L^2(t, T; \mathbb{R}^{m \times n})$. Define a bounded linear operator $\mathcal{A} : \mathcal{U}[t, T] \rightarrow \mathcal{U}[t, T]$ by

$$\mathcal{A}u = u - \Theta(X_0^u - \mathbb{E}[X_0^u]).$$

Then \mathcal{A} is bijective and its inverse \mathcal{A}^{-1} is given by

$$\mathcal{A}^{-1}u = u + \Theta \left(\tilde{X}_0^u - \mathbb{E}[\tilde{X}_0^u] \right),$$

where $\tilde{X}_0^u(\cdot)$ is the solution of

$$\begin{cases} d\tilde{X}_0^u(s) = \left\{ (A + B\Theta)\tilde{X}_0^u + (\bar{A} - B\Theta)\mathbb{E}[\tilde{X}_0^u] + Bu + \bar{B}\mathbb{E}[u] \right\} ds \\ \quad + \left\{ (C + D\Theta)\tilde{X}_0^u + (\bar{C} - D\Theta)\mathbb{E}[\tilde{X}_0^u] + Du + D\mathbb{E}[u] \right\} dW(s), & s \in [t, T], \\ \tilde{X}_0^u(t) = 0. \end{cases}$$

By the bounded inverse theorem, \mathcal{A}^{-1} is bounded with norm $\|\mathcal{A}^{-1}\| > 0$. Thus,

$$\begin{aligned} \mathbb{E} \int_t^T |u(s)|^2 ds &= \mathbb{E} \int_t^T |(\mathcal{A}^{-1}\mathcal{A}u)(s)|^2 ds \leq \|\mathcal{A}^{-1}\| \mathbb{E} \int_t^T |(\mathcal{A}u)(s)|^2 ds \\ &= \|\mathcal{A}^{-1}\| \mathbb{E} \int_t^T |u(s) - \Theta(s)(X_0^u(s) - \mathbb{E}[X_0^u(s)])|^2 ds, & \forall u(\cdot) \in \mathcal{U}[t, T], \end{aligned}$$

which implies the first inequality in (5.2) with $\gamma = \|\mathcal{A}^{-1}\|^{-1}$.

To prove the second, for any $v(\cdot) \in L^2(t, T; \mathbb{R}^{m \times n})$, let $y^v(\cdot)$ be the solution to the following ODE:

$$\begin{cases} \dot{y}^v(s) = [A(s) + \bar{A}(s)]y^v(s) + [B(s) + \bar{B}(s)]v(s), & s \in [t, T], \\ y^v(t) = 0. \end{cases} \quad (5.3)$$

For $\bar{\Theta}(\cdot) \in L^2(t, T; \mathbb{R}^{m \times n})$, we define a bounded linear operator $\mathcal{B} : L^2(t, T; \mathbb{R}^m) \rightarrow L^2(t, T; \mathbb{R}^m)$ by

$$\mathcal{B}v = v - \bar{\Theta}y^v.$$

Similar to the previous argument, one can show that \mathcal{B} is invertible and

$$\int_t^T |v(s) - \bar{\Theta}(s)y^v(s)|^2 ds \geq \frac{1}{\|\mathcal{B}^{-1}\|} \int_t^T |v(s)|^2 ds, \quad \forall v(\cdot) \in L^2(t, T; \mathbb{R}^{m \times n}).$$

Observe that $\mathbb{E}[X_0^u(\cdot)]$ satisfies (5.3) with $v(\cdot) = \mathbb{E}[u(\cdot)]$. The result therefore follows. \square

Now we present the main result of this section, which gives a characterization for the uniform convexity of the cost functional as well as a feedback representation of the optimal control.

Theorem 5.2. *Let (H1)–(H2) hold. Then the map $u(\cdot) \mapsto J^0(t, 0; u(\cdot))$ is uniformly convex if and only if the Riccati equation (4.7) admits a strongly regular solution $P(\cdot)$ such that*

$$\Sigma \equiv R + \bar{R} + (D + \bar{D})^\top P(D + \bar{D}) \gg 0, \quad (5.4)$$

and the corresponding Riccati equation (4.16) admits a solution $\Pi(\cdot)$. In this case, the unique optimal $u^*(\cdot)$ of Problem (MF-LQ) at (t, ξ) is given by

$$u^* = \Theta(X^* - \mathbb{E}[X^*]) + \bar{\Theta}\mathbb{E}[X^*] + \varphi - \mathbb{E}[\varphi] + \bar{\varphi}, \quad (5.5)$$

where

$$\begin{cases} \Theta = -(R + D^\top PD)^{-1}(B^\top P + D^\top PC + S), \\ \bar{\Theta} = -\Sigma^{-1}[(B + \bar{B})^\top \Pi + (D + \bar{D})^\top P(C + \bar{C}) + (S + \bar{S})], \\ \varphi = -(R + D^\top PD)^{-1}[B^\top \eta + D^\top (\zeta + P\sigma) + \rho], \\ \bar{\varphi} = -\Sigma^{-1}\{(B + \bar{B})^\top \bar{\eta} + (D + \bar{D})^\top (\mathbb{E}[\zeta] + P\mathbb{E}[\sigma]) + \mathbb{E}[\rho] + \bar{\rho}\}, \end{cases} \quad (5.6)$$

with $(\eta(\cdot), \zeta(\cdot))$ and $\bar{\eta}(\cdot)$ being the (adapted) solutions to the following BSDE

$$\begin{cases} d\eta(s) = -[(A + B\Theta)^\top \eta + (C + D\Theta)^\top \zeta + (C + D\Theta)^\top P\sigma \\ \quad + \Theta^\top \rho + Pb + q] ds + \zeta dW(s), & s \in [t, T], \\ \eta(T) = g, \end{cases} \quad (5.7)$$

and ordinary differential equation

$$\begin{cases} \dot{\bar{\eta}} + [(A + \bar{A}) + (B + \bar{B})\bar{\Theta}]^\top \bar{\eta} + \bar{\Theta}^\top \left\{ (D + \bar{D})^\top (P\mathbb{E}[\sigma] + \mathbb{E}[\zeta]) + \mathbb{E}[\rho] + \bar{\rho} \right\} \\ + (C + \bar{C})^\top (P\mathbb{E}[\sigma] + \mathbb{E}[\zeta]) + \mathbb{E}[q] + \bar{q} + \Pi\mathbb{E}[b] = 0, & \text{a.e. } s \in [t, T], \\ \bar{\eta}(T) = \mathbb{E}[g] + \bar{g}, \end{cases} \tag{5.8}$$

respectively, and $X^*(\cdot)$ is the solution of the closed-loop system

$$\begin{cases} dX^*(s) = \left\{ (A + B\Theta)(X^* - \mathbb{E}[X^*]) + [(A + \bar{A}) + (B + \bar{B})\bar{\Theta}]\mathbb{E}[X^*] \right. \\ \quad \left. + B(\varphi - \mathbb{E}[\varphi]) + (B + \bar{B})\bar{\varphi} + b \right\} ds \\ \quad + \left\{ (C + D\Theta)(X^* - \mathbb{E}[X^*]) + [(C + \bar{C}) + (D + \bar{D})\bar{\Theta}]\mathbb{E}[X^*] \right. \\ \quad \left. + D(\varphi - \mathbb{E}[\varphi]) + (D + \bar{D})\bar{\varphi} + \sigma \right\} dW(s), \quad s \in [t, T], \\ X^*(t) = \xi. \end{cases} \tag{5.9}$$

Moreover, the value $V(t, \xi)$ is given by

$$\begin{aligned} V(t, \xi) &= \mathbb{E} \langle P(t)(\xi - \mathbb{E}[\xi]) + 2\eta(t), \xi - \mathbb{E}[\xi] \rangle + \langle \Pi(t)\mathbb{E}[\xi] + 2\bar{\eta}(t), \mathbb{E}[\xi] \rangle \\ &\quad + \mathbb{E} \int_t^T \left\{ \langle P\sigma, \sigma \rangle + 2\langle \eta, b - \mathbb{E}[b] \rangle + 2\langle \zeta, \sigma \rangle + 2\langle \bar{\eta}, \mathbb{E}[b] \rangle \right. \\ &\quad \left. - \langle \Sigma_0(\varphi - \mathbb{E}[\varphi]), \varphi - \mathbb{E}[\varphi] \rangle - \langle \Sigma\bar{\varphi}, \bar{\varphi} \rangle \right\} ds, \end{aligned} \tag{5.10}$$

where $\Sigma_0 = R + D^\top PD$.

Proof. The “only if” part has been proved in Section 4. Let us now show the “if” part. For any $\xi \in L^2_{\mathcal{F}_t}(\Omega; \mathbb{R}^n)$ and $u(\cdot) \in \mathcal{U}[t, T]$, let $X(\cdot) \equiv X(\cdot; t, \xi, u(\cdot))$ be the corresponding solution of (1.1). Set

$$z(\cdot) = X(\cdot) - \mathbb{E}[X(\cdot)], \quad v(\cdot) = u(\cdot) - \mathbb{E}[u(\cdot)], \quad y(\cdot) = \mathbb{E}[X(\cdot)].$$

Then

$$\begin{cases} dz(s) = \left\{ Az + Bv + b - \mathbb{E}[b] \right\} ds \\ \quad + \left\{ Cz + Dv + \sigma + (C + \bar{C})y + (D + \bar{D})\mathbb{E}[u] \right\} dW(s), \quad s \in [t, T], \\ z(t) = \xi - \mathbb{E}[\xi], \end{cases} \tag{5.11}$$

and

$$\begin{cases} \dot{y} = (A + \bar{A})y + (B + \bar{B})\mathbb{E}[u] + \mathbb{E}[b], & s \in [t, T], \\ y(t) = \mathbb{E}[\xi]. \end{cases} \tag{5.12}$$

Now we rewrite the cost functional as follows:

$$\begin{aligned} J(t, \xi; u(\cdot)) &= \mathbb{E} \left\{ \langle Gz(T) + 2g, z(T) \rangle + \int_t^T \left[\left\langle \begin{pmatrix} Q & S^\top \\ S & R \end{pmatrix} \begin{pmatrix} z \\ v \end{pmatrix}, \begin{pmatrix} z \\ v \end{pmatrix} \right\rangle + 2 \left\langle \begin{pmatrix} q \\ \rho \end{pmatrix}, \begin{pmatrix} z \\ v \end{pmatrix} \right\rangle \right] ds \right\} \\ &\quad + \langle (G + \bar{G})y(T) + 2(\mathbb{E}[g] + \bar{g}), y(T) \rangle \\ &\quad + \int_t^T \left[\left\langle \begin{pmatrix} Q + \bar{Q} & (S + \bar{S})^\top \\ S + \bar{S} & R + \bar{R} \end{pmatrix} \begin{pmatrix} y \\ \mathbb{E}[u] \end{pmatrix}, \begin{pmatrix} y \\ \mathbb{E}[u] \end{pmatrix} \right\rangle + 2 \left\langle \begin{pmatrix} \mathbb{E}[q] + \bar{q} \\ \mathbb{E}[\rho] + \bar{\rho} \end{pmatrix}, \begin{pmatrix} y \\ \mathbb{E}[u] \end{pmatrix} \right\rangle \right] ds. \end{aligned} \tag{5.13}$$

Applying Itô's formula to $s \mapsto \langle P(s)z(s) + 2\eta(s), z(s) \rangle$, we have (noting $\mathbb{E}[z] \equiv 0, \mathbb{E}[v] \equiv 0$)

$$\begin{aligned}
 & \mathbb{E}\langle Gz(T) + 2g, z(T) \rangle - \mathbb{E}\langle P(t)(\xi - \mathbb{E}[\xi]) + 2\eta(t), \xi - \mathbb{E}[\xi] \rangle \\
 & + \mathbb{E} \int_t^T \left[\langle Qz, z \rangle + 2\langle Sz, v \rangle + \langle Rv, v \rangle + 2\langle q, z \rangle + 2\langle \rho, v \rangle \right] ds \\
 = & \mathbb{E} \int_t^T \left[\langle \dot{P}z, z \rangle + \langle P(Az + Bv + b - \mathbb{E}[b]), z \rangle + \langle Pz, (Az + Bv + b - \mathbb{E}[b]) \rangle \right. \\
 & + \langle P\{Cz + Dv + \sigma + (C + \bar{C})y + (D + \bar{D})\mathbb{E}[u]\}, \\
 & \quad Cz + Dv + \sigma + (C + \bar{C})y + (D + \bar{D})\mathbb{E}[u] \rangle \\
 & - 2\langle (A + B\Theta)^\top \eta + (C + D\Theta)^\top \zeta + (C + D\Theta)^\top P\sigma + \Theta^\top \rho + Pb + q, z \rangle \\
 & + 2\langle \eta, Az + Bv + b - \mathbb{E}[b] \rangle \\
 & \left. + 2\langle \zeta, Cz + Dv + \sigma + (C + \bar{C})y + (D + \bar{D})\mathbb{E}[u] \rangle \right] ds \\
 & + \mathbb{E} \int_t^T \left[\langle Qz, z \rangle + 2\langle Sz, v \rangle + \langle Rv, v \rangle + 2\langle q, z \rangle + 2\langle \rho, v \rangle \right] ds \\
 = & \mathbb{E} \int_t^T \left[\langle (\dot{P} + PA + A^\top P + C^\top PC + Q)z, z \rangle + 2\langle (PB + C^\top PD + S^\top)v, z \rangle \right. \\
 & + \langle (R + D^\top PD)v, v \rangle + 2\langle B^\top \eta + D^\top \zeta + D^\top P\sigma + \rho, v - \Theta z \rangle \\
 & + 2\langle P\mathbb{E}[\sigma] + \mathbb{E}[\zeta], (C + \bar{C})y + (D + \bar{D})\mathbb{E}[u] \rangle \\
 & + \langle P\{(C + \bar{C})y + (D + \bar{D})\mathbb{E}[u]\}, (C + \bar{C})y + (D + \bar{D})\mathbb{E}[u] \rangle \\
 & \left. + \langle P\sigma, \sigma \rangle + 2\langle \eta, b - \mathbb{E}[b] \rangle + 2\langle \zeta, \sigma \rangle \right] ds \\
 = & \mathbb{E} \int_t^T \left[\langle \Theta^\top \Sigma_0 \Theta z, z \rangle - 2\langle \Theta^\top \Sigma_0 v, z \rangle + \langle \Sigma_0 v, v \rangle - 2\langle \Sigma_0 \varphi, v - \Theta z \rangle \right. \\
 & + \langle (C + \bar{C})^\top P(C + \bar{C})y, y \rangle + 2\langle (C + \bar{C})^\top P(D + \bar{D})\mathbb{E}[u], y \rangle \\
 & + \langle (D + \bar{D})^\top P(D + \bar{D})\mathbb{E}[u], \mathbb{E}[u] \rangle + 2\langle (C + \bar{C})^\top (P\mathbb{E}[\sigma] + \mathbb{E}[\zeta]), y \rangle \\
 & \left. + 2\langle (D + \bar{D})^\top (P\mathbb{E}[\sigma] + \mathbb{E}[\zeta]), \mathbb{E}[u] \rangle + \langle P\sigma, \sigma \rangle + 2\langle \eta, b - \mathbb{E}[b] \rangle + 2\langle \zeta, \sigma \rangle \right] ds \\
 = & \mathbb{E} \int_t^T \left[\langle \Sigma_0(v - \Theta z - \varphi), v - \Theta z - \varphi \rangle - \langle \Sigma_0 \varphi, \varphi \rangle \right. \\
 & + \langle (C + \bar{C})^\top P(C + \bar{C})y, y \rangle + 2\langle (C + \bar{C})^\top P(D + \bar{D})\mathbb{E}[u], y \rangle \\
 & + \langle (D + \bar{D})^\top P(D + \bar{D})\mathbb{E}[u], \mathbb{E}[u] \rangle + 2\langle (C + \bar{C})^\top (P\mathbb{E}[\sigma] + \mathbb{E}[\zeta]), y \rangle \\
 & \left. + 2\langle (D + \bar{D})^\top (P\mathbb{E}[\sigma] + \mathbb{E}[\zeta]), \mathbb{E}[u] \rangle + \langle P\sigma, \sigma \rangle + 2\langle \eta, b - \mathbb{E}[b] \rangle + 2\langle \zeta, \sigma \rangle \right] ds.
 \end{aligned} \tag{5.14}$$

Applying the integration by parts formula to $s \mapsto \langle \Pi(s)y(s) + 2\bar{\eta}(s), y(s) \rangle$, we have

$$\begin{aligned}
 & \langle (G + \bar{G})y(T) + 2(\mathbb{E}[g] + \bar{g}), y(T) \rangle - \langle \Pi(t)\mathbb{E}[\xi] + 2\bar{\eta}(t), \mathbb{E}[\xi] \rangle \\
 & + \int_t^T \left[\left\langle \begin{pmatrix} Q + \bar{Q} & (S + \bar{S})^\top \\ S + \bar{S} & R + \bar{R} \end{pmatrix} \begin{pmatrix} y \\ \mathbb{E}[u] \end{pmatrix}, \begin{pmatrix} y \\ \mathbb{E}[u] \end{pmatrix} \right\rangle + 2 \left\langle \begin{pmatrix} \mathbb{E}[q] + \bar{q} \\ \mathbb{E}[\rho] + \bar{\rho} \end{pmatrix}, \begin{pmatrix} y \\ \mathbb{E}[u] \end{pmatrix} \right\rangle \right] ds \\
 = & \int_t^T \left[\langle \dot{H}y, y \rangle + \langle \Pi \{ (A + \bar{A})y + (B + \bar{B})\mathbb{E}[u] + \mathbb{E}[b] \}, y \rangle \right. \\
 & \left. + \langle \Pi y, (A + \bar{A})y + (B + \bar{B})\mathbb{E}[u] + \mathbb{E}[b] \rangle + 2\langle \dot{\eta}, y \rangle + 2\langle \bar{\eta}, (A + \bar{A})y + (B + \bar{B})\mathbb{E}[u] + \mathbb{E}[b] \rangle \right] ds \\
 & + \int_t^T \left[\langle (Q + \bar{Q})y, y \rangle + 2\langle (S + \bar{S})y, \mathbb{E}[u] \rangle + \langle (R + \bar{R})\mathbb{E}[u], \mathbb{E}[u] \rangle + 2\langle \mathbb{E}[q] + \bar{q}, y \rangle + 2\langle \mathbb{E}[\rho] + \bar{\rho}, \mathbb{E}[u] \rangle \right] ds \\
 = & \int_t^T \left\{ \langle [\dot{H} + \Pi(A + \bar{A}) + (A + \bar{A})^\top \Pi + Q + \bar{Q}]y, y \rangle \right. \\
 & + 2\langle [\Pi(B + \bar{B}) + (S + \bar{S})^\top] \mathbb{E}[u], y \rangle + 2\langle \dot{\eta} + (A + \bar{A})^\top \bar{\eta} + \mathbb{E}[q] + \bar{q} + \Pi \mathbb{E}[b], y \rangle \\
 & \left. + 2\langle (B + \bar{B})^\top \bar{\eta} + \mathbb{E}[\rho] + \bar{\rho}, \mathbb{E}[u] \rangle + \langle (R + \bar{R})\mathbb{E}[u], \mathbb{E}[u] \rangle + 2\langle \bar{\eta}, \mathbb{E}[b] \rangle \right\} ds. \tag{5.15}
 \end{aligned}$$

Adding (5.14) and (5.15) together and noting (5.13), we obtain

$$\begin{aligned}
 & J(t, \xi; u(\cdot)) - \mathbb{E}\langle P(t)(\xi - \mathbb{E}[\xi]) + 2\eta(t), \xi - \mathbb{E}[\xi] \rangle - \langle \Pi(t)\mathbb{E}[\xi] + 2\bar{\eta}(t), \mathbb{E}[\xi] \rangle \\
 & = \mathbb{E} \int_t^T \left\{ \langle \Sigma_0(v - \Theta z - \varphi), v - \Theta z - \varphi \rangle - \langle \Sigma_0 \varphi, \varphi \rangle \right. \\
 & \quad + \langle P\sigma, \sigma \rangle + 2\langle \eta, b - \mathbb{E}[b] \rangle + 2\langle \zeta, \sigma \rangle + 2\langle \bar{\eta}, \mathbb{E}[b] \rangle \left. \right\} ds \\
 & \quad + \int_t^T \left\{ \langle [\dot{H} + \Pi(A + \bar{A}) + (A + \bar{A})^\top \Pi + Q + \bar{Q} + (C + \bar{C})^\top P(C + \bar{C})]y, y \rangle \right. \\
 & \quad + 2\langle [\Pi(B + \bar{B}) + (C + \bar{C})^\top P(D + \bar{D}) + (S + \bar{S})^\top] \mathbb{E}[u], y \rangle \\
 & \quad + \langle [R + \bar{R} + (D + \bar{D})^\top P(D + \bar{D})] \mathbb{E}[u], \mathbb{E}[u] \rangle \\
 & \quad + 2\langle \dot{\eta} + (A + \bar{A})^\top \bar{\eta} + (C + \bar{C})^\top (P\mathbb{E}[\sigma] + \mathbb{E}[\zeta]) + \mathbb{E}[q] + \bar{q} + \Pi \mathbb{E}[b], y \rangle \\
 & \quad \left. + 2\langle (B + \bar{B})^\top \bar{\eta} + (D + \bar{D})^\top (P\mathbb{E}[\sigma] + \mathbb{E}[\zeta]) + \mathbb{E}[\rho] + \bar{\rho}, \mathbb{E}[u] \rangle \right\} ds \\
 = & \mathbb{E} \int_t^T \left\{ \langle \Sigma_0(v - \Theta z - \varphi), v - \Theta z - \varphi \rangle - \langle \Sigma_0 \varphi, \varphi \rangle \right. \\
 & \quad + \langle P\sigma, \sigma \rangle + 2\langle \eta, b - \mathbb{E}[b] \rangle + 2\langle \zeta, \sigma \rangle + 2\langle \bar{\eta}, \mathbb{E}[b] \rangle \left. \right\} ds \\
 & \quad + \int_t^T \left\{ \langle \bar{\Theta}^\top \Sigma \bar{\Theta} y, y \rangle - 2\langle \bar{\Theta}^\top \Sigma \mathbb{E}[u], y \rangle \right. \\
 & \quad \left. + \langle \Sigma \mathbb{E}[u], \mathbb{E}[u] \rangle - 2\langle \Sigma \bar{\varphi}, \mathbb{E}[u] - \bar{\Theta} y \rangle \right\} ds \tag{5.16}
 \end{aligned}$$

$$\begin{aligned}
 &= \mathbb{E} \int_t^T \left\{ \langle \Sigma_0(v - \Theta z - \varphi), v - \Theta z - \varphi \rangle - \langle \Sigma_0 \varphi, \varphi \rangle \right. \\
 &\quad + \langle \Sigma(\mathbb{E}[u] - \bar{\Theta} y - \bar{\varphi}), \mathbb{E}[u] - \bar{\Theta} y - \bar{\varphi} \rangle - \langle \Sigma \bar{\varphi}, \bar{\varphi} \rangle \\
 &\quad \left. + \langle P\sigma, \sigma \rangle + 2\langle \eta, b - \mathbb{E}[b] \rangle + 2\langle \zeta, \sigma \rangle + 2\langle \bar{\eta}, \mathbb{E}[b] \rangle \right\} ds \\
 &= \mathbb{E} \int_t^T \left\{ \langle P\sigma, \sigma \rangle + 2\langle \eta, b - \mathbb{E}[b] \rangle \right. \\
 &\quad + 2\langle \zeta, \sigma \rangle + 2\langle \bar{\eta}, \mathbb{E}[b] \rangle - \langle \Sigma \bar{\varphi}, \bar{\varphi} \rangle \\
 &\quad - \langle \Sigma_0(\varphi - \mathbb{E}[\varphi]), \varphi - \mathbb{E}[\varphi] \rangle + \langle \Sigma_0(v - \Theta z - \varphi + \mathbb{E}[\varphi]), v - \Theta z - \varphi + \mathbb{E}[\varphi] \rangle \\
 &\quad \left. + \langle \Sigma(\mathbb{E}[u] - \bar{\Theta} y - \bar{\varphi}), \mathbb{E}[u] - \bar{\Theta} y - \bar{\varphi} \rangle \right\} ds.
 \end{aligned}$$

Since $\Sigma_0, \Sigma \gg 0$, (5.16) implies that

$$\begin{aligned}
 J(t, \xi; u(\cdot)) &\geq \mathbb{E} \langle P(t)(\xi - \mathbb{E}[\xi]) + 2\eta(t), \xi - \mathbb{E}[\xi] \rangle + \langle \Pi(t)\mathbb{E}[\xi] + 2\bar{\eta}(t), \mathbb{E}[\xi] \rangle \\
 &\quad + \mathbb{E} \int_t^T \left\{ \langle P\sigma, \sigma \rangle + 2\langle \eta, b - \mathbb{E}[b] \rangle + 2\langle \zeta, \sigma \rangle + 2\langle \bar{\eta}, \mathbb{E}[b] \rangle \right. \\
 &\quad \left. - \langle \Sigma_0(\varphi - \mathbb{E}[\varphi]), \varphi - \mathbb{E}[\varphi] \rangle - \langle \Sigma \bar{\varphi}, \bar{\varphi} \rangle \right\} ds,
 \end{aligned} \tag{5.17}$$

with the equality holding if and only if

$$\begin{cases} u - \mathbb{E}[u] = v = \Theta z + \varphi - \mathbb{E}[\varphi] = \Theta(X - \mathbb{E}[X]) + \varphi - \mathbb{E}[\varphi], \\ \mathbb{E}[u] = \bar{\Theta} y + \bar{\varphi} = \bar{\Theta}\mathbb{E}[X] + \bar{\varphi}, \end{cases}$$

which is also equivalent to

$$u = \Theta(X - \mathbb{E}[X]) + \bar{\Theta}\mathbb{E}[X] + \varphi - \mathbb{E}[\varphi] + \bar{\varphi}.$$

In particular, when $b(\cdot), \sigma(\cdot), g(\cdot), \bar{g}(\cdot), q(\cdot), \bar{q}(\cdot), \rho(\cdot), \bar{\rho}(\cdot) = 0$, we have

$$(\eta(\cdot), \zeta(\cdot)) = (0, 0), \quad \bar{\eta}(\cdot) = 0, \quad \varphi(\cdot) = \bar{\varphi}(\cdot) = 0.$$

Take $\xi = 0$. Then $X(\cdot)$ satisfies

$$\begin{cases} dX(s) = \left\{ A(s)X(s) + \bar{A}(s)\mathbb{E}[X(s)] + B(s)u(s) + \bar{B}(s)\mathbb{E}[u(s)] \right\} ds \\ \quad + \left\{ C(s)X(s) + \bar{C}(s)\mathbb{E}[X(s)] + D(s)u(s) + \bar{D}(s)\mathbb{E}[u(s)] \right\} dW(s), & s \in [t, T], \\ X(t) = 0, \end{cases}$$

and (5.16) becomes

$$\begin{aligned}
 J^0(t, 0; u(\cdot)) &= \mathbb{E} \int_t^T \left\{ \langle \Sigma_0[u - \mathbb{E}[u] - \Theta(X - \mathbb{E}[X])], u - \mathbb{E}[u] - \Theta(X - \mathbb{E}[X]) \rangle \right. \\
 &\quad \left. + \langle \Sigma(\mathbb{E}[u] - \bar{\Theta}\mathbb{E}[X]), \mathbb{E}[u] - \bar{\Theta}\mathbb{E}[X] \rangle \right\} ds.
 \end{aligned} \tag{5.18}$$

Noting that $\Sigma_0, \Sigma \geq \delta I$ for some $\delta > 0$ and making use of Lemma 5.1, we have

$$\begin{aligned} J^0(t, 0; u(\cdot)) &\geq \delta \mathbb{E} \int_t^T \left\{ |u - \mathbb{E}[u] - \Theta(X - \mathbb{E}[X])|^2 + |\mathbb{E}[u] - \bar{\Theta}\mathbb{E}[X]|^2 \right\} ds \\ &\geq \delta \mathbb{E} \int_t^T |u - \Theta(X - \mathbb{E}[X])|^2 - 2\langle u - \Theta(X - \mathbb{E}[X]), \mathbb{E}[u] \rangle + (1 + \gamma)|\mathbb{E}[u]|^2 ds \\ &\geq \frac{\delta\gamma}{1 + \gamma} \mathbb{E} \int_t^T |u - \Theta(X - \mathbb{E}[X])|^2 ds \\ &\geq \frac{\delta\gamma^2}{1 + \gamma} \mathbb{E} \int_t^T |u(s)|^2 ds, \quad \forall u(\cdot) \in \mathcal{U}[t, T], \end{aligned}$$

for some $\gamma > 0$. The uniform convexity of $u(\cdot) \mapsto J^0(t, 0; u(\cdot))$ follows immediately. □

Note that for Problem (MF-LQ)⁰ (where $b(\cdot), \sigma(\cdot), g(\cdot), \bar{g}(\cdot), q(\cdot), \bar{q}(\cdot), \rho(\cdot), \bar{\rho}(\cdot) = 0$), under the uniform convexity condition (H4), the value at (t, ξ) is given by

$$V^0(t, \xi) = \mathbb{E} \langle P(t)(\xi - \mathbb{E}[\xi]), \xi - \mathbb{E}[\xi] \rangle + \langle \Pi(t)\mathbb{E}[\xi], \mathbb{E}[\xi] \rangle,$$

where $P(\cdot)$ and $\Pi(\cdot)$ are the solutions to the Riccati equations (4.7) and (4.16), respectively. The unique optimal $u^*(\cdot)$ is given by

$$u^* = \Theta(X^* - \mathbb{E}[X^*]) + \bar{\Theta}\mathbb{E}[X^*],$$

where $\Theta(\cdot), \bar{\Theta}(\cdot)$ are defined by (5.6) and $X^*(\cdot)$ is the solution of

$$\begin{cases} dX^*(s) = \left\{ (A + B\Theta)(X^* - \mathbb{E}[X^*]) + [(A + \bar{A}) + (B + \bar{B})\bar{\Theta}]\mathbb{E}[X^*] \right\} ds \\ \quad + \left\{ (C + D\Theta)(X^* - \mathbb{E}[X^*]) + [(C + \bar{C}) + (D + \bar{D})\bar{\Theta}]\mathbb{E}[X^*] \right\} dW(s), & s \in [t, T], \\ X^*(t) = \xi. \end{cases}$$

To conclude this section, we present a sufficient condition for the uniform convexity of the cost functional. From the following result, we will see that (1.4) implies the uniform convexity condition (H4). However, the converse fails. A counterexample will be present in the next section (see Example 6.1).

Proposition 5.3. *Let (H1)–(H2) hold and $t \in [0, T]$ be given. If there exists a constant $\delta > 0$ such that*

$$\begin{cases} G, G + \bar{G} \geq 0, & R(s), R(s) + \bar{R}(s) \geq \delta I, & Q(s) - S(s)^\top R(s)^{-1} S(s) \geq 0, \\ Q(s) + \bar{Q}(s) - [S(s) + \bar{S}(s)]^\top [R(s) + \bar{R}(s)]^{-1} [S(s) + \bar{S}(s)] \geq 0, & \text{a.e. } s \in [t, T], \end{cases} \quad (5.19)$$

then the map $u(\cdot) \mapsto J^0(t, 0; u(\cdot))$ is uniformly convex.

Proof. For any $u(\cdot) \in \mathcal{U}[t, T]$, let $X_0^u(\cdot)$ be the solution of (5.1). Then

$$\begin{aligned} J^0(t, 0; u(\cdot)) &= \mathbb{E} \left\{ \langle G(X_0^u(T) - \mathbb{E}[X_0^u(T)]), X_0^u(T) - \mathbb{E}[X_0^u(T)] \rangle \right. \\ &\quad + \int_t^T \left\langle \begin{pmatrix} Q & S^\top \\ S & R \end{pmatrix} \begin{pmatrix} X_0^u - \mathbb{E}[X_0^u] \\ u - \mathbb{E}[u] \end{pmatrix}, \begin{pmatrix} X_0^u - \mathbb{E}[X_0^u] \\ u - \mathbb{E}[u] \end{pmatrix} \right\rangle ds \Big\} \\ &\quad + \langle (G + \bar{G})\mathbb{E}[X_0^u(T)], \mathbb{E}[X_0^u(T)] \rangle \\ &\quad + \int_t^T \left\langle \begin{pmatrix} Q + \bar{Q} & (S + \bar{S})^\top \\ S + \bar{S} & R + \bar{R} \end{pmatrix} \begin{pmatrix} \mathbb{E}[X_0^u] \\ \mathbb{E}[u] \end{pmatrix}, \begin{pmatrix} \mathbb{E}[X_0^u] \\ \mathbb{E}[u] \end{pmatrix} \right\rangle ds \end{aligned}$$

$$\begin{aligned}
 &\geq \mathbb{E} \int_t^T \left\{ \langle Q(X_0^u - \mathbb{E}[X_0^u]), X_0^u - \mathbb{E}[X_0^u] \rangle + 2 \langle S(X_0^u - \mathbb{E}[X_0^u]), u - \mathbb{E}[u] \rangle + \langle R(u - \mathbb{E}[u]), u - \mathbb{E}[u] \rangle \right\} ds \\
 &\quad + \int_t^T \left\{ \langle (Q + \bar{Q})\mathbb{E}[X_0^u], \mathbb{E}[X_0^u] \rangle + 2 \langle (S + \bar{S})\mathbb{E}[X_0^u], \mathbb{E}[u] \rangle + \langle (R + \bar{R})\mathbb{E}[u], \mathbb{E}[u] \rangle \right\} ds \\
 &= \mathbb{E} \int_t^T \left\{ \langle (Q - S^\top R^{-1}S)(X_0^u - \mathbb{E}[X_0^u]), X_0^u - \mathbb{E}[X_0^u] \rangle \right. \\
 &\quad \left. + \langle R[u - \mathbb{E}[u] + R^{-1}S(X_0^u - \mathbb{E}[X_0^u])], u - \mathbb{E}[u] + R^{-1}S(X_0^u - \mathbb{E}[X_0^u]) \rangle \right\} ds \\
 &\quad + \int_t^T \left\{ \langle [Q + \bar{Q} - (S + \bar{S})^\top (R + \bar{R})^{-1} (S + \bar{S})]\mathbb{E}[X_0^u], \mathbb{E}[X_0^u] \rangle \right. \\
 &\quad \left. + \langle (R + \bar{R})\left(\mathbb{E}[u] + (R + \bar{R})^{-1}(S + \bar{S})\mathbb{E}[X_0^u]\right), \mathbb{E}[u] + (R + \bar{R})^{-1}(S + \bar{S})\mathbb{E}[X_0^u] \rangle \right\} ds \\
 &\geq \delta \mathbb{E} \int_t^T |u - \mathbb{E}[u] + R^{-1}S(X_0^u - \mathbb{E}[X_0^u])|^2 ds + \delta \int_t^T |\mathbb{E}[u] + (R + \bar{R})^{-1}(S + \bar{S})\mathbb{E}[X_0^u]|^2 ds.
 \end{aligned}$$

Consequently, by Lemma 5.1 (taking $\Theta = -R^{-1}S$ and $\bar{\Theta} = -(R + \bar{R})^{-1}(S + \bar{S})$), we have

$$\begin{aligned}
 J^0(t, 0; u(\cdot)) &\geq \delta \mathbb{E} \int_t^T \left\{ |u - \mathbb{E}[u] + R^{-1}S(X_0^u - \mathbb{E}[X_0^u])|^2 + \gamma |\mathbb{E}[u]|^2 \right\} ds \\
 &\geq \frac{\delta\gamma}{1 + \gamma} \mathbb{E} \int_t^T |u + R^{-1}S(X_0^u - \mathbb{E}[X_0^u])|^2 ds \\
 &\geq \frac{\delta\gamma^2}{1 + \gamma} \mathbb{E} \int_t^T |u(s)|^2 ds, \quad \forall u(\cdot) \in \mathcal{U}[t, T],
 \end{aligned}$$

for some $\gamma > 0$. This completes the proof. □

6. EXAMPLES

In this section we present two illustrative examples. In the first example, the condition (1.4) does not hold, but the corresponding Riccati equations are still solvable. Thus, by Theorem 5.2, the cost functional is uniformly convex. This example shows that the uniform convexity condition (H4) is indeed weaker than (1.4).

Example 6.1. Consider the following Problem (MF-LQ)⁰ with one-dimensional state equation

$$\begin{cases} dX(s) = \{ \mathbb{E}[X(s)] + u(s) + \mathbb{E}[u(s)] \} ds + \sqrt{2}u(s)dW(s), & s \in [t, 1], \\ X(t) = \xi, \end{cases}$$

and cost functional

$$J(t, \xi; u(\cdot)) = \mathbb{E} \left\{ G|X(1)|^2 + \bar{G}|\mathbb{E}[X(1)]|^2 + \int_t^1 \left(R(s)|u(s)|^2 + \bar{R}(s)|\mathbb{E}[u(s)]|^2 \right) ds \right\},$$

where

$$\begin{cases} G = 8, & \bar{G} = -\alpha - 8 \quad \text{with} \quad 0 < \alpha < \frac{1}{2(e^2 - 1)}, \\ R(s) = (s + 1)^3 - 4(s + 1)^2, & \bar{R}(s) = 1 - (s + 1)^3, \quad s \in [0, 1]. \end{cases}$$

The Riccati equations for the above problem are

$$\begin{cases} \dot{P}(s) - \frac{P(s)^2}{R(s) + 2P(s)} = 0, \\ P(1) = 8, \end{cases}$$

and

$$\begin{cases} \dot{\Pi}(s) + 2\Pi(s) - \frac{4\Pi(s)^2}{R(s) + \bar{R}(s) + 2P(s)} = 0, \\ \Pi(1) = -\alpha. \end{cases}$$

Clearly,

$$G + \bar{G} = -\alpha < 0, \quad R(s) = (s + 1)^2(s - 3) \leq -2, \quad R(s) + \bar{R}(s) = 1 - 4(s + 1)^2 \leq -3, \quad s \in [0, 1].$$

Hence, the condition (1.4) does not hold. However, one can verify that the above Riccati equations are solvable on the whole interval $[0, 1]$ with solutions given by

$$P(s) = 2(s + 1)^2, \quad \Pi(s) = \frac{\alpha e^{2(1-s)}}{2\alpha[e^{2(1-s)} - 1] - 1} < 0, \quad s \in [0, 1].$$

Note that

$$\begin{cases} R(s) + D(s)^\top P(s)D(s) = (s + 1)^3 \geq 1, \\ R(s) + \bar{R}(s) + [D(s) + \bar{D}(s)]^\top P(s)[D(s) + \bar{D}(s)] = 1, \end{cases} \quad s \in [0, 1].$$

By Theorem 5.2, the cost functional $J(t, \xi; u(\cdot))$ is uniformly convex in $u(\cdot)$, and for any initial pair $(t, \xi) \in [0, 1] \times L^2_{\mathcal{F}_t}(\Omega; \mathbb{R})$, the problem admits a unique optimal control $u^*(\cdot)$ given by

$$u^*(s) = -\frac{2}{s + 1}X^*(s) + \left[\frac{2}{s + 1} - 2\Pi(s) \right] \mathbb{E}[X^*(s)], \quad s \in [t, 1]$$

with $X^*(\cdot)$ being the solution to the following closed-loop system:

$$\begin{cases} dX^*(s) = \left\{ -\frac{2}{s + 1}X^*(s) + \left[\frac{s + 3}{s + 2} - 4\Pi(s) \right] \mathbb{E}[X^*(s)] \right\} ds \\ \quad + \left\{ -\frac{2\sqrt{2}}{s + 2}X^*(s) + 2\sqrt{2} \left[\frac{1}{s + 1} - \Pi(s) \right] \mathbb{E}[X^*(s)] \right\} dW(s), \quad s \in [t, 1], \\ X^*(t) = \xi. \end{cases}$$

Now we present an example in which the mean-field LQ problem is open-loop solvable, but the cost functional is not uniformly convex. Hence, the optimal control cannot be constructed directly in terms of the Riccati equations. However, an optimal could still be found by making use of Theorems 3.2 and 5.2.

Example 6.2. Consider the following Problem (MF-LQ)⁰ with one-dimensional state equation

$$\begin{cases} dX(s) = \{X(s) - \mathbb{E}[X(s)] + \mathbb{E}[u(s)]\} ds + u(s)dW(s), \quad s \in [t, T], \\ X(t) = \xi, \end{cases}$$

and cost functional

$$J(t, \xi; u(\cdot)) = \mathbb{E} \left\{ 2|X(T)|^2 + |\mathbb{E}[X(T)]|^2 + \int_t^T \left(-4|X(s)|^2 - |u(s)|^2 + 4|\mathbb{E}[X(s)]|^2 - |\mathbb{E}[u(s)]|^2 \right) ds \right\}.$$

In this example,

$$\begin{cases} A = 1, \quad \bar{A} = -1, \quad B = 0, \quad \bar{B} = 1, \quad C = \bar{C} = 0, \quad D = 1, \quad \bar{D} = 0, \\ G = 2, \quad \bar{G} = 1, \quad Q = -4, \quad \bar{Q} = 4, \quad S = \bar{S} = 0, \quad R = \bar{R} = -1. \end{cases}$$

Clearly, the condition (1.4) does not hold. The Riccati equations for the problem are

$$\begin{cases} \dot{P}(s) + 2P(s) - 4 = 0, & s \in [t, T], \\ P(T) = 2, \end{cases} \tag{6.1}$$

and

$$\begin{cases} \dot{\Pi}(s) - \frac{\Pi(s)^2}{P(s) - 2} = 0, & s \in [t, T], \\ \Pi(T) = 3. \end{cases} \tag{6.2}$$

It is easy to see that $P(\cdot) \equiv 2$ is the unique solution of (6.1). However, since

$$R(s) + \bar{R}(s) + [D(s) + \bar{D}(s)]^\top P(s)[D(s) + \bar{D}(s)] = P(s) - 2 = 0, \quad s \in [t, T],$$

we cannot use (6.2) to solve the problem directly. To investigate the open-loop solvability of the above problem, let us now consider the following cost functionals for $\varepsilon > 0$:

$$\begin{aligned} J_\varepsilon(t, \xi; u(\cdot)) &= J(t, \xi; u(\cdot)) + \varepsilon \mathbb{E} \int_t^T |u(s)|^2 ds \\ &= \mathbb{E} \left\{ 2|X(T)|^2 + |\mathbb{E}[X(T)]|^2 + \int_t^T \left(-4|X(s)|^2 + (\varepsilon - 1)|u(s)|^2 + 4|\mathbb{E}[X(s)]|^2 - |\mathbb{E}[u(s)]|^2 \right) ds \right\}. \end{aligned}$$

We denote the corresponding mean-field LQ problem and value function by Problem (MF-LQ) $_\varepsilon^0$ and $V_\varepsilon^0(\cdot, \cdot)$, respectively. The Riccati equations for Problem (MF-LQ) $_\varepsilon^0$ are

$$\begin{cases} \dot{P}_\varepsilon(s) + 2P_\varepsilon(s) - 4 = 0, & s \in [t, T], \\ P_\varepsilon(T) = 2, \end{cases}$$

and

$$\begin{cases} \dot{\Pi}_\varepsilon(s) - \frac{\Pi_\varepsilon(s)^2}{\varepsilon - 2 + P_\varepsilon(s)} = 0, & s \in [t, T], \\ \Pi_\varepsilon(T) = 3. \end{cases}$$

A straightforward calculation leads to

$$P_\varepsilon(s) = 2, \quad \Pi_\varepsilon(s) = \frac{3\varepsilon}{\varepsilon + 3(T - s)}; \quad s \in [t, T].$$

Since

$$R + \varepsilon + D^\top P_\varepsilon D = 1 + \varepsilon, \quad R + \varepsilon + \bar{R} + (D + \bar{D})^\top P_\varepsilon (D + \bar{D}) = \varepsilon,$$

by Theorem 5.2, the map $u(\cdot) \mapsto J_\varepsilon(t, 0; u(\cdot))$ is uniformly convex for all $\varepsilon > 0$ and hence $u(\cdot) \mapsto J(t, 0; u(\cdot))$ is convex. Moreover,

$$V_\varepsilon^0(t, \xi) = \mathbb{E} \langle P_\varepsilon(t)(\xi - \mathbb{E}[\xi]), \xi - \mathbb{E}[\xi] \rangle + \langle \Pi_\varepsilon(t) \mathbb{E}[\xi], \mathbb{E}[\xi] \rangle, \tag{6.3}$$

and the unique optimal control of Problem (MF-LQ) $_\varepsilon^0$ at (t, ξ) is given by

$$u_\varepsilon^*(s) = -\frac{\Pi_\varepsilon(s)}{\varepsilon} \mathbb{E}[X_\varepsilon^*(s)], \quad s \in [t, T],$$

with $X_\varepsilon^*(\cdot)$ being the solution to the following closed-loop system:

$$\begin{cases} dX_\varepsilon^*(s) = \left\{ X_\varepsilon^*(s) - \left(1 + \frac{\Pi_\varepsilon(s)}{\varepsilon} \right) \mathbb{E}[X_\varepsilon^*(s)] \right\} ds - \frac{\Pi_\varepsilon(s)}{\varepsilon} \mathbb{E}[X_\varepsilon^*(s)] dW(s), & s \in [t, T], \\ X_\varepsilon^*(t) = \xi. \end{cases}$$

Letting $\varepsilon \rightarrow 0$ in (6.3), we have from Theorem 3.2 that

$$V^0(t, \xi) = \lim_{\varepsilon \rightarrow 0} V_\varepsilon^0(t, \xi) = \begin{cases} 2\text{var}[\xi], & 0 \leq t < T, \\ 2\mathbb{E}[\xi^2] + (\mathbb{E}[\xi])^2, & t = T. \end{cases}$$

Note that

$$\begin{cases} d\mathbb{E}[X_\varepsilon^*(s)] = -\frac{\Pi_\varepsilon(s)}{\varepsilon} \mathbb{E}[X_\varepsilon^*(s)] ds, & s \in [t, T], \\ \mathbb{E}[X_\varepsilon^*(t)] = \mathbb{E}[\xi]. \end{cases}$$

Hence,

$$\mathbb{E}[X_\varepsilon^*(s)] = \mathbb{E}[\xi] \exp \left\{ -\int_t^s \frac{\Pi_\varepsilon(r)}{\varepsilon} dr \right\} = \frac{\varepsilon + 3(T-s)}{\varepsilon + 3(T-t)} \mathbb{E}[\xi], \quad s \in [t, T],$$

and

$$u_\varepsilon^*(s) = -\frac{\Pi_\varepsilon(s)}{\varepsilon} \mathbb{E}[X_\varepsilon^*(s)] = -\frac{3\mathbb{E}[\xi]}{\varepsilon + 3(T-t)}, \quad s \in [t, T].$$

It is clear that for $t \in [0, T]$, $u_\varepsilon^*(s)$ converges uniformly to

$$u^*(s) \equiv -\frac{\mathbb{E}[\xi]}{T-t}, \quad s \in [t, T],$$

which, by Theorem 3.2, is an optimal control of the original problem at (t, ξ) .

Acknowledgements. The author wishes to thank Prof. Jiongmin Yong for his valuable comments, which have helped to improve the quality of the manuscript. The author also would like to thank Dr. Xun Li for his useful suggestions and financial support.

REFERENCES

- [1] N.U. Ahmed, Nonlinear diffusion governed by McKean–Vlasov equation on Hilbert space and optimal control. *SIAM J. Control Optim.* **46** (2007) 356–378.
- [2] N.U. Ahmed and X. Ding, A semilinear McKean–Vlasov stochastic evolution equation in Hilbert space. *Stoch. Proc. Appl.* **60** (1995) 65–85.
- [3] N.U. Ahmed and X. Ding, Controlled McKean–Vlasov equations. *Commun. Appl. Anal.* **5** (2001) 183–206.
- [4] D. Andersson and B. Djehiche, A maximum principle for SDEs of mean-field type. *Appl. Math. Optim.* **63** (2011) 341–356.
- [5] V.S. Borkar and K.S. Kumar, McKean–Vlasov limit in portfolio optimization. *Stoch. Anal. Appl.* **28** (2010) 884–906.
- [6] R. Buckdahn, B. Djehiche and J. Li, A general maximum principle for SDEs of mean-field type. *Appl. Math. Optim.* **64** (2011) 197–216.
- [7] R. Buckdahn, B. Djehiche, J. Li and S. Peng, Mean-field backward stochastic differential equations: A limit approach. *Ann. Probab.* **37** (2009) 1524–1565.
- [8] R. Buckdahn, J. Li and S. Peng, Mean-field backward stochastic differential equations and related partial differential equations. *Stoch. Process. Appl.* **119** (2009) 3133–3154.
- [9] T. Chan, Dynamics of the McKean–Vlasov equation. *Ann. Probab.* **22** (1994) 431–441.
- [10] T. Chiang, McKean–Vlasov equations with discontinuous coefficients. *Soochow J. Math.* **20** (1994) 507–526.
- [11] D. Crisan and J. Xiong, Approximate McKean–Vlasov representations for a class of SPDEs. *Stochastics* **82** (2010) 53–68.
- [12] D.A. Dawson, Critical dynamics and fluctuations for a mean-field model of cooperative behavior. *J. Statist. Phys.* **31** (1983) 29–85.
- [13] D.A. Dawson and J. Gärtner, Large deviations from the McKean–Vlasov limit for weakly interacting diffusions. *Stochastics* **20** (1987) 247–308.
- [14] J. Gärtner, On the McKean–Vlasov limit for interacting diffusions. *Math. Nachr.* **137** (1988) 197–248.
- [15] C. Graham, McKean–Vlasov Ito-Skorohod equations, and nonlinear diffusions with discrete jump sets. *Stoch. Proc. Appl.* **40** (1992) 69–82.
- [16] J. Huang, X. Li, and J. Yong, A linear-quadratic optimal control problem for mean-field stochastic differential equations in infinite horizon. *Math. Control Relat. Fields* **5** (2015) 97–139.
- [17] M. Huang, R.P. Malhamé, and P.E. Caines, Large population stochastic dynamic games: Closed-loop McKean–Vlasov systems and the Nash certainty equivalence principle. *Commun. Inf. Syst.* **6** (2006) 221–251.

- [18] M. Kac, Foundations of kinetic theory. *Proc. of 3rd Berkeley Sympos. Math. Statist. Prob.* **3** (1956) 171–197.
- [19] P.M. Kotelenetz and T.G. Kurtz, Macroscopic limit for stochastic partial differential equations of McKean–Vlasov type. *Prob. Theory Relat. Fields* **146** (2010) 189–222.
- [20] N. I. Mahmudov and M.A. McKibben, On a class of backward McKean–Vlasov stochastic equations in Hilbert space: Existence and convergence properties. *Dynam. Syst. Appl.* **16** (2007) 643–664.
- [21] H.P. McKean, A class of Markov processes associated with nonlinear parabolic equations. *Proc. Natl. Acad. Sci. USA* **56** (1966) 1907–1911.
- [22] T. Meyer-Brandis, B. Øksendal and X. Y. Zhou, A mean-field stochastic maximum principle via Malliavin calculus. *Stochastics* **84** (2012) 643–666.
- [23] J.Y. Park, P. Balasubramaniam and Y.H. Kang, Controllability of McKean–Vlasov stochastic integrodifferential evolution equation in Hilbert spaces. *Numer. Funct. Anal. Optim.* **29** (2008) 1328–1346.
- [24] M. Scheutzow, Uniqueness and non-uniqueness of solutions of Vlasov-McKean equations. *J. Austral. Math. Soc., Ser. A* **43** (1987) 246–256.
- [25] J. Sun and J. Yong, Linear Quadratic Stochastic Differential Games: Open-Loop and Closed-Loop Saddle Points. *SIAM J. Control Optim.* **52** (2014) 4082–4121.
- [26] J. Sun, X. Li and J. Yong, Open-Loop and Closed-Loop Solvabilities for Stochastic Linear Quadratic Optimal Control Problems. *SIAM J. Control Optim.* **54** (2016) 2274–2308.
- [27] A. Yu. Veretennikov, On ergodic measures for McKean–Vlasov stochastic equations, in Monte Carlo and quasi-Monte Carlo methods 2004. Springer, Berlin (2006) 471–486.
- [28] J. Yong, Linear-Quadratic Optimal Control Problems for Mean-Field Stochastic Differential Equations. *SIAM J. Control Optim.* **51** (2013) 2809–2838.
- [29] J. Yong and X.Y. Zhou, Stochastic Controls: Hamiltonian Systems and HJB Equations. Springer-Verlag, New York (1999).