

SHARP SUMMABILITY FOR MONGE TRANSPORT DENSITY VIA INTERPOLATION

LUIGI DE PASCALE¹ AND ALDO PRATELLI²

Abstract. Using some results proved in De Pascale and Pratelli [*Calc. Var. Partial Differ. Equ.* **14** (2002) 249-274] (and De Pascale *et al.* [*Bull. London Math. Soc.* **36** (2004) 383-395]) and a suitable interpolation technique, we show that the transport density relative to an L^p source is also an L^p function for any $1 \leq p \leq +\infty$.

Mathematics Subject Classification. 41A05, 49N60, 49Q20, 90B06.

Received May 5, 2003.

This paper is concerned with the transport problem, which consists in minimizing

$$\int_{\Omega} |x - t(x)| \, df^+(x) \tag{1}$$

among the *transports*, which are the measurable functions $t : \text{spt}(f^+) \rightarrow \text{spt}(f^-)$ such that $t_{\#}f^+ = f^-$, *i.e.* for any Borel set B it is $f^+(t^{-1}(B)) = f^-(B)$; here $f = f^+ - f^-$ is a L^1 function on Ω with $\int f = 0$, while Ω is a convex and bounded subset of \mathbb{R}^N (to find more general descriptions of the transport problem, see [1, 9]). To each optimal – *i.e.* minimizing (1) – transport t it is possible to associate a positive measure σ on Ω defined by

$$\langle \sigma, \varphi \rangle := \int_{\Omega} \left(\int_{\Omega} \varphi(z) \, d\mathcal{H}_{t(x)}^1(z) \right) \, df^+(x) \tag{2}$$

where φ is any function in $C_0(\Omega)$ and \mathcal{H}_{xy}^1 is the one-dimensional Hausdorff measure on the segment xy . It has been proved (see [1, 8]) that there always exist (in this setting) optimal transports and in particular there are invertible optimal transports whose inverse is also an optimal transport for $-f$. A fundamental result, due to [1, 8], is that, even if there can be many different optimal transports, all define *via* (2) the same measure σ , which is then called *transport density* relative to f . This measure is very interesting for the transport problem (for example it plays an important role in [7]), and moreover it represents the connection between this problem and some shape optimization problem (see [3, 4]), which can be reduced to the research of a positive measure σ

Keywords and phrases. Transport density, interpolation, summability.

¹ Dipartimento di Matematica Applicata, Università di Pisa, via Bonanno Pisano 25/B, 56126 Pisa, Italy; e-mail: depascal@dm.unipi.it

² Scuola Normale Superiore, Piazza dei Cavalieri 7, 56126 Pisa, Italy; e-mail: a.pratelli@sns.it

and a 1-Lipschitz function u solving

$$\begin{cases} -\operatorname{div}(\sigma Du) = f & \text{on } \Omega \\ |Du| = 1 & \sigma - \text{a.e.} \end{cases} \tag{3}$$

The relationship between the two problems relies on the fact that the (unique) transport density is also the unique solution of (3) (see [1,3,6]); the functions u solving (3) together with σ are also meaningful in the context of the transport problem, they are referred to as *Kantorovich potentials*. Equation (3) is often referred to as *Monge-Kantorovich equation*. Thus the study of the regularity of σ is useful both for the transport problem and for the shape optimization problem. It was proved (see [1,6,8]) that the fact that $f \in L^1$ implies also that $\sigma \in L^1$.

In this paper we will show some sharp relationship between the summability of f and that of σ . The problem to derive regularity of σ from that of f has already been studied in [5,6] following two different methods: in [6] we used a geometric construction starting from the definition (2), while in [5] the proofs used PDE tools starting from the equivalent definition (3). In the first work it was proved that

$$\begin{aligned} f \in L^1 &\implies \sigma \in L^1, & f \in L^\infty &\implies \sigma \in L^\infty, \\ f \in L^p &\implies \sigma \in L^{p-\epsilon} \text{ for any } \epsilon > 0, \end{aligned} \tag{4}$$

and some examples were given in which $f \in L^p$ and $\sigma \notin L^q$ for any $q > p$. Thus it was left open the problem whether or not it is true that $f \in L^p$ implies $\sigma \in L^p$ for $p \neq 1, +\infty$. In the second work this problem was partially solved, since it was proved that

$$f \in L^p \implies \sigma \in L^p \text{ for any } 2 \leq p < +\infty. \tag{5}$$

Since the cases $p = 1, \infty$ had already been solved in the first work, it was left open only the case with $1 < p < 2$. In this work we will show how the classic Marcinkiewicz interpolation result can be used to infer from the results already mentioned the general property for any p . Note that this is not trivial since the map associating the transport density σ to any function f with $\int f = 0$ is far from being linear or sublinear, as easy examples show; however, this map is 1-homogeneous, as one can hope in view of (6).

The result of this paper is the following

Theorem A. *For any $1 \leq p \leq +\infty$, if $f \in L^p$ is a function with $\int f = 0$, then the associated transport density σ is also in L^p . More precisely, there exist a constant C_p , depending only on Ω , such that*

$$\|\sigma\|_{L^p} \leq C_p \|f\|_{L^p}. \tag{6}$$

To prove the theorem, we first of all recall the known results we use, the regularity results proved in [5,6] and the Marcinkiewicz interpolation result, which can be found for example in [10].

Theorem 1. *For $p = 1$ and for any $p \geq 2$ there exists a constant C_p depending on Ω such that, for any $f \in L^p$ with $\int f = 0$,*

$$\|\sigma\|_{L^p} \leq C_p \|f\|_{L^p} \tag{7}$$

where σ is the transport density associated to f .

Theorem 2 (Marcinkiewicz). *If $T : L^1 \rightarrow L^1$ is a linear mapping such that, for two suitable constants M_p and M_q with $1 \leq p < q \leq +\infty$,*

$$\|T(g)\|_{L^p} \leq M_p \|g\|_{L^p} \quad \text{and} \quad \|T(g)\|_{L^q} \leq M_q \|g\|_{L^q}, \tag{8}$$

then it is also true that for any $s \in (p, q)$

$$\|T(g)\|_{L^s} \leq C M_p^{\frac{p(q-s)}{s(q-p)}} M_q^{\frac{q(s-p)}{s(q-p)}} \|g\|_{L^s}, \tag{9}$$

where C is a geometric constant depending only on Ω .

To prove our result, let us fix now a function $f \in L^p$ with $\int f = 0$ and $1 \leq p \leq +\infty$. We can assume $p \neq 1, +\infty$, since otherwise we already know that $\sigma \in L^p$. Fix also an invertible optimal transport t for f (as we said, this always exists when $f \in L^1$, even though it is not unique). For any function $g \in L^1$, there are of course two uniquely determined measurable functions λ and ν supported respectively on $\text{spt}(f^+)$ and on $\Omega \setminus \text{spt}(f^+)$ such that

$$g = \lambda f^+ + \nu. \tag{10}$$

Let us finally define the operator $T : L^1 \rightarrow \mathcal{M}(\Omega)$ to which we will apply later the Marcinkiewicz theorem: given any $g \in L^1$ and following the notations of (10), we define

$$T(g) := \int_{\Omega} \lambda(x) f^+(x) \mathcal{H}_{xt(x)}^1 dx, \tag{11}$$

which can also be rewritten as

$$\langle T(g), \varphi \rangle = \int_{x \in \Omega} \left(\int_{z \in \Omega} \varphi(z) d\mathcal{H}_{xt(x)}^1(z) \right) \lambda(x) f^+(x) dx$$

for any $\varphi \in C_0(\Omega)$. Notice that $T(g)$ is *a priori* a measure, and that the definition of T depends on the function f we fixed; moreover, we point out that of course $T(f)$ is the transport density σ associated to f (just recall (2) and (11)).

We define now σ_1, σ_2 and f_1, f_2 (depending on f and g) as follows:

$$\begin{aligned} \sigma_1 &:= \int_{\Omega} \lambda^+(x) f^+(x) \mathcal{H}_{xt(x)}^1 dx & f_1 &:= \lambda^+ f^+ - (\lambda^+ \circ t^{-1}) f^- \\ \sigma_2 &:= \int_{\Omega} \lambda^-(x) f^+(x) \mathcal{H}_{xt(x)}^1 dx & f_2 &:= \lambda^- f^+ - (\lambda^- \circ t^{-1}) f^-; \end{aligned} \tag{12}$$

note that also these definitions depend on f and g , and that $T(g) = \sigma_1 + \sigma_2$. First we prove the

Lemma 3. *The function $t : \text{spt}(f^+) \rightarrow \text{spt}(f^-)$ is defined $f_i^+ - a.e.$ and it is an optimal transport for the functions $f_i, i = 1, 2$ defined in (12); moreover, each σ_i is the transport density associated to f_i .*

Proof. The optimality of a transport is equivalent to the cyclical monotonicity of its graph (see [2, 9] to find the definition of the cyclical monotonicity and the proof of this assert). Then the fact that t is optimal for f assures that its graph is monotonically cyclic; thus, given any function h on Ω with 0 mean and such that $h^+ \ll f^+, t$ is defined $h^+ - a.e.$ and it is an optimal transport for h if and only if it is a transport. Then to prove the first part of the assert it is enough to check that $t_{\#} f_i^+ = f_i^-$ for $i = 1, 2$, which is a straightforward consequence of the fact that $t_{\#} f^+ = f^-$ and of the properties of the push-forward. Finally, the fact that each σ_i is the transport density associated to f_i follows comparing (12) with the definition (2) of the transport density (replace f and σ in (2) by f_i and σ_i). \square

We can then prove the following

Lemma 4. *$T : L^1 \rightarrow L^1$ is a linear operator.*

Proof. The fact that T is linear follows immediately from the definition (11); moreover $T(g)$ is a L^1 function (recall that *a priori* we knew it only to be a measure) since it is the sum of the two transport densities σ_i thanks to the preceding Lemma, and thanks to Theorem 1 each of these densities is in L^1 since so is each f_i – recall (12) and that $g \in L^1$. \square

We prove now the validity of (8) with $p = 1$ and $q = +\infty$ in order to apply the Marcinkiewicz Theorem.

Lemma 5. *The inequalities (8) hold for T with $p = 1$ and $q = +\infty$; in particular, $M_1 = 2C_1$ and $M_\infty = 2C_\infty$, where the C_i 's are the constants of (7).*

Proof. In view of Lemma 3 and Theorem 1, σ_1 is the transport density relative to f_1 and then $\|\sigma_1\|_{L^1} \leq C_1\|f_1\|_{L^1}$; but since $t_{\#}f_1^+ = f_1^-$, then $\|f_1^+\|_{L^1} = \|f_1^-\|_{L^1}$ and we infer

$$\|\sigma_1\|_{L^1} \leq 2C_1\|f_1^+\|_{L^1}.$$

In the same way we deduce also $\|\sigma_2\|_{L^1} \leq 2C_1\|f_2^-\|_{L^1}$. Using now the fact that the supports of the f_i 's are essentially disjoint – that is clear from (12) –, we have

$$\begin{aligned} \|T(g)\|_{L^1} &= \|\sigma_1 + \sigma_2\|_{L^1} \leq \|\sigma_1\|_{L^1} + \|\sigma_2\|_{L^1} \leq 2C_1(\|f_1^+\|_{L^1} + \|f_2^+\|_{L^1}) \\ &= 2C_1\|f_1^+ + f_2^+\|_{L^1} = 2C_1\|\lambda^+f^+ + \lambda^-f^+\|_{L^1} = 2C_1\|\lambda f^+\|_{L^1} \\ &\leq 2C_1\|g\|_{L^1}, \end{aligned}$$

which gives the first estimate.

On the other hand, to show the L^∞ inequality we note that, thanks to (10) and (12), it is $\|f_i\|_{L^\infty} \leq \|g\|_{L^\infty}$ for each i . Since $T(g) = \sigma_1 + \sigma_2$, from Lemma 3 and Theorem 1 we infer

$$\|T(g)\|_{L^\infty} \leq 2C_\infty\|g\|_{L^\infty},$$

and then also the L^∞ inequality follows. \square

Thanks to Lemmas 4 and 5, we can apply Theorem 2 to prove (6), recalling that $T(f)$ is the transport density σ associated to f . Recall now that the function $f \in L^p$ was fixed at the beginning, but the constants C_p we obtained do not depend on f , but only on p and Ω . Then the estimate (6) is true, with the same constants, for any function $f \in L^p$.

REFERENCES

- [1] L. Ambrosio, Mathematical Aspects of Evolving Interfaces. *Lect. Notes Math.* **1812** (2003) 1-52.
- [2] L. Ambrosio and A. Pratelli, Existence and stability results in the L^1 theory of optimal transportation. *Lect. Notes Math.* **1813** (2003) 123-160.
- [3] G. Bouchitté and G. Buttazzo, Characterization of optimal shapes and masses through Monge-Kantorovich equation. *J. Eur. Math. Soc.* **3** (2001) 139-168.
- [4] G. Bouchitté, G. Buttazzo and P. Seppecher, Shape Optimization Solutions via Monge-Kantorovich Equation. *C. R. Acad. Sci. Paris I* **324** (1997) 1185-1191.
- [5] L. De Pascale, L.C. Evans and A. Pratelli, Integral Estimates for Transport Densities. *Bull. London Math. Soc.* **36** (2004) 383-395.
- [6] L. De Pascale and A. Pratelli, Regularity properties for Monge transport density and for solutions of some shape optimization problem. *Calc. Var. Partial Differ. Equ.* **14** (2002) 249-274.
- [7] L.C. Evans and W. Gangbo, Differential Equations Methods for the Monge-Kantorovich Mass Transfer Problem. *Mem. Amer. Math. Soc.* **137** (1999).
- [8] M. Feldman and R. McCann, Uniqueness and transport density in Monge's mass transportation problem. *Calc. Var. Partial Differ. Equ.* **15** (2002) 81-113.
- [9] W. Gangbo and R.J. McCann, The geometry of optimal transportation. *Acta Math.* **177** (1996) 113-161.
- [10] M. Giaquinta, *Introduction to regularity theory for nonlinear elliptic systems*. Birkhäuser Verlag (1993).