

MÉMOIRES DE LA S. M. F.

ANDRZEJ PROSZYNSKI

Some functors related to polynomial theory. II

Mémoires de la S. M. F., tome 59 (1979), p. 125-129

http://www.numdam.org/item?id=MSMF_1979__59__125_0

© Mémoires de la S. M. F., 1979, tous droits réservés.

L'accès aux archives de la revue « Mémoires de la S. M. F. » (<http://smf.emath.fr/Publications/Memoires/Presentation.html>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

SOME FUNCTORS RELATED TO POLYNOMIAL THEORY, II

by

Andrzej PRÓSZYŃSKI

1. Introduction. We consider the following natural transformation :

$$T^m : \mathcal{P}_R^m(X, Y) \rightarrow \text{Map}(X, Y), \quad T^m(f) = f_R$$

where R denotes a commutative ring with 1, X, Y - R -modules, and $\mathcal{P}_R^m(X, Y)$ is the R -module of all forms of degree m on the pair (X, Y) (in the sense of N. Roby [2]).

An element of $\mathcal{P}_R^m(X, Y)$ is a system $f = (f_A)$ indexed by all commutative R -algebras A , where $f_A : X \otimes A \rightarrow Y \otimes A$ are mappings satisfying the following conditions :

(i) $(1 \otimes u) \circ f_A = f_B \circ (1 \otimes u)$ for any R -algebra homomorphism $u : A \rightarrow B$,

(ii) $f_A(xa) = f_A(x)a^m$ for any R -algebra A , any $x \in X \otimes A$ and $a \in A$.

It is proved in [1] that in the case $X = R^n, Y = R$ we obtain :

$$T^m : R[T_1, \dots, T_n]_m \rightarrow \text{Map}(R^n, R), \quad T^m(F)(x_1, \dots, x_n) = F(x_1, \dots, x_n).$$

It is well-known that the above homomorphism is not always injective ; this is the starting point and the motivation of the following considerations.

It is known from [2] that the functor $\mathcal{P}_R^m(X, -)$ is represented by the m -th divided power $\Gamma_R^m(X)$ of the module X . Similarly, it is proved in [1] that $\mathcal{P}_R^m(X, -) = \text{Ker } T^m$ is represented by $\hat{\Gamma}_R^m(X)$ where :

$$\hat{\Gamma}_R^m(X) = \Gamma_R^m(X) / R\{x^{(m)}; x \in X\}.$$

The above module is generated by the classes of elements :

$$y_{m_1, \dots, m_k}^{(x_1, \dots, x_k)} = x_1^{(m_1)} \dots x_k^{(m_k)}, \quad m_i \geq 0, \quad m_1 + \dots + m_k = m, \quad x_1, \dots, x_k \in X,$$

which are denoted by $\tilde{y}_{m_1, \dots, m_k}^{(x_1, \dots, x_k)}$.

It is easy to see that $\hat{\Gamma}_R^m$ is an endo-functor of the category $R\text{-Mod}$.

We recall the following results contained in [1] :

Lemma 1.1. $\hat{\Gamma}_R^m$ commutes with direct limits.

Lemma 1.2. $\hat{\Gamma}_R^m(X)$ is finitely generated if so is X .

Theorem 1.3. There exist the natural isomorphisms :

- (1) $\widehat{\Gamma}_R^m(X_S) \approx \widehat{\Gamma}_R^m(X)_S$ for any multiplicative set S in R
 (2) $\widehat{\Gamma}_{R/I}^m(X/IX) \approx \widehat{\Gamma}_R^m(X)/I \widehat{\Gamma}_R^m(X)$ for any ideal I in R
 (3) $\widehat{\Gamma}_{R \times R'}^m(X \times X') \approx \widehat{\Gamma}_R^m(X) \times \widehat{\Gamma}_{R'}^m(X')$.

Theorem 1.4. For a finitely generated R-module X, the following conditions are equivalent :

- (i) $\widehat{\Gamma}_R^m(X) = 0$
 (ii) $\widehat{\Gamma}_{R/P}^m(X/PX) = 0$ for any $P \in \text{Max}(R)$
 (iii) For any $P \in \text{Max}(R)$: either $\dim_{R/P}(X/PX) \leq 1$ or $m \leq |R/P|$.
 In particular, $\widehat{\Gamma}_R^m = 0$ iff $m \leq d(R) : \inf \{|R/P| ; P \in \text{Max}(R)\}$.

2. The structure of $\widehat{\Gamma}_R^m(X)$. We shall give some structural informations on $\widehat{\Gamma}_R^m(X)$ which generalize results contained in [1]. The first step is the following

Lemma 2.1. If $P \in \text{Spec}(R) - \text{Max}(R)$ then $\widehat{\Gamma}_R^m(X)_P = 0$ for any R-module X. Moreover, is X is finitely generated then $\text{Ann}(\widehat{\Gamma}_R^m(X)) \not\subset P$.

Proof : Observe that R/P is an infinite domain (it is not a field!) and hence $d(R_P) = \infty$. It follows from Theorem 1.3 and 1.4 that $\widehat{\Gamma}_R^m(X)_P = \widehat{\Gamma}_{R_P}^m(X_P) = 0$. Then the second part of the lemma follows from Lemma 1.2.

Corollary 2.2. If $\dim(R) > 0$ then :

- (1) $\widehat{\Gamma}_R^m(X)$ are torsion modules.
 (2) $\widehat{\Gamma}_R^m(X)$ is free iff it is zero.
If $\dim(R_P) > 0$ for any $P \in \text{Max}(R)$ then :
 (3) $\widehat{\Gamma}_R^m(X)$ is projective iff it is zero.

Now we explain the structure of $\widehat{\Gamma}_R^m(X)$ over Noetherian rings.

Theorem 2.3. Let R be a Noetherian ring and let X be a finitely generated R-module. Then there exists a natural R-isomorphism :

$$\widehat{\Gamma}_R^m(X) \approx \bigoplus_{P \in \text{Max}(R)} \widehat{\Gamma}_{R/P}^m(X/P \cdot k_P)$$

induced by $X \rightarrow X/P \cdot k_P X$, for all sufficiently large k_P .

Proof : We can assume that $\text{Ann}(\widehat{\Gamma}_R^m(X)) \neq R$. Let $\text{Ann}(\widehat{\Gamma}_R^m(X)) = Q_1 \cap \dots \cap Q_s$ be a primary decomposition, and let $P_i = \text{rad}(Q_i)$. Observe that $P_i \subset Q_i$ for all sufficiently large k_i . Denote $I = P_1^{k_1} \dots P_s^{k_s} \subset \text{Ann}(\widehat{\Gamma}_R^m(X))$. Since $P_1, \dots, P_s \in \text{Max}(R)$ by Lemma 2.1, it follows that $R/I \approx \prod_{i=1}^s R/P_i^{k_i}$ and hence :

$$\widehat{\Gamma}_R^m(X) = \widehat{\Gamma}_R^m(X) / \widehat{\Gamma}_R^m(X) \approx \widehat{\Gamma}_{R/I}^m(X/IX) \approx \bigoplus_{i=1}^s \widehat{\Gamma}_{R/P_i}^m(X/P_i^k X).$$

If $P \in \text{Max}(R) - \{P_1, \dots, P_s\}$ then $I + P^k = R$ for each natural k , and hence :

$$\widehat{\Gamma}_{R/P^k}^m(X/P^k X) = \widehat{\Gamma}_R^m(X)/P^k \widehat{\Gamma}_R^m(X) = 0.$$

This completes the proof.

Corollary 2.4. If R is a Noetherian ring then there exists a natural R -isomorphism

$$\widehat{\Gamma}_R^m(X) \approx \bigoplus_{P \in \text{Max}(R)} \widehat{\Gamma}_{R/P}^m(X_P)$$

induced by $X \rightarrow X_P$.

Proof: Compare the decompositions from Theorem 2.3 for X and X_P in the case if X is finitely generated. Next apply Lemma 1.1.

The same argument prove the following

Corollary 2.5. If R is a local Noetherian ring then there exists a natural R -isomorphism :

$$\widehat{\Gamma}_R^m(X) \approx \widehat{\Gamma}_{\hat{R}}^m(X \otimes \hat{R})$$

induced by $X \rightarrow X \otimes \hat{R}$.

Observe that the above two corollaries reduce the computation of $\widehat{\Gamma}_R^m(X)$ for Noetherian R to the case when R is local and complete. Theorem 2.3 reduces this problem (for finitely generated X) to the case when R is local Artinian. This case will be studied in the next section.

3. The Artinian case. Let (R, P) be an Artinian local ring. Then $P^k = 0$ for some natural k . Observe that $r^2 = 0$ for any $r \in P^{k-1}$ (if $k > 1$). This is the motivation of the following.

Proposition 3.1. If $r^2=0$ in R and $m \leq 5$ then $r\widehat{\Gamma}_R^m(X) = 0$ for any R -module X .

Proof : To start with, we give some general formulas. It follows from [1] that :

$$\sum_{m_i > 0} \widetilde{\gamma}_{m_1, \dots, m_n}^{m_i}(x_1, \dots, x_n) = 0 \text{ for any } x_1, \dots, x_n \in X.$$

Denote $\Sigma/m_1, \dots, m_n/ = \widetilde{\gamma}_{m_1, \dots, m_n}^{m_i}(x_1, \dots, x_n)$ for $m_i > 0$, $\Sigma m_i = m$. We must prove that r annihilates all this generators. We have :

$$(1) \quad \Sigma/m_1, \dots, m_n/ = 0.$$

Replacing x_1 by rx_1 and $(1+r)x_1$ we get :

$$(2) \quad r\Sigma/1, m_2, \dots, m_n/ = 0$$

$$(2') \quad \Sigma(1+rm_1)/m_1, \dots, m_n/ = 0$$

since $r^2 = 0$ and $(1+r)^k = 1+kr$. In view of (1) and (2) we get from (2') :

$$(3) \quad r \sum_{k=3}^{m-n+1} (k-2) \Sigma/k, m_2, \dots, m_n / = 0.$$

In particular, it follows that :

- (a) $/1, \dots, 1/ = 0$ by (1) ($n=m$)
- (b) $r/2, 1, \dots, 1/ = 0$ by (1) and (2) ($n=m-1$)
- (c) $r/3, 1, \dots, 1/ = 0$ by (3) ($n=m-2$)
- (d) $r/1, m-1/ = 0$ by (2) ($n=2$).

For $m \leq 2$ there is nothing to prove. For $m=3$ we utilize (a),(b). For $m=4$ we get $r/3, 1/ = r/1, 3/ = r/2, 1, 1/ = r/1, 2, 1/ = r/1, 1, 2/ = r/1, 1, 1, 1/ = 0$. Hence also $r/2, 2/ = 0$ by (1). For $m=5$ we have $r/1, 4/ = r/3, 1, 1/ = r/2, 1, 1, 1/ = r/1, 1, 1, 1, 1/ = 0$ and analogously for any permutation. Then (2) and (3) get us $r/1, 2, 2/ = r/3, 2/ = 0$. This completes the proof.

Remark 3.2. Using the same formulas (when we also replace x_2 by $-x_2$) we can prove the above proposition for $m \leq 7$ with the assumption that 2 is invertible in R.

Corollary 3.3. Let R be a Noetherian ring and $m \leq 5$ (or $m \leq 7$ and 2 is invertible in R). Then there exists a natural R-isomorphism :

$$\widehat{\Gamma}_R^m(X) \approx \bigoplus_{P \in \text{Max}(R)} \widehat{\Gamma}_{R/P}^m(X/PX)$$

induced by $X \rightarrow X/PX$.

Proof : It can be assumed that X is finitely generated. In view of Theorem 2.3, it suffices to prove that $\widehat{\Gamma}_R^m(X) \approx \widehat{\Gamma}_{R/P}^m(X/PX)$ for any Artinian local (R,P). If $P^k = 0, P^{k-1} \neq 0$ and $k > 1$ (i.e. R is not a field) then :

$$\widehat{\Gamma}_R^m(X) = \widehat{\Gamma}_R^m(X) / P^{k-1} \widehat{\Gamma}_R^m(X) \approx \widehat{\Gamma}_{R/P^{k-1}}^m(X/P^{k-1}X)$$

by Proposition 3.1 and Remark 3.2. Induction on k completes the proof.

Remark 3.4. The assumptions of the above corollary are necessary. In fact, it can be computed that :

$$\widehat{\Gamma}_{Z_4}^6(Z_4^2) = Z_2 \oplus Z_2 \oplus Z_2 \oplus Z_4, \quad \widehat{\Gamma}_{Z_9}^6(Z_9^2) = Z_3 \oplus Z_3 \oplus Z_3 \oplus Z_3 \oplus Z_9.$$

Remark 3.5. Since the dimensions of $\widehat{\Gamma}^m(X)$ over fields are known (see [1]), Corollary 3.3 solves the problem of computation of $\widehat{\Gamma}^m(X)$ over Noetherian rings for small m. For example, it can be proved that :

$$\begin{aligned} \widehat{\Gamma}_Z^3(Z^n) &= \binom{n}{2} Z_2 \\ \widehat{\Gamma}_Z^4(Z^n) &= 2 \binom{n+1}{3} Z_2 \oplus \binom{n}{2} Z_3 \end{aligned}$$

$$\tilde{\Gamma}_Z^5(Z^n) = (3 \binom{n}{2} + 5 \binom{n}{3} + 3 \binom{n}{4})Z_2 \oplus 2 \binom{n+1}{3}Z_3$$

where $\binom{n}{k} = 0$ for $n < k$. However, the problem is open for large m .

REFERENCES

- [1] A. PROSZYNSKI, Some functors related to polynomial theory, *Fundamenta Math.* 98, (1978), 219-229.
- [2] N. ROBY, Lois polynomes et lois formelles en théorie des modules, *Ann. Scient. Ec. Norm. Sup.* 80 (1963) 213-348.

Andrzej PRÓSZYŃSKI
 Institute of Mathematics
 Polish Academy of Sciences
 ul. Chopina 12
 87-100 TORUŃ, POLAND
