

MÉMOIRES DE LA S. M. F.

REMI SENTIS

Discretized feedback for differential games

Mémoires de la S. M. F., tome 60 (1979), p. 153-159

http://www.numdam.org/item?id=MSMF_1979__60__153_0

© Mémoires de la S. M. F., 1979, tous droits réservés.

L'accès aux archives de la revue « Mémoires de la S. M. F. » (<http://smf.emath.fr/Publications/Memoires/Presentation.html>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

DISCRETIZED FEEDBACK FOR DIFFERENTIAL GAMES

Rémi SENTIS

§1. Introduction

Let us recall a result of minimization for optimal control (see SENTIS [1]). For any initial condition (t, x) of $[0, T] \times \mathbb{R}^d$, we call $\mathcal{V}_{t,x}^2$ the set of the controls b of $L^\infty(0, T; \mathbb{R}^d)$ such that (1) admits a solution (which is denoted y_b):

$$(1) \quad \begin{aligned} y'(s) &= b(s) & b(s) &\in B(s, y(s)) & \text{a.e. } s \in [t, T] \\ y(t) &= x \end{aligned}$$

with the hypothesis:

$$(2) \quad \begin{aligned} B &\text{ is a Lipschitzian multivalued mapping from } [0, T] \times \mathbb{R}^d \\ &\text{with convex compact values in the sphere of radius } Q_0. \end{aligned}$$

For fixed (t, x) we will minimize on $\mathcal{V}_{t,x}^2$ the following cost

$$(3) \quad J_{t,x}(b) + F(y_b(T))$$

where F is Lipschitzian and has no propriety of convexity. This problem admits an optimal open-loop control, but we look for a feedback which approaches the optimum for any initial condition. For that purpose we discretize the interval of time defining:

$$(4) \quad \begin{cases} h_n = T/n \\ t_n^k = kh_n & \forall k \in \mathbb{N} \\ \theta_n^t \text{ the unique integer such that } t \in [t_n^k, t_n^{k+1}[\end{cases}$$

And there exist multivalued (m.v.) mappings $v_n^0, v_n^1, \dots, v_n^{n-1}$ from \mathbb{R}^d to \mathbb{R}^d such that

$$v_n^k(z) \subset B(t_n^k, z) \quad \forall z \in \mathbb{R}^d$$

and such that for any initial condition (t, x) , if we define a trajectory y_n (linear on any interval $[t_n^k, t_n^{k+1}[$) by $y_n(t_n^k) = x_n^k$: with

$$\begin{cases} x_n^{\theta t} = x \\ x_n^{k+1} \in x_n^k + v_n^k(x_n^k) h_n \end{cases} \quad k \geq \theta t$$

then any accumulation point y in $C^0(0, T; \mathbb{R}^d)$ of $(y_n)_n$ is a solution of (1) and is optimal, that is to say:

$$F(y(T)) = J_{t,x}(y') = \underset{b \in \mathcal{V}_{t,x}}{\text{Min}} J_{t,x}(b)$$

Let us now consider the following differential game. For any initial condition (t, x) the admissible trajectories are the solutions of

$$(7) \begin{cases} y'(s) \in A(s, y(s)) + B(s, y(s)) & \text{a.e. } s \in [t, T] \\ y(t) = x \end{cases}$$

where A and B satisfy (2). Let F be a Lipschitzian function on \mathbb{R}^d .

Heuristically, if u and v are two sections of A and B such that there exists a solution (denoted $y_{u,v}$) of:

$$(8) \begin{cases} y'(s) = u(s, y(s)) + v(s, y(s)) \\ y(t) = x \end{cases}$$

then we look for u^* and v^* , sections of A and B , such that

$$(9) \quad F(y_{u,v^*}(T)) \leq F(y_{u^*,v^*}(T)) \leq F(y_{u^*,v}(T))$$

for any u and v section of A and B .

In general, there do not exist sections u^* and v^* verifying (9) and such that u^* and v^* are continuous with respect to the state variable. (Obviously there do not exist open-loop controls u^* and v^* verifying (9).) The topic of this paper is to find a couple of strategies which is a saddle-point for the differential game in a certain class of strategies. For that purpose we must first define the class of admissible strategies (we use the notations (4) except $h_n = T/2^n$ and we write \bar{n} for 2^n)

Definition 1. An admissible strategy for the player U [or V] is a sequence $(u_n)_n$ [or $(v_n)_n$] of elements u_n [or v_n] (which are called discretized feedbacks) with:

$$(10) \left\{ \begin{array}{l} u_n = \{u_n^0, u_n^1, u_n^2, \dots, u_n^{\bar{n}-1}\} \in \prod_{k=0}^{\bar{n}-1} \mathcal{U}_n^k \\ \text{where } \mathcal{U}_n^k \text{ is the set of the m.v. mappings } u \text{ on } \mathbb{R}^d \text{ verifying :} \\ u(z) \subset A(t_n^k, z) \end{array} \right.$$

and :

$$(10') \left\{ \begin{array}{l} v_n = \{v_n^0, v_n^1, v_n^2, \dots, v_n^{\bar{n}-1}\} \in \prod_{k=0}^{\bar{n}-1} \mathcal{V}_n^k \\ \text{where } \mathcal{V}_{n,k} \text{ is the set of the m.v. mappings } v \text{ on } \mathbb{R}^d \text{ verifying :} \\ v(z) \subset B(t_n^k, z) \quad \forall z \in \mathbb{R}^d \end{array} \right.$$

In §2, we exhibit particular discretized feedbacks associated to each h_n and in §3, let n go to infinity, to show that the sequences of such discretized feedbacks constitutes a saddle point in the class of admissible strategies (for detailed proofs, see SENTIS [2]).

§2. Definition of the discretized feedbacks \tilde{u}_n and \tilde{v}_n .

The following proposition justifies the term admissible in definition 1.

Proposition 1. Let us fix (t, x) . If (u_n) and (v_n) are admissible strategies and if we define a trajectory y_n linear on each interval $[t_n^k, t_n^{k+1}[$ by $y_n(t_n^k) = x_n^k$ and x_n^k given by (11)

$$(11) \left\{ \begin{array}{l} x_n^0 = x \\ x_n^{k+1} \in x_n^k + h_n (u_n^k(x_n^k) + v_n^k(x_n^k)) \end{array} \right. \quad k \geq \theta t$$

then any accumulation point y in C^0 verifies (7).

Now let us give two definitions for the cost of a game with initial conditions (t, x) .

Definition 2. The cost of the game for the two discretized feedbacks u_n and v_n is the subset of \mathbb{R} defined by :

$$J_{t,x}(u_n, v_n) = \{F(x_n^k)\} \text{ such that there exists } (x_n^k)_k \text{ verifying (11)}$$

Definition 3. The cost of the game for the two admissible strategies (u_n) and (v_n) is the subset of R denoted by $J_{t,x}((u_n), (v_n))$ and containing the accumulation points of all the sequences (a_n) verifying $a_n \in J_{t,x}((u_n), (v_n))$. Let us yet define the lower and upper optimal cost-functions \bar{W}_n^k and \hat{W}_n^k as FRIEDMAN [1] by decreasing induction :

$$(12) \quad \begin{cases} \bar{W}_n^k(x) = F(x) \\ \bar{W}_n^k(x) = \text{Max}_{u \in A(t_n^k, x)} \bar{Z}_n^k(x, u) \text{ and } \bar{Z}_n^k(x, u) = \text{Min}_{v \in B(t_n^k, x)} \bar{W}_n^{k+1}(x+(u+v)h_n) \end{cases}$$

and :

$$(12') \quad \begin{cases} \hat{W}_n^k(x) = F(x) \\ \hat{W}_n^k(x) = \text{Min}_{v \in B(t_n^k, x)} \hat{Z}_n^k(x, v) \text{ and } \hat{Z}_n^k(x, v) = \text{Max}_{u \in A(t_n^k, x)} \hat{W}_n^{k+1}(x+(u+v)h_n) \end{cases}$$

Now we can exhibit the m.v. mappings \hat{u}_n^k and \hat{v}_n^k , which do not depend on the initial conditions.

$$(13) \quad \begin{cases} \hat{u}_n^k(x) = \text{Arg Max}_{u \in A(t_n^k, x)} \hat{Z}_n^k(x, u) \\ \hat{v}_n^k(x) = \text{Arg Min}_{v \in B(t_n^k, x)} \hat{Z}_n^k(x, v) \end{cases}$$

We can prove easily by induction the following :

Proposition 2. All the mappings \bar{W}_n^k , \bar{Z}_n^k , \hat{W}_n^k , \hat{Z}_n^k are Lipschitzian (with respect to x) with constant K (independent of n and k).

§3. Saddle point theorem

Proposition 3 We have when n goes to infinity:

$$\overline{W}_n^{\theta t}(x) \rightarrow W^-(t,x) \quad \widehat{W}_n^{\theta t}(x) \rightarrow W^+(t,x)$$

moreover:

$$W^-(t,x) \leq W^+(t,x)$$

principle of the proof.

First we show by decreasing induction on k that

$$\overline{W}_n^k(x) - \overline{W}_{n+1}^{2k}(x) \leq (\overline{n-k}) C_0 (h_n)^2 \quad \forall k$$

And as $\theta_{n+1}t$ is equal to $(2\theta_n t)$ or $(2\theta_n t+1)$ we have according to proposition 2:

$$(14) \quad \overline{W}_n^{\theta t}(x) - \overline{W}_n^{\theta_{n+1}t}(x) \leq C_1 h_n \quad \text{with } C_1 = C_0 T + 2KQ_0$$

Hence if we denote:

$$W^-(t,x) = \lim_n \sup \overline{W}_n^{\theta t}(x)$$

we can show easily according to (14) that $\overline{W}_n^{\theta t}(x) \rightarrow W^-(t,x)$. We show exactly the same way that $\widehat{W}_n^{\theta t}(x) \rightarrow W^+(t,x)$. The end of the proposition is a consequence of the following fact:

$$\overline{W}_n^k(x) \leq \widehat{W}_n^k(x) \quad \forall x, n, k \quad \text{Q.E.D.}$$

The following proposition is fundamental and is proved in FRIEDMAN [1], using the m.v. mappings:

$$\text{Arg Min}_{v \in B(t_n^k, x)} \bar{W}_n^{k+1}(x+(u+v)h_n) \quad \text{and} \quad \text{Arg Max}_{u \in A(t_n^k, x)} \hat{W}_n^{k+1}(x+(u+v)h_n)$$

Proposition 4We have

$$\bar{W}^-(t, x) = \hat{W}^+(t, x)$$

We write thus $W(t, x)$ instead of $\bar{W}^-(t, x)$. This number is called the value of the game.

Proposition 5

For any $u_n \in \prod_{k=0}^{\bar{n}-1} \mathcal{U}_n^k$, we have

$$(15) \quad J_{t,x}(u_n, \tilde{v}_n) \leq \hat{W}_n^{\theta t}(x)$$

(This means that any element of the left-hand side is smaller than the right-hand side.)

Proof

Using the notations (11) (changing v_n^k into \tilde{v}_n^k), we note that there exist $q_n^k \in \tilde{v}_n(x_n^k)$ such that

$$x_n^{k+1} \in x_n^k + h_n(u_n^k(x_n^k) + q_n^k) \quad \forall k \geq \theta t$$

Thus we have:

$$\hat{W}_n^k(x_n^k) = \hat{Z}_n^k(x_n^k, q_n^k) \geq \hat{W}_n^{k+1}(x_n^{k+1})$$

Rewriting this inequality for k from θt to \bar{n} , we obtain (15). Q.E.D.

We have evidently also:

$$(16) \quad J_{t,x}(\tilde{u}_n, v_n) \geq \bar{W}_n^{\theta t}(x)$$

Let n go to infinity in (15) and (16), we deduce immediately from the propositions 3 and 4 the following:

Theorem

For any admissible strategy (u_n) and (v_n) , we have:

$$J_{t,x}((u_n), (\tilde{v}_n)) \leq W(t,x) \leq J_{t,x}((\tilde{u}_n), (v_n))$$

Thus we have:

$$W(t,x) = \min_{(v_n)} \max_{(u_n)} J_{t,x}((u_n), (v_n)) = \max_{(u_n)} \min_{(v_n)} J_{t,x}((u_n), (v_n))$$

And if y is an accumulation point in C^0 of trajectories y_n associated to \tilde{u}_n and \tilde{v}_n we have

$$W(t,x) = J_{t,x}((\tilde{u}_n), (\tilde{v}_n)) = F(y(T))$$

References:

- A. FRIEDMAN [1] Differential games, Wiley Interscience, N.Y., 1971.
- R. SENTIS [1] Discretized feedback for non-stationary and stationary optimal control problem. To be published.
- [2] Feedbacks discretisés en jeux différentiels, Cahiers de Mathématiques de la Décision, No. 780, Université Paris IX - Dauphine.

Rémy S E N T I S
28 rue du Fief
92100 BOULOGNE