

ANNALES DE L'INSTITUT FOURIER

JERRY M. LODDER

Leibniz cohomology for differentiable manifolds

Annales de l'institut Fourier, tome 48, n° 1 (1998), p. 73-95

http://www.numdam.org/item?id=AIF_1998__48_1_73_0

© Annales de l'institut Fourier, 1998, tous droits réservés.

L'accès aux archives de la revue « Annales de l'institut Fourier » (<http://annalif.ujf-grenoble.fr/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

LEIBNIZ COHOMOLOGY FOR DIFFERENTIABLE MANIFOLDS

by Jerry M. LODDER*

Introduction.

The goal of this paper is to extend Jean-Louis Loday's Leibniz cohomology $[L, P]$ from a Lie algebra invariant to a new invariant for differentiable manifolds. The extension is achieved by using a cochain complex of tensors from classical differential geometry, and as such, generalizes the de Rham complex of differential forms (i.e. alternating tensors). The exterior derivative of an n -tensor, however, is not necessarily an $(n + 1)$ -tensor, but more generally an operator on vector fields. The Leibniz cochain complex thus becomes a non-commutative version of Gelfand-Fuks cohomology for smooth vector fields. The Leibniz cohomology groups, $HL^*(M)$, are diffeomorphism invariants, but fail to be homotopy invariants. In fact the first obstruction to the homotopy invariance of $HL^*(\mathbf{R}^n)$ is the universal Godbillon-Vey invariant in dimension $2n + 1$. The main calculational result of the paper is then the computation of $HL^*(\mathbf{R}^n)$ in terms of (i) certain universal invariants of foliations, and (ii) Loday's product structure on HL^* . Unlike Lie algebra or Gelfand-Fuks cohomology, the Leibniz cohomology of vector fields on \mathbf{R}^n contains infinite families of elements which support non-trivial products.

* Supported by the National Science Foundation, grant no. DMS-9704891.

Key words: Leibniz cohomology – Foliation – Differentiable manifolds – Gelfand-Fuks cohomology.

Math. classification: 17A30 – 57R30 – 57R32.

The paper begins with a brief review of Loday's definition of HL^* , and proceeds with a conceptual discussion of HL^* for differentiable manifolds. The calculational results for $HL^*(\mathbf{R}^n)$ appear in §2 along with the necessary background material on foliations needed to state these results. Finally §3 concludes with a conjectural characteristic map in the setting of Leibniz cohomology.

1. Non-commutative cohomologies.

We begin by reviewing Loday's definition of Leibniz (co)homology [L1], [L2], [L,P]. Let k be a commutative ring and \mathfrak{g} a k -module together with a bilinear map $[\cdot, \cdot] : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathfrak{g}$. Then, by definition, \mathfrak{g} is a Leibniz algebra if the bracket satisfies the "Leibniz identity"

$$(1.1) \quad [x, [y, z]] = [[x, y], z] - [[x, z], y]$$

for all x, y, z in \mathfrak{g} . If in addition to (1.1), the bracket is skew-symmetric, $[x, y] = -[y, x]$, then (1.1) is equivalent to the Jacobi identity, and \mathfrak{g} itself becomes a Lie algebra. In this sense, a Leibniz algebra is a non-commutative version of a Lie algebra. We also need the notion of a representation of a Leibniz algebra [L,P], which is a k -module N equipped with left and right actions of \mathfrak{g}

$$[\cdot, \cdot] : \mathfrak{g} \times N \rightarrow N, \quad [\cdot, \cdot] : N \times \mathfrak{g} \rightarrow N$$

which are bilinear and satisfy the three properties

$$(1.2) \quad \begin{aligned} [m, [x, y]] &= [[m, x], y] - [[m, y], x] \\ [x, [m, y]] &= [[x, m], y] - [[x, y], m] \\ [x, [y, m]] &= [[x, y], m] - [[x, m], y] \end{aligned}$$

for all $x, y \in \mathfrak{g}$ and $m \in N$. Of course, if N is a representation of a Lie algebra \mathfrak{g} , then the above three conditions are equivalent, since a left representation of a Lie algebra, $[\cdot, \cdot] : \mathfrak{g} \times N \rightarrow N$ determines a right representation of the same Lie algebra $[\cdot, \cdot] : N \times \mathfrak{g} \rightarrow N$ via $[x, m] = -[m, x]$.

Returning to the setting of a Leibniz algebra \mathfrak{g} and a representation N , the Leibniz cohomology of \mathfrak{g} with coefficients in N , written $HL^*(\mathfrak{g}; N)$, is the homology of the cochain complex

$$C^m(\mathfrak{g}; N) = \text{Hom}_k(\mathfrak{g}^{\otimes m}, N), \quad m \geq 0.$$

To streamline the notation for the coboundary map

$$d : C^n(\mathfrak{g}; N) \rightarrow C^{n+1}(\mathfrak{g}; N)$$

let (g_1, g_2, \dots, g_n) denote the element $g_1 \otimes g_2 \otimes \dots \otimes g_n \in \mathfrak{g}^{\otimes n}$. For $f \in C^n(\mathfrak{g}; N)$, we have

$$(1.3) \quad \begin{aligned} & (df)(g_1, g_2, \dots, g_{n+1}) \\ &= \sum_{1 \leq i < j \leq n+1} (-1)^{j+1} f(g_1, \dots, g_{i-1}, [g_i, g_j], g_{i+1}, \dots, \hat{g}_j, \dots, g_{n+1}) \\ & \quad + [g_1, f(g_2, \dots, g_{n+1})] + \sum_{i=2}^{n+1} (-1)^i [f(g_1, \dots, \hat{g}_i, \dots, g_{n+1}), g_1]. \end{aligned}$$

Loday and Pirashvili [LP] prove that $d \circ d = 0$ in (1.3), thus establishing that $C^*(\mathfrak{g}; N)$ is a cochain complex. For a Lie algebra \mathfrak{h} and a representation N , the projection to the exterior product $\mathfrak{h}^{\otimes n} \rightarrow \mathfrak{h}^{\wedge n}$ induces a natural homomorphism

$$H_{\text{Lie}}^*(\mathfrak{h}; N) \rightarrow HL^*(\mathfrak{h}; N),$$

where H_{Lie}^* is Lie-algebra cohomology.

We wish to extend Leibniz cohomology to an invariant for differentiable manifolds so that in a certain sense HL^* is a non-commutative version of de Rham cohomology. In particular the non-commutativity arises by considering a cochain complex of tensors (from differential geometry) which are not necessarily skew-symmetric. Let M be a differentiable manifold and α an n -tensor on M . Then, by definition, α is a differentiable section

$$\alpha : M \rightarrow (T^*M)^{\otimes n},$$

where T^*M is the cotangent bundle of M . To define the de Rham cohomology groups, $H_{DR}^*(M)$, however, one uses forms $\omega \in \Omega^n(M)$, where ω is a differentiable section of the n -th exterior power of T^*M ,

$$\omega : M \rightarrow (T^*M)^{\wedge n}.$$

Any tensor α determines a form ω_α by first symmetrizing over the symmetric group Σ_n , i.e.

$$\omega_\alpha(v_1 \wedge v_2 \wedge \dots \wedge v_n) = \sum_{\sigma \in \Sigma_n} (\text{sgn } \sigma) \alpha(v_{\sigma(1)} \otimes v_{\sigma(2)} \otimes \dots \otimes v_{\sigma(n)}).$$

This is the step which yields a graded commutative ring structure on

$$\Omega^*(M) = \sum_{n \geq 0} \Omega^n(M).$$

Recall E. Cartan's global formulation for the exterior derivative of a differential form in terms of vector fields (see Spivak [S], p. 289 for a modern treatment). Let X_1, X_2, \dots, X_{n+1} be differentiable vector fields on M and let $\omega \in \Omega^n(M)$. Then there is a unique $(n+1)$ -form $d\omega$ such that

$$(1.4) \quad d\omega(X_1 \wedge X_2 \wedge \cdots \wedge X_{n+1}) = \sum_{i=1}^{n+1} (-1)^{i+1} X_i(\omega(X_1 \wedge \cdots \wedge \hat{X}_i \cdots \wedge X_{n+1})) \\ + \sum_{1 \leq i < j \leq n+1} (-1)^{j+1} \omega(X_1 \wedge \cdots \wedge X_{i-1} \wedge [X_i, X_j] \wedge X_{i+1} \wedge \cdots \\ \cdots \wedge \hat{X}_j \cdots \wedge X_{n+1}),$$

where $[,]$ is the Lie bracket of vector fields and

$$X_i(\omega(X_1 \wedge \cdots \wedge \hat{X}_i \cdots \wedge X_{n+1}))$$

is the directional derivative of the function

$$\omega(X_1 \wedge \cdots \wedge \hat{X}_i \cdots \wedge X_{n+1}) : M \rightarrow \mathbf{R}.$$

Although $d\omega$ is defined on vector fields in (1.4), the value of the exterior derivative

$$d\omega(X_1(p) \wedge X_2(p) \wedge \cdots \wedge X_{n+1}(p))$$

depends only on one point $p \in M$, since $d\omega$ is linear over C^∞ functions on M . Thus, $d\omega$ determines a well-defined $(n+1)$ -form. See Spivak [S], pp. 162, 289, for more details. The following lemma is immediate.

1.5. LEMMA. — *If in (1.4) ω is an n -tensor and the exterior product \wedge is replaced with the tensor product \otimes , then $(d \circ d)(\omega) = 0$.*

Proof. — Let $\chi(M)$ be the vector space of all differentiable vector fields on M . Then $\chi(M)$ is a Lie algebra under the Lie bracket of vector fields, and, hence a Leibniz algebra. Let $C^\infty(M)$ be the vector space of C^∞ functions $f : M \rightarrow \mathbf{R}$. Clearly $C^\infty(M)$ is a representation of $\chi(M)$ by

$$[X, f] = X(f),$$

where $X(f)$ is the Lie derivative of f in the direction X . Setting $[f, X] = -[X, f]$, we see that $C^\infty(M)$ becomes a representation of the Leibniz algebra $\chi(M)$. The lemma follows since $d \circ d = 0$ in (1.3). \square

It must be noted, however, that when using tensors, $d\omega$ is **no longer linear over C^∞ functions**. The obstruction to $C^\infty(M)$ -linearity is precisely the failure of a tensor to be skew-symmetric. This is the essential distinction between the commutative and non-commutative cases. As a

consequence $d\omega$ is not necessarily an $(n+1)$ -tensor, but more generally an \mathbf{R} -linear mapping

$$d\omega : \chi(M)^{\otimes(n+1)} \rightarrow C^\infty(M).$$

We propose the following definition.

1.6. DEFINITION. — *The Leibniz cohomology of a differentiable manifold M , written*

$$HL^*(\chi(M); C^\infty(M)),$$

is the homology of the complex of continuous cochains

$$\text{Hom}_{\mathbf{R}}^{\text{cont}}(\chi(M)^{\otimes n}, C^\infty(M)), n \geq 0,$$

in the C^∞ topology, where $\chi(M)$ is the Lie algebra of differentiable vector fields on M , and $C^\infty(M)$ denotes the ring of C^∞ real-valued functions on M . The coboundary map d is given in (1.3).

Several observations are in order.

1.7. — The continuous Lie-algebra cohomology of $\chi(M)$ is called Gelfand-Fuks cohomology, $H_{GF}^*(\chi(M); \mathbf{R})$, and is essentially the subject of the book *Cohomology of Infinite Dimensional Lie Algebras* [F].

1.8. — There is a commutative diagram

$$\begin{array}{ccc} H_{DR}^*(M) & \xrightarrow{i} & H_{GF}^*(\chi(M); C^\infty(M)) \\ \pi \circ i \downarrow & & \swarrow \pi \\ HL^*(\chi(M); C^\infty(M)) & & \end{array}$$

which arises from the fact that $H_{DR}^*(M)$ is the homology of the cochain complex

$$\text{Hom}_{C^\infty(M)}^{\text{cont}}(\chi(M)^{\wedge n}, C^\infty(M)), n \geq 0,$$

with boundary map given by (1.4). In fact there is a natural inclusion of cochain complexes

$$\Omega^n(M) \rightarrow \text{Hom}_{C^\infty(M)}^{\text{cont}}(\chi(M)^{\wedge n}, C^\infty(M))$$

given by

$$\omega \mapsto \omega(X_1 \wedge X_2 \wedge \cdots \wedge X_n) : M \rightarrow \mathbf{R}$$

$$\omega(X_1 \wedge X_2 \wedge \cdots \wedge X_n)(p) = \omega(X_1(p) \wedge X_2(p) \wedge \cdots \wedge X_n(p)).$$

The map $i : H_{DR}^*(M) \rightarrow H_{GF}^*(\chi(M); C^\infty(M))$ is then induced by the inclusion

$$\text{Hom}_{C^\infty(M)}^{\text{cont}}(\chi(M)^{\wedge n}, C^\infty(M)) \rightarrow \text{Hom}_{\mathbf{R}}^{\text{cont}}(\chi(M)^{\wedge n}, C^\infty(M)).$$

See [F], p. 21 for further details. The map

$$\pi : H_{GF}^*(\chi(M); C^\infty(M)) \rightarrow HL^*(\chi(M); C^\infty(M))$$

is induced by the projection $\chi(M)^{\otimes n} \rightarrow \chi(M)^{\wedge n}$.

1.9. — If $M = G$ is a Lie group, then $\chi(G)^G$, the left-invariant vector fields on G form the Lie algebra \mathfrak{g} of G . For invariant functions on G , we have $C^\infty(G)^G = \mathbf{R}$. Thus,

$$T(\mathfrak{g}) = \sum_{n \geq 0} \mathfrak{g}^{\otimes n}$$

is a subcomplex of

$$\{C^\infty(G) \otimes \chi(G)^{\otimes n}\}_{n \geq 0}.$$

In fact $T(\mathfrak{g})$ is the original complex proposed by Loday [L1], p. 324 for the Leibniz homology of a Lie algebra. Of course, for G compact and connected, there is the Chevalley-Eilenberg [C,E] isomorphism

$$H_{\text{Lie}}^*(\mathfrak{g}; \mathbf{R}) \simeq H_{DR}^*(G; \mathbf{R}).$$

Such an isomorphism appears not to hold in the setting of Leibniz cohomology however.

1.10. LEMMA. — *Leibniz cohomology is a diffeomorphism invariant, i.e. if $f : M \rightarrow N$ is a diffeomorphism, then there is an induced isomorphism*

$$f^* : HL^*(\chi(N); C^\infty(N)) \rightarrow HL^*(\chi(M); C^\infty(M)).$$

Proof. — Note that if X is a C^∞ vector field on M , then $f_*(X)$ is a C^∞ vector field on N , where

$$(f_*(X))_q = f_*(X_{f^{-1}(q)}) \quad \text{for } q \in N.$$

See Spivak [S], p. 186. To define

$$f^\# : \text{Hom}_{\mathbf{R}}^c(\chi(N)^{\otimes n}, C^\infty(N)) \rightarrow \text{Hom}_{\mathbf{R}}^c(\chi(M)^{\otimes n}, C^\infty(M)),$$

let $\beta \in \text{Hom}_{\mathbf{R}}^c(\chi(N)^{\otimes n}, C^\infty(N))$, and $X_i \in \chi(M)$, $i = 1, 2, \dots, n$. Then

$$f^\#(\beta)(X_1 \otimes X_2 \otimes \cdots \otimes X_n) : M \rightarrow \mathbf{R}$$

is given by

$$f^\#(\beta)(X_1 \otimes \cdots \otimes X_n)(p) = \beta(f_*(X_1) \otimes \cdots \otimes f_*(X_n))(f(p)).$$

Of course for $n = 0$, $\text{Hom}_{\mathbf{R}}^c(\mathbf{R}, C^\infty(N)) = C^\infty(N)$, and for $\beta \in C^\infty(N)$,

$$(f^\#(\beta))(p) = \beta(f(p)).$$

Two properties are needed to show that $f^\#$ is a cochain map. First,

$$f_*([X_1, X_2]) = [f_*(X_1), f_*(X_2)].$$

To state the second, recall that $C^\infty(M)$ is a left representation of the Lie algebra $\chi(M)$ via

$$[X, \varphi] = X(\varphi), \varphi \in C^\infty(M), X \in \chi(M).$$

For $\gamma \in C^\infty(N)$, the composition $\gamma \circ f \in C^\infty(M)$, and the second needed property is

$$[X, \gamma \circ f] = [f_*(X), \gamma] \circ f.$$

This is simply “the chain rule.”

Since both compositions $(f^{-1})^* \circ f^*$ and $f^* \circ (f^{-1})^*$ are the identity, it follows that

$$f^* : HL^*(\chi(N); C^\infty(N)) \rightarrow HL^*(\chi(M); C^\infty(M))$$

is an isomorphism. Clearly if using trivial coefficients, then

$$f^* : HL^*(\chi(N); \mathbf{R}) \rightarrow HL^*(\chi(M); \mathbf{R})$$

is also an isomorphism. □

1.11. — We wish now to compare, at least philosophically, Leibniz cohomology with cyclic homology, both of which serve as certain non-commutative versions of de Rham cohomology. For cyclic homology, what has been generalized to a non-commutative setting is the coefficient ring, i.e. $C^\infty(M)$ is replaced with a non-commutative algebra. For Leibniz cohomology the differential operator (exterior derivative) has been cast in a non-commutative framework. That these are in fact genuinely different generalizations can be seen from the calculation of $HP_*(C^\infty(M))$, the periodic cyclic homology of $C^\infty(M)$. We have [C]

$$HP_*(C^\infty(M)) \simeq H_{DR}^*(M)$$

for M σ -compact.

The notions of non-commutativity can be summarized in the following diagram, where the column headings describe the coefficients, and the row headings describe the operators.

| | COMMUTATIVE | NON-COMMUTATIVE |
|-----------------|---|---|
| COMMUTATIVE | de Rham (singular cohomology) | cyclic homology (K-theory) |
| NON-COMMUTATIVE | Leibniz cohomology of Lie algebras (foliations) | Leibniz cohomology of Leibniz algebras (quantum field theory) |

The parenthetical remarks refer to the type of invariants which can be computed by each theory. For the relation between cyclic homology and K-theory, see for example Goodwillie's paper [Gw]. In this paper we capture the Godbillon-Vey invariant for foliations via a Leibniz cohomology computation of a certain Lie algebra. In the next remark we observe how a Leibniz algebra arises naturally in V. Kač's work [K] on vertex operator algebras. The Leibniz cohomology groups of this algebra would then provide new invariants of quantum field theory.

1.12. — We now review a construction of Kač [K] concerning vertex algebras, and show that a Lie algebra occurring in [K]—formed by a certain quotient—is genuinely a Leibniz (non-Lie) algebra if the quotient is not formed. Let U be an associative, but not necessarily commutative algebra, such as $\text{End}(V)$ for a vector space V . Let $a(z)$ be a formal power series in z and z^{-1} with coefficients in U , i.e. $a(z)$ is a formal distribution. For

$$a(z), b(z) \in U[[z, z^{-1}]],$$

the 0-th order product, $a(z)_{(0)}b(z)$, is defined as

$$a(z)_{(0)}b(z) = \text{Res}_z (a(z)b(w) - b(w)a(z)),$$

where for $c(z) = \sum_{k \in \mathbf{Z}} c_k z^k$, we set $\text{Res}_z(c(z)) = c_{-1}$. Then

$$a(z)_{(0)}b(z) = \sum_{k \in \mathbf{Z}} (a_{-1}b_k - b_k a_{-1})z^k \in U[[z, z^{-1}]].$$

By the work of Kač, the quotient $\mathcal{A}/\partial\mathcal{A}$ is a Lie algebra with respect to the 0-th order product, where $\mathcal{A} = U[[z, z^{-1}]]$. Setting $[a(z), b(z)] = a(z)_{(0)}b(z)$, we see that this bracket is not skew-symmetric on \mathcal{A} , but satisfies what Loday and Pirashvili [LP] call the left Leibniz identity:

$$(1.13) \quad [[a(z), b(z)], c(z)] = [a(z), [b(z), c(z)]] - [b(z), [a(z), c(z)]].$$

Thus, $U[[z, z^{-1}]]$ is a left Leibniz algebra. There is a one-to-one correspondence between left Leibniz algebras $(\mathfrak{g}, [,]_L)$ for which the bracket satisfies (1.13) and right Leibniz algebras $(\mathfrak{g}, [,]_R)$ for which the bracket satisfies (1.1). The correspondence is simply given by

$$\mathfrak{g} \longleftrightarrow \mathfrak{g} \quad [y, x]_L \longleftrightarrow [x, y]_R.$$

Also, if V is a $\mathbf{Z}/2$ -graded vector space (i.e. a super-space), and $U = \text{End}(V)$, then there are corresponding statements for $\mathbf{Z}/2$ -graded Lie algebras and $\mathbf{Z}/2$ -graded Leibniz algebras. A different Leibniz algebra in the setting of vertex algebras has been found by Kosmann-Schwarzbach [K-S] by using the residue of a dot product of $a(z)$ and $b(z)$ after multiplication

by z^{-1} . This differs from our example where we use the residue of a commutator of $a(z)$ and $b(z)$ (after multiplication by $z^0 = 1$).

2. Leibniz cohomology of formal vector fields.

In this section we begin the calculation of $HL^*(\chi(\mathbf{R}^n); \mathbf{R})$, the Leibniz cohomology of \mathbf{R}^n with trivial coefficients. These groups are isomorphic to the continuous Leibniz cohomology of formal vector fields W_n , which we now describe. Let E be the real vector space with basis

$$\left\{ \frac{\partial}{\partial x_1}, \frac{\partial}{\partial x_2}, \dots, \frac{\partial}{\partial x_n} \right\},$$

and let $E' = \text{Hom}_{\mathbf{R}}(E, \mathbf{R})$ be the dual space with dual basis $\{x_1, x_2, \dots, x_n\}$. Then [F],[Gb]

$$W_n = \left(\prod_{k \geq 0} S^k(E') \right) \otimes E,$$

where S^k denotes the k -symmetric power, and W_n becomes a Lie algebra with the bracket given by

$$\left[\sum_{i=1}^n P_i \frac{\partial}{\partial x_i}, \sum_{i=1}^n Q_i \frac{\partial}{\partial x_i} \right] = \sum_{k=1}^n \left(\sum_{j=1}^n P_j \frac{\partial Q_k}{\partial x_j} - Q_j \frac{\partial P_k}{\partial x_j} \right) \frac{\partial}{\partial x_k}$$

where P_i and Q_i are formal power series in x_1, x_2, \dots, x_n . Furthermore W_n is a topological space in the \mathcal{M} -adic topology, where

$$\|x_i^k\| = c^{-k}, i = 1, 2, \dots, n,$$

for some fixed integer $c > 1$. By $HL^*(W_n)$ we mean the continuous Leibniz cohomology with coefficients in \mathbf{R} . We now define a Taylor series map $\phi : \chi(\mathbf{R}^n) \rightarrow W_n$. For $X \in \chi(\mathbf{R}^n)$, let $X = \sum_{i=1}^n f_i e_i$, where the $f_i : \mathbf{R}^n \rightarrow \mathbf{R}$ are C^∞ functions and the e_i are the canonical vector fields on \mathbf{R}^n (the unit vector fields following the coordinate axes). Then

$$\phi(X) = \sum_{i=1}^n \sum_J \frac{1}{j_1! j_2! \dots j_n!} \frac{\partial^J (f_i)}{\partial x^J} (0) x_1^{j_1} x_2^{j_2} \dots x_n^{j_n} \frac{\partial}{\partial x_i},$$

where J is the multi-index $j_1, j_2, \dots, j_n \geq 0$. It can be checked that ϕ is a continuous homomorphism of Lie algebras.

2.1. LEMMA. — *The induced map $\phi^* : HL^*(W_n) \rightarrow HL^*(\chi(\mathbf{R}^n); \mathbf{R})$ is an isomorphism.*

Proof. — The proof follows essentially from the work of Bott and Segal [BS] who prove that for continuous Lie algebra cohomology

$$\phi^* : H_{\text{Lie}}^*(W_n) \rightarrow H_{GF}^*(\chi(\mathbf{R}^n); \mathbf{R})$$

is an isomorphism. We do not offer complete details of their proof here, but instead state how a few ideas are extended from Lie cochains to Leibniz cochains. Let

$$\chi = \chi(\mathbf{R}^n), C^q = \text{Hom}_{\mathbf{R}}^{\text{cont}}(\chi^{\otimes q}, \mathbf{R}),$$

and define P_q to be the subspace of $\chi^{\otimes q}$ spanned by \mathbf{R} -linear combinations of

$$f_{i_1} e_{i_1} \otimes f_{i_2} e_{i_2} \otimes \cdots \otimes f_{i_q} e_{i_q},$$

where each f_{i_j} is a polynomial and

$$\sum_{j=1}^q \deg(f_{i_j}) \leq q.$$

Let $C_P^q = \text{Hom}_{\mathbf{R}}^{\text{cont}}(P_q, \mathbf{R})$. Then $d : C_P^q \rightarrow C_P^{q+1}$, and we have a short exact sequence

$$0 \longrightarrow B^* \longrightarrow C^* \longrightarrow C_P^* \longrightarrow 0,$$

where B^* is defined by $C_P^* = C^*/B^*$. As in the case for Lie algebra cohomology [BS], we show that B^* is acyclic, thus proving that $C^* \rightarrow C_P^*$ induces an isomorphism on HL^* .

Let $T_t : \mathbf{R}^n \rightarrow \mathbf{R}^n$ be the contraction defined by $T_t(x) = tx$, $0 < t \leq 1$. Then T_t acts on C^q by

$$(T_t \alpha)(X_1 \otimes \cdots \otimes X_q) = t^{-q} \alpha(T_t^*(X_1) \otimes \cdots \otimes T_t^*(X_q)),$$

where $\alpha \in C^q$ and $X_i \in \chi$. Since the one-parameter family of diffeomorphisms T_t is generated by a vector field ρ on \mathbf{R}^n , we have

$$T_t^{-1} \cdot t(d/dt)T_t = \theta(\rho) : C^* \rightarrow C^*,$$

where for the Leibniz cochain complex

$$\begin{aligned} \theta(\rho)(\alpha)(X_1 \otimes X_2 \otimes \cdots \otimes X_q) \\ = - \sum_{i=1}^q \alpha(X_1 \otimes \cdots \otimes X_{i-1} \otimes [X_i, \rho] \otimes X_{i+1} \otimes \cdots \otimes X_q). \end{aligned}$$

From the work of Loday and Pirashvili [LP]

$$\theta(\rho) = i(\rho)d + di(\rho),$$

where $i(\rho) : C^{q+1} \rightarrow C^q$ is given by

$$i(\rho)\alpha(X_1 \otimes \cdots \otimes X_q) = (-1)^{q+1}\alpha(X_1 \otimes X_2 \otimes \cdots \otimes X_q \otimes \rho).$$

It now follows as in [BS] that for $\alpha \in B^*$,

$$\alpha = Kd\alpha + dK\alpha,$$

where

$$K(\alpha) = i(\rho) \int_0^1 t^{-1} T_t(\alpha) dt.$$

Thus K is a contracting chain homotopy for B^* regardless of whether α is skew-symmetric. \square

We now turn to the calculation of $HL^*(W_n)$ in terms of certain universal properties of foliations. Let \mathfrak{F} be a codimension n , C^∞ foliation on a manifold M with trivial normal bundle. Then there is a characteristic map associated to \mathfrak{F} [B] [F] [H]

$$(2.2) \quad \text{char}_{\mathfrak{F}} : H_{\text{Lie}}^*(W_n) \rightarrow H_{DR}^*(M).$$

If \mathfrak{F}_t is a smooth one-parameter family of such foliations, then one can compute the derivative

$$\frac{d}{dt} [\text{char}_{\mathfrak{F}_t}(\alpha)|_{t=0} \in H_{DR}^q(M; \mathbf{R})].$$

The class $\alpha \in H_{\text{Lie}}^q(W_n)$ is called variable if there exists a family \mathfrak{F}_t for which

$$\frac{d}{dt} [\text{char}_{\mathfrak{F}_t}(\alpha)|_{t=0} \neq 0.$$

Otherwise, α is called rigid. From Fuks [F] a necessary condition for the variability of α is that $\text{var}(\alpha) \neq 0$, where var is the homomorphism

$$(2.3) \quad \text{var} : H_{\text{Lie}}^q(W_n) \rightarrow H_{\text{Lie}}^{q-1}(W_n; W'_n)$$

given on the cochain level by

$$\text{var}(\alpha)(g_1, g_2, \dots, g_{q-1})(g_0) = (-1)^{q-1}\alpha(g_0, g_1, g_2, \dots, g_{q-1}).$$

Here and further $H^*(W_n; W'_n)$ denotes the continuous Lie algebra cohomology with coefficients in the co-adjoint representation

$$W'_n = \text{Hom}_{\mathbf{R}}^{\text{cont}}(W_n, \mathbf{R}).$$

Our calculation of $HL^*(W_n)$ is in terms of the homomorphism var and the product structure on Leibniz cohomology.

Recall that for a Leibniz algebra \mathfrak{g} , Loday [L3] has defined a non-commutative, non-associative product on $HL^*(\mathfrak{g})$ which affords $HL^*(\mathfrak{g})$

the structure of a dual Leibniz algebra, where dual is taken in the sense of Koszul duality for operads [G,K], [L4]. More specifically, let

$$C^p = \text{Hom}(\mathfrak{g}^{\otimes p}, \mathbf{R}), \alpha \in C^p, \beta \in C^q,$$

and define $\alpha \cdot \beta \in C^{p+q}$ by

$$(2.4) \quad (\alpha \cdot \beta)(g_1 \otimes g_2 \otimes \cdots \otimes g_{p+q}) = \sum_{\sigma \in Sh_{p-1,q}} \text{sgn}(\sigma) \alpha(g_1 \otimes g_{\sigma(2)} \otimes \cdots \\ \cdots \otimes g_{\sigma(p)}) \cdot \beta(g_{\sigma(p+1)} \otimes \cdots \otimes g_{\sigma(p+q)}),$$

where the sum is over all $(p-1, q)$ -shuffles of the symmetric group Σ_{p+q-1} , i.e.

$$\sigma(2) < \sigma(3) < \cdots < \sigma(p),$$

$$\sigma(p+1) < \sigma(p+2) < \cdots < \sigma(p+q).$$

Then from [L3] $d(\alpha \cdot \beta) = (d\alpha) \cdot \beta + (-1)^{|\alpha|} \alpha \cdot (d\beta)$, and for $x \in HL^p(\mathfrak{g})$, $y \in HL^q(\mathfrak{g})$, $z \in HL^r(\mathfrak{g})$, we have

$$(x \cdot y) \cdot z = x \cdot (y \cdot z) + (-1)^{qr} x \cdot (z \cdot y).$$

In the sequel we will need the following formula for products of homogeneous elements in $HL^*(\mathfrak{g})$

$$(2.5) \quad \{a_0 \cdot (a_1 \cdot (a_2 \cdots (a_{p-1} \cdot a_p) \cdots))\} \cdot \{a_{p+1} \cdot (a_{p+2} \cdot (\cdots (a_{p+q-1} \cdot a_{p+q}) \cdots))\} \\ = \sum_{\sigma \in Sh_{p,q}} \pm a_0 \cdot (a_{\sigma^{-1}(1)} \cdot (a_{\sigma^{-1}(2)} \cdot (\cdots (a_{\sigma^{-1}(p+q-1)} \cdot a_{\sigma^{-1}(p+q)}) \cdots))).$$

Note that we are using σ^{-1} in the above formula. Related to this formula is Loday's observation [L3] that for a vector space V , the reduced tensor module

$$\bar{T}(V) = \sum_{k \geq 1} V^{\otimes k}$$

carries the structure of a free dual Leibniz algebra for which

$$(2.6) \quad (v_0, v_1, \dots, v_p) \cdot (v_{p+1}, \dots, v_{p+q}) \\ = \sum_{\sigma \in Sh_{p,q}} (v_0, v_{\sigma^{-1}(1)}, v_{\sigma^{-1}(2)}, \dots, v_{\sigma^{-1}(p+q)})$$

for homogeneous elements v_i . Signs can be used in (2.6) if $\bar{T}(V)$ is given a grading.

An essential technique for the calculation of $HL^*(\mathfrak{g})$ in the special case of a Lie algebra is the Pirashvili spectral sequence [P] which uses $H_{\text{Lie}}^*(\mathfrak{g})$ and $H_{\text{Lie}}^*(\mathfrak{g}; \mathfrak{g}')$ to glean information about $HL^*(\mathfrak{g})$. In [P], Pirashvili provides the details for Leibniz homology. Here we outline the construction

for cohomology and show that Loday's product is defined on the filtration which yields the cohomology spectral sequence. To begin, let \mathfrak{g} be a Lie algebra and define $\Omega^n(\mathfrak{g})$ to be the real vector space of skew-symmetric (alternating) homomorphisms $\mathfrak{g}^{\otimes n} \rightarrow \mathbf{R}$. If \mathfrak{g} has a topology, then all morphisms are taken to be continuous. For $\alpha \in \Omega^n(\mathfrak{g})$, it is evident that

$$\alpha(g_1, \dots, g_i, \dots, g_j, \dots, g_n) = -\alpha(g_1, \dots, g_j, \dots, g_i, \dots, g_n),$$

and $\Omega^*(\mathfrak{g})$ is the cochain complex for $H_{\text{Lie}}^*(\mathfrak{g})$. We have a short exact sequence

$$0 \longrightarrow \Omega^*(\mathfrak{g}) \longrightarrow C^*(\mathfrak{g}) \longrightarrow C^*(\mathfrak{g})/\Omega^*(\mathfrak{g}) \longrightarrow 0,$$

where $C^n(\mathfrak{g}) = \text{Hom}_{\mathbf{R}}(\mathfrak{g}^{\otimes n}, \mathbf{R})$. Clearly $\Omega^0(\mathfrak{g}) = C^0(\mathfrak{g})$ and $\Omega^1(\mathfrak{g}) = C^1(\mathfrak{g})$. Following Pirashvili's grading, set

$$C_{\text{rel}}^*[2] = C^*(\mathfrak{g})/\Omega^*(\mathfrak{g}).$$

We then have a long exact sequence

$$(2.7) \quad \cdots \longrightarrow H_{\text{Lie}}^n(\mathfrak{g}) \longrightarrow HL^n(\mathfrak{g}) \longrightarrow H_{\text{rel}}^{n-2}(\mathfrak{g}) \longrightarrow H_{\text{Lie}}^{n+1}(\mathfrak{g}) \longrightarrow \cdots$$

Define a left representation of \mathfrak{g} on $\mathfrak{g}' = \text{Hom}_{\mathbf{R}}(\mathfrak{g}, \mathbf{R})$ by

$$(g\gamma)(h) = \gamma([h, g])$$

where $g, h \in \mathfrak{g}$ and $\gamma \in \mathfrak{g}'$. Let $\Omega^n(\mathfrak{g}; \mathfrak{g}')$ denote the vector space of skew-symmetric homomorphisms $\alpha : \mathfrak{g}^{\otimes n} \rightarrow \mathfrak{g}'$. Thus, $\Omega^*(\mathfrak{g}; \mathfrak{g}')$ is the cochain complex for $H_{\text{Lie}}^*(\mathfrak{g}; \mathfrak{g}')$. There are inclusions of cochain complexes

$$i_1 : \Omega^{n+1}(\mathfrak{g}) \rightarrow \Omega^n(\mathfrak{g}; \mathfrak{g}'), \quad i_2 : \Omega^n(\mathfrak{g}; \mathfrak{g}') \rightarrow C^{n+1}(\mathfrak{g})$$

given by

$$(i_1(\alpha)(g_1, g_2, \dots, g_n))(g_0) = (-1)^n \alpha(g_0, g_1, g_2, \dots, g_n)$$

$$(i_2(\beta))(g_0, g_1, \dots, g_n) = (-1)^n \beta(g_1, \dots, g_n)(g_0).$$

Note that i_1 has occurred as var in (2.3), and i_2 allows us to consider an element of $\Omega^n(\mathfrak{g}; \mathfrak{g}')$ as an element of $C^{n+1}(\mathfrak{g})$ which is alternating on the last n tensor factors of $\mathfrak{g}^{\otimes(n+1)}$. In [F] $H_{\text{Lie}}^*(\mathfrak{g}; \mathfrak{g}')$ is given the structure of a right module over $H_{\text{Lie}}^*(\mathfrak{g})$, which is induced by a map of cochain complexes

$$\bullet : \Omega^p(\mathfrak{g}; \mathfrak{g}') \otimes \Omega^q(\mathfrak{g}) \rightarrow \Omega^{p+q}(\mathfrak{g}; \mathfrak{g}').$$

For $\alpha \in \Omega^p(\mathfrak{g}; \mathfrak{g}')$ and $\beta \in \Omega^q(\mathfrak{g})$,

$$\begin{aligned} (\alpha \bullet \beta)(g_1, g_2, \dots, g_{p+q})(g_0) \\ = \sum_{\sigma \in Sh_{p,q}} \text{sgn}(\sigma) \left(\alpha(g_{\sigma(1)}, \dots, g_{\sigma(p)})(g_0) \right) \left(\beta(g_{\sigma(p+1)}, \dots, g_{\sigma(p+q)}) \right). \end{aligned}$$

Then $d(\alpha \bullet \beta) = (d\alpha) \bullet \beta + (-1)^p \alpha \bullet d\beta$, and

$$i_2(\alpha \bullet \beta) = (i_2(\alpha)) \cdot \beta,$$

where the small dot, \cdot , is the product defined in (2.4).

The short exact sequence

$$0 \longrightarrow \Omega^*(\mathfrak{g}) \longrightarrow \Omega^{*-1}(\mathfrak{g}; \mathfrak{g}') \longrightarrow CR^*(\mathfrak{g})[1] \longrightarrow 0,$$

for which $CR^*(\mathfrak{g})[1] = \Omega^{*-1}(\mathfrak{g}; \mathfrak{g}')/\Omega^*(\mathfrak{g})$, yields the long exact sequence

$$(2.8) \quad \cdots \longrightarrow H_{\text{Lie}}^n(\mathfrak{g}) \longrightarrow H_{\text{Lie}}^{n-1}(\mathfrak{g}; \mathfrak{g}') \longrightarrow HR^{n-2}(\mathfrak{g}) \longrightarrow H_{\text{Lie}}^{n+1}(\mathfrak{g}) \longrightarrow \cdots.$$

Consider now the decreasing filtration of C_{rel}^* given by $F_*^0 = C_{\text{rel}}^*$, and for $s \geq 1$, $F_*^s = \{f \in C^*(\mathfrak{g}) \mid f \text{ is alternating in the last } (s+1)\text{-many tensor factors} / \Omega^*(\mathfrak{g})\}$.

2.9. LEMMA. — *If $f : \mathfrak{g}^{\otimes(n+s)} \rightarrow \mathbf{R}$ is alternating in the last s -many factors, then*

$$df : \mathfrak{g}^{\otimes(n+1+s)} \rightarrow \mathbf{R}$$

is alternating in the last s -many factors.

Proof. — This can be checked by hand. Compare with [P]. \square

2.10. THEOREM. — *There is a spectral sequence converging to $H_{\text{rel}}^*(\mathfrak{g})$ with*

$$E_2^{s,n} \simeq HL^n(\mathfrak{g}) \otimes HR^s(\mathfrak{g})$$

provided that $HL^(\mathfrak{g})$ and $HR^*(\mathfrak{g})$ are finite dimensional vector spaces in each dimension. The completed tensor product can be used if this finiteness condition is not satisfied.*

Proof. — Let $\hat{\otimes}$ denote the completed tensor product. Then

$$(F^s/F^{s+1})_n \simeq A/B$$

$$A = C^n(\mathfrak{g}) \hat{\otimes} \{f \in C^{s+2}(\mathfrak{g}) \mid f \text{ is alternating in last } (s+1) \text{ tensor factors}\}$$

$$B = C^n(\mathfrak{g}) \hat{\otimes} \Omega^{s+2}(\mathfrak{g}).$$

Thus, $E_0^{s,n} \simeq C^n(\mathfrak{g}) \hat{\otimes} CR^s(\mathfrak{g})$, and $E_1^{s,n} \simeq HL^n(\mathfrak{g}) \hat{\otimes} CR^s(\mathfrak{g})$. Since the co-adjoint action of \mathfrak{g} on $HL^*(\mathfrak{g})$ is trivial [L1, 10.1.7], we conclude that

$$E_2^{s,n} \simeq HL^n(\mathfrak{g}) \hat{\otimes} HR^s(\mathfrak{g}).$$

If $HL^*(\mathfrak{g})$ and $HR^*(\mathfrak{g})$ are finitely generated in each dimension, then

$$E_2^{s,n} \simeq HL^n(\mathfrak{g}) \otimes HR^s(\mathfrak{g}).$$

\square

2.11. THEOREM. — *If $\alpha : \mathfrak{g}^{\otimes p} \rightarrow \mathbf{R}$ and $\beta : \mathfrak{g}^{\otimes q} \rightarrow \mathbf{R}$ are alternating in the last s factors, where $s \leq p$ and $s \leq q$, then $\alpha \cdot \beta : \mathfrak{g}^{\otimes(p+q)} \rightarrow \mathbf{R}$ is alternating in the last s factors, i.e. $F^{s-1} \cdot F^{s-1} \subset F^{s-1}$.*

Proof. — It suffices to show that for $p+q-s < i < p+q$, we have

$$(\alpha \cdot \beta)(g_1, \dots, g_i, g_{i+1}, \dots, g_{p+q}) = -(\alpha \cdot \beta)(g_1, \dots, g_{i+1}, g_i, \dots, g_{p+q}).$$

Let σ be a $(p-1, q)$ -shuffle of $\{2, 3, \dots, p+q\}$. There are four possibilities to consider.

(i) Suppose that both i and $i+1$ appear in the sequence

$$\sigma(p+1), \sigma(p+2), \dots, \sigma(p+q).$$

Since there are at most $p+q-(i+1)$ -many possible choices for j so that $i+1 < \sigma(j)$, the integers i and $i+1$ must appear among the last s -many terms of the increasing sequence

$$\sigma(p+1), \sigma(p+2), \dots, \sigma(p+q).$$

(ii) Suppose that i appears in the sequence

$$\sigma(2), \sigma(3), \dots, \sigma(p)$$

and $i+1$ appears in the sequence

$$\sigma(p+1), \sigma(p+2), \dots, \sigma(p+q).$$

Let $\sigma(x) = i$ and $\sigma(y) = i+1$. Then $\hat{\sigma}$ defined by

$$\hat{\sigma}(j) = \begin{cases} \sigma(j), & j \neq x, j \neq y \\ \sigma(x), & j = y \\ \sigma(y), & j = x, \end{cases}$$

is a $(p-1, q)$ -shuffle of $\{2, 3, \dots, p+q\}$ having the opposite sign of σ .

(iii) If i appears in the sequence

$$\sigma(p+1), \sigma(p+2), \dots, \sigma(p+q),$$

and $i+1$ appears in the sequence $\sigma(2), \sigma(3), \dots, \sigma(p)$, then proceed as in (ii).

(iv) If both i and $i+1$ appear in the sequence $\sigma(2), \sigma(3), \dots, \sigma(p)$, then argue as in (i). \square

It may well be that certain terms in the sum for $\alpha \cdot \beta$ lie in filtration degree greater than s , which is useful in the computation of higher differentials for the Pirashvili spectral sequence.

For the reader's convenience we record the results of the calculation of $H_{\text{Lie}}^*(W_n)$ [B] [F] [Gb]. First observe that there is a monomorphism of Lie algebras

$$\varphi : \mathfrak{gl}_n(\mathbf{R}) \rightarrow W_n$$

given by $\varphi(E_{ij}) = x_i \frac{\partial}{\partial x_j}$, where E_{ij} is the elementary matrix with 1 in the i -th row and j -th column, and zeroes everywhere else. The image of φ consists of the one jets in W_n . Using the Hochschild-Serre spectral sequence associated to a subalgebra of a Lie algebra [HS] and invariant theory [W], one has

2.12. THEOREM ([B] [F] [Gb]). — *The E_2 -term of the Hochschild-Serre spectral sequence converging to $H_{\text{Lie}}^*(W_n)$ has form*

$$E_2^{*,*} \simeq H_{\text{Lie}}^*(\mathfrak{gl}_n(\mathbf{R})) \otimes (\mathbf{R}[P_1, \dots, P_n]/I),$$

where $\mathbf{R}[P_1, \dots, P_n]$ is the polynomial algebra with $\deg(P_j) = 2j$, and I is the ideal of $\mathbf{R}[P_1, \dots, P_n]$ generated by polynomials of degree greater than $2n$. Moreover, letting

$$H_{\text{Lie}}^*(\mathfrak{gl}_n(\mathbf{R})) = \Lambda(u_1, u_2, \dots, u_n),$$

for certain generators u_i with $\deg(u_i) = 2i - 1$, the differentials are determined by

$$d_{2r}(u_r) = P_r, r = 1, 2, \dots, n.$$

2.13. COROLLARY ([Gb]). — *We have $H_{\text{Lie}}^i(W_n) = 0$ for $1 \leq i \leq 2n$ and $i > n^2 + 2n$.*

2.14. COROLLARY ([Gb]). — *The Vey basis for $H_{\text{Lie}}^*(W_n)$ is given by the monomials*

$$u_{i_1} u_{i_2}, \dots, u_{i_r} P_{j_1} P_{j_2}, \dots, P_{j_s}$$

where

$$1 \leq i_1 < i_2 < \dots < i_r \leq n,$$

$$1 \leq j_1 \leq j_2 \leq \dots \leq j_s \leq n,$$

$$j_1 + j_2 + \dots + j_s \leq n,$$

$$i_1 + j_1 + j_2 + \dots + j_s > n,$$

$$i_1 \leq j_1.$$

The above monomial is then in degree

$$2(i_1 + i_2 + \dots + i_r + j_1 + j_2 + \dots + j_s) - r.$$

The following computation of Feigin, Fuks and Gelfand [F] is quite useful:

2.15. THEOREM ([F]). — *There is a natural isomorphism*

$$H_{\text{Lie}}^q(W_n; W'_n) \xrightarrow{\simeq} H_{\text{Lie}}^{2n+1}(W_n) \otimes H_{\text{Lie}}^{q-2n}(\mathfrak{gl}_n(\mathbf{R})),$$

and

$$\text{var} : H_{\text{Lie}}^{2n+1}(W_n) \rightarrow H_{\text{Lie}}^{2n}(W_n; W'_n)$$

is an isomorphism, where var is given in (2.3).

For its applications to the Godbillon-Vey invariant, we prove the following:

2.16. LEMMA. — *The natural map*

$$H_{\text{Lie}}^{2n+1}(W_n) \rightarrow HL^{2n+1}(W_n)$$

is an isomorphism.

Proof. — From the Vey basis, we see that $H_{\text{Lie}}^{2n+2}(W_n) = 0$. Since

$$\text{var} : H_{\text{Lie}}^{2n+1}(W_n) \rightarrow H_{\text{Lie}}^{2n}(W_n; W'_n)$$

is an isomorphism, it follows from (2.8) that $HR^{2n-1}(W_n) = 0$. Actually,

$$HR^i(W_n) = 0 \quad \text{for } i = 0, 1, 2, \dots, 2n-1,$$

and, therefore, $H_{\text{rel}}^i(W_n) = 0$ for $i = 0, 1, \dots, 2n-1$. The lemma follows from (2.7). \square

2.17. COROLLARY. — *If α_{GV} is a cochain representing the universal Godbillon-Vey invariant in $H_{\text{Lie}}^{2n+1}(W_n)$, then α_{GV} represents a non-zero class in $HL^{2n+1}(W_n)$.*

Proof. — The corollary follows from (2.16). \square

We may now state the main calculational result of the paper.

2.18. THEOREM. — *Letting T denote the tensor algebra, there is an isomorphism of dual Leibniz algebras (n any positive integer)*

$$HL^*(W_n) \simeq \mathbf{R} \oplus [\text{Im} \oplus A] \otimes T(A \oplus B),$$

$$\text{Im} = \text{Im}(\text{var})$$

$$A = \text{coker}(\text{var})[1]$$

$$B \simeq \ker(\text{var})[-1],$$

where \mathbf{R} is in dimension zero.

Proof. — We illustrate several techniques germane to the proof by considering the special case $n = 1$ for which

$$\overline{H}_{\text{Lie}}^q(W_1) = \begin{cases} \mathbf{R}, & q = 3 \\ 0, & \text{otherwise,} \end{cases}$$

and $\ker(\text{var}) = 0$. Note that $HR^2(W_1)$ is generated (as a vector space) by a tensor $h : W_1^{\otimes 4} \rightarrow \mathbf{R}$,

$$h \in \text{coker}[\text{var} : \Omega^4(W_1) \rightarrow \Omega^3(W_1; W_1')],$$

and $HR^i(W_1) = 0$ for $i \neq 2$. Moreover the alternating tensor $f : W_1^{\otimes 3} \rightarrow \mathbf{R}$ corresponding to the Godbillon-Vey invariant generates $HL^3(W_1)$. In the Pirashvili spectral sequence $d_r = 0$ for $r \geq 2$, whence as vector spaces

$$(2.19) \quad HL^*(W_1) \simeq \mathbf{R} \oplus [HL^3(W_1) \oplus HR^2(W_1)] \otimes T(HR^2(W_1)).$$

Certainly $f^2 = 0$ in $HL^*(W_1)$, since $HL^6(W_1) = 0$, but also since $f^2 \in F^4$ (the fourth filtration degree), and

$$H_*(F^4/F^5) = 0.$$

We now show that $h \otimes h$ is represented by h^2 (i.e. $h \cdot h$) in $HL^*(W_1)$. This would follow by showing that $h \otimes h$ is represented by h^2 in $H_*(F^2/F^3)$. From (2.11) we conclude that $h^2 - h \otimes h \in F^2$. By omitting the identity permutation in the definition of Loday's product, we have

$$(2.20) \quad h^2 - h \otimes h \in F^3.$$

The proof of this requires four cases as in (2.11) and the fact that no permutation shuffles a factor into the first tensor position, where h is not alternating. (See (2.4) for the formula of the shuffles.) Thus, $h^2 - h \otimes h = 0$ in E_1 and in $HL^*(W_1)$. By a similar argument,

$$h \cdot (h \cdot h), \quad h \otimes h^2, \quad \text{and} \quad h \otimes h \otimes h$$

represent the same element in $HL^*(W_1)$. By induction, the cochains

$$\underbrace{h \cdot (h \cdot (h \cdots (h \cdot h) \cdots))}_{n\text{-many}} \quad \text{and} \quad \underbrace{h \otimes h \otimes \cdots \otimes h}_{n\text{-many}}$$

represent the same element in $HL^*(W_1)$. A similar statement is valid for

$$f \cdot (h \cdot (h \cdots (h \cdot h) \cdots)) \quad \text{and} \quad f \otimes h \otimes h \otimes \cdots \otimes h.$$

Of course, hf represents zero in $HL^*(W_1)$, since $hf \in F^5$. For $n > 1$, this type of filtration argument can be used to determine products between elements in $T(A \oplus B)$ and Im .

For any positive integer n and $q > n^2 + 2n$, we have $H_{\text{Lie}}^q(W_n) = 0$, and, therefore,

$$HL^q(W_n) \simeq H_{\text{rel}}^{q-2}(W_n).$$

Since $H_{\text{Lie}}^q(W_n) = 0$ for $q = 1, 2, \dots, 2n$, and $H^q(W_n; W'_n) = 0$ for $q = 0, 1, 2, \dots, 2n - 1$, we see that $HL^q(W_n) = 0$ for $q = 1, 2, \dots, 2n$. Also, $HR^q(W_n) = 0$ for $q = 0, 1, 2, \dots, 2n - 1$ and for $q \geq n^2 + 2n$. It follows that for $0 \leq q \leq 4n$

$$H_{\text{rel}}^q(W_n) \simeq HR^q(W_n) \simeq \left(\ker(\text{var})[-3] \oplus \text{coker}(\text{var})[-1] \right)_q.$$

Thus, for $1 \leq q \leq 4n + 2$

$$HL^q(W_n) \simeq \text{Im}(\text{var}) \oplus \text{coker}(\text{var})[+1].$$

We now show that for $r \geq 2$, $d_r(\alpha \otimes \beta) = 0$, where $\alpha \in HL^q(W_n)$, $1 \leq q \leq 4n + 2$, and

$$\beta \in HR^j(W_n) \simeq \ker(\text{var})[-3] \oplus \text{coker}(\text{var})[-1], \quad j \geq 0.$$

In the E^2 term represent $\alpha \otimes \beta$ as $\alpha \cdot \beta$ in a fashion similar to (2.20). Then

$$d(\alpha \cdot \beta) = (d\alpha) \cdot \beta + (-1)^{|\alpha|} \alpha \cdot (d\beta).$$

Necessarily $d\alpha = 0$. If β corresponds to an element of $\text{coker}(\text{var})$, then $d\beta = 0$. If β corresponds to an element of $\ker(\text{var})$, then

$$d\beta \in \Omega^*(W_n),$$

and $d\beta$ represents zero in $HL^*(W_n)$. Thus, $\alpha \cdot d\beta$ is zero in $HL^*(W_n)$, and the map

$$HL^*(W_n) \rightarrow H_{\text{rel}}^{*-2}(W_n)$$

sends the Leibniz cohomology class of $\alpha \cdot d\beta$ to zero. It follows that $d_r(\alpha \otimes \beta) = 0$, $r \geq 2$.

To compute the boundary map

$$\partial : H_{\text{rel}}^q(W_n) \rightarrow H_{\text{Lie}}^{q+3}(W_n), \quad q \geq 4n + 1,$$

first consider elements in $H_{\text{rel}}^q(W_n)$ represented by $\alpha \cdot \beta$, where either $\alpha \in \text{Im}(\text{var})$ or $\alpha \in \text{coker}(\text{var})$, and either $\beta \in \text{coker}(\text{var})$ or β corresponds to an element of $\ker(\text{var})$. Then as cochains

$$\partial(\alpha \cdot \beta) = (-1)^{k_1} \alpha \cdot (d\beta) \in \Omega^{q+3}(W_n),$$

where $k_1 = |\alpha|$, $k_2 = |\beta|$. Either $d\beta = 0$ or $d\beta \in \Omega^*(W_n)$ as above. The notation $S(\alpha \cdot (d\beta))$ is introduced to denote the cochain $\alpha \cdot (d\beta)$ symmetrized over the symmetric group, i.e.

$$S(\alpha \cdot (d\beta))(g_1 \otimes \cdots \otimes g_{q+3}) = (\alpha \cdot (d\beta)) \left(\sum_{\sigma \in \Sigma_{q+3}} \text{sgn}(\sigma) g_{\sigma(1)} \otimes \cdots \otimes g_{\sigma(q+3)} \right)$$

for $g_1, \dots, g_{q+3} \in W_n$. On one hand,

$$S(\alpha \cdot (d\beta)) = (-1)^{k_1} S(\partial(\alpha \cdot \beta)) = (-1)^{k_1} (q+3)! \partial(\alpha \cdot \beta).$$

On the other hand,

$$S(\alpha \cdot (d\beta)) = \binom{k_1 + k_2}{k_1 - 1} (S\alpha) \wedge (Sd\beta),$$

where \wedge denotes the usual Lie algebra product of cochains. Certainly $Sd\beta$ is a Lie algebra cocycle. We argue that $S\alpha$ is as well, and use the vanishing of all products in $H_{\text{Lie}}^*(W_n)$ [F], p. 79, to conclude that the class of

$$(S\alpha) \wedge (Sd\beta)$$

is zero in $H_{\text{Lie}}^*(W_n)$. Since α is not necessarily skew-symmetric (as an element of $C^*(W_n)$), we cannot conclude that

$$d(S\alpha) = \frac{S(d\alpha)}{k_1 + 1}.$$

However, using (2.15), α may be written as $\rho_1 \otimes \rho_2$, where

$$\rho_1 \in \text{Im} \left(\text{var} : \Omega^{2n+1}(W_n) \rightarrow \Omega^{2n}(W_n; W'_n) \right),$$

and $\rho_2 \in \Omega^*(W_n)$. Furthermore, in the Hochschild-Serre spectral sequence calculation for $H_{\text{Lie}}^*(W_n; W'_n)$ [F], p. 100, $\rho_1 \otimes \rho_2$ can be represented as $\rho_1 \bullet \rho_2$ when considering $\Omega^*(W_n; W'_n)$ as a right module over $\Omega^*(W_n)$. Let $p_1 = |\rho_1|$ and $p_2 = |\rho_2|$, where degree is computed as elements of $\Omega^*(W_n)$. Necessarily $d\rho_1 = 0$. We have

$$0 = d\alpha = (-1)^{p_1} \rho_1 \bullet (d\rho_2)$$

$$0 = S(\rho_1 \bullet (d\rho_2)) = \binom{p_1 + p_2 - 1}{p_1 - 1} (S\rho_1) \wedge (Sd\rho_2).$$

Since ρ_1 and $d\rho_2$ are skew-symmetric, $0 = \rho_1 \wedge (d\rho_2)$. Now,

$$\begin{aligned} S\alpha &= S(\rho_1 \bullet \rho_2) = \binom{p_1 + p_2 - 1}{p_1 - 1} (S\rho_1) \wedge (S\rho_2) \\ &= p_1(p_1 + p_2 - 1)! \rho_1 \wedge \rho_2, \end{aligned}$$

and

$$d(S\alpha) = (-1)^{p_1} p_1(p_1 + p_2 - 1)! \rho_1 \wedge (d\rho_2) = 0.$$

Thus, $S\alpha$ is a Lie-algebra cocycle.

Technically the proof proceeds by induction on the number of tensor factors in a monomial in $HL^*(W_n)$, since the Pirashvili spectral sequence for $H_{\text{rel}}^*(W_n)$ requires knowledge of $HL^*(W_n)$ in lower dimensions. Having begun this induction, we show that

$$d_r(\alpha_1 \cdot (\alpha_2 \cdot (\dots (\alpha_{k-1} \cdot \alpha_k) \dots))) = 0, r \geq 2,$$

where $\alpha_1 \in \text{Im}(\text{var})$ or $\alpha_1 \in \text{coker}(\text{var})$, and for $i \geq 2$, either $\alpha_i \in \text{coker}(\text{var})$ or α_i corresponds to an element in $\text{ker}(\text{var})$. This follows from

$$\begin{aligned} d(\alpha_1 \cdot (\alpha_2 \cdot (\cdots (\alpha_{k-1} \cdot \alpha_k) \cdots))) \\ = (d\alpha_1) \cdot (\alpha_2 \cdot (\cdots (\alpha_{k-1} \cdot \alpha_k) \cdots)) \\ + (-1)^{|\alpha_1|} \alpha_1 \cdot d(\alpha_2 \cdot (\alpha_3 \cdots (\alpha_{k-1} \cdot \alpha_k) \cdots)). \end{aligned}$$

As a cochain, $d\alpha_1 = 0$. Letting

$$\gamma = \alpha_2 \cdot (\alpha_3 \cdot (\cdots (\alpha_{k-1} \cdot \alpha_k) \cdots)),$$

we note that $d\gamma$ represents zero in $HL^*(W_n)$. Thus, $\alpha_1 \cdot d\gamma$ is zero in $HL^*(W_n)$, and the map

$$HL^*(W_n) \rightarrow H_{\text{rel}}^{*-2}(W_n)$$

sends the class of $\alpha_1 \cdot d\gamma$ to zero.

To conclude the computation of $\partial : H_{\text{rel}}^q(W_n) \rightarrow H_{\text{Lie}}^{q+3}(W_n)$, note that

$$\begin{aligned} \partial(\alpha_1 \cdot (\alpha_2 \cdot (\cdots (\alpha_{k-1} \cdot \alpha_k) \cdots))) \\ = \pm \alpha_1 \cdot (d\alpha_2 \cdot (\cdots (\alpha_{k-1} \cdot \alpha_k) \cdots)) \\ \vdots \\ \pm \alpha_1 \cdot (\alpha_2 \cdot (\cdots ((d\alpha_{k-1}) \cdot \alpha_k) \cdots)) \\ \pm \alpha_1 \cdot (\alpha_2 \cdot (\cdots (\alpha_{k-1} \cdot (d\alpha_k)) \cdots)) \in \Omega^*(W_n). \end{aligned}$$

Thus, $\partial(\alpha_1 \cdot (\alpha_2 \cdot (\cdots (\alpha_{k-1} \cdot \alpha_k) \cdots)))$ is a linear combination of the cochains

$$\begin{aligned} S\alpha_1 \wedge S(d\alpha_2) \wedge S\alpha_3 \wedge \cdots \wedge S\alpha_k, \\ S\alpha_1 \wedge S\alpha_2 \wedge S(d\alpha_3) \wedge \cdots \wedge S\alpha_k, \\ \vdots \\ S\alpha_1 \wedge S\alpha_2 \wedge S\alpha_3 \wedge \cdots \wedge S(d\alpha_k). \end{aligned}$$

Letting $\beta = \alpha_i, i \geq 2$, note that $\alpha_1 \cdot \beta$ is a Leibniz cocycle. Thus,

$$0 = d(\alpha_1 \cdot \beta) = \alpha_1 \cdot (d\beta).$$

From $S(\alpha_1 \cdot (d\beta)) = 0$, it follows that over \mathbf{R} , $S\alpha_1 \wedge (Sd\beta) = 0$. Thus, each summand in the expression for

$$\partial(\alpha_1 \cdot (\alpha_2 \cdot (\cdots (\alpha_{k-1} \cdot \alpha_k) \cdots)))$$

is zero. □

3. The characteristic map for Leibniz cohomology.

Recall that for a C^∞ , codimension n foliation \mathfrak{F} on M with trivial normal bundle, there is a characteristic homomorphism [B] [F] [H]

$$\text{char}_{\mathfrak{F}} : H_{\text{Lie}}^*(W_n) \rightarrow H_{DR}^*(M).$$

In this section we conjecture the existence of a homomorphism

$$L_{\mathfrak{F}} : HL^*(W_n) \rightarrow HL^*(\chi(M); C^\infty(M))$$

so that the following diagram commutes

$$(3.1) \quad \begin{array}{ccc} H_{\text{Lie}}^*(W_n) & \xrightarrow{\text{char}_{\mathfrak{F}}} & H_{DR}^*(M) \\ \downarrow & & \downarrow \\ HL^*(W_n) & \xrightarrow{L_{\mathfrak{F}}} & HL^*(\chi(M); C^\infty(M)), \end{array}$$

where $H_{DR}^*(M) \rightarrow HL^*(\chi(M); C^\infty(M))$ is given in (1.8). A fixed foliation determines a splitting of tangent bundle

$$(3.2) \quad TM \simeq T_L M \oplus T_\eta M,$$

where $T_L M$ is the tangent space to the leaves and $T_\eta M$ is the normal bundle. For the special case $n = 1$, let ω be a determining one-form for the foliation \mathfrak{F} . Then $\omega(v_p) = 0$ for $v_p \in T_L M$, and $\omega(v_p) = 1$, where v_p is chosen to be a unit vector with positive orientation in $T_\eta M$. Moreover, the splitting of the tangent bundle (3.2) yields a vector space isomorphism

$$\chi(M) \simeq \chi_L(M) \oplus \chi_\eta(M),$$

where $\chi_L(M)$ are vector fields along the leaves and $\chi_\eta(M)$ are vector fields perpendicular to the leaves. Given any $X \in \chi(M)$, we may write X uniquely as a sum $X_L + X_\eta$, where

$$X_L \in \chi_L(M) \text{ and } X_\eta \in \chi_\eta(M).$$

Define $L_\omega : \chi(M) \rightarrow C^\infty(M)$ by $L_\omega(X) = f$, where $f(p) = \omega(X_\eta(p))$ for all $p \in M$. Then L_ω is simply the image of ω under the inclusion

$$\Omega^1(M) \rightarrow \text{Hom}_{\mathbf{R}}^{\text{cont}}(\chi(M), C^\infty(M))$$

given in (1.8). In this way, all canonical one-forms used to define $\text{char}_{\mathfrak{F}}$ in the commutative framework occur as elements of

$$\text{Hom}_{\mathbf{R}}^{\text{cont}}(\chi(M), C^\infty(M)).$$

Acknowledgements. — The author would like to express his sincere gratitude to the Centre National de la Recherche Scientifique whose support permitted a productive collaboration with Jean-Louis Loday at the Institut de Recherche Mathématique Avancée in Strasbourg.

BIBLIOGRAPHY

- [B] R. BOTT, Lectures on Characteristic Classes and Foliations, Lecture Notes in Mathematics, 279 (1972), 1–94.
- [BS] R. BOTT, G. SEGAL, The Cohomology of the Vector Fields on a Manifold, *Topology*, 16 (1977), 285–298.
- [CE] C. CHEVALLEY, S. EILENBERG, Cohomology Theory of Lie groups and Lie algebras, *Trans. Amer. Math. Soc.*, 63 (1948), 85–124.
- [C] A. CONNES, *Géométrie Non Commutative*, Inter Editions, Paris, 1990.
- [F] D.B. FUKS, *Cohomology of Infinite-Dimensional Lie Algebras*, Consultants Bureau, 1986 (A.B. Sosinskii translator).
- [GK] V. GINZBURG, M.M. KAPRANOV, Koszul Duality for Operads, *Duke Jour. Math.*, 76-1 (1994), 203–272.
- [Gb] C. GODBILLON, Cohomologies d'algèbres de Lie de champs de vecteurs formels, *Séminaire Bourbaki*, 421 (1972).
- [Gw] T.G. GOODWILLIE, Relative Algebraic K-Theory and Cyclic Homology, *Annals of Math.*, 124 (1986), 347–402.
- [H] A. HAEFLIGER, Sur les classes caractéristiques des feuilletages, *Séminaire Bourbaki*, 412 (1972).
- [HS] G. HOCHSCHILD J.-P. SERRE, Cohomology of Lie Algebras, *Ann. of Math.*, 57 (1953), 591–603 .
- [K] V. KAČ, *Vertex Algebras for Beginners*, Am. Math. Soc., Univ. Lecture Series, 10 (1996).
- [K-S] Y. KOSMANN-SCHWARZBACH, From Poisson Algebras to Gerstenhaber Algebra, *Ann. Inst. Fourier, Grenoble*, 46-5 (1996), 1243–1274.
- [L1] J.-L. LODAY, *Cyclic Homology*, *Grund. Math. Wissen.* 301, Springer Verlag, 1992.
- [L2] J.-L. LODAY, Une version non commutative des algèbres de Lie: les algèbres de Leibniz, *L'Enseignement Math.*, 39 (1993), 269–293.
- [L3] J.-L. LODAY, Cup product for Leibniz cohomology and dual Leibniz algebras, *Math. Scand.*, 77-2 (1995), 189–196.
- [L4] J.-L. LODAY, La Renaissance des Opérades, *Séminaire Bourbaki*, 792 (1994–95).
- [LP] J.-L. LODAY, T. PIRASHVILI, Universal Enveloping Algebras of Leibniz Algebras and (Co)-homology, *Math. Annalen*, 296 (1993), 139–158.
- [P] T. PIRASHVILI, On Leibniz Homology, *Ann. Inst. Fourier, Grenoble*, 44-2 (1994), 401–411.
- [S] M. SPIVAK, *A Comprehensive Introduction to Differential Geometry*, Vol. I, Publish or Perish, Inc. (1979).
- [W] H. WEYL, *The Classical Groups*, Princeton University Press, 1946.

Manuscrit reçu le 24 mars 1997,
 accepté le 8 septembre 1997.

Jerry M. LODDER,
 New Mexico State University
 Mathematical Sciences
 Box MB
 Las Cruces, NM 88003 (USA).
 jlodder@nmsu.edu