

ANDRZEJ PLIS

**On sets filled by asymptotic solutions of
differential equations**

Annales de l'institut Fourier, tome 14, n° 1 (1964), p. 191-194

http://www.numdam.org/item?id=AIF_1964__14_1_191_0

© Annales de l'institut Fourier, 1964, tous droits réservés.

L'accès aux archives de la revue « Annales de l'institut Fourier » (<http://annalif.ujf-grenoble.fr/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

ON SETS FILLED BY ASYMPTOTIC SOLUTIONS OF DIFFERENTIAL EQUATIONS

A. PLIS (Krakow)

Consider an ordinary differential equation

$$(1) \quad \begin{aligned} x' &= f(t, x) \\ x &= (x_1, \dots, x_n), \quad f = (f_1, \dots, f_n). \end{aligned}$$

Assumption I. Suppose that the domain D of $f(t, x)$ is open, $f(t, x)$ is continuous on D and through each point

$$A: t = a_0, \quad x = a = (a_1, a_2, \dots, a_n)$$

of D passes only one integral $x = x(t, A)$ of (1).

Denote by $(\alpha(A), \beta(A))$ the maximal interval on which there exists the integral passing through A . We shall denote

$$X(t, A) = (t, x(t, A)) \quad \text{for} \quad t \in (\alpha(A), \beta(A)).$$

Let E be an open subset of D . In the following we shall deal with the set $Z(E)$ of such points A , that $X(t, A) \in E$ for $a_0 \leq t < \infty$. Obviously set $Z(E)$ depends on both set E and system (1). It is evident that $E \subset F$ implies $Z(E) \subset Z(F)$. Let φ be a family of subsets F of D . We shall consider the following properties of equation (1).

PROPERTY I (of equation (1) in respect to E and φ). — *For every $F \in \varphi$ $Z(E) \cap F$ is empty or consists of one point.*

PROPERTY II. — *For every $F \in \varphi$ $Z(E) \cap F$ is not empty.*

Let $I^+(A)$ denote the set of all points $B = X(t, A)$ for $t \geq a_0$.

We say that the point $A \in P(G) \cap D$, where $P(G)$ denotes the boundary of an open set G , is the point of egress from G (with respect to equation (1) and set D) if there exists such an integral $x(t)$ of (1) and a positive number $\varepsilon > 0$ that

$$x(a_0) = a \quad \text{and} \quad (t, x(t)) \in G$$

for $a_0 - \varepsilon < t < a_0$ (under Assumption I, $X(t, A) \in G$ for $a_0 - \varepsilon < t < a_0$). If no point of $P(G)$ is a point of egress from G then $A \in G$ implies $I^+(A) \subset G$. If Property I is satisfied and $B \in Z(E) \cap F$ then $(F - B) \cap Z(E) = \emptyset$, where $F - B$ denotes the set of all points of the set F except the point B . It follows that for every $A \in F$, $A \neq B$ either $I^+(A) \sim \in E$ or $\beta(A) < \infty$. Let G be such a set that $\bar{G} \cap E$ has no common point with a plane $t = c > a_0$, where \bar{G} denotes the closure of G , then $I^+(A) \subset \bar{G}$ implies $A \sim \in Z(E)$.

LEMMA. — *Suppose Assumption I and the following conditions. For each set $G_i (i = 1, \dots) G_i \cap E$ is contained in a halfspace $t < c_i$. No point of $P(G_i)$ is a point of egress. Set F satisfies inclusion $F - O \subset \bigcup_{i=1}^{\infty} G_i$.*

Then $(F - O) \cap Z(E) = \emptyset$.

THEOREM 1. — *Suppose Assumption I and the following conditions. The intersection $E(s)$ of a given set E and the plane $t = s$ satisfies the inequality $\text{diam}(E(t)) < p(t)$, where $p(t)$ is a positive function continuous on $(-\infty, \infty)$. No point of $P(G_i)$ is a point of egress in respect to the equation*

$$x' = f(t + a_0, x + a(t)) - f(t + a_0, a(t)),$$

where a_0 is a real number and $x = a(t)$ is such a Lipchitzian function that the right side of the equation is defined. Set F satisfies inclusion $F - O \subset \bigcup_{i=1}^{\infty} G_i$. For any i and s there exists a constant $c(i, s)$ that $\text{dist}(G_i(t), 0) \geq p(t + s)$ for $t \geq c(i, s)$, where $G_i(s)$ is the intersection of G_i and the plane $t = s$.

Under these assumptions if $A \in Z(E)$, then

$$(F(A) - A) \cap Z(E) = \emptyset,$$

where $F(A)$ denotes set obtained from A by translation of R^{n+1} transforming O on A .

THEOREM 2. — If assumptions of Theorem 1 are satisfied and F is a plane then equation (1) possesses property I in respect to E and the family of planes parallel to F (and of the same dimension).

Suppose now that set F is a plane and in the coordinate system t , $x = (u, \nu)$, $u = (u_1, \dots, u_k)$, $\nu = (\nu_1, \dots, \nu_{n-k})$ it has the equation $t = 0$, $u = 0$. Now Property I (for the family of planes $t = c_0$, $u = (c_1, \dots, c_k)$, c_i arbitrary) is necessary and sufficient for set $Z(E)$ to be the graph of a single-valued function $\nu = q(t, u)$. Putting $g = (f_1, \dots, f_k)$, $h = (f_{k+1}, \dots, f_n)$ system (1) takes the form

$$(2) \quad u' = g(t, u, \nu), \quad \nu' = h(t, u, \nu).$$

The following result formulated in terms of inequalities can be obtained from Theorem 1 formulated in terms of sets ⁽¹⁾

THEOREM 3. — Suppose that system (2) satisfies Assumption I and that the functions $g(t, u, \nu)$, $h(t, u, \nu)$ for

$$(t, u, \nu) \in D, \quad (t, \bar{u}, \bar{\nu}) \in D$$

satisfy inequalities

$$(3) \quad (g(t, u, \nu) - g(t, \bar{u}, \bar{\nu})) (u - \bar{u}) \leq \gamma(t) (u - \bar{u})^2$$

for $|\nu - \bar{\nu}| = |u - \bar{u}|$, where $|z|$ denotes Euclidean distance of point z from 0,

$$(4) \quad (h(t, u, \nu) - h(t, \bar{u}, \bar{\nu})) (\nu - \bar{\nu}) \geq \gamma(t) (\nu - \bar{\nu})^2,$$

for

$$|u - \bar{u}| \leq |\nu - \bar{\nu}|,$$

where $\gamma(t)$ is a function summable in every finite interval, and such that

$$\int_0^\infty \gamma(s) ds = \infty,$$

then set Z of points A lying on the integrals of (2) (remaining in D) bounded for $a_0 \leq t < \infty$ is a graph of a single-valued function $\nu = q(t, u)$ defined in a certain set $S (S \subset \mathbb{R}^{k+1})$ satisfying the Lipschitz condition with respect to all the variables

⁽¹⁾ Such kind of formulation was suggested by T. Wazewski.

and in particular the condition

$$|q(t, u) - q(t, \bar{u})| \leq |u - \bar{u}|$$

in the set S or the set Z is an empty set.

Theorem 3 is a particular case of theorem 2 in [1].

Now for illustration of Property II we present a variant of an example from [2].

Let system (2) satisfy Assumption I on a neighbourhood D of the set $H: |u| \leq 1, |\varphi| \leq 1, -\infty < t < \infty$. Moreover suppose that $g(t, u, \varphi)u < 0$ for $|u| = 1, |\varphi| \leq 1$ and arbitrary t , $h(t, u, \varphi) > 0$ for $|\varphi| = 1, |u| \leq 1$ and arbitrary t .

Under these assumptions for every $\bar{u}, |\bar{u}| < 1$ and arbitrary \bar{t} , there exists $\bar{\varphi}$, that $I^+(\bar{t}, \bar{u}, \bar{\varphi}) \in H$.

BIBLIOGRAPHY

- [1] A. PLIS, Sets filled by asymptotic integrals of ordinary differential equations, *Bull. Acad. Polon. Sci. Cl. III*, **4** (1956), 749-752.
- [2] T. WAZEWSKI, Sur un principe topologique de l'examen de l'allure asymptotique des intégrales des équations différentielles ordinaires, *Ann. Soc. Polon. Math.*, **20** (1947), 279.