



ANNALES

DE

L'INSTITUT FOURIER

Karl-Hermann NEEB

Non-abelian extensions of infinite-dimensional Lie groups

Tome 57, n° 1 (2007), p. 209-271.

http://aif.cedram.org/item?id=AIF_2007__57_1_209_0

© Association des Annales de l'institut Fourier, 2007, tous droits réservés.

L'accès aux articles de la revue « Annales de l'institut Fourier » (<http://aif.cedram.org/>), implique l'accord avec les conditions générales d'utilisation (<http://aif.cedram.org/legal/>). Toute reproduction en tout ou partie cet article sous quelque forme que ce soit pour tout usage autre que l'utilisation à fin strictement personnelle du copiste est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

cedram

Article mis en ligne dans le cadre du
Centre de diffusion des revues académiques de mathématiques
<http://www.cedram.org/>

NON-ABELIAN EXTENSIONS OF INFINITE-DIMENSIONAL LIE GROUPS

by Karl-Hermann NEEB

ABSTRACT. — In this article we study non-abelian extensions of a Lie group G modeled on a locally convex space by a Lie group N . The equivalence classes of such extension are grouped into those corresponding to a class of so-called smooth outer actions S of G on N . If S is given, we show that the corresponding set $\text{Ext}(G, N)_S$ of extension classes is a principal homogeneous space of the locally smooth cohomology group $H_{ss}^2(G, Z(N))_S$. To each S a locally smooth obstruction class $\chi(S)$ in a suitably defined cohomology group $H_{ss}^3(G, Z(N))_S$ is defined. It vanishes if and only if there is a corresponding extension of G by N . A central point is that we reduce many problems concerning extensions by non-abelian groups to questions on extensions by abelian groups, which have been dealt with in previous work. An important tool is a Lie theoretic concept of a smooth crossed module $\alpha: H \rightarrow G$, which we view as a central extension of a normal subgroup of G .

RÉSUMÉ. — Dans cet article nous étudions les extensions non abéliennes d'un groupe de Lie G modelé sur un espace localement convexe par un groupe de Lie N . Les classes d'équivalence de telles extensions sont groupées en celles qui correspondent à la classe des actions dites des actions extérieures S de G sur N . Si S est donné, nous montrons que l'ensemble correspondant $\text{Ext}(G, N)_S$ des classes d'extensions est un espace homogène principal du groupe de cohomologie localement lisse $H_{ss}^2(G, Z(N))_S$. Pour chaque S une obstruction localement lisse $\chi(S)$ dans un groupe de cohomologie $H_{ss}^3(G, Z(N))_S$ est définie. Elle s'annule si et seulement si il existe une extension correspondante de G par N . Un point central est que nous ramenons plusieurs problèmes concernant des extensions par des groupes non abéliens à des questions sur des extensions par des groupes abéliens, qui ont été étudiées dans des travaux antérieurs. Un outil important est une notion de module croisé lisse, relevant de la théorie de Lie, $\alpha: H \rightarrow G$, que nous voyons comme une extension centrale d'un sous-groupe normal de G .

Introduction

In the present paper we continue our investigation of extensions of infinite-dimensional Lie groups. In this sense it is a sequel to [18] and [19],

Keywords: Lie group extension, smooth outer action, crossed module, Lie group cohomology, automorphisms of group extension.

Math. classification: 22E65, 57T10, 22E15.

where we studied central, resp. abelian extensions. We now turn to extensions by groups which are not necessarily abelian.

The concept of a (not necessarily finite-dimensional) Lie group used here is that a *Lie group* G is a manifold modeled on a locally convex space endowed with a group structure for which the group operations are smooth (cf. [15, 8]). An *extension* of G by N is an exact sequence of Lie groups $N \hookrightarrow \widehat{G} \twoheadrightarrow G$ which defines a locally trivial smooth N -principal bundle over the Lie group G . It is called *abelian*, resp. *central* if N is abelian, resp. central in \widehat{G} . This setup implies in particular that for each extension $q: \widehat{G} \rightarrow G$ of G by N there exists a section $\sigma: G \rightarrow \widehat{G}$ of q mapping $\mathbf{1}$ to $\mathbf{1}$ which is smooth in an identity neighborhood of G . Starting with such a section, we consider the functions

$$S: G \rightarrow \text{Aut}(N), \quad S(g)(n) := \sigma(g)n\sigma(g)^{-1},$$

$$\omega: G \times G \rightarrow N, \quad (g, g') \mapsto \sigma(g)\sigma(g')\sigma(gg')^{-1}.$$

Then the map $N \times G \rightarrow \widehat{G}$, $(n, g) \mapsto n\sigma(g)$ is a bijection, and transferring the group structure from \widehat{G} to $N \times G$ yields

$$(\dagger) \quad (n \cdot g)(n'g') = (nS(g)(n')\omega(g, g'), gg').$$

In the first half of the present paper we are mainly concerned with the appropriate smoothness requirements for the functions S and ω under which the product set $N \times G$ with the group structure given by (\dagger) carries a Lie group structure for which $N \times G \rightarrow G$, $(n, g) \mapsto g$ is an extension of Lie groups, and when such Lie group extensions are equivalent. The natural context to deal with such questions is a non-abelian locally smooth Lie group cohomology introduced in Section 1. In Section 2 the cohomological setup is linked to the classification of Lie group extensions of G by N . In particular we show that all such extensions are constructed from pairs of maps (S, ω) as above, where

- (1) $S(g)S(g') = c_{\omega(g, g')}S(gg')$ for $g, g' \in G$ and $c_n(n') := nn'n^{-1}$ denotes conjugation.
- (2) $S(g)(\omega(g', g''))\omega(g, g'g'') = \omega(g, g')\omega(gg', g'')$ for $g, g', g'' \in G$.
- (3) The map $G \times N \rightarrow N$, $(g, n) \mapsto S(g)n$ is smooth on a set of the form $U \times N$, where U is an identity neighborhood of G .
- (4) ω is smooth in an identity neighborhood.
- (5) For each $g \in G$ the map $\omega_g: G \rightarrow N$, $x \mapsto \omega(g, x)\omega(gxg^{-1}, g)^{-1}$ is smooth in an identity neighborhood of G .

Conditions (1) and (2) ensure that $N \times G$ is a group with the multiplication defined by (\dagger) , and (3)–(5) imply that $N \times G$ can be endowed with a

manifold structure turning it into a Lie group. In general this manifold will not be diffeomorphic to the product manifold $N \times G$, but the projection onto G will define a smooth N -principal bundle. A map $S: G \rightarrow \text{Aut}(N)$ satisfying (3) and for which there is an ω satisfying (1) is called a *smooth outer action* of G on N .

In the setting of abstract groups, the corresponding results are due to Eilenberg and MacLane ([7]). In this context it is more convenient to work with the homomorphism $s: G \rightarrow \text{Out}(N), g \mapsto [S(g)]$ defined by S . They show that if there is some extension corresponding to s , then all N -extensions of G corresponding to s form a principal homogeneous space of the cohomology group $H^2(G, Z(N))_s$, where $Z(N)$ is endowed with the G -module structure determined by s . In loc.cit. it is also shown that, for a given s , there is a characteristic class in $H^3(G, Z(N))_s$ vanishing if and only if s corresponds to a group extension. In Section 2 we adapt these results to the Lie group setting in the sense that for a smooth outer action S , we show that a certain characteristic class $\chi(S)$ in the Lie group cohomology $H_{ss}^3(G, Z(N))_S$ determines whether there exists a corresponding Lie group extension, *i.e.*, there exists another choice of ω such that also (2) is satisfied. We further show that in the latter case the set $\text{Ext}(G, N)_S$ of N -extensions of G corresponding to S is a principal homogeneous space of the cohomology group $H_{ss}^2(G, Z(N))_S$ defined in Section 1.

A particular subtlety entering the picture in our Lie theoretic context is that in general the center $Z(N)$ of N need not be a Lie group. For the results of Sections 1 and 2 it will suffice that $Z(N)$ carries the structure of an initial Lie subgroup, which is a rather weak requirement. For the definition of the $Z(N)$ -valued cochain spaces this property is not needed because we use a definition of $H_{ss}^2(G, Z(N))_S$ and $H_{ss}^3(G, Z(N))_S$ that does not require a manifold structure on $Z(N)$ since we consider $Z(N)$ -valued cochains as N -valued functions to specify smoothness requirements.

In Section 3 we introduce crossed modules for Lie groups and discuss their relation to group extensions. A crossed module is a morphism $\alpha: H \rightarrow G$ of Lie groups for which $\text{im}(\alpha)$ and $\ker(\alpha)$ are split Lie subgroups, together with an action of G on H lifting the conjugation action of G and extending the conjugation action of H on itself. Then $\ker(\alpha) \hookrightarrow H \twoheadrightarrow \text{im}(\alpha)$ is a central extension of $N := \text{im}(\alpha)$ by $Z := \ker(\alpha)$ and the question arises whether this Z -extension of N can be enlarged in an equivariant fashion to an extension of G by Z . As in Section 2, it turns out that such an enlargement exists if and only if a certain cohomology class in $H_{ss}^3(G, Z)_\alpha$ vanishes (Theorem 3.8).

If the group N is abelian, then the data given by a smooth outer action S of G on N is the same as a smooth G -module structure on N . In Section 4 we start with a smooth outer action and the assumption that $Z(N)$ is a split Lie subgroup of N , so that we have a central Lie group extension $Z(N) \hookrightarrow N \twoheadrightarrow N_{\text{ad}} := N/Z(N)$. From that we construct a Lie group G^S which is an extension of G by N_{ad} , depending on S , which has the interesting property that all N -extensions of G corresponding to S are extensions of the Lie group G^S by $Z(N)$. Moreover, the map $N \rightarrow N_{\text{ad}} \subseteq G^S$ together with the natural action of G^S on N defines a crossed module of Lie groups whose characteristic class coincides with the characteristic class $\chi(S) \in H_{ss}^3(G, Z(N))_S$.

These observations reduce many questions on general Lie group extensions to the special case of abelian extensions which has been treated in detail in [19]. In particular, the question whether a topologically split extension $\mathfrak{n} \hookrightarrow \widehat{\mathfrak{g}} \twoheadrightarrow \mathfrak{g}$ of Lie algebras is integrable if we assume that we already have groups G and N , together with a compatible smooth outer action of G on N , can be reduced to the question whether the corresponding abelian extension $\mathfrak{z}(\mathfrak{n}) \hookrightarrow \mathfrak{g}^S \twoheadrightarrow \mathfrak{g}$ is integrable, which is characterized in [19] by the discreteness of the image of a period homomorphism $\pi_2(G^S) \rightarrow \mathfrak{z}(\mathfrak{n})$. In [19] we also describe an exact sequence which explains how the cohomology group $H_s^2(G, Z(N))_S$ is given in terms of topological data associated to the group G and the corresponding Lie algebra cohomology space $H_c^2(\mathfrak{g}, \mathfrak{z}(\mathfrak{n}))_S$.

Non-abelian extensions of Lie groups occur quite naturally in the context of smooth principal bundles over compact manifolds M . If $q: P \rightarrow M$ is a smooth K -principal bundle over the compact manifold, where K is a finite-dimensional Lie group, then the group $\text{Aut}(P)$ is a Lie group extension

$$\text{Gau}(P) \hookrightarrow \text{Aut}(P) \twoheadrightarrow \text{Diff}(M)_{[P]},$$

where $\text{Gau}(P)$ is the gauge group of P and $\text{Diff}(M)_{[P]}$ is the open subgroup of $\text{Diff}(M)$ consisting of all diffeomorphisms preserving the bundle class $[P]$ under pull-backs. If K is abelian, then $\text{Gau}(P) \cong C^\infty(M, K)$ and we have an abelian extension.

In an appendix we collect several results on automorphisms of group extensions and on smooth Lie group actions on Lie group extensions. Although only very specific results of this appendix are used in the present paper, we felt that they are very useful in many different contexts, so that it makes good sense to collect them in such an appendix. For a discussion of automorphisms of abstract groups from a cohomological point of view, we refer to Huebschmann's paper [10].

The problem to parameterize all extensions of a given group G by a group N is a core problem in group theory. It seems that the first systematic treatment using a parameterization of the extended group \widehat{G} as a product set $N \times G$ is due to Schreier ([23, 24]). He mainly works with presentations of the groups involved. Baer discovered the first examples of outer actions for which no group extensions exist ([1]), and Turing refined the methods introduced by Baer for abelian extensions to extensions by non-abelian groups ([25]). In particular he discusses extensions of a finite cyclic group by a group N . Cohomological methods to study group extensions first appear in the work of Eilenberg and MacLane ([6], [7]) who coin the term G -kernel for a pair (N, s) consisting of a group N and a homomorphism $s: G \rightarrow \text{Out}(N)$. They show that all extensions corresponding to a given kernel can be parametrized by the cohomology group $H^2(G, Z(N))_s$ and that the obstruction of a kernel to correspond to a group extension is an element in $H^3(G, Z(N))_s$. They also prove for a given G -module Z and each cohomology class $c \in H^3(G, Z)$ the existence of some G -kernel (N, s) with $Z(N) = Z$ whose obstruction class is c . In loc.cit. one even finds the reduction to abelian extensions in the sense that N -extensions of G corresponding to s can also be described as $Z(N)$ -extensions of a group G^s , depending only on s .

Non-abelian extensions of topological groups have been studied by Brown under the assumptions that G and N are polonais ([3]). Then the appropriate group cohomology is defined by measurable cocycles, and there is a natural action of the group $C^1(G, N)$ of measurable cochains on a certain set of pairs (S, ω) , also satisfying certain measurability conditions, such that the orbits $[(S, \omega)]$ of $C^1(G, N)$ are in one-to-one correspondence with the equivalence classes of topological extensions of G by N (cf. our Corollary 2.11).

In [16] C. C. Moore defines group cohomology for second countable locally compact groups in terms of cochains which are Borel measurable. This is a natural context to parametrize abelian group extensions with Borel measurable cross sections ([12]). For finite-dimensional Lie groups measurable cocycles are equivalent to locally smooth cocycles, hence lead to analytic group structures on the extensions ([26, Th. 7.21]; see also [4] and [12]). Varadarajan's book [26] also contains a wealth of information on the connection between projective unitary representations and central extensions of Lie groups.

For a survey of the relations between group cohomology and non-abelian group extensions with twisted actions and crossed product algebras we refer

to [21]. Non-abelian group extensions also arise naturally in mathematical physics ([5]), where the groups involved are mostly infinite-dimensional Lie groups, although they are often dealt with by ad hoc methods, and not in the framework of a general theory, to which we contribute with the present paper.

In a subsequent paper we plan to return to this topic and address the question how to describe elements of $H_s^3(G, Z)$, where Z is a smooth G -module, in terms of Lie algebra cohomology classes in $H_c^3(\mathfrak{g}, \mathfrak{z})$ and topological data associated to G and Z , a topic that has been dealt with in degree 2 in [19]. Another interesting question is how to realize classes in $H_s^3(G, Z)$ as characteristic classes of smooth crossed modules $\alpha: H_1 \rightarrow H_2$ with $\ker(\alpha) \cong Z$ and $\operatorname{coker}(\alpha) \cong G$. Both questions are of particular interest if G is a group of diffeomorphisms of a compact manifold M and \mathfrak{z} is the quotient of the space $\Omega^p(M, \mathbb{R})$ of smooth p -forms on M modulo the subspace of exact forms. Further, it should be possible to describe the criteria for the integrability of the abelian Lie algebra extensions $\mathfrak{z}(\mathfrak{n}) \hookrightarrow \widehat{\mathfrak{g}} \twoheadrightarrow \mathfrak{g}^S$ arising in Section 4 directly in terms of data associated to G , N and the smooth outer action S .

We are most grateful to Ch. Wockel for a thorough proofreading of this article.

1. Lie groups and their cohomology

In this section we introduce notation and terminology used throughout this paper. In particular we describe a natural cohomology for Lie groups also used in [19] in the context of abelian extensions and provide some basic results and calculations that will be crucial for the extension theory.

1.1. Infinite-dimensional Lie groups

Infinite-dimensional manifolds. In this paper $\mathbb{K} \in \{\mathbb{R}, \mathbb{C}\}$ denotes the field of real or complex numbers. Let X and Y be topological \mathbb{K} -vector spaces, $U \subseteq X$ open and $f: U \rightarrow Y$ a map. Then the *derivative of f at x in the direction of h* is defined as

$$df(x)(h) := \lim_{t \rightarrow 0} \frac{1}{t} (f(x + th) - f(x))$$

whenever the limit exists. The function f is called *differentiable at x* if $df(x)(h)$ exists for all $h \in X$. It is called *continuously differentiable or C^1*

if it is continuous, differentiable at all points of U , and

$$df: U \times X \rightarrow Y, \quad (x, h) \mapsto df(x)(h)$$

is a continuous map. It is called a C^n -map if f is C^1 and df is a C^{n-1} -map, and C^∞ (smooth) if it is C^n for all $n \in \mathbb{N}$. This is the notion of differentiability used in [15] and [8], where the latter reference deals with the modifications necessary for incomplete spaces. Since we have a chain rule for C^1 -maps between locally convex spaces, we can define smooth manifolds as in the finite-dimensional case.

Lie groups. A Lie group G is a smooth manifold modeled on a locally convex space for which the group multiplication and the inversion are smooth maps. We refer to [15] for basic results on Lie groups in this context. We write $\mathbf{1} \in G$ for the identity element,

$$\lambda_g(x) := gx, \quad \rho_g(x) := xg \text{ and } c_g(x) := C_G(g)(x) := gxg^{-1}$$

for left and right translations, resp. conjugation on G . The group of all Lie group automorphisms of G is denoted $\text{Aut}(G)$, we write $C_G(G) = \{c_g : g \in G\}$ for the normal subgroup of inner automorphisms, and $\text{Out}(G) := \text{Aut}(G)/C_G(G)$ for the quotient group of outer automorphisms. The conjugation action of G is the homomorphism $C_G: G \rightarrow \text{Aut}(G)$ with range $C_G(G)$. For a normal subgroup $N \trianglelefteq G$ we write $C_N: G \rightarrow \text{Aut}(N)$ with $C_N(g) := C_G(g)|_N$. If G is a connected Lie group, then we write $q_G: \tilde{G} \rightarrow G$ for its universal covering Lie group and identify $\pi_1(G)$ with the kernel of q_G .

Lie subgroups. A subgroup $H \leq G$ of a Lie group G is called an initial Lie subgroup if it carries a Lie group structure for which the inclusion map $\iota: H \hookrightarrow G$ is an immersion in the sense that the tangent map $T\iota: TH \rightarrow TG$ is fiberwise injective, and for each smooth map $f: M \rightarrow G$ from a smooth manifold M to G with $\text{im}(f) \subseteq H$ the corresponding map $\iota^{-1} \circ f: M \rightarrow H$ is smooth. The latter property implies in particular that the initial Lie group structure on H is unique. In fact, assume that H' is H , endowed with another initial Lie group structure and $\iota': H' \hookrightarrow G$ is the corresponding injective immersion with $\iota'(H') = H$. Then the maps $\iota^{-1} \circ \iota': H' \rightarrow H$ and $\iota'^{-1} \circ \iota: H \rightarrow H'$ are smooth morphism of Lie groups. Since these maps are each others inverse, the Lie groups H and H' are isomorphic.

Lie algebras of Lie groups. Each $x \in T_{\mathbf{1}}(G)$ corresponds to a unique left invariant vector field x_l with $g \cdot x := x_l(g) := d\lambda_g(\mathbf{1}) \cdot x, g \in G$. The space of left invariant vector fields is closed under the Lie bracket of vector

fields, hence inherits a Lie algebra structure. In this sense we obtain on the locally convex space $T_1(G)$ a continuous Lie bracket which is uniquely determined by the relation $[x, y]_l = [x_l, y_l]$.

We call a Lie algebra \mathfrak{g} which is a topological vector space such that the Lie bracket is continuous a *topological Lie algebra* \mathfrak{g} . In this sense the Lie algebra $\mathbb{L}(G) := (T_1(G), [\cdot, \cdot])$ of a Lie group G is a locally convex topological Lie algebra.

Topological splitting. A linear subspace W of a topological vector space V is called (*topologically*) *split* if it is closed and there is a continuous linear map $\sigma: V/W \rightarrow V$ for which the map

$$W \times (V/W) \rightarrow V, \quad (w, x) \mapsto w + \sigma(x)$$

is an isomorphism of topological vector spaces. Note that the closedness of W guarantees that the quotient topology turns V/W into a Hausdorff space which is a topological vector space with respect to the induced vector space structure. A continuous linear map $f: V \rightarrow W$ between topological vector spaces is said to be (*topologically*) *split* if the subspaces $\ker(f) \subseteq V$ and $\text{im}(f) \subseteq W$ are topologically split.

1.2. Locally smooth cohomology of Lie groups

DEFINITION 1.1. — *If G and N are Lie groups, then we put $C_s^0(G, N) := N$ and for $p \in \mathbb{N}$ we call a map $f: G^p \rightarrow N$ locally smooth if there exists an open identity neighborhood $U \subseteq G^p$ such that $f|_U$ is smooth. We say that f is normalized if*

$$(1.1) \quad (\exists j) \ g_j = \mathbf{1} \Rightarrow f(g_1, \dots, g_p) = \mathbf{1}.$$

We write $C_s^p(G, N)$ for the space of all normalized locally smooth maps $G^p \rightarrow N$, the so-called (*locally smooth*) p -cochains.

We shall see below that for $p = 2$ and non-connected Lie groups, we sometimes have to require additional smoothness: We write $C_{ss}^2(G, N)$ for the set of all elements $\omega \in C_s^2(G, N)$ with the additional property that for each $g \in G$ the map

$$\omega_g: G \rightarrow N, \quad x \mapsto \omega(g, x)\omega(gxg^{-1}, g)^{-1}$$

is smooth in an identity neighborhood of G . Note that $\omega_g(\mathbf{1}) = \mathbf{1}$.

For $f \in C_s^1(G, N)$ we define

$$\delta_f: G \times G \rightarrow N, \quad \delta_f(g, g') := f(g)f(g')f(gg')^{-1}$$

and observe that

$$\begin{aligned}
 (\delta_f)_g(x) &= \delta_f(g, x)\delta_f(gxg^{-1}, g)^{-1} \\
 &= f(g)f(x)f(gx)^{-1}f(gx)f(g)^{-1}f(gxg^{-1})^{-1} \\
 &= f(g)f(x)f(g^{-1})f(gxg^{-1})^{-1}
 \end{aligned}$$

is smooth in an identity neighborhood of G , so that $\delta_f \in C_{ss}^2(G, N)$. Note that δ_f measures the failure of f for being a homomorphism.

If G and N are abstract groups, we write $C^p(G, N)$ for the set of all functions $G^p \rightarrow N$, normalized as in (1.1).

For the following we also observe that for $h, f \in C_s^1(G, N)$ we have

$$\begin{aligned}
 (1.2) \quad \delta_{hf}(g, g') &= h(g)f(g)h(g')f(g')f(gg')^{-1}h(gg')^{-1} \\
 &= h(g)c_{f(g)}(h(g'))\delta_f(g, g')h(gg')^{-1}.
 \end{aligned}$$

For cochains with values in abelian groups, resp. smooth modules, the following definition recalls the natural Lie group cohomology setting from [19].

DEFINITION 1.2. — Let G be a Lie group and A a smooth G -module, i.e., a pair (A, S) of an abelian Lie group A and a smooth G -action $G \times A \rightarrow A$ given by the homomorphism $S: G \rightarrow \text{Aut}(A)$. We call the elements of $C_s^p(G, A)$ (locally smooth normalized) p -cochains and write

$$d_S: C_s^p(G, A) \rightarrow C_s^{p+1}(G, A)$$

for the group differential given by

$$\begin{aligned}
 (d_S f)(g_0, \dots, g_p) &:= S(g_0)(f(g_1, \dots, g_p)) \\
 &+ \sum_{j=1}^p (-1)^j f(g_0, \dots, g_{j-1}g_j, \dots, g_p) + (-1)^{p+1} f(g_0, \dots, g_{p-1}).
 \end{aligned}$$

It is easy to verify that $d_S(C_s^p(G, A)) \subseteq C_s^{p+1}(G, A)$. We thus obtain a sub-complex of the standard group cohomology complex $(C^\bullet(G, A), d_S)$. In view of $d_S^2 = 0$, the space $Z_s^p(G, A)_S := \ker d_S|_{C_s^p(G, A)}$ of p -cocycles contains the space $B_s^p(G, A)_S := d_S(C_s^{p-1}(G, A))$ of p -coboundaries. The quotient

$$H_s^p(G, A)_S := Z_s^p(G, A)_S / B_s^p(G, A)_S$$

is the p^{th} locally smooth cohomology group of G with values in the G -module A . We write $[f] \in H_s^p(G, A)_S$ for the cohomology class of a cocycle $f \in Z_s^p(G, A)_S$.

In the following we also need generalizations of Lie group cohomology to settings where the target groups are non-abelian. For 1-cocycles the generalization is quite straight forward:

DEFINITION 1.3. — *Let G and N be Lie groups and $S: G \rightarrow \text{Aut}(N)$ a homomorphism defining a smooth action of G on N . A function $f \in C_s^1(G, N)$ is called a 1-cocycle (or a crossed homomorphism) if*

$$f(gg') = f(g)S(g)(f(g')) \text{ for all } g, g' \in G.$$

We write $Z_s^1(G, N)_S$ for the set of all 1-cocycles $G \rightarrow N$ with respect to the action of G on N given by S . The group $N = C_s^0(G, N)$ acts on $Z_s^1(G, N)_S$ by

$$(n \cdot f)(g) := nf(g)S(g)(n)^{-1},$$

and the set of orbits of N in $Z_s^1(G, N)_S$ is the cohomology set

$$H_s^1(G, N)_S := Z_s^1(G, N)_S / N.$$

For the extension theory of Lie groups we also need non-abelian 2-cocycles, but for the appropriate concept we have to weaken the setting of a smooth action of G on N as follows.

DEFINITION 1.4. — *Let G and N be Lie groups. We define $C_s^1(G, \text{Aut}(N))$ as the set of all maps $S: G \rightarrow \text{Aut}(N)$ with $S(\mathbf{1}) = \text{id}_N$ and for which there exists an open identity neighborhood $U \subseteq G$ such that the map*

$$U \times N \rightarrow N, \quad (g, n) \mapsto S(g)(n)$$

is smooth.

We call a map $S \in C_s^1(G, \text{Aut}(N))$ a smooth outer action of G on N if there exists $\omega \in C_s^2(G, N)$ with $\delta_S = C_N \circ \omega$. It is called strongly smooth if ω can be chosen in the smaller set $C_{ss}^2(G, N)$.

On the set of smooth outer actions we define an equivalence relation by

$$S \sim S' \iff (\exists h \in C_s^1(G, N)) S' = (C_N \circ h) \cdot S.$$

In Lemma 1.10 below we shall see that for each (strongly) smooth outer action S the outer action $S_h := (C_N \circ h) \cdot S$ also is a (strongly) smooth outer action because $\delta_S = C_N \circ \omega$ implies

$$\delta_{S_h}(g, g') = C_N \left(h(g)S(g)(h(g'))\omega(g, g')h(gg')^{-1} \right) \text{ for } g, g' \in G.$$

We write $[S]$ for the equivalence class of S and call $[S]$ a smooth G -kernel.

Remark 1.5.

(a) A smooth outer action $S: G \rightarrow \text{Aut}(N)$ need not be a group homomorphism, but $\delta_S = C_N \circ \omega$ implies that S induces a group homomorphism $s := Q_N \circ S: G \rightarrow \text{Out}(N)$, where $Q_N: \text{Aut}(N) \rightarrow \text{Out}(N)$ denotes the quotient homomorphism. Clearly s depends only on the class $[S]$ of S because $Q_N \circ C_N = \mathbf{1}$.

(b) If G and N are discrete groups, then for each homomorphism $s: G \rightarrow \text{Out}(N)$ there exists a map $S: G \rightarrow \text{Aut}(N)$ with $S(\mathbf{1}) = \text{id}_N$ and $Q_N \circ S = s$ and a map $\omega: G \times G \rightarrow N$ with $\delta_S = C_N \circ \omega$. In this case all outer actions are smooth, the smoothness conditions on S and ω are vacuous, and $S \sim S'$ is equivalent to $Q_N \circ S = Q_N \circ S'$, so that a smooth G -kernel is nothing but a homomorphism $s: G \rightarrow \text{Out}(N)$.

(c) If $S \in C_s^1(G, \text{Aut}(N))$ is a homomorphism of groups, then we may choose $\omega = \mathbf{1}$ and S defines a smooth action of G on N . In fact, the action is smooth on a set of the form $U \times N$, so that the assertion follows from the fact that all automorphisms $S(g)$ are smooth.

(d) Suppose that $S_1, S_2 \in C_s^1(G, \text{Aut}(N))$ and that the map

$$U \times N \rightarrow N^2, \quad (g, n) \mapsto (S_1(g)(n), S_2(g)(n))$$

is smooth. Then the map $U \times N \rightarrow N, (g, n) \mapsto S_1(g)S_2(g)(n)$ is smooth because it is a composition of smooth maps. Therefore $C_s^1(G, \text{Aut}(N))$ is closed under pointwise products.

(e) If $S: G \rightarrow \text{Aut}(N)$ is a smooth outer action, then we obtain in particular

$$C_N(\omega(g, g^{-1})) = \delta_S(g, g^{-1}) = S(g)S(g^{-1}),$$

which leads to

$$S(g)^{-1} = S(g^{-1})C_N(\omega(g, g^{-1}))^{-1}.$$

Hence there also exists an identity neighborhood $U \subseteq G$ for which the map

$$U \times N \rightarrow N, \quad (g, n) \mapsto S(g)^{-1}(n) = S(g^{-1})\omega(g, g^{-1})^{-1}n\omega(g, g^{-1})$$

is smooth.

LEMMA 1.6. — *Each smooth outer action $S: G \rightarrow \text{Aut}(N)$ defines on $Z(N)$ the structure of a G -module by $g \cdot z := S_Z(g)(z) := S(g)(z)$. If, in addition, $Z(N)$ is an initial Lie subgroup of N , then $Z(N)$ is a smooth G -module.*

Proof. — First we observe that inner automorphisms act trivially on the center $Z(N)$ of N , which implies that $g \cdot z := S(g)(z)$ defines indeed a G -action on $Z(N)$.

Let $U \subseteq G$ be an open identity neighborhood such that the map $U \times N \rightarrow N, (g, n) \mapsto S(g) \cdot n$ is smooth. If $Z(N)$ is an initial Lie subgroup of N , restriction and co-restriction define a smooth map

$$(1.3) \quad U \times Z(N) \rightarrow Z(N), \quad (g, z) \mapsto g \cdot z = S_Z(g)(z).$$

As each element of g acts as a Lie group automorphism on $Z(N)$, the smoothness of the action $G \times Z(N) \rightarrow Z(N)$ follows from the smoothness of the map in (1.3). \square

The following lemma sheds some more light on the concept of an initial subgroup of a Lie group. In the present paper we shall mostly use it for the center of a Lie group N which is not always a Lie group, but mostly carries an initial Lie subgroup structure.

LEMMA 1.7. — *Let G be a Lie group with Lie algebra \mathfrak{g} and $H \leq G$ a subgroup. Then the following assertions hold:*

- (a) *If H is an initial normal subgroup, then the conjugation action of G on H is smooth.*
- (b) *If H is totally disconnected, then the discrete group H_d (the group H endowed with the discrete topology) defines on H via the inclusion map $\iota: H_d \rightarrow H \subseteq G$ the structure of an initial Lie subgroup. The same conclusion holds if all smooth arcs in H are constant, i.e., if the smooth arc-components in H are trivial.*
- (c) *If $\dim G < \infty$, then any subgroup $H \leq G$ carries an initial Lie group structure.*
- (d) *Let G be a Lie group with a smooth exponential function and*

$$\mathfrak{z}(\mathfrak{g})^G := \{x \in \mathfrak{z}(\mathfrak{g}) : (\forall g \in G) \operatorname{Ad}(g) \cdot x = x\}.$$

We assume that the group $\Gamma_Z := \exp^{-1}(\mathbf{1}) \cap \mathfrak{z}(\mathfrak{g})$ is discrete and that $\mathfrak{z}(\mathfrak{g})^G$ is Mackey complete. Then $Z(G)$ carries an initial Lie group structure $Z(G)^L$ with $\mathbb{L}(Z(G)^L) = \mathfrak{z}(\mathfrak{g})^G$.

Proof.

(a) The conjugation map $G \times H \rightarrow G, (g, h) \mapsto ghg^{-1}$ is smooth with values in H , hence smooth as a map $G \times H \rightarrow H$.

(b) Since every discrete space carries a natural 0-dimensional smooth manifold structure, the group H_d is a Lie group with $\mathbb{L}(H_d) = \{0\}$. If M is a smooth manifold and $f: M \rightarrow G$ a smooth map with $f(M) \subseteq H$, then f maps each connected component of M into a connected component of H , so that f is locally constant because H is totally disconnected. Hence the map $\iota^{-1} \circ f: M \rightarrow H_d$ is locally constant and therefore smooth.

(c) Let $H_a \subseteq H$ be the arc-component of H , viewed as a topological subgroup of G . According to Yamabe's Theorem ([9]), the arc-component H_a of G is analytic, i.e., there exists a Lie subalgebra $\mathfrak{h} \subseteq \mathfrak{g}$ such that $H_a = \langle \exp_G \mathfrak{h} \rangle$. Let H_a^L denote the group H_a endowed with its intrinsic Lie group topology for which $\exp: \mathfrak{h} \rightarrow H_a^L$ is a local diffeomorphism in $\mathbf{0}$. Then $H \subseteq \{g \in G: \text{Ad}(g)\mathfrak{h} = \mathfrak{h}\}$ implies that H acts by conjugation smoothly on H_a^L , so that H carries a Lie group structure for which H_a^L is an open subgroup (apply Theorem B.1 with $K = H_a^L$). Let H^L denote this Lie group. Now the inclusion map $\iota: H^L \rightarrow G$ is an immersion whose differential in $\mathbf{1}$ is the inclusion $\mathfrak{h} \hookrightarrow \mathfrak{g}$.

We claim that $\iota: H^L \rightarrow G$ is an initial Lie subgroup. In fact, let $f: M \rightarrow G$ be a smooth map from the smooth manifold M to G with $f(M) \subseteq H$. We have to show that f is smooth, i.e., that f is smooth in a neighborhood of each point $m \in M$. Replacing f by $f \cdot f(m)^{-1}$ and observing that the group operations in H and G are smooth, we may w.l.o.g. assume that $f(m) = \mathbf{1}$.

Let $U \subseteq \mathfrak{h}$ be an open $\mathbf{0}$ -neighborhood, $\mathfrak{m} \subseteq \mathfrak{g}$ be a vector space complement to \mathfrak{h} and $V \subseteq \mathfrak{m}$ an open $\mathbf{0}$ -neighborhood for which the map

$$\Phi: U \times V \rightarrow G, \quad (x, y) \mapsto \exp x \exp y$$

is a diffeomorphism onto an open subset of G . Then

$$H_a \cap (\exp U \exp V) = \bigcup_{\exp y \in H_a} \exp U \exp y$$

and each set $\exp U \exp y$ contained in H_a also is an open subset of H^L . Since the topology of H_a^L is second countable, the set $\exp^{-1}(H_a) \cap \mathfrak{m}$ is countable. Every smooth arc $\gamma: I \rightarrow \exp U \exp V$ is of the form $\gamma(t) = \exp \alpha(t) \exp \beta(t)$ with smooth arcs $\alpha: I \rightarrow U$ and $\beta: I \rightarrow V$, and for every smooth arc contained in H_a the arc β is constant.

We conclude that if $W \subseteq M$ is an open connected neighborhood of m with $f(W) \subseteq \exp U \exp V$, then $f(W) \subseteq \exp U$. Then the map

$$\exp|_U^{-1} \circ f|_W: W \rightarrow \mathfrak{h}$$

is smooth, so that the corresponding map

$$\iota^{-1} \circ f|_W = \exp_{H^L} \circ \exp|_U^{-1} \circ f|_W: W \rightarrow H^L$$

is also smooth. This proves that the map $\iota^{-1} \circ f: M \rightarrow H^L$ is smooth, and hence that $\iota: H^L \rightarrow H$ is an initial Lie subgroup of G .

(d) First we observe that the exponential function $\exp_Z: \mathfrak{z}(\mathfrak{g})^G \rightarrow Z(G)$ is a group homomorphism, and our condition implies that $\Gamma := \ker \exp_Z$ is a discrete subgroup of $\mathfrak{z}(\mathfrak{g})^G$, so that $Z := \mathfrak{z}(\mathfrak{g})^G / \Gamma$ carries a natural Lie

group structure, and the smooth function $\exp_Z: \mathfrak{z}(\mathfrak{g})^G \rightarrow G$ factors through an injective immersion of Lie groups $\iota: Z \rightarrow G$ whose image is the subgroup $\exp(\mathfrak{z}(\mathfrak{g})^G)$ of $Z(G)$. We now endow $Z(G)$ with the Lie group structure for which $\exp \mathfrak{z}(\mathfrak{g})^G$ is an open subgroup isomorphic to Z (Theorem B.1) and write $Z(G)^L$ for this Lie group. Then the inclusion map $\iota: Z(G)^L \rightarrow G$ is an injective immersion of Lie groups.

To see that this defines an initial Lie group structure on $Z(G)$, let $f: M \rightarrow G$ be a smooth map with $f(M) \subseteq Z(G)$. We may w.l.o.g. assume that M is connected and fix some $m \in M$. Then the left logarithmic derivative

$$\delta(f) \in \Omega^1(M, \mathfrak{z}(\mathfrak{g})^G) \text{ with } \delta(f)(m) = f(m)^{-1}df(m)$$

is a smooth $\mathfrak{z}(\mathfrak{g})^G$ -valued 1-form on M . In fact, for each smooth curve $\gamma: I \rightarrow Z(G)$ we have $\gamma(t)^{-1}\gamma'(t) \in \mathfrak{z}(\mathfrak{g})^G$ for each $t \in I$ because for each $g \in G$ the relation $c_g \circ \gamma = \gamma$ leads to

$$\gamma(t)^{-1}.\gamma'(t) = (c_g \circ \gamma)(t)^{-1}.(c_g \circ \gamma)'(t) = \text{Ad}(g).(\gamma(t)^{-1}.\gamma'(t)).$$

The 1-form $\delta(f)$ satisfies the Maurer–Cartan equation

$$d(\delta(f)) + \frac{1}{2}[\delta(f), \delta(f)] = 0$$

([11]), so that the fact that $\mathfrak{z}(\mathfrak{g})$ is abelian implies that $\delta(f)$ is closed. Let $q_M: \widetilde{M} \rightarrow M$ be a universal covering map. As $\mathfrak{z}(\mathfrak{g})$ is Mackey complete, the Poincaré Lemma holds for $\mathfrak{z}(\mathfrak{g})$ -valued 1-forms, so that there exists a smooth function $f_1: \widetilde{M} \rightarrow \mathfrak{z}(\mathfrak{g})$ with $df_1 = q_M^*\delta(f)$ (cf. [18, Prop. III.4]). If $\widetilde{m} \in \widetilde{M}$ is a point with $q_M(\widetilde{m}) = m$, we may assume, in addition, that $f_1(\widetilde{m}) = 0$. Then the smooth function

$$f_2 := \exp_Z \circ f_1: \widetilde{M} \rightarrow Z(G)^L \text{ satisfies } \delta(f_2) = df_1 = q_M^*\delta(f) = \delta(f \circ q_M),$$

which in turn leads to

$$f \circ q_M = f(m) \cdot f_2$$

since the solutions of the Maurer–Cartan equation are uniquely determined by their values in \widetilde{m} . This implies that the function $f \circ q_M: \widetilde{M} \rightarrow Z(G)^L$ is smooth, and hence that the function $\iota^{-1} \circ f: M \rightarrow Z(G)^L$ is smooth. \square

The following example shows that in general $Z(G)$, considered as a topological group, is not a Lie group because it may be a totally disconnected non-discrete subgroup. Since all known Lie groups modeled on complete locally convex spaces satisfy the requirements from Lemma 1.7(d), for all these Lie groups G the center $Z(G)$ carries an initial Lie group structure. Therefore the assumption that $Z(G)$ is an initial Lie subgroup is not restrictive.

Example 1.8. — We consider the Lie group

$$N := \mathbb{C}^{\mathbb{N}} \rtimes \mathbb{R}^{\mathbb{N}} \text{ with } ((z_n), (t_n)) * ((z'_n), (t'_n)) := ((z_n + e^{t_n} z'_n), (t_n + t'_n)),$$

where the locally convex structure on $\mathbb{C}^{\mathbb{N}}$ and $\mathbb{R}^{\mathbb{N}}$ is the product topology. We observe that N is an infinite topological product of groups isomorphic to $\mathbb{C} \rtimes \mathbb{R}$, endowed with the multiplication

$$(z, t) * (z', t') = (z + e^t z', t + t').$$

Therefore the center of N is

$$Z(N) = \{0\} \times (2\pi i \mathbb{Z})^{\mathbb{N}} \cong \mathbb{Z}^{\mathbb{N}},$$

and this group is totally disconnected, but not discrete, hence not a Lie group. Therefore the center of a Lie group need not be a Lie group.

Since $Z(N)$ is totally disconnected, Lemma 1.7 implies that the inclusion map $Z(N)_d \rightarrow N$ of the discrete group $Z(N)_d$ defines on $Z(N)$ the structure of an initial Lie subgroup.

If S is a smooth outer action of G on N and $Z(N)$ is an initial Lie subgroup of N , then it carries the structure of a smooth G -module (Lemma 1.6) and the definitions below are consistent with Definition 1.2, but in general we do not want to assume that $Z(N)$ is an initial Lie subgroup and it is not necessary for the results in Section 2 below.

DEFINITION 1.9. — *If S is a smooth outer action of G on the Lie group N , then we define*

$$\begin{aligned} C_s^p(G, Z(N)) &:= \{f \in C_s^p(G, N) : \text{im}(f) \subseteq Z(N)\}, \\ C_{ss}^2(G, Z(N)) &:= C_{ss}^2(G, N) \cap C_s^2(G, Z(N)) \end{aligned}$$

and accordingly

$$Z_s^p(G, Z(N))_S := \{f \in C_s^p(G, Z(N)) : d_S f = \mathbf{1}\}$$

and $Z_{ss}^2(G, Z(N))_S := Z_s^2(G, Z(N))_S \cap C_{ss}^2(G, Z(N))$ (cf. Lemma 1.6). Note that on $f \in C_s^p(G, Z(N))$ the group differential d_S for the G -module structure on $Z(N)$ can be written, in additive notation, as

$$\begin{aligned} (d_S f)(g_0, \dots, g_p) &= S(g_0)(f(g_1, \dots, g_n)) \\ &+ \sum_{j=1}^n (-1)^j f(g_0, \dots, g_{j-1} g_j, \dots, g_n) + (-1)^{n+1} f(g_0, \dots, g_{n-1}). \end{aligned}$$

This implies that $d_S(C_s^p(G, Z(N))) \subseteq C^{p+1}(G, Z(N))$, so that we obtain a differential complex $(C_s^\bullet(G, Z(N)), d_S)$ whose cohomology groups are denoted by

$$H_s^p(G, Z(N))_S := Z_s^p(G, Z(N))_S / d_S(C_s^p(G, Z(N))).$$

For $\alpha \in C_s^1(G, Z(N))$ and $g \in G$ we then have

$$\begin{aligned} (d_S\alpha)_g(x) &= (d_S\alpha)(g, x) - (d_S\alpha)(g x g^{-1}, g) \\ &= g.\alpha(x) - \alpha(gx) + \alpha(g) - (g x g^{-1}).\alpha(g) + \alpha(gx) - \alpha(g x g^{-1}) \\ &= g.\alpha(x) + \alpha(g) - (g x g^{-1}).\alpha(g) - \alpha(g x g^{-1}), \end{aligned}$$

which is smooth in an identity neighborhood of G . Hence

$$B_s^2(G, Z(N))_S := d_S(C_s^1(G, Z(N))) \subseteq Z_{ss}^2(G, Z(N))_S,$$

and we define

$$H_{ss}^2(G, Z(N))_S := Z_{ss}^2(G, Z(N))_S / B_s^2(G, Z(N))_S,$$

and

$$H_{ss}^3(G, Z(N))_S := Z_s^3(G, Z(N))_S / d_S(C_{ss}^2(G, Z(N))).$$

We observe that the action of G on $Z(N)$ defined by a smooth outer action S only depends on the equivalence class $[S]$. In this sense we also write

$$\begin{aligned} Z_s^p(G, Z(N))_{[S]} &:= Z^p(G, Z(N))_S, \\ H_s^p(G, Z(N))_{[S]} &:= H_s^p(G, Z(N))_S, \text{ etc.} \end{aligned}$$

In Definition 1.3 above we have defined $H_s^1(G, N)$ as the set of orbits of $N = C_s^0(G, N)$ on the set $Z_s^1(G, N)$. The following lemma describes an action of $C_s^1(G, N)$ that in some sense is a generalization of the N -action on $Z_s^1(G, N)$ for degree 1 to degree 2.

LEMMA 1.10. — *Let G and N be Lie groups and consider the group $C_s^1(G, N)$ with respect to pointwise multiplication. This group acts on the set $C_s^1(G, \text{Aut}(N))$ by $h.S := (C_N \circ h) \cdot S$ and on the product set*

$$(1.4) \quad C_s^1(G, \text{Aut}(N)) \times C_s^2(G, N) \text{ by } h.(S, \omega) := (h.S, h *_S \omega)$$

with

$$(h *_S \omega)(g, g') := h(g)S(g)(h(g'))\omega(g, g')h(gg')^{-1}.$$

The stabilizer of (S, ω) is given by

$$C_s^1(G, N)_{(S, \omega)} = Z_s^1(G, Z(N))_S$$

which depends only on $[S]$, but not on ω . Moreover, the following assertions hold:

- (1) The subset $\{(S, \omega) : \delta_S = C_N \circ \omega\}$ is invariant.
- (2) The subset $\{(S, \omega) \in C_s^1(G, \text{Aut}(N)) \times C_{ss}^2(G, N) : \delta_S = C_N \circ \omega\}$ is also invariant.
- (3) If $\delta_S = C_N \circ \omega$, then $\text{im}(d_S\omega) \subseteq Z(N)$ holds for

$$(d_S\omega)(g, g', g'') := S(g)(\omega(g', g''))\omega(g, g'g'')\omega(gg', g'')^{-1}\omega(g, g')^{-1}.$$

- (4) If $\delta_S = C_N \circ \omega$ and $h.(S, \omega) = (S', \omega')$, then $d_{S'}\omega' = d_S\omega$.
- (5) If $\delta_S = C_N \circ \omega$, then $d_S\omega \in Z_s^3(G, Z(N))_S$.
- (6) For a smooth outer action S with $\delta_S = C_N \circ \omega$, the cohomology class $[d_S\omega] \in H_s^3(G, Z(N))_S$ depends only on the equivalence class $[S]$. Similarly, for a strongly smooth outer action and $\omega \in C_{ss}^2(G, N)$ with $\delta_S = C_N \circ \omega$ the class $[d_S\omega] \in H_{ss}^3(G, Z(N))_S$ depends only on the equivalence class $[S]$.

Proof. — That (1.4) defines a group action follows from the trivial relation $(hh').S = h.(h'.S)$, and

$$\begin{aligned} &(h *_{h'.S} (h' *_S \omega))(g, g') \\ &= h(g)(h'.S)(g)(h(g'))(h' *_S \omega)(g, g')h(gg')^{-1} \\ &= h(g)h'(g)S(g)(h(g'))h'(g)^{-1}h'(g)S(g)(h'(g'))\omega(g, g')h'(gg')^{-1}h(gg')^{-1} \\ &= (hh')(g)S(g)(hh'(g'))\omega(g, g')(hh')(gg')^{-1} \\ &= ((hh') *_S \omega)(g, g'). \end{aligned}$$

To calculate the stabilizer of the pair (S, ω) , we observe that the condition $h.S = S$ is equivalent to $h \in C_s^1(G, Z(N))$. Then $h *_S \omega = \omega \cdot d_S h$, and this equals ω if and only if h is a cocycle. Therefore $C_s^1(G, N)_{(S, \omega)} = Z_s^1(G, Z(N))_S$.

(1) If the pair (S, ω) satisfies $\delta_S = C_N \circ \omega$, then we obtain for $h \in C_s^1(G, N)$ with Formula (1.2) (after Definition 1.1), the relation

$$\begin{aligned} \delta_{h.S}(g, g') &= \delta_{(C_N \circ h).S}(g, g') \\ &= C_N(h(g))c_{S(g)}(C_N(h(g')))\delta_S(g, g')C_N(h(gg')^{-1}) \\ &= C_N(h(g))C_N(S(g)(h(g'))C_N(\omega(g, g'))C_N(h(gg')^{-1}) \\ &= C_N(h(g)S(g)(h(g'))\omega(g, g')h(gg')^{-1}) \\ &= (C_N \circ (h *_S \omega))(g, g'). \end{aligned}$$

(2) Suppose that $\omega \in C_{ss}^2(G, N)$ satisfies $\delta_S = C_N \circ \omega$. We have to show that $h *_S \omega \in C_{ss}^2(G, N)$. In fact, for $g, x \in G$ we have

$$\begin{aligned} (h *_S \omega)_g(x) &= (h *_S \omega)(g, x)(h *_S \omega)(g x g^{-1}, g)^{-1} \\ &= h(g)S(g)(h(x))\omega(g, x)h(gx)^{-1}(h(gxg^{-1})S(gxg^{-1})(h(g)) \\ &\quad \cdot \omega(gxg^{-1}, g)h(gx)^{-1})^{-1} \\ &= h(g)S(g)(h(x))\omega(g, x)\omega(gxg^{-1}, g)^{-1}S(gxg^{-1})(h(g)^{-1}) \\ &\quad \cdot h(gxg^{-1})^{-1} \\ &= h(g)S(g)(h(x))\omega_g(x)S(gxg^{-1})(h(g)^{-1})h(gxg^{-1})^{-1}. \end{aligned}$$

This expression is a product of factors which are smooth functions of x in an identity neighborhood of G . Therefore $h *_S \omega \in C_{ss}^2(G, N)$.

(3) This follows from

$$\begin{aligned} C_N((d_S \omega)(g, g', g'')) &= C_N(S(g)(\omega(g', g''))\omega(g, g'g'')\omega(gg', g'')^{-1}\omega(g, g')^{-1}) \\ &= S(g)(C_N \circ \omega)(g', g'')S(g)^{-1}(C_N \circ \omega)(g, g'g'')(C_N \circ \omega)(gg', g'')^{-1} \\ &\quad \cdot (C_N \circ \omega)(g, g')^{-1} \\ &= S(g)\delta_S(g', g'')S(g)^{-1}\delta_S(g, g'g'')\delta_S(gg', g'')^{-1}\delta_S(g, g')^{-1} \\ &= S(g)(S(g')S(g'')S(g'g'')^{-1})S(g)^{-1}S(g)S(g'g'')S(gg'g'')^{-1} \\ &\quad \cdot S(gg'g'')S(g'')^{-1}S(gg')^{-1}S(gg')S(g')^{-1}S(g)^{-1} = \mathbf{1}. \end{aligned}$$

(4) In the following calculation we frequently use that $d_S \omega$ and $d_{S'} \omega'$ have values in $Z(N)$ to see that they commute with all elements of N . We use under-braces to indicate which parts of the formula cancel each other.

We calculate

$$\begin{aligned} (d_{S'} \omega')(g, g', g'') &= S'(g)(\omega'(g', g''))\omega'(g, g'g'')\omega'(gg', g'')^{-1}\omega'(g, g')^{-1} \\ &= (C_N(h(g)) \circ S(g)(\omega'(g', g'')))\omega'(g, g'g'')\omega'(gg', g'')^{-1}\omega'(g, g')^{-1} \\ &= \underbrace{h(g)}_1 S(g)[h(g')S(g')(h(g''))\omega(g', g'') \underbrace{h(g'g'')^{-1}}_5] \underbrace{h(g)^{-1}}_2 \underbrace{h(g)}_2 \\ &\quad \underbrace{S(g)(h(g'g''))}_5 \omega(g, g'g'') \underbrace{h(gg'g'')^{-1}}_3 \underbrace{h(gg'g'')}_3 \omega(gg', g'')^{-1} \\ &\quad S(gg')(h(g''))^{-1} \underbrace{h(gg')^{-1}}_4 \underbrace{h(gg')}_4 \omega(g, g')^{-1} S(g)(h(g'))^{-1} \underbrace{h(g)^{-1}}_1 \end{aligned}$$

$$\begin{aligned}
 &= \underbrace{S(g)(h(g'))}_1 S(g) \circ S(g')(h(g'')) [S(g)(\omega(g', g''))\omega(g, g'g'')\omega(gg', g'')^{-1}] \\
 &\quad S(gg')(h(g''))^{-1}\omega(g, g')^{-1} \underbrace{S(g)(h(g'))^{-1}}_1 \\
 &= [\omega(g, g')S(gg')(h(g''))\omega(g, g')^{-1}] [(d_S\omega)(g, g', g'')\omega(g, g')] \\
 &\quad S(gg')(h(g''))^{-1}\omega(g, g')^{-1} \\
 &= (d_S\omega)(g, g', g'')
 \end{aligned}$$

because the values of $d_S\omega$ are central.

(5) It is easily seen that $d_S\omega$ vanishes if one of its three arguments is $\mathbf{1}$, so that, in view of (3), we have $d_S\omega \in C_s^3(G, Z(N))$. It remains to show that it is a cocycle.

In the following calculation we write the group structure on $Z(N)$ multiplicatively. We shall use that the values of $d_S\omega$ are central, so that they commute with all values of ω . The corresponding equalities are marked by (*). We simply write $g.z := S(g)(z)$ for the action of G on $Z(N)$ induced by the smooth outer action S . We have to show that for $g, g', g'', g''' \in G$ the following expression vanishes:

$$\begin{aligned}
 &d_S(d_S\omega)(g, g', g'', g''') \\
 &= g.(d_S\omega)(g', g'', g''')(d_S\omega)(gg', g'', g''')^{-1}(d_S\omega)(g, g'g'', g''') \\
 &\quad (d_S\omega)(g, g', g''g''')^{-1}(d_S\omega)(g, g', g'') \\
 &= g.(d_S\omega)(g', g'', g''')(d_S\omega)(gg', g'', g''')^{-1}(d_S\omega)(g, g'g'', g''') \\
 &\quad (d_S\omega)(g, g', g'')(d_S\omega)(g, g', g''g''')^{-1} \\
 &= S(g)[S(g')(\omega(g'', g'''))\omega(g', g'g''g''')\omega(g'g'', g''')^{-1}\omega(g', g'')^{-1}] \\
 &\quad \omega(gg', g'')\omega(gg'g'', g''')\omega(gg', g''g''')^{-1}S(gg')(\omega(g'', g''')^{-1}) \\
 &\quad S(g)(\omega(g'g'', g'''))\omega(g, g'g''g''')\omega(gg'g'', g''')^{-1}\omega(g, g'g'')^{-1} \\
 &\quad S(g)(\omega(g', g''))\omega(g, g'g'')\omega(gg', g'')^{-1}\omega(g, g')^{-1} \\
 &\quad \omega(g, g')\omega(gg', g''g''')\omega(g, g'g''g''')^{-1}S(g)(\omega(g', g''g''')^{-1}) \\
 &= [\omega(g, g')S(gg')(\omega(g'', g'''))\omega(g, g')^{-1}]S(g)[\omega(g', g''g''')\omega(g'g'', g''')^{-1} \\
 &\quad \omega(g', g'')^{-1}][\omega(gg', g'')\omega(gg'g'', g''')\omega(gg', g''g''')^{-1}S(gg')(\omega(g'', g''')^{-1})] \\
 &\quad [S(g)(\omega(g'g'', g'''))\omega(g, g'g''g''')\omega(gg'g'', g''')^{-1}\omega(g, g'g'')^{-1}] \\
 &\quad S(g)(\omega(g', g''))\omega(g, g'g'')\omega(gg', g'')^{-1}\omega(gg', g''g''')\omega(g, g'g''g''')^{-1} \\
 &\quad S(g)(\omega(g', g''g''')^{-1})
 \end{aligned}$$

$$\begin{aligned}
&= \omega(g, g') [\omega(gg', g'') \omega(gg'g'', g''') \omega(gg', g''g''')^{-1} S(gg') (\omega(g'', g''')^{-1})] \\
&\quad S(gg') (\omega(g'', g''')) \omega(g, g')^{-1} S(g) [\omega(g', g''g''') \omega(g'g'', g''')^{-1} \omega(g', g'')^{-1}] \\
&\quad S(g) (\omega(g', g'')) \omega(g, g'g'') \omega(gg', g'')^{-1} \omega(gg', g''g''') \omega(g, g'g''g''')^{-1} \\
&\quad S(g) (\omega(g', g''g''')^{-1}) [S(g) (\omega(g'g'', g''')) \omega(g, g'g''g''') \omega(gg'g'', g''')^{-1} \\
&\quad \omega(g, g'g'')^{-1}] \\
&= \omega(g, g') \omega(gg', g'') \omega(gg'g'', g''') \omega(gg', g''g''')^{-1} \omega(g, g')^{-1} S(g) [\omega(g', g''g''') \\
&\quad \omega(g'g'', g''')^{-1}] \omega(g, g'g'') \omega(gg', g'')^{-1} \omega(gg', g''g''') \omega(g, g'g''g''')^{-1} \\
&\quad S(g) (\omega(g', g''g''')^{-1}) [S(g) (\omega(g'g'', g''')) \omega(g, g'g''g''') \\
&\quad \omega(gg'g'', g''')^{-1} \omega(g, g'g'')^{-1}] \\
&= \omega(g, g') \omega(gg', g'') \omega(gg'g'', g''') \omega(gg', g''g''')^{-1} \omega(g, g')^{-1} S(g) \\
&\quad [\omega(g', g''g''') \omega(g'g'', g''')^{-1}] [S(g) (\omega(g'g'', g''')) \omega(g, g'g''g''') \\
&\quad \omega(gg'g'', g''')^{-1} \omega(g, g'g'')^{-1}] \omega(g, g'g'') \omega(gg', g'')^{-1} \omega(gg', g''g''') \\
&\quad \omega(g, g'g''g''')^{-1} S(g) (\omega(g', g''g''')^{-1}) \\
&= \omega(g, g') \omega(gg', g'') \omega(gg'g'', g''') \omega(gg', g''g''')^{-1} \omega(g, g')^{-1} \\
&\quad \{S(g) (\omega(g', g''g''')) \omega(g, g'g''g''') [\omega(gg'g'', g''')^{-1} \omega(gg', g'')^{-1} \\
&\quad \omega(gg', g''g''')] \omega(g, g'g''g''')^{-1} S(g) (\omega(g', g''g''')^{-1})\} \\
&= \omega(g, g') \omega(gg', g'') \omega(gg'g'', g''') \omega(gg', g''g''')^{-1} \omega(g, g')^{-1} \\
&\quad c_{S(g) [\omega(g', g''g''')] \omega(g, g'g''g''')} [\omega(gg'g'', g''')^{-1} \omega(gg', g'')^{-1} \omega(gg', g''g''')] \\
&\stackrel{(*)}{=} \omega(g, g') \omega(gg', g'') \omega(gg'g'', g''') \omega(gg', g''g''')^{-1} \omega(g, g')^{-1} \\
&\quad c_{\omega(g, g') \omega(gg', g''g''')} [\omega(gg'g'', g''')^{-1} \omega(gg', g'')^{-1} \omega(gg', g''g''')] \\
&= \omega(g, g') \omega(gg', g'') \omega(gg'g'', g''') \omega(gg', g''g''')^{-1} \omega(g, g')^{-1} \omega(g, g') \\
&\quad \omega(gg', g''g''') [\omega(gg'g'', g''')^{-1} \omega(gg', g'')^{-1} \omega(gg', g''g''')] \\
&\quad \omega(gg', g''g''')^{-1} \omega(g, g')^{-1} = \mathbf{1}.
\end{aligned}$$

(6) We prove the assertion for strongly smooth outer actions. The other case is proved similarly by using elements $\omega \in C_s^2(G, N)$ instead.

We fix some $\omega \in C_{ss}^2(G, N)$ with $\delta_S = C_N \circ \omega$. If $\omega' \in C_{ss}^2(G, N)$ also satisfies $\delta_S = C_N \circ \omega'$, then $\beta := \omega' \cdot \omega^{-1} \in C_{ss}^2(G, Z(N))$ and thus $d_S \omega' = d_S \omega \cdot d_S \beta$, hence $[d_S \omega'] = [d_S \omega]$ and the cohomology class $[d_S \omega]$ does not depend on the choice of ω .

If $S' \sim S$, then there exists an $h \in C_s^1(G, N)$ with $S' = (C_N \circ h) \cdot S$, and we have seen in (2) that $\omega' := h *_S \omega \in C_{ss}^2(G, N)$ satisfies $\delta_{S'} = C_N \circ \omega'$.

Now $d_{S'}\omega' = d_S\omega$ implies in particular that $[d_{S'}\omega'] = [d_S\omega]$ and that this cohomology class neither depends on the representative S' of $[S]$ nor on the choice of ω . □

DEFINITION 1.11. — *Let $S \in C_s^1(G, \text{Aut}(N))$ be a strongly smooth outer action and pick $\omega \in C_{ss}^2(G, N)$ with $\delta_S = C_N \circ \omega$. In view of Lemma 1.10(6), the cohomology class*

$$\chi_{ss}(S) := [d_S\omega] \in H_{ss}^3(G, Z(N))_S$$

does not depend on the choice of ω and is constant on the equivalence class of S , so that we may also write $\chi_{ss}([S]) := \chi_{ss}(S)$. We call $\chi_{ss}(S)$ the characteristic class of S .

For a smooth outer action we likewise obtain a characteristic class

$$\chi_s(S) = \chi_s([S]) = [d_S\omega] \in H_s^3(G, Z(N))_S.$$

2. Extensions of Lie groups

In this section we discuss the basic concepts related to extensions of Lie groups. Since every discrete group can be viewed as a Lie group, our discussion includes in particular the algebraic context of group extensions ([13], [7]).

DEFINITION 2.1. — *Let G be a Lie group. A subgroup H of G is called a split Lie subgroup if it carries a Lie group structure for which the inclusion map $i_H: H \hookrightarrow G$ is a morphism of Lie groups and the right action of H on G defined by restricting the multiplication map of G to a map $G \times H \rightarrow G$ defines a smooth H -principal bundle. This means that the coset space G/H is a smooth manifold and that the quotient map $\pi: G \rightarrow G/H$ has smooth local sections.*

Example 2.2. — Since the Lie algebra \mathfrak{h} of a Lie subgroup H of a Lie group G need not have a closed complement in \mathfrak{g} , not every Lie subgroup is split. A simple example is the subgroup $H := c_0(\mathbb{N}, \mathbb{R})$ in $G := \ell^\infty(\mathbb{N}, \mathbb{R})$ (cf. [27, Satz IV.6.5]).

LEMMA 2.3. — *Each split Lie subgroup of G is initial.*

Proof. — The condition that H is a split Lie subgroup implies that there exists an open subset U of some locally convex space V and a smooth map $\sigma: U \rightarrow G$ such that the map

$$U \times H \rightarrow G, \quad (x, h) \mapsto \sigma(x)h$$

is a diffeomorphism onto an open subset of G . We conclude that for every manifold M a map $f: M \rightarrow H$ is smooth whenever it is smooth as a map to G , resp. the open subset $\sigma(U)H$ of G . \square

DEFINITION 2.4. — An extension of Lie groups is a short exact sequence

$$\mathbf{1} \rightarrow N \xrightarrow{\iota} \widehat{G} \xrightarrow{q} G \rightarrow \mathbf{1}$$

of Lie group morphisms, for which $N \cong \ker q$ is a split Lie subgroup. This means that \widehat{G} is a smooth N -principal bundle over G and $G \cong \widehat{G}/N$. In the following we shall identify N with the subgroup $\iota(N) \trianglelefteq \widehat{G}$.

We call two extensions $N \hookrightarrow \widehat{G}_1 \twoheadrightarrow G$ and $N \hookrightarrow \widehat{G}_2 \twoheadrightarrow G$ of the Lie group G by the Lie group N equivalent if there exists a Lie group morphism $\varphi: \widehat{G}_1 \rightarrow \widehat{G}_2$ such that the following diagram commutes:

$$\begin{array}{ccccc} N & \hookrightarrow & \widehat{G}_1 & \twoheadrightarrow & G \\ \downarrow \text{id}_N & & \downarrow \varphi & & \downarrow \text{id}_G \\ N & \hookrightarrow & \widehat{G}_2 & \twoheadrightarrow & G. \end{array}$$

It is easy to see that any such φ is in particular an isomorphism of Lie groups. We write $\text{Ext}(G, N)$ for the set of equivalence classes of Lie group extensions of G by N .

We call an extension $q: \widehat{G} \twoheadrightarrow G$ with $\ker q = N$ split if there exists a Lie group morphism $\sigma: G \rightarrow \widehat{G}$ with $q \circ \sigma = \text{id}_G$. In this case the map $N \times_S G \rightarrow \widehat{G}, (n, g) \mapsto n\sigma(g)$ is an isomorphism, where the semidirect product is defined by the homomorphism

$$S := C_N \circ \sigma: G \rightarrow \text{Aut}(N),$$

where $C_N: \widehat{G} \rightarrow \text{Aut}(N)$ is the conjugation action of \widehat{G} on N . In view of Lemmas 1.7 and 2.3, S defines a smooth action of G on N .

Remark 2.5. — We give a description of Lie group extensions $N \hookrightarrow \widehat{G} \twoheadrightarrow G$ in terms of data associated to G and N . Let $q: \widehat{G} \twoheadrightarrow G$ be a Lie group extension of G by N . By assumption, the map q has a smooth local section. Hence there exists a locally smooth global section $\sigma: G \rightarrow \widehat{G}$ which is normalized in the sense that $\sigma(\mathbf{1}) = \mathbf{1}$, i.e., $\sigma \in C_s^1(G, \widehat{G})$. Then the map

$$\Phi: N \times G \rightarrow \widehat{G}, \quad (n, g) \mapsto n\sigma(g)$$

is a bijection which restricts to a local diffeomorphism on an identity neighborhood. In general Φ is not continuous, but we may nevertheless use it to identify \widehat{G} with the product set $N \times G$ endowed with the multiplication

$$(2.1) \quad (n, g)(n', g') = (nS(g)(n')\omega(g, g'), gg'),$$

where

$$(2.2) \quad S := C_N \circ \sigma : G \rightarrow \text{Aut}(N) \text{ for } C_N : \widehat{G} \rightarrow \text{Aut}(N), \quad C_N(g) = gng^{-1},$$

and

$$(2.3) \quad \omega : G \times G \rightarrow N, \quad (g, g') \mapsto \sigma(g)\sigma(g')\sigma(gg')^{-1}.$$

Note that ω is smooth in an identity neighborhood and the map $G \times N \rightarrow N, (g, n) \mapsto S(g).n$ is smooth in a set of the form $U \times N$, where U is an identity neighborhood of G . The maps S and ω satisfy the relations

$$(2.4) \quad \sigma(g)\sigma(g') = \omega(g, g')\sigma(gg'),$$

$$(2.5) \quad S(g)S(g') = C_N(\omega(g, g'))S(gg'), \text{ i.e., } \delta_S = C_N \circ \omega,$$

and

$$(2.6) \quad \omega(g, g')\omega(gg', g'') = S(g)(\omega(g', g''))\omega(g, g'g''), \text{ i.e., } d_S\omega = \mathbf{1}.$$

DEFINITION 2.6. — *The elements of the set*

$$Z_{ss}^2(G, N) := \{(S, \omega) \in C_s^1(G, \text{Aut}(N)) \times C_{ss}^2(G, N) : \delta_S = C_N \circ \omega, d_S\omega = \mathbf{1}\}$$

are called smooth factor systems for the pair (G, N) or locally smooth 2-cocycles. Note that for

$$Z_{ss}^2(G, N)_S := \{\omega \in C_{ss}^2(G, N) : \delta_S = C_N \circ \omega, d_S\omega = \mathbf{1}\}$$

we may write the set $Z_{ss}^2(G, N)$ as the disjoint union

$$Z_{ss}^2(G, N) = \bigcup_S Z_{ss}^2(G, N)_S,$$

which explains the shift in notation from 2-cocycles as functions ω to pairs (S, ω) . If N is abelian, then one can fix S and deal with each set $Z_{ss}^2(G, N)_S$ separately, but this is impossible for nonabelian groups N .

LEMMA 2.7. — *Let $S \in C^1(G, \text{Aut}(N))$ and $\omega \in C^2(G, N)$ with $\delta_S = C_N \circ \omega$. We define a product on $N \times G$ by*

$$(n, g)(n', g') = (nS(g)(n')\omega(g, g'), gg').$$

Then $N \times G$ is a group if and only if $d_S\omega = \mathbf{1}$. Inversion in this group is given by

$$\begin{aligned} (n, g)^{-1} &= (S(g)^{-1}(n^{-1}\omega(g, g^{-1})^{-1}), g^{-1}) \\ &= (\omega(g^{-1}, g)^{-1}S(g^{-1})(n^{-1}), g^{-1}) \end{aligned}$$

and in particular

$$(\mathbf{1}, g)^{-1} = (S(g)^{-1} \cdot (\omega(g, g^{-1})^{-1}), g^{-1}) = (\omega(g^{-1}, g)^{-1}, g^{-1}).$$

Conjugation is given by

$$(n, g)(n', g')(n, g)^{-1} = \left(nS(g)(n')\omega_g(g')S(gg'g^{-1})^{-1}(n^{-1}), gg'g^{-1} \right),$$

so that $(n, \mathbf{1})(n', g')(n, \mathbf{1})^{-1} = (nn'S(g')(n^{-1}), g')$ and

$$(2.7) \quad (\mathbf{1}, g)(n', g')(\mathbf{1}, g)^{-1} = (S(g)(n')\omega_g(g'), gg'g^{-1}).$$

Proof. — The associativity of the multiplication on $N \times G$ is equivalent to the equality of

$$\begin{aligned} ((n, g)(n', g'))(n'', g'') &= (nS(g)(n')\omega(g, g'), gg')(n'', g'') \\ &= (nS(g)(n')\omega(g, g')S(gg')(n'')\omega(gg', g''), gg'g'') \end{aligned}$$

and

$$\begin{aligned} (n, g)((n', g')(n'', g'')) &= (n, g)(n'S(g')(n'')\omega(g', g''), g'g'') \\ &= (nS(g)(n'S(g')(n'')\omega(g', g''))\omega(g, g'g''), gg'g'') \\ &= (nS(g)(n')(S(g)S(g')(n''))S(g)(\omega(g', g''))\omega(g, g'g''), gg'g'') \end{aligned}$$

for all $g, g', g'' \in G$ and $n, n', n'' \in N$. This means that

$$\omega(g, g')S(gg')(n'')\omega(gg', g'') = (S(g)S(g')(n''))S(g)(\omega(g', g''))\omega(g, g'g'').$$

For $n'' = \mathbf{1}$ this leads to $d_S\omega = \mathbf{1}$. If, conversely, $d_S\omega = \mathbf{1}$, then the associativity condition is equivalent to

$$\omega(g, g')S(gg')(n'')\omega(gg', g'') = (S(g)S(g')(n''))\omega(g, g')\omega(gg', g'')$$

and hence to $\delta_S = C_N \circ \omega$. Therefore the conditions $\delta_S = C_N \circ \omega$ and $d_S\omega = \mathbf{1}$ are equivalent to the associativity of the multiplication on $N \times G$.

To see that we actually obtain a group, we first observe that $S(\mathbf{1}) = \text{id}_N$ and $\omega(g, \mathbf{1}) = \omega(\mathbf{1}, g) = \mathbf{1}$ imply that $\mathbf{1} := (\mathbf{1}, \mathbf{1})$ is an identity element of $\widehat{G} := N \times G$, so that $(\widehat{G}, \mathbf{1})$ is a monoid. For $(n, g) \in \widehat{G}$ the element

$$(S(g)^{-1}(n^{-1}\omega(g, g^{-1})^{-1}), g^{-1})$$

is a right inverse and likewise

$$(\omega(g^{-1}, g)^{-1}S(g^{-1})(n^{-1}), g^{-1})$$

is a left inverse. Now the associativity of \widehat{G} implies that left and right inverse are equal, hence an inverse of (n, g) . Therefore \widehat{G} is a group.

The formula for the inversion has already been obtained above. For the conjugation we first observe that for $g, g' \in G$ we have

$$\begin{aligned} \mathbf{1} &= (d_S\omega)(gg', g^{-1}, g) \\ &= S(gg')(\omega(g^{-1}, g))\omega(gg', \mathbf{1})\omega(gg'g^{-1}, g)^{-1}\omega(gg', g^{-1})^{-1} \\ &= S(gg')(\omega(g^{-1}, g))\omega(gg'g^{-1}, g)^{-1}\omega(gg', g^{-1})^{-1}, \end{aligned}$$

so that

$$S(gg')(\omega(g^{-1}, g))^{-1}\omega(gg', g^{-1}) = \omega(gg'g^{-1}, g)^{-1}.$$

We now obtain

$$\begin{aligned} (n, g)(n', g')(n, g)^{-1} &= \left(nS(g)(n')\omega(g, g'), gg' \right) \left(\omega(g^{-1}, g)^{-1}S(g^{-1})(n^{-1}), g^{-1} \right) \\ &= \left(nS(g)(n')\omega(g, g')S(gg')(\omega(g^{-1}, g)^{-1}S(g^{-1})(n^{-1})) \right. \\ &\quad \left. \cdot \omega(gg', g^{-1}), gg'g^{-1} \right) \\ &= \left(nS(g)(n')\omega(g, g')\omega(gg'g^{-1}, g)^{-1}\omega(gg', g^{-1})^{-1} \right. \\ &\quad \left. \cdot (S(gg')S(g^{-1})(n^{-1}))\omega(gg', g^{-1}), gg'g^{-1} \right) \\ &= \left(nS(g)(n')\omega_g(g')C_N(\omega(gg', g^{-1})^{-1})(S(gg')S(g^{-1})(n^{-1})), gg'g^{-1} \right) \\ &= \left(nS(g)(n')\omega_g(g')\delta_S(gg', g^{-1})^{-1}(S(gg')S(g^{-1})(n^{-1})), gg'g^{-1} \right) \\ &= \left(nS(g)(n')\omega_g(g')S(gg'g^{-1})^{-1}(n^{-1}), gg'g^{-1} \right). \end{aligned}$$

□

For a smooth factor system (S, ω) we write $N \times_{(S, \omega)} G$ for the set $N \times G$ endowed with the group multiplication (2.1).

PROPOSITION 2.8.

(1) *If (S, ω) is a smooth factor system, then $N \times_{(S, \omega)} G$ carries a unique structure of a Lie group for which the map*

$$N \times G \rightarrow N \times_{(S, \omega)} G, \quad (n, g) \mapsto (n, g)$$

is smooth on a set of the form $N \times U$, where U is an open identity neighborhood in G . Then

$$q: N \times_{(S, \omega)} G \rightarrow G, \quad (n, g) \mapsto g$$

is a Lie group extension of G by N .

(2) *Each Lie group extension $q: \widehat{G} \rightarrow G$ of G by N gives rise to smooth factor systems by choosing a locally smooth normalized section $\sigma: G \rightarrow$*

\widehat{G} and defining $(S, \omega) := (C_N \circ \sigma, \delta_\sigma)$. Then the extension $q: \widehat{G} \rightarrow G$ is equivalent to $N \times_{(S, \omega)} G$.

Proof.

(1) Let $U_G \subseteq G$ be an open symmetric **1**-neighborhood such that the maps

$$U_G \times N \rightarrow N, \quad (g, n) \mapsto S(g).n \text{ and } \omega|_{U_G \times U_G}$$

are smooth. We consider the subset

$$U := N \times U_G = q^{-1}(U_G) \subseteq \widehat{G} := N \times_{(S, \omega)} G.$$

Then $U = U^{-1}$ because q is a group homomorphism. We endow U with the product manifold structure from $N \times U_G$. Since the multiplication $m_G|_{U_G \times U_G}: U_G \times U_G \rightarrow G$ is continuous, there exists an open symmetric identity neighborhood $V_G \subseteq U_G$ with $V_G V_G \subseteq U_G$. Then the set $V := N \times V_G$ is an open subset of U such that the multiplication map

$$V \times V \rightarrow U, \quad ((n, g), (n', g')) \mapsto (nS(g)(n')\omega(g, g'), gg')$$

is smooth. The inversion

$$U \rightarrow U, \quad (n, g) \mapsto (\omega(g^{-1}, g)^{-1}S(g^{-1})(n^{-1}), g^{-1})$$

(cf. Lemma 2.7) is also smooth.

For $(n, g) \in \widehat{G}$ let $V_g \subseteq U_G$ be an open identity neighborhood with $c_g(V_g) \subseteq U_G$. Then $c_{(n, g)}(q^{-1}(V_g)) \subseteq U$. That the conjugation map $c_{(n, g)}: q^{-1}(V_g) \rightarrow U$ is smooth in an identity neighborhood follows immediately from formula (2.7) in Lemma 2.7 because $\omega \in C_{ss}^2(G, N)$ ensures that the function ω_g is smooth in an identity neighborhood of G .

Eventually Theorem B.1 implies that \widehat{G} carries a unique Lie group structure for which the inclusion map $U = N \times U_G \hookrightarrow \widehat{G}$ is a local diffeomorphism onto an identity neighborhood. It is clear that with respect to the Lie group structure on \widehat{G} , the map $q: \widehat{G} \rightarrow G$ is a smooth N -principal bundle because the map $V_G \rightarrow \widehat{G}, g \mapsto (0, g)$ defines a section of q which is smooth on an identity neighborhood in G that might be smaller than V_G .

(2) Assume, conversely, that $q: \widehat{G} \rightarrow G$ is a Lie group extension of G by N . Then there exists an open **1**-neighborhood $U_G \subseteq G$ and a smooth section $\sigma: U_G \rightarrow \widehat{G}$ of the map $q: \widehat{G} \rightarrow G$. We extend σ to a global section $G \rightarrow \widehat{G}$ and put $(S, \omega) := (C_N \circ \sigma, \delta_\sigma)$. We have already seen in Definition 1.1 that $\omega = \delta_\sigma \in C_{ss}^2(G, N)$. Therefore (S, ω) is a smooth factor system, so that we can use (1) to obtain a Lie group structure on $N \times_{(S, \omega)} G$ for which there exists an identity neighborhood $V_G \subseteq G$ for which the product map

$$N \times V_G \rightarrow N \times_{(S, \omega)} G, \quad (n, v) \mapsto (n, \mathbf{1})(0, v) = (n, v)$$

is smooth. This implies that the group isomorphism $N \times_{(S,\omega)} G \rightarrow \widehat{G}$ is a local diffeomorphism, hence an isomorphism of Lie groups. \square

If the group G is connected, then we can weaken the condition on smooth factor systems by replacing $\omega \in C_{ss}^2(G, N)$ by the weaker condition $\omega \in C_s^2(G, N)$.

LEMMA 2.9. — *If G is connected and $S \in C_s^1(G, \text{Aut}(N))$ and $\omega \in C_s^2(G, N)$ satisfy*

$$\delta_S = C_N \circ \omega \text{ and } d_S \omega = \mathbf{1},$$

then $\omega \in C_{ss}^2(G, N)$ and in particular $(S, \omega) \in Z_{ss}^2(G, N)$.

Proof. — That the map $\omega_g : G \rightarrow N$ is smooth in an identity neighborhood means that conjugation with $(\mathbf{1}, g)$ is smooth in an identity neighborhood because

$$(\mathbf{1}, g)(n', g')(\mathbf{1}, g)^{-1} = \left(S(g)(n')\omega_g(g'), gg'g^{-1} \right)$$

by formula (2.7).

The set of all elements $x \in N \times_{(S,\omega)} G$ for which the conjugation c_x is smooth in an identity neighborhood is a subsemigroup containing N (cf. Lemma 2.7) and $\{\mathbf{1}\} \times W$ for some identity neighborhood W in G . If G is connected, it is generated by W as a semigroup, so that $N \times_{(S,\omega)} G$ is generated by $N \times W$. Therefore the local smoothness requirement for the functions ω_g is redundant. \square

In the following proposition we show that equivalence classes of extensions correspond to orbits of the group $C_s^1(G, N)$ in the set of smooth factor systems.

PROPOSITION 2.10. — *For two smooth factor systems $(S, \omega), (S', \omega') \in Z_{ss}^2(G, N)$ the following are equivalent*

- (1) $N \times_{(S,\omega)} G$ and $N \times_{(S',\omega')} G$ are equivalent extensions of G by N .
- (2) There exists an $h \in C_s^1(G, N)$ with $h.(S, \omega) = (S', \omega')$.

If these conditions are satisfied, then the map

$$\varphi : N \times_{(S',\omega')} G \rightarrow N \times_{(S,\omega)} G, \quad (n, g) \mapsto (nh(g), g)$$

is an equivalence of extensions and all equivalences of extensions $N \times_{(S',\omega')} G \rightarrow N \times_{(S,\omega)} G$ are of this form.

Proof. — Let $\varphi : N \times_{(S',\omega')} G \rightarrow N \times_{(S,\omega)} G$ be an equivalence of extensions. Then there exists a function $h \in C_s^1(G, N)$ such that φ has the form

$\varphi(n, g) = (nh(g), g)$ with respect to the product coordinates on $N \times_{(S, \omega)} G$, resp. $N \times_{(S', \omega')} G$. We then have

$$\begin{aligned} \varphi(n, g)\varphi(n', g') &= (nh(g), g)(n'h(g'), g') = (nh(g)S(g)(n'h(g'))\omega(g, g'), gg') \\ &= (n[C_N(h(g))(S(g)(n'))]h(g)S(g)(h(g'))\omega(g, g'), gg') \\ &= (n(h.S)(g)(n')(h *_S \omega)(g, g')h(gg'), gg') \\ &= \varphi(n(h.S)(g)(n')(h *_S \omega)(g, g'), gg'). \end{aligned}$$

Since φ is an equivalence of extensions, it is an injective morphism of groups, which implies that the product in $N \times_{(S', \omega')} G$ is given by

$$\begin{aligned} (nS'(g)(n')\omega'(g, g'), gg') &= (n, g)(n', g') \\ &= (n(h.S)(g)(n')(h *_S \omega)(g, g'), gg'). \end{aligned}$$

This implies that $\omega' = h *_S \omega$ and further that $S' = h.S$, so that “(1) \Rightarrow (2)” is proved.

If, conversely, $(S', \omega') = h.(S, \omega) = (h.S, h *_S \omega)$, then we define

$$\varphi: N \times_{(S', \omega')} G \rightarrow N \times_{(S, \omega)} G, \quad \varphi(n, g) := (nh(g), g),$$

and the calculation above shows that φ is a homomorphisms of groups. Since φ is smooth in an identity neighborhood, it is a morphisms of Lie groups. Clearly $\varphi|_N = \text{id}_N$ and φ induces the identity map on G , so that it is an equivalence of extensions. This completes the proof. \square

COROLLARY 2.11. — *The map*

$$Z_{ss}^2(G, N) \rightarrow \text{Ext}(G, N), \quad (S, \omega) \mapsto [N \times_{(S, \omega)} G]$$

induces a bijection

$$H_{ss}^2(G, N) := Z_{ss}^2(G, N)/C_s^1(G, N) \rightarrow \text{Ext}(G, N).$$

The preceding proposition implies in particular that

$$N \times_{(S, \omega)} G \sim N \times_{(S', \omega')} G$$

implies $[S] = [S']$, i.e., equivalent extensions correspond to the same smooth G -kernel. In the following we write $\text{Ext}(G, N)_{[S]}$ for the set of equivalence classes of N -extensions of G corresponding to the G -kernel $[S]$.

THEOREM 2.12. — *Let S be a smooth outer action of G on N with $\text{Ext}(N, G)_{[S]} \neq \emptyset$. Then each extension class in $\text{Ext}(G, N)_{[S]}$ can be represented by a Lie group of the form $N \times_{(S, \omega)} G$. Any other Lie group extension $N \times_{(S, \omega')} G$ representing an element of $\text{Ext}(G, N)_{[S]}$ satisfies*

$$\omega' \cdot \omega^{-1} \in Z_{ss}^2(G, Z(N))_{[S]},$$

and the Lie groups $N \times_{(S,\omega)} G$ and $N \times_{(S,\omega')} G$ define equivalent extensions if and only if

$$\omega' \cdot \omega^{-1} \in B_s^2(G, Z(N))_{[S]}.$$

Proof. — From Proposition 2.8 we know that each Lie group extension \widehat{G} of G by N is equivalent to one of the form $N \times_{(S',\omega')} G$. If $[S] = [S']$ and $h \in C_s^1(G, N)$ satisfies $h.S = S'$, then $h^{-1}.(S', \omega') = (S, h^{-1} *_S \omega')$, so that $\omega'' := h^{-1} *_S \omega'$ satisfies $N \times_{(S',\omega')} G \cong N \times_{(S,\omega'')} G$. This means that each extension class in $\text{Ext}(G, N)_{[S]}$ can be represented by a group of the form $N \times_{(S,\omega'')} G$.

If (S, ω) and (S, ω') are smooth factor systems, then $C_N \circ \omega = \delta_S = C_N \circ \omega'$ implies $\beta := \omega' \cdot \omega^{-1} \in C_s^2(G, Z(N))$. Further $\text{im}(\beta) \subseteq Z(N)$ leads to $\beta_g = \omega'_g \omega_g^{-1}$ for each $g \in G$:

$$\begin{aligned} \beta_g(x) &= \beta(g, x)\beta(gxg^{-1}, g)^{-1} \\ &= \omega'(g, x)\omega(g, x)^{-1}\omega(gxg^{-1}, g)\omega'(gxg^{-1}, g)^{-1} \\ &= \omega'_g(x)C_N(\omega'(gxg^{-1}, g))(\omega(g, x)^{-1}\omega(gxg^{-1}, g)) \\ &= \omega'_g(x)C_N(\omega(gxg^{-1}, g))(\omega(g, x)^{-1}\omega(gxg^{-1}, g)) \\ &= \omega'_g(x)\omega_g(x)^{-1}. \end{aligned}$$

This shows that β_g is smooth in an identity neighborhood, so that $\beta \in C_{ss}^2(G, Z(N))$. Moreover, $d_S \omega' = d_S \omega = \mathbf{1}$, so that

$$\mathbf{1} = d_S \omega' = d_S \omega \cdot d_S \beta = d_S \beta$$

implies $\beta \in Z_{ss}^2(G, Z(N))_{[S]}$. This means that $\omega' \in \omega \cdot Z_{ss}^2(G, Z(N))_S$.

In view of Proposition 2.10, the equivalence of the extensions $N \times_{(S,\omega')} G$ and $N \times_{(S,\omega)} G$ is equivalent to the existence of an $h \in C_s^1(G, N)$ with

$$S = h.S = (C_N \circ h) \cdot S \text{ and } \omega' = h *_S \omega.$$

Then $C_N \circ h = \text{id}_N$ implies $h \in C_s^1(G, Z(N))$ which further leads to $\omega' = h *_S \omega = (d_S h) \cdot \omega$, i.e., $\omega' \omega^{-1} \in B_s^2(G, Z(N))_S$. If, conversely, $\omega' \omega^{-1} = d_S h$ for some $h \in C_s^1(G, Z(N))$, then $h.S = S$ and $h *_S \omega = \omega'$. \square

COROLLARY 2.13. — *For a smooth G -kernel $[S]$ with $\text{Ext}(G, N)_{[S]} \neq \emptyset$, the map*

$$\begin{aligned} H_{ss}^2(G, Z(N))_{[S]} \times \text{Ext}(G, N)_{[S]} &\rightarrow \text{Ext}(G, N)_{[S]}, \\ (\beta, [N \times_{(S,\omega)} G]) &\mapsto [N \times_{(S,\omega \cdot \beta)} G] \end{aligned}$$

is a well-defined simply transitive action.

Remark 2.14 (Abelian extensions). — Suppose that $N = A$ is an abelian Lie group. Then the adjoint representation of A is trivial and a smooth factor system (S, ω) for (G, A) consists of a smooth module structure $S: G \rightarrow \text{Aut}(A)$ and an element $\omega \in C_{ss}^2(G, A)$. In this case d_S is the Lie group differential corresponding to the module structure on A (Definition 1.2). Therefore (S, ω) defines a Lie group $N \times_{(S, \omega)} G$ if and only if $d_S \omega = \mathbf{1}$, i.e., $\omega \in Z_{ss}^2(G, A)$. In this case we write $A \times_\omega G$ for this Lie group, which is $A \times G$, endowed with the multiplication

$$(a, g)(a', g') = (a + g.a' + \omega(g, g'), gg').$$

Further $S \sim S'$ if and only if $S = S'$. Hence a smooth G -kernel $[S]$ is the same as a smooth G -module structure S on A and $\text{Ext}(G, A)_S := \text{Ext}(G, A)_{[S]}$ is the class of all A -extensions of G for which the associated G -module structure on A is S .

According to Corollary 2.13, the equivalence classes of extensions correspond to cohomology classes of cocycles, so that the map

$$H_{ss}^2(G, A)_S \rightarrow \text{Ext}(G, A)_S, \quad [\omega] \mapsto [A \times_\omega G]$$

is a well-defined bijection. This was also shown directly in [19] in the context of abelian extensions.

Remark 2.15. — The result of Corollary 2.13 can also be visualized on the level of extensions as follows. For that we assume that $Z(N)$ is an initial Lie subgroup of N , so that it carries its own Lie group structure and the smooth outer action S induces on $Z(N)$ the structure S_Z of a smooth G -module (Lemma 1.6). In view of Remark 2.14, we then have

$$H_{ss}^2(G, Z(N))_S \cong \text{Ext}(G, Z(N))_{S_Z}.$$

Let $Z(N) \rightarrow \widehat{G}_1 \xrightarrow{q_1} G$ be an extension of G by the smooth G -module $Z(N)$ and $N \rightarrow \widehat{G}_2 \xrightarrow{q_2} G$ an extension of G by N corresponding to the G -kernel $[S]$. Then we consider the group

$$H := q_1^* \widehat{G}_2 \cong \{(g_1, g_2) \in \widehat{G}_1 \times \widehat{G}_2 : q_1(g_1) = q_2(g_2)\}$$

which is a Lie group extension of G by $Z(N) \times N$, and the subgroup $\Delta_Z := \{(z, z^{-1}) : z \in Z(N)\}$ is a central split Lie subgroup of H . Therefore the Baer product $\widehat{G} := H/\Delta_Z$ is a Lie group extension of G by $N \cong (Z(N) \times N)/\Delta_Z$.

If $\widehat{G}_1 \cong Z(N) \times_f G$ and $\widehat{G}_2 \cong N \times_{(S, \omega)} G$, then

$$H \cong (Z(N) \times N) \times_{((S_Z, S), (f, \omega))} G$$

and $\widehat{G} \cong N \times_{(S, f \cdot \omega)} G$. This implies that the action of the abelian group $H_{ss}^2(G, Z(N))_S$ on $\text{Ext}(G, N)_{[S]}$ corresponds to the Baer multiplication of extensions of G by $Z(N)$ with N -extensions of G .

Remark 2.16. — The preceding results show that the fibers of the map $[(S, \omega)] \mapsto [S]$ that assigns to an extension of G by N , represented by a smooth factor system (S, ω) , the class $[S]$ of the corresponding smooth outer action, are given by $Q^{-1}([S]) = \text{Ext}(G, N)_{[S]}$, and the group $H_{ss}^2(G, Z(N))_S$ acts simply transitively on the fibers. Clearly the action of G on $Z(N)$ depends on $[S]$ and so does the cohomology group $H_{ss}^2(G, Z(N))_S$.

In the example described below, we shall see that for an abelian smooth G -module $A = N = Z(N)$ the group $H_{ss}^2(G, A)_S$ very much depends on the action S of G on A . For $G = \mathbb{R}^2$ and $A = \mathbb{R}$ we have $H_s^2(G, A) \cong H_c^2(\mathfrak{g}, \mathfrak{a})$ (cf. [19, Th. 7.2]). On the other hand $C_c^2(\mathfrak{g}, \mathfrak{a})$ is the space of alternating bilinear maps $\mathfrak{g} \times \mathfrak{g} \rightarrow \mathfrak{a}$, hence 1-dimensional. Further $\dim \mathfrak{g} = 2$ implies that each 2-cochain is a cocycle. Since $B_c^2(\mathfrak{g}, \mathfrak{a}) = d_{\mathfrak{g}}\mathfrak{a}$ vanishes if the module \mathfrak{a} is trivial and coincides with $Z_c^2(\mathfrak{g}, \mathfrak{a})$ otherwise, we have

$$H_{ss}^2(G, A) \cong H_s^2(G, A) \cong H_c^2(\mathfrak{g}, \mathfrak{a}) \cong \begin{cases} \mathbb{R} & \text{for } \mathfrak{g} \cdot \mathfrak{a} = \{0\} \\ \{0\} & \text{for } \mathfrak{g} \cdot \mathfrak{a} \neq \{0\}. \end{cases}$$

The following theorem provides a Lie theoretic criterion for the non-emptiness of the set $\text{Ext}(G, N)_{[S]}$.

THEOREM 2.17. — *If S is a strongly smooth outer action of G on N , then*

$$\chi_{ss}(S) = \mathbf{1} \iff \text{Ext}(G, N)_{[S]} \neq \emptyset.$$

If G is connected and S is a smooth outer action, then

$$\chi_s(S) = \mathbf{1} \iff \text{Ext}(G, N)_{[S]} \neq \emptyset.$$

Proof. — If there exists a Lie group extension \widehat{G} corresponding to $[S]$, then we may w.l.o.g. assume that it is of the form $N \times_{(S, \omega)} G$ for a smooth factor system (S, ω) (Theorem 2.12). This implies in particular that S is strongly smooth, and $d_S\omega = \mathbf{1}$ (Lemma 2.7) leads to $\chi_{ss}(S) = [d_S\omega] = \mathbf{1}$.

Suppose, conversely, that S is strongly smooth with $\chi_{ss}(S) = \mathbf{1}$. Then there exists $\omega \in C_{ss}^2(G, N)$ with $\delta_S = C_N \circ \omega$ and some $h \in C_{ss}^2(G, Z(N))$ with $d_S\omega = d_S h^{-1}$, so that $\omega' := \omega \cdot h \in C_{ss}^2(G, N)$ satisfies $d_S\omega' = d_S\omega \cdot d_S h = \mathbf{1}$ and $\delta_S = C_N \circ \omega'$. Hence (S, ω') is a smooth factor system, and Proposition 2.8 implies the existence of a Lie group extension $N \times_{(S, \omega)} G$ of G by N corresponding to $[S]$.

If G is connected and S is a smooth outer action with $\chi_s(S) = \mathbf{1}$, then a similar argument provides some $\omega' \in C_s^2(G, N)$ with $\delta_S = C_N \circ \omega'$ and

$d_S \omega' = \mathbf{1}$. In view of Lemma 2.9, $\omega' \in C_{ss}^2(G, N)$, and we thus obtain a Lie group extension $N \times_{(S, \omega')} G$ of G by N . \square

Split extensions

We conclude this section with a characterization of the *split extensions*, i.e., those extensions $q: \widehat{G} \rightarrow G$ for which there exists a smooth homomorphism $\sigma: G \rightarrow \widehat{G}$ with $q \circ \sigma = \text{id}_G$, which implies that $\widehat{G} \cong N \rtimes_S G$ with respect to the smooth action of G on N given by $S := C_N \circ \sigma$.

PROPOSITION 2.18. — *The extension $\widehat{G} = N \times_{(S, \omega)} G$ splits if and only if there exists an $h \in C_s^1(G, N)$ with*

$$(2.8) \quad h *_S \omega = \mathbf{1}.$$

Assume, in addition, that $\omega \in C_{ss}^2(G, Z(N))$. Then S is an action of G on N . With respect to this action, (2.8) is equivalent to

$$d_S h = \omega^{-1}.$$

Consider the set

$$Z_s^1(G, N, Z(N))_S := \{f \in C_s^1(G, N) : \text{im}(d_S f) \subseteq Z(N)\}$$

and the action of G on $\text{Aut}(N)$ by $S'(g)(\varphi) := S(g) \circ \varphi \circ S(g)^{-1}$. Then the abelian group $C_s^1(G, Z(N))$ acts on $C_s^1(G, N)$ by pointwise multiplication and we have an injective map

$$Z_s^1(G, N, Z(N))_S / C_s^1(G, Z(N)) \rightarrow Z^1(G, \text{Aut}(N))_S, \quad [f] \mapsto C_N \circ f$$

of the set of $C_s^1(G, Z(N))$ -orbits in $Z_s^1(G, N, Z(N))_S$ onto the set of all 1-cocycles $G \rightarrow C_N(N) \subseteq \text{Aut}(N)$ that can be lifted to elements of $C_s^1(G, N)$. Moreover,

$$[\omega] \cdot [N \times_{(S, \mathbf{1})} G] = [N \times_{(S, \omega)} G] \in \text{Ext}(G, N)_{[S]} \text{ splits} \iff [\omega^{-1}] \in \text{im}(\delta),$$

where δ is the natural map

$$\delta: Z_s^1(G, N, Z(N))_S / C_s^1(G, Z(N)) \rightarrow H_{ss}^2(G, Z(N))_S, \quad [f] \mapsto [d_S f].$$

Proof. — Since the orbit of the factor system (S, ω) under the group $C_s^1(G, N)$ consists of all other factor systems describing the same extension, i.e., corresponding to different choices of sections $G \rightarrow \widehat{G}$ (Proposition 2.10), the extension \widehat{G} splits if and only if this orbit contains a factor system $(S', \omega') = h \cdot (S, \omega)$ for which the corresponding section $\sigma': G \rightarrow \widehat{G}, g \mapsto (h(g), g)$ is a group homomorphism, i.e., $\delta_{\sigma'} = \mathbf{1}$. Hence the first assertion follows from $\delta_{\sigma'} = h *_S \omega$.

Suppose now that ω has values in $Z(N)$. Then $\delta_S = C_N \circ \omega = \mathbf{1}$ shows that S is a representation. Moreover, $h *_S \omega = d_S h \cdot \omega$ vanishes if and only if $d_S h = \omega^{-1}$.

To verify the remaining assertions, we first observe that for $f \in Z_s^1(G, N, Z(N))_S$ and $h \in C_s^1(G, Z(N))$ we have

$$\text{im}(d_S(fh)) = \text{im}(d_S(f)d_S(h)) \subseteq Z(N),$$

so that the subset $Z_s^1(G, N, Z(N))_S$ of $C_s^1(G, N)$ is invariant under the action of $C_s^1(G, Z(N))$ by pointwise multiplication. It is clear that for two elements $f, f' \in Z_s^1(G, N, Z(N))_S$ the condition $C_N \circ f = C_N \circ f'$ is equivalent to $f' \in f \cdot C_s^1(N, Z(N))$, which leads to the asserted injectivity statement. Moreover, for each $f \in Z_s^1(G, N, Z(N))_S$ the map $C_N \circ f: G \rightarrow \text{Aut}(N)$ is a 1-cocycle because $d_{S'}(C_N \circ f) = C_N \circ d_S f = \mathbf{1}$, and the latter relation shows that $Z_s^1(G, N, Z(N))_S$ is precisely the set of all elements of $C_s^1(G, N)$ for which $C_N \circ f$ is a 1-cocycle.

Finally we observe that the map δ is well-defined and that $[\omega^{-1}] \in \text{im}(\delta)$ is equivalent to the existence of some $h \in Z_s^1(G, N, Z(N))_S$ with $d_S h = \omega^{-1}$. Since ω has values in $Z(N)$, the observations from above show that this is equivalent to the splitting of $N \times_{(S, \omega)} G$. □

Remark 2.19. — In general the image of the map δ is a quite complicated subset of the cohomology group $H_{ss}^2(G, Z(N))_S$. To understand this subtle point, we note that $d_S h = \omega^{-1}$ does not seem to imply $d_S h^{-1} = \omega$. It is true that for $S_h = (C_N \circ h) \cdot S$ we have

$$d_{S_h} h^{-1} = h^{-1} *_S h \mathbf{1} = h^{-1} *_S (h *_S \omega) = \omega$$

because $(S, \omega) = h^{-1} \cdot (h \cdot (S, \omega)) = h^{-1} \cdot (S_h, \mathbf{1})$. The condition $d_S h^{-1} = d_{S_h} h^{-1}$ is equivalent to

$$S(g)(h(g')) = h(g)S(g)(h(g'))h(g)^{-1}$$

for $g, g' \in G$, which means that the two functions h and $S \cdot h$ commute pointwise.

For the special case $S = \mathbf{1}$ we have $d_S h = \delta_h$ and $\text{im}(\delta_h) \subseteq Z(N)$ means that the map $C_N \circ h: G \rightarrow \text{Aut}(N)$ is a homomorphism. Fix some $g > 1$ and let us consider the special case, where $N = \widetilde{\text{SL}}_2(\mathbb{R})$ is the universal covering group of $\text{SL}_2(\mathbb{R})$ and G is the discrete group with $2g$ generators $\alpha_1, \dots, \alpha_{2g}$ subject to the commutator relation

$$[\alpha_1, \alpha_2] \cdots [\alpha_{2g-1}, \alpha_{2g}] = \alpha_1 \alpha_2 \alpha_1^{-1} \alpha_2^{-1} \cdots \alpha_{2g-1} \alpha_{2g} \alpha_{2g-1}^{-1} \alpha_{2g}^{-1} = \mathbf{1}.$$

In this case $Z(N) \cong \mathbb{Z}$ is discrete and $N_{\text{ad}} := N/Z(N) \cong \text{PSL}_2(\mathbb{R})$. Then each homomorphism $h: G \rightarrow N_{\text{ad}}$ corresponds to a $(2g)$ -tuple of points

$(x_1, \dots, x_{2g}) \in \mathrm{PSL}_2(\mathbb{R})^{2g}$ satisfying

$$[x_1, x_2] \cdots [x_{2g-1}, x_{2g}] = \mathbf{1}.$$

Since G and $Z(N)$ are discrete, we have

$$H_{ss}^2(G, Z(N))_S = H^2(G, Z(N)).$$

Using the fact that G is the fundamental group $\pi_1(\Sigma_g)$ of a compact orientable surface Σ_g of genus g , it is known that

$$H^2(G, Z(N)) \cong H^2(\Sigma_g, Z(N)) \cong \mathrm{Hom}(H_2(\Sigma_g), Z(N)) \cong Z(N)$$

because $H_2(\Sigma_g) \cong \mathbb{Z}$ ([14]). The isomorphism

$$\Phi: H^2(G, Z(N)) \rightarrow Z(N)$$

can be obtained by choosing for each central extension $Z(N) \hookrightarrow \widehat{G} \rightarrow G$ a section $\sigma: G \rightarrow \widehat{G}$ and then putting

$$\delta([\widehat{G}]) := [\sigma(\alpha_1), \sigma(\alpha_2)] \cdots [\sigma(\alpha_{2g-1}), \sigma(\alpha_{2g})].$$

From this observation it follows that $\mathrm{im}(\delta) \subseteq Z(N)$ coincides with the set

$$\{[y_1, y_2] \cdots [y_{2g-1}, y_{2g}]: (y_1, \dots, y_{2g}) \in N, [y_1, y_2] \cdots [y_{2g-1}, y_{2g}] \in Z(N)\}.$$

In [14] it is shown that, as a subset of $\mathbb{Z} \cong Z(N)$, this set coincides with $\{n \in \mathbb{Z}: |n| < g\}$. In particular it is not a subgroup.

Remark 2.20. — Suppose that the extension $\widehat{G} = N \times_{(S,\omega)} G$ splits, so that we may w.l.o.g. assume that $\omega = \mathbf{1}$.

(a) For $h \in C_s^1(G, N)$ we then have

$$h.(S, \mathbf{1}) = (h.S, h *_S \mathbf{1}) = (h.S, d_S h)$$

with $d_S h(g, g') := h(g)S(g)(h(g'))h(gg')^{-1}$. We conclude that

$$\{h \in C_s^1(G, N): h *_S \mathbf{1} = \mathbf{1}\} = Z_s^1(G, N)_S$$

is the set of 1-cocycles for the smooth action of G on N defined by the homomorphism $S: G \rightarrow \mathrm{Aut}(N)$ (cf. Definition 1.3). Therefore the orbit of $(S, \mathbf{1})$ under $C_s^1(G, N)$ may contain different elements of the form $(S', \mathbf{1})$. The representations $S': G \rightarrow \mathrm{Aut}(N)$ arising that way have the form

$$S_f := (C_N \circ f) \cdot S, \quad f \in Z_s^1(G, N)_S.$$

If N is abelian, all representations S_f are the same, so that the $C_s^1(G, N)$ -orbit of $(S, \mathbf{1})$ contains no other element of the form $(S', \mathbf{1})$. But if $f \in Z_s^1(G, N)_S$ is a cocycle whose values do not lie in the center $Z(N)$ of N , then $S_f \neq S$. Nevertheless, the two split extensions

$$N \rtimes_S G \text{ and } N \rtimes_{S_f} G$$

are equivalent.

(b) A condition that is slightly weaker than the condition $d_S h = \mathbf{1}$ is that $S_h = (C_N \circ h) \cdot S$ is a homomorphism, which, in view of $\delta_{h,S} = C_N \circ (d_S h)$ (cf. proof of Lemma 1.10(1)) is equivalent to $\text{im}(d_S h) \subseteq Z(N)$, which in turn means that $C_N \circ h \in Z^1(G, \text{Aut}(N))$ with respect to the action of G on $\text{Aut}(N)$ given by $g \cdot \varphi := S(g)\varphi S(g)^{-1}$. Then $d_S d_S h = d_{S_h} d_S h = d_{S_h}(h *_S \mathbf{1}) = \mathbf{1}$ (Lemma 1.10(4)) implies that $d_S h \in Z_{ss}^2(G, Z(N))_S$, so that on the level of extension classes we have

$$[N \rtimes_S G] = [N \times_{(h,S,d_S h)} G] = [d_S h].[N \times_{(S_h, \mathbf{1})} G] = [d_S h].[N \rtimes_{S_h} G].$$

This shows that the semidirect products

$$N \rtimes_S G \text{ and } N \rtimes_{S_h} G$$

are equivalent extensions of G by N if and only if the class $[d_S h] \in H_{ss}^2(G, Z(N))_S$ vanishes.

COROLLARY 2.21. — *Suppose that $Z(N) \trianglelefteq N$ is a split Lie subgroup, so that $N_{\text{ad}} := N/Z(N)$ carries a canonical Lie group structure for which $q_N: N \rightarrow N_{\text{ad}} = N/Z(N)$ defines a central extension of N_{ad} by $Z(N)$. Further let $S: G \rightarrow \text{Aut}(N)$ be a smooth action of G on N and consider the induced action \bar{S} of G on N_{ad} . Then we have an exact sequence of groups, resp. pointed spaces,*

$$Z(N)^G \hookrightarrow N^G \rightarrow N_{\text{ad}}^G \rightarrow H_s^1(G, Z(N))_S \rightarrow H_s^1(G, N)_S \rightarrow H_s^1(G, N_{\text{ad}})_{\bar{S}} \xrightarrow{\delta} H_{ss}^2(G, Z(N))_S,$$

where for $f \in Z_s^1(G, N_{\text{ad}})_S$ we have

$$\delta([f]) = [d_S \hat{f}]$$

for some $\hat{f} \in Z_s^1(G, N, Z(N))_S$ satisfying $q_N \circ \hat{f} = f$ and

$$\text{Ext}(G, N)_{[S], \text{split}} = -\text{im}(\delta).[N \rtimes_S G].$$

Proof. — For the 7-term exact sequence and the connecting map δ corresponding to the central Lie group extension $Z(N) \hookrightarrow N \twoheadrightarrow N_{\text{ad}}$, we refer to [17, Sect. II].

If the cocycle $f: G \rightarrow N$ has values in $Z(N)$, then $f \cdot (S, \mathbf{1}) = (S, \mathbf{1})$, so that $Z_s^1(G, Z(N))_S$ corresponds to the stabilizer of $(S, \mathbf{1}) \in Z_{ss}^2(G, N)$ and the map

$$Z_s^1(G, N, Z(N))_S \rightarrow Z_{ss}^2(G, N), \quad f \mapsto f \cdot (S, \mathbf{1})$$

factors through an injective map

$$Z_s^1(G, N, Z(N))_S / Z_s^1(G, Z(N))_S \cong Z_s^1(G, N_{\text{ad}}), \quad [f] \mapsto f \cdot (S, \mathbf{1})$$

because for each $h \in Z_s^1(G, N_{\text{ad}})_S$ there is an $\widehat{h} \in Z_s^1(G, N, Z(N))_S$ with $q_N \circ \widehat{h} = h$ and $\widehat{h} \cdot (S, \mathbf{1}) = (\widehat{h} \cdot S, d_S \widehat{h})$.

Since $d_S \widehat{h} \in Z_{ss}^2(G, Z(N))_S$, this factor system corresponds to the extension class

$$[d_S \widehat{h}].[N \rtimes_S G] = \delta(h).[N \rtimes_S G] \in \text{Ext}(G, N)_{[S]}.$$

This class equals $[N \rtimes_S G]$ if and only if $\delta(h) = 0$ which is equivalent to $[h]$ being contained in the image of $H_s^1(G, N)_S$ under the connecting map δ .

The remaining assertion follows from Remark 2.20(b) and Proposition 2.18. □

In [2], Borovoi deals with non-abelian H^2 -sets in the context of Galois cohomology, where he calls the cohomology classes in $H_{ss}^2(G, N)$ corresponding to split extensions *neutral*. Our Corollary 2.21 is a Lie group version of Borovoi’s Proposition 2.3.

3. Smooth crossed modules

In this section we introduce the notion of a smooth crossed module for a Lie group G . Our point of view is that a smooth crossed module is a central extension $\widehat{N} \rightarrow N$ of a normal split Lie subgroup $N \trianglelefteq G$ for which the conjugation action of G on N lifts to a smooth action on \widehat{N} . It turns out that smooth crossed modules of Lie groups provide the natural framework to reduce problems related to general extensions of Lie groups to abelian extensions, which is carried out in Section 4.

DEFINITION 3.1. — *A morphism $\alpha: H \rightarrow G$ of Lie groups, together with a homomorphism $\widehat{S}: G \rightarrow \text{Aut}(H)$ defining a smooth action $\widehat{S}: G \times H \rightarrow H, (g, h) \mapsto g.h = \widehat{S}(g)(h)$ of G on H , is called a smooth crossed module if the following conditions are satisfied:*

- (CM1) $\alpha \circ \widehat{S}(g) = c_{\alpha(g)} \circ \alpha$ for $g \in G$.
- (CM2) $\widehat{S} \circ \alpha = C_H: H \rightarrow \text{Aut}(H)$ is the conjugation action.
- (CM3) $\ker(\alpha)$ is a split Lie subgroup of H and $\text{im}(\alpha)$ is a split Lie subgroup of G for which α induces an isomorphism $H/\ker(\alpha) \rightarrow \text{im}(\alpha)$.

The conditions (CM1/2) express the compatibility of the G -action on H with the conjugation actions of G and H .

LEMMA 3.2. — *If $\alpha: H \rightarrow G$ is a smooth crossed module, then the following assertions hold:*

- (1) $\text{im}(\alpha)$ is a normal subgroup of G .
- (2) $\ker(\alpha) \subseteq Z(H)$.
- (3) $\ker(\alpha)$ is G -invariant and G acts smoothly on $\ker(\alpha)$.

Proof.

(1) follows from (CM1), (2) from (CM2), and (3) from (CM1), Lemma 1.7(a) and Lemma 2.3. □

Crossed modules for which α is injective are inclusions of normal split Lie subgroups and surjective crossed modules are central extensions. In this sense the concept of a crossed module interpolates between split normal subgroups and central extensions.

In the following we shall adopt the following perspective on crossed modules. Let $\alpha: H \rightarrow G$ be a smooth crossed module. Then $N := \text{im}(\alpha)$ is a split normal Lie subgroup of G and $\alpha: H \rightarrow N$ is a central Lie group extension of N by $Z := \ker(\alpha)$. In this sense a smooth crossed module can be viewed as a central Lie group extension $\alpha: H \rightarrow N$ of a split normal subgroup N of G for which there exists a smooth G -action on H satisfying (CM1/2).

If, conversely, N is a split normal subgroup of G and $\alpha: \widehat{N} \rightarrow N$ a central Lie group extension of N by Z , then we have a natural smooth N -module structure on \widehat{N} given by $n.n' := \widehat{C}_N(n)(n') := nn'n^{-1}$. To obtain the structure of a smooth crossed module for $\alpha: \widehat{N} \rightarrow G$, one has to extend the action of N on \widehat{N} to a smooth G -action $G \times \widehat{N} \rightarrow \widehat{N}$, $(g, n') \mapsto g.n'$ for which α is equivariant.

From now on we assume that this data is given. We may w.l.o.g. assume that $\widehat{N} = Z \times_f N$ for some $f \in Z_{ss}^2(N, Z)$ (Remark 2.14) and write the G -module structure on Z as $(g, z) \mapsto g.z$. Then the G -module structure on $\widehat{N} = Z \times_f N$ can be written as

$$(3.1) \quad g.(z, n) = (g.z + \theta(g)(g.n), g.n),$$

where the function $\theta: G \rightarrow C_s^1(N, Z)$ is a map for which $\tilde{\theta}: G \times N \rightarrow Z, (g, n) \mapsto \theta(g)(n)$ is smooth on an open neighborhood of $G \times \{\mathbf{1}\}$ in $G \times N$. That (3.1) defines a group action of G on \widehat{N} means that θ is a cocycle with respect to the natural action of G on $C_s^1(N, Z)$ by $(g.\alpha)(n) := g.\alpha(g^{-1}ng)$, and that with respect to the natural action of G on $C_s^2(N, Z)$ by $(g.\beta)(n_1, n_2) := g.\beta(g^{-1}n_1g, g^{-1}n_2g)$ and we have with respect to the trivial action of N on Z :

$$(3.2) \quad d_N(\theta(g)) = g.f - f = d_S(f)(g) \text{ for all } g \in G.$$

(cf. Corollary A.4 and Lemma A.11).

Therefore the data we are now working with is given by $f \in Z_{ss}^2(N, Z)$ with $\widehat{N} = Z \times_f N$ and $\theta \in Z_{ss}^1(G, C_s^1(N, Z))_f$ for which the corresponding smooth action \widehat{S} of G on \widehat{N} turns the homomorphism $\alpha: \widehat{N} \rightarrow G, (z, n) \mapsto n$ into a smooth crossed module.

LEMMA 3.3. — *For a cocycle $\theta: G \rightarrow C_s^1(N, Z)$ satisfying (3.2), the following are equivalent:*

- (1) *The corresponding action of G on \widehat{N} is smooth.*
- (2) *The function $\widetilde{\theta}: G \times N \rightarrow Z, (g, n) \mapsto \theta(g)(n)$ is smooth in an identity neighborhood and for each $n \in N$ the function*

$$(3.3) \quad \theta_n: G \rightarrow Z, \quad g \mapsto \theta(g)(g.n) + f(g.n, n^{-1})$$

is smooth in an identity neighborhood of G .

Proof. — This is an immediate consequence of Proposition A.13. □

DEFINITION 3.4. — *In the following we write $C_s^1(G, C_s^1(N, Z))$ for the set of all maps $\alpha: G \rightarrow C_s^1(N, Z)$ for which the function $\widetilde{\alpha}: G \times N \rightarrow Z, (g, n) \mapsto \alpha(g)(n)$ is smooth in an identity neighborhood. We write $C_{ss}^1(G, C_s^1(N, Z))_f$ for the set of all elements with the additional property that for each $n \in N$ the function*

$$\alpha_n: G \rightarrow Z, \quad g \mapsto \alpha(g)(g.n) + f(g.n, n^{-1})$$

is smooth in an identity neighborhood. The set of all cocycles in $C_s^1(G, C_s^1(N, Z))$ is denoted $Z_s^1(G, C_s^1(N, Z))$ and likewise

$$Z_{ss}^1(G, C_s^1(N, Z))_f := C_{ss}^1(G, C_s^1(N, Z))_f \cap Z_s^1(G, C_s^1(N, Z)).$$

The natural action of N on \widehat{N} is given by

$$n.(z', n') := (z, n)(z', n')(z, n)^{-1} = (z' + f(n, n') - f(nn'n^{-1}, n), nn'n^{-1}),$$

where we note that the right hand side does not depend on $z \in Z(\widehat{N})$. We define

$$(3.4) \quad \widetilde{f}(n)(n') := f(n', n) - f(n, n^{-1}n'n),$$

and observe that (3.1) defines a smooth action extending the given action of N if $\theta|_N = \widetilde{f}$, i.e., that the pair (f, θ) is an element of the group

$$D := \{(f, \theta) \in Z_{ss}^2(N, Z) \times Z_s^1(G, C_s^1(N, Z)) : d_N \circ \theta = d_S(f), \theta|_N = \widetilde{f}, \\ \theta \in C_{ss}^1(G, C_s^1(N, Z))_f\}.$$

That the set of all these elements is indeed a subgroup follows directly from Lemma 3.3. We also write

$$D(f) := \{\theta \in Z_{ss}^1(G, C_s^1(N, Z))_f : (f, \theta) \in D\} = \text{pr}_2(D)$$

for the set of all smooth actions of G on \widehat{N} extending the given action of N .

The following lemma provides a convenient description of the set $D(f)$ as a homogeneous space.

LEMMA 3.5. — Let $q: G \rightarrow G/N$ denote the quotient map, $\widehat{N} = Z \times_f N$ a central Lie group extension of N by Z and a smooth G -action on \widehat{N} turning $\alpha: \widehat{N} \rightarrow N \subseteq G$ into a smooth crossed module. For any $(f, \theta) \in D$ and $\beta \in Z_{ss}^1(G/N, Z_s^1(N, Z))_0$ we have

$$\beta * (f, \theta) := (f, \theta + q^* \beta) \in D.$$

This defines for each $f \in \text{pr}_1(D) \subseteq Z_{ss}^2(N, Z)$ a simply transitive action of $Z_{ss}^1(G/N, Z_s^1(N, Z))_0$ on the set $D(f)$.

Proof. — That $\beta * (f, \theta) \in D$ follows from

$$d_N \circ (\theta + q^* \beta) = d_N \circ \theta + d_N \circ q^* \beta = d_N \circ \theta + q^*(d_N \circ \beta) = d_N \circ \theta = d_S(f).$$

If, conversely, $(f, \theta), (f, \theta') \in D$, then the difference $\theta' - \theta$ has values in $Z_s^1(N, Z)$ and vanishes on N , hence is of the form $q^* \beta$ for some $\beta \in Z_{ss}^1(G/N, Z_s^1(N, Z))_0$ because for each $n \in N$ the map

$$g \mapsto (\theta' - \theta)(g)(g.n) = \theta'(g)(g.n) + f(g.n, n^{-1}) - (\theta(g)(g.n) - f(g.n, n^{-1}))$$

is smooth in an identity neighborhood of G . This proves the lemma. \square

We want to construct a map $Q: D \rightarrow H_s^3(G/N, Z)$ such that $Q(f, \theta) = 0$ characterizes the extendability of f to a cocycle $\tilde{f} \in Z_s^2(G, Z)$ (where we assume that we are given a smooth action of G on Z) satisfying $\tilde{f}|_{G \times N} = \tilde{\theta}$. In the following we fix a pair $(f, \theta) \in D$ and consider the corresponding smooth action of G on \widehat{N} given by

$$\widehat{S}(g)(z, n) := (g.z + \theta(g)(g.n), g.n)$$

(cf. Corollary A.4 and Proposition A.13) extending the given action on Z . To define the map Q , let $\sigma: G/N \rightarrow G$ be a normalized locally smooth section of the quotient map and define

$$S := \widehat{S} \circ \sigma: G/N \rightarrow \text{Aut}(\widehat{N}).$$

There exists an $\omega \in C_{ss}^2(G/N, \widehat{N})$ with $q_N \circ \omega = \delta_\sigma$, where $q_N: \widehat{N} \rightarrow N$ is the quotient map. Then

$$(3.5) \quad \delta_S = \widehat{S} \circ \delta_\sigma = \widehat{S} \circ q_N \circ \omega = C_{\widehat{N}} \circ \omega,$$

which shows that S is a smooth outer action of G/N on \widehat{N} .

If we even find an $\omega \in C_{ss}^2(G/N, \widehat{N})$ with $\delta_S = C_N \circ \omega$, we have a strongly smooth outer action, but this does not follow from the definition

of a smooth crossed module. Using Lemma 1.10, it is easy to see that this additional smoothness condition for S does not depend on the choice of the section σ , hence is a property of the pair (f, θ) . We write $D_{ss} \subseteq D$ for the set of all pairs $(f, \theta) \in D$ for which S is a strongly smooth outer action. In terms of the product coordinates $\widehat{N} = Z \times_f N$, this means that there exists $\omega_Z \in C_{ss}^2(G/N, Z)$ such for $\omega := (\omega_Z, \delta_\sigma) \in C_s^2(G/N, \widehat{N})$ the functions $\omega_g: G/N \rightarrow \widehat{N}$ given by

$$\begin{aligned} \omega_g(x) &= \omega(g, x)\omega(gxg^{-1}, g)^{-1} \\ &= (\omega_Z(g, x), \delta_\sigma(g, x))(\omega_Z(gxg^{-1}, g), \delta_\sigma(gxg^{-1}, g))^{-1} \\ &= (\omega_Z(g, x), \delta_\sigma(g, x))(-\omega_Z(gxg^{-1}, g) \\ &\quad - f(\delta_\sigma(gxg^{-1}, g)^{-1}, \delta_\sigma(gxg^{-1}, g)), \delta_\sigma(gxg^{-1}, g)^{-1}) \\ &= (\omega_{Z,g}(x) - f(\delta_\sigma(gxg^{-1}, g)^{-1}, \delta_\sigma(gxg^{-1}, g)) \\ &\quad + f(\delta_\sigma(g, x), \delta_\sigma(gxg^{-1}, g)^{-1}), \delta_{\sigma,g}(x)) \end{aligned}$$

are smooth in an identity neighborhood. For $\delta_{\sigma,g}$ this follows as in Definition 1.1, but for the Z -component this depends on the values of the section σ in elements far from the identity, so that the smoothness assumptions in the identity do not imply anything.

LEMMA 3.6. — *The function*

$$\begin{aligned} \widetilde{Q}(f, \theta, \sigma, \omega) &:= d_S\omega: (G/N)^3 \rightarrow \widehat{N}, \quad (g, g', g'') \\ &\mapsto S(g)(\omega(g', g''))\omega(g, g'g'')\omega(gg', g'')^{-1}\omega(g, g')^{-1} \end{aligned}$$

has values in Z and is a 3-cocycle in $Z_s^3(G/N, Z)$. Its cohomology class $[d_S\omega] \in H_s^3(G/N, Z)$ does not depend on the choices of ω and σ . Likewise, for $(f, \theta) \in D_{ss}$ the cohomology class of $d_S\omega$ in $H_{ss}^3(G/N, Z)$ does not depend on ω and σ .

Proof. — First we observe that $(d_S\omega)(g, g', g'') \in Z$ follows from

$$\begin{aligned} q_N((d_S\omega)(g, g', g'')) &= C_N(\sigma(g))(\delta_\sigma(g', g''))\delta_\sigma(g, g'g'')\delta_\sigma(gg', g'')^{-1}\delta_\sigma(g, g')^{-1} \\ &= \sigma(g)(\sigma(g')\sigma(g'')\sigma(g'g'')^{-1})\sigma(g)^{-1}\sigma(g)\sigma(g'g'')\sigma(gg'g'')^{-1} \\ &\quad \sigma(gg'g'')\sigma(g'')^{-1}\sigma(gg')^{-1}\sigma(gg')\sigma(g')^{-1}\sigma(g)^{-1} = \mathbf{1}. \end{aligned}$$

It is easy to verify that $d_S\omega$ vanishes if one of its three arguments is $\mathbf{1}$, so that $d_S\omega \in C_s^3(G/N, Z)$. Since S is a smooth outer action of G/N on \widehat{N} and $\delta_S = C_{\widehat{N}} \circ \omega$, Lemma 1.10(5) implies that $d_S\omega \in Z_s^3(G/N, Z(\widehat{N}))$. Since its values lie in the subgroup Z of $Z(\widehat{N})$, we have $d_S\omega \in Z_s^3(G/N, Z)$.

It remains to show that the cohomology class of $[d_S\omega]$ in $H_s^3(G/N, Z)$, resp. $H_{ss}^3(G/N, Z)$ does not depend on the choices of ω and σ . We only discuss the case $(\theta, f) \in D_{ss}$. First we show that it does not depend on the choice of ω if σ is fixed. So let $\tilde{\omega} \in C_{ss}^2(G/N, \hat{N})$ be another map with $q_N \circ \tilde{\omega} = \delta_\sigma$. Then there exists a function $\beta \in C_{ss}^2(G/N, Z)$ with $\tilde{\omega} = \beta \cdot \omega$. As all values of β commute with all values of ω , this leads to

$$d_S\tilde{\omega} = d_S\beta + d_S\omega = d_{G/N}\beta + d_S\omega \in d_S\omega + d_{G/N}(C_{ss}^2(G/N, Z)).$$

Therefore the choice of ω for a fixed section σ has no effect on the cohomology class $[d_S\omega]$ in $H_{ss}^3(G/N, Z)$.

Now we consider another section $\tilde{\sigma} \in C_s^1(G/N, G)$ of q . Then there exists a function $h \in C_s^1(G/N, N)$ with $\tilde{\sigma} = h \cdot \sigma$ and

$$\begin{aligned} \delta_{\tilde{\sigma}}(g, g') &= h(g)\sigma(g)h(g')\sigma(g')(h(gg')\sigma(gg'))^{-1} \\ &= h(g)c_{\sigma(g)}(h(g'))\delta_\sigma(g, g')h(gg')^{-1}. \end{aligned}$$

Let $\hat{h} \in C_s^1(G/N, \hat{N})$ with $q_N \circ \hat{h} = h$. Then the prescription

$$\tilde{\omega}(g, g') = \hat{h}(g)S(g)(\hat{h}(g'))\omega(g, g')\hat{h}(gg')^{-1} = (\hat{h} *_S \omega)(g, g')$$

defines an element $\tilde{\omega} \in C_{ss}^2(G/N, \hat{N})$ for which $\tilde{S} := \hat{S} \circ \tilde{\sigma} = (C_{\hat{N}} \circ \hat{h}) \cdot S$ satisfies $\delta_{\tilde{S}} = C_{\hat{N}} \circ \tilde{\omega}$ (Lemma 1.10(1)), and we conclude from Lemma 1.10(4) that $d_{\tilde{S}}\tilde{\omega} = d_S\omega$. □

The preceding lemma implies that the cohomology class

$$Q(f, \theta) := [\tilde{Q}(f, \theta, \sigma, \omega)] = [d_S\omega]$$

in $H_s^3(G/N, Z)$ only depends on the pair (f, θ) , hence defines a map $Q: D \rightarrow H_s^3(G/N, Z)$. In the strongly smooth case we obtain a map $Q_{ss}: D_{ss} \rightarrow H_{ss}^3(G/N, Z)$.

LEMMA 3.7. — *The maps $Q: D \rightarrow H_s^3(G/N, Z)$ and $Q_{ss}: D_{ss} \rightarrow H_{ss}^3(G/N, Z)$ are group homomorphisms.*

Proof. — Again we consider only the case $(\theta, f) \in D_{ss}$. We choose $\omega \in C_{ss}^2(G/N, \hat{N})$ with $q_N \circ \omega = \delta_\sigma$. We have to see how $Q(f, \theta, \sigma, \omega)$ depends on (f, θ) . Writing $\hat{N} := Z \times_f N$, any ω with $q_N \circ \omega = \delta_\sigma$ can be written as $\omega = (\omega_Z, \delta_\sigma)$, independently of f and θ .

Writing $g.n = gng^{-1}$ for the conjugation action of G on N , we now obtain

$$\begin{aligned} S(g).\omega(g', g'') &= \hat{S}(\sigma(g))(\omega_Z(g', g''), \delta_\sigma(g', g'')) \\ &= (g.\omega_Z(g', g'') + \theta(\sigma(g))(\sigma(g).\delta_\sigma(g', g'')), \sigma(g).\delta_\sigma(g', g'')). \end{aligned}$$

Therefore

$$\begin{aligned} & (S(g).\omega(g', g''))\omega(g, g'g'') \\ &= \left(g.\omega_Z(g', g'') + \omega_Z(g, g'g'') + \theta(\sigma(g))(\sigma(g).\delta_\sigma(g', g'')) \right. \\ & \quad \left. + f(\sigma(g).\delta_\sigma(g', g''), \delta_\sigma(g, g'g'')), (\sigma(g).\delta_\sigma(g', g''))\delta_\sigma(g, g'g'') \right), \end{aligned}$$

and similarly

$$\begin{aligned} \omega(g, g')\omega(gg', g'') &= (\omega_Z(g, g') + \omega_Z(gg', g'')) \\ & \quad + f(\delta_\sigma(g, g'), \delta_\sigma(gg', g'')), \delta_\sigma(g, g')\delta_\sigma(gg', g'')). \end{aligned}$$

Since $\text{im}(d_S\omega) \subseteq Z$, this implies that

$$\begin{aligned} (d_S\omega)(g, g', g'') &= (d_S\omega_Z)(g, g', g'') + \theta(\sigma(g))(\sigma(g).\delta_\sigma(g', g'')) \\ & \quad + f(\sigma(g).\delta_\sigma(g', g''), \delta_\sigma(g, g'g'')) - f(\delta_\sigma(g, g'), \delta_\sigma(gg', g'')). \end{aligned}$$

Hence the cohomology class of $d_S\omega$ depends additively on the pair (f, θ) , i.e., Q_{ss} is a group homomorphism $D_{ss} \rightarrow H_{ss}^3(G/N, Z)$. \square

The following theorem explains the meaning of the condition $Q(f, \theta) = \mathbf{1}$. It is the main result of the present section and establishes an important link between smooth crossed modules and the extendability of central extensions of normal subgroups to the whole group.

THEOREM 3.8. — *Let $f \in Z_{ss}^2(N, Z)$ with $\widehat{N} = Z \times_f N$ and $\theta \in Z_{ss}^1(G, C_s^1(N, Z))_f$ describe a smooth action \widehat{S} of G on \widehat{N} such that the homomorphism $\alpha: \widehat{N} \rightarrow G, (z, n) \mapsto n$ is a smooth crossed module. Further choose a locally smooth section $\sigma \in C_s^1(G/N, G)$ of $q: G \rightarrow G/N$ and $\omega \in C_{ss}^2(G/N, \widehat{N})$ with $\delta_\sigma = q_N \circ \omega$, and $S = \widehat{S} \circ \sigma$. Then the following are equivalent:*

- (1) $Q_{ss}(f, \theta) = [d_S\omega] = \mathbf{1}$ in $H_{ss}^3(G/N, Z)$.
- (2) There exists a Lie group extension $Z \hookrightarrow \widehat{G} \xrightarrow{q} G$ and a G -equivariant equivalence $\widehat{N} \rightarrow q^{-1}(N)$ of Z -extensions of N .
- (3) There exists a cocycle $f_G \in Z_{ss}^2(G, Z)$ with $f_G|_{N \times N} = f$ and $\widehat{f}_G|_{G \times N} = \theta$.

Proof.

(1) \Rightarrow (2): Suppose that $[d_S\omega] = \mathbf{1}$, i.e., $d_S\omega = d_S\beta$ for some $\beta \in C_{ss}^2(G/N, Z)$. Then $\omega' := \omega \cdot \beta^{-1}$ satisfies $q_N \circ \omega' = \delta_\sigma$ and $d_S\omega' = d_S\omega - d_S\beta = 0$. Since $\omega' \in C_{ss}^2(G/N, \widehat{N})$ follows from $\beta \in C_{ss}^2(G/N, Z)$, the pair (S, ω') is a smooth factor system for G/N and \widehat{N} , so that we obtain a group

$\widehat{G} = \widehat{N} \times_{(S, \omega')} G/N$ with the product

$$(n, g)(n', g') := (nS(g)(n')\omega'(g, g'), gg')$$

and a natural Lie group structure (Proposition 2.8). Then \widehat{G} is a Lie group extension of G/N by \widehat{N} . On the other hand, we can view $(C_N \circ \sigma, \delta_\sigma)$ as a smooth factor system for G/N and N , which leads to the description of G as $N \times (G/N)$, endowed with the product

$$(n, g)(n', g') := (nC_N(\sigma(g))(n')\delta_\sigma(g, g'), gg').$$

Hence the map

$$\begin{aligned} q_G: \widehat{G} = \widehat{N} \times (G/N) &= (Z \times_f N) \times (G/N) \\ &\rightarrow G = N \times (G/N), \quad ((z, n), g) \mapsto (n, g) \end{aligned}$$

defines a Lie group extension of G by Z containing \widehat{N} . It remains to verify that the action of G on \widehat{N} induced by the conjugation action of \widehat{G} , for which Z acts trivially, coincides with the action given by θ .

It follows from the construction, that the conjugation action of \widehat{G} on $\widehat{N} = Z \times_f N$ is given by

$$\begin{aligned} (n, g)(z, n')(n, g)^{-1} &= c_{\widehat{N}}(n)S(g)(z, n') \\ &= c_{\widehat{N}}(n)(g.z + \theta(g)(C_N(g)(n')), C_N(g)(n')) \\ &= (g.z + \theta(g)(C_N(g)(n')) + \theta(n)(C_N(n)C_N(g)(n')), C_N(n)C_N(g)(n')) \\ &= (g.z + \theta((n, g))(C_N(n)C_N(g)(n')), C_N((n, g))(n')) \\ &= (n, g).(z, n'). \end{aligned}$$

This completes the proof of (2).

(2) \Rightarrow (3): Let $q: \widehat{G} \rightarrow G$ be an extension as in (2). We may w.l.o.g. assume that $\widehat{G} = Z \times_{f_G} G$ for some $f_G \in Z_{ss}^2(G, Z)$, where f_G extends f , i.e., $\widehat{N} = Z \times_f N$ is contained in \widehat{G} . The condition that the inclusion $\widehat{N} \hookrightarrow \widehat{G}$ is G -equivariant means that $\widetilde{f}_G|_{G \times N} = \theta$.

(3) \Rightarrow (1): Let $f_G \in Z_{ss}^2(G, Z)$ be a cocycle as in (3) and

$$q_G: \widehat{G} := Z \times_{f_G} G \rightarrow G$$

the corresponding Lie group extension. The condition $f_G|_{N \times N} = f$ means that the group $\widehat{N} = Z \times_f N$ is a split Lie subgroup of \widehat{G} , and the second condition $\widetilde{f}_G|_{G \times N} = \theta$ ensures that the conjugation action $C_{\widehat{N}}$ of \widehat{G} on \widehat{N} induces the action of G on \widehat{N} defined by θ .

We consider the locally smooth normalized section $\widehat{\sigma}: G/N \rightarrow \widehat{G}, g \mapsto (0, \sigma(g))$. Then $\delta_{\widehat{\sigma}}$ satisfies $q_N \circ \delta_{\widehat{\sigma}} = \delta_{\sigma}$. In view of $d_S \omega = \mathbf{1}$ and $S = C_{\widehat{N}} \circ \widehat{\sigma}$, we now have

$$Q(f, \theta, \sigma, \delta_{\widehat{\sigma}}) = d_S \delta_{\widehat{\sigma}} = \mathbf{1}.$$

It follows in particular that $Q(f, \theta) = [d_S \delta_{\widehat{\sigma}}] = \mathbf{1}$ (Lemma 3.6). □

Remark 3.9. — Suppose that G is connected and that $(f, \theta) \in D$. If $Q(f, \theta) = \mathbf{1}$ in $H_s^3(G/N, Z)$, then the same argument as in the proof of Theorem 3.8 provides $\omega' \in C_s^2(G/N, \widehat{N})$ with $\delta_S = C_{\widehat{N}} \circ \omega'$ and $d_S \omega' = \mathbf{1}$. Now Lemma 2.9 implies that $\omega' \in C_{ss}^2(G/N, \widehat{N})$ and that (2) still holds. All other implications do not require the strong smoothness of S . Hence Theorem 3.8 remains true for connected groups G if we replace the condition in (1) by the weaker requirement $Q(f, \theta) = \mathbf{1}$ in $H_s^3(G/N, Z)$.

4. Applications to non-abelian extensions of Lie groups

Let G and N be Lie groups, and assume that $Z(N)$ is a split Lie subgroup of N , so that $N_{\text{ad}} := N/Z(N)$ carries a natural Lie group structure. Further let $S \in C_s^1(G, \text{Aut}(N))$ be a strongly smooth outer action. In this section we describe the set $\text{Ext}(G, N)_{[S]}$ of N -extensions of G corresponding to $[S]$ in terms of abelian Lie group extensions. The main point is that S defines a Lie group extension

$$\mathbf{1} \rightarrow N_{\text{ad}} := N/Z(N) \rightarrow G^S \xrightarrow{qs} G \rightarrow \mathbf{1}$$

and that for each N -extension \widehat{G} of G corresponding to S , we have $\widehat{G}/Z(N) \cong G^S$, so that we may consider \widehat{G} as an extension of G^S by $Z(N)$. This relation reduces several problems concerning general Lie group extensions to the abelian case for which we refer to [19].

Pick $\omega \in C_{ss}^2(G, N)$ with $\delta_S = C_N \circ \omega$. Since $N_{\text{ad}} \cong C_N(N) \subseteq \text{Aut}(N)$, we write the elements of this group as $C_N(n)$ and not as cosets $nZ(N)$. On the product set

$$G^S := N_{\text{ad}} \times G$$

we define the product

$$\begin{aligned} (C_N(n), g)(C_N(n'), g') &:= \left(C_N(nS(g)(n')\omega(g, g')), gg' \right) \\ &= \left(C_N(n)c_{S(g)}(C_N(n'))\delta_S(g, g'), gg' \right). \end{aligned}$$

Note that the second form of the product shows in particular that it does not depend on ω . We observe that

$$S_1: G \rightarrow \text{Aut}(N_{\text{ad}}), \quad S_1(g)(C_N(n)) = C_N(S(g).n)$$

is a smooth outer action with $\delta_{S_1} = C_{N_{\text{ad}}} \circ \delta_S$ if we consider δ_S as N_{ad} -valued function and G acts on $N_{\text{ad}} \cong C_N(N)$ by conjugation.

LEMMA 4.1. — *If $Z(N)$ is a split Lie subgroup of N , then G^S is a group with the following properties:*

- (1) *It carries a natural Lie group structure for which $q_S: G^S \rightarrow G, (C_N(n), g) \mapsto g$ is a Lie group extension of G by N_{ad} . The equivalence class of this extension only depends on the G -kernel $[S]$.*
- (2) *The map $\rho: G^S \rightarrow \text{Aut}(N), (C_N(n), g) \mapsto C_N(n)S(g)$ defines on N a smooth G^S -action.*
- (3) *The map $\alpha: N \rightarrow G^S, n \mapsto (C_N(n), \mathbf{1})$ defines a smooth crossed module with*

$$\ker \alpha = Z(N) \text{ and } \text{coker}(\alpha) = G^S/N_{\text{ad}} \cong G,$$

and S is the corresponding strongly smooth outer action of $G \cong G^S/N_{\text{ad}}$ on N .

Proof.

(1) First we have to show that (S_1, δ_S) is a smooth factor system for G and N_{ad} : Since we define the Lie group structure on $N_{\text{ad}} = C_N(N)$ in such a way that $C_N: N \rightarrow N_{\text{ad}}$ is the quotient map, the relation $\delta_S = C_N \circ \omega$ implies $\delta_S \in C_{ss}^2(G, N_{\text{ad}})$, and we have $\delta_{S_1} = C_{N_{\text{ad}}} \circ \delta_S$.

If $U \subseteq G$ is an open identity neighborhood such that the map $U \times N \rightarrow N, (g, n) \mapsto S(g).n$ is smooth, then the map

$$\begin{aligned} U \times N_{\text{ad}} &\rightarrow N_{\text{ad}}, \quad (g, C_N(n)) \mapsto C_N(S(g).n) = S(g)C_N(n)S(g)^{-1} \\ &= S_1(g)(C_N(n)) \end{aligned}$$

is also smooth.

The local smoothness of $\delta_S = C_N \circ \omega$ follows from the local smoothness of ω . Moreover,

$$\begin{aligned} \delta_{S,g}(g') &= \delta_S(g, g')\delta_S(gg'g^{-1}, g)^{-1} = C_N(\omega(g, g')\omega(gg'g^{-1}, g)^{-1}) \\ &= C_N(\omega_g(g')) \end{aligned}$$

is smooth in an identity neighborhood because $\omega \in C_{ss}^2(G, N)$.

For $g, g', g'' \in G$ we have

$$\begin{aligned}
& S_1(g)(\delta_S(g', g''))\delta_S(g, g'g'') \\
&= S(g)\left(S(g')S(g'')S(g'g'')^{-1}\right)S(g)^{-1}S(g)S(g'g'')S(gg'g'')^{-1} \\
&= S(g)S(g')S(g'')S(gg'g'')^{-1} \\
&= S(g)S(g')S(gg')^{-1}S(gg')S(g'')S(gg'g'')^{-1} \\
&= \delta_S(g, g')\delta_S(gg', g'').
\end{aligned}$$

This shows that (S_1, δ_S) is a smooth factor system, so that G^S is a group and Proposition 2.8 provides a natural Lie group structure for which $q_S: G^S \rightarrow G$ is a Lie group extension of G by N_{ad} .

To see that the equivalence class of the extension $G^S = N_{\text{ad}} \times_{(S_1, \delta_S)} G$ of G by N_{ad} depends only on the equivalence class $[S]$, let $\alpha \in C_s^1(G, N)$ and $S' := (C_N \circ \alpha) \cdot S$. Then

$$S'_1 := (C_{N_{\text{ad}}} \circ h) \cdot S_1 = h \cdot S_1 \text{ for } h := C_N \circ \alpha \in C_s^1(G, N_{\text{ad}}),$$

and formula (1.2) after Definition 1.1 leads to

$$\delta_{S'}(g, g') = C_N\left(\alpha(g)S(g)(\alpha(g'))\right)\delta_S(g, g')C_N(\alpha(gg')^{-1}) = (h *_{S_1} \delta_S)(g, g').$$

According to Proposition 2.10, the map

$$\psi: G^{S'} \rightarrow G^S, \quad (C_N(n), g) \mapsto (C_N(n\alpha(g)), g) = (C_N(n)h(g), g)$$

is an equivalence of extensions of G by N_{ad} .

(2) First we show that ρ is a representation:

$$\begin{aligned}
& \rho(C_N(n), g)\rho(C_N(n'), g') \\
&= C_N(n)S(g)C_N(n')S(g') = C_N(n)C_N(S(g).n')S(g)S(g') \\
&= C_N(n(S(g).n'))\delta_S(g, g')S(gg') \\
&= \rho\left(C_N(n(S(g).n'))\delta_S(g, g'), gg'\right) = \rho((C_N(n), g)(C_N(n'), g')).
\end{aligned}$$

Since $\rho(G^S)$ consists of smooth automorphisms of N , the smoothness of the representation follows from the local smoothness and the smoothness of the orbit maps; local smoothness is a consequence of the local smoothness of the map

$$(N_{\text{ad}} \times G) \times N \rightarrow N, \quad ((C_N(n), S(g)), n') \rightarrow C_N(n)S(g).n'$$

which, in addition, shows that the action of G^S on N is a map $G^S \times N \rightarrow N$ that is smooth on a set of the form $U^S \times N$, where U^S is an identity neighborhood in G^S . Since G^S acts on N by smooth maps, this implies the smoothness of the action.

(3) is an immediate consequence of (1) and (2). □

Remark 4.2. — If the group G is connected, then we can use Lemma 2.9, instead of Proposition 2.8, in the preceding proof and draw the same conclusions if S is not strongly smooth.

LEMMA 4.3. — *The map*

$$\psi = (\rho, q_S): G^S \rightarrow \text{Aut}(N) \times G$$

is injective and yields an isomorphism of groups

$$G^S \cong \{(\varphi, g) \in \text{Aut}(N) \times G: S(g) \in \varphi \cdot C_N(N)\}.$$

Proof. — Since $\ker q_S = N_{\text{ad}}$ and $(\ker \rho) \cap N_{\text{ad}} = \{\mathbf{1}\}$, the map ψ is an injective homomorphism of groups.

For each element $(C_N(n), g) \in G^S$ we have

$$\psi(C_N(n), g) = (C_N(n)S(g), g),$$

which proves “ \subseteq ,” and for any pair $(\varphi, g) \in \text{Aut}(N) \times G$ with $\varphi \in S(g) \cdot C_N(N)$ we find an element $n \in N$ with $\varphi = C_N(n)S(g)$, which means that $(\varphi, g) = \psi(C_N(n), g)$. This proves the lemma. □

PROPOSITION 4.4. — *Let $q: \widehat{G} \rightarrow G$ be a Lie group extension of G by N , corresponding to the smooth G -kernel $[S]$, and C_N the corresponding representation of \widehat{G} on N . Assume further that $Z(N)$ is a split Lie subgroup of N . Then $\widehat{G}/Z(N) \cong G^S$ and the map*

$$\gamma = (C_N \cdot (S \circ q)^{-1}, q): \widehat{G} \rightarrow \text{Aut}(N) \times G, \quad g \mapsto (C_N(g)S(q(g))^{-1}, q(g))$$

defines a Lie group extension

$$Z(N) \hookrightarrow \widehat{G} \xrightarrow{\gamma} G^S.$$

This assignment has the following properties:

(1) *If $q_j: \widehat{G}_j \rightarrow G, j = 1, 2$, are equivalent extensions of G by N , then $\gamma_j: \widehat{G}_j \rightarrow G^S$ are equivalent extensions of G^S by the smooth G^S -module $Z(N)$. We thus obtain a map*

$$\tau: \text{Ext}(G, N)_{[S]} \rightarrow \text{Ext}(G^S, Z(N))_{[\rho]}.$$

(2) *A Lie group extension $\gamma: \widehat{G}^S \rightarrow G^S$ of G^S by $Z(N)$ comes from an extension of G by N corresponding to $[S]$ if and only if there exists a G^S -equivariant equivalence*

$$\alpha: N \rightarrow \gamma^{-1}(N_{\text{ad}}) \subseteq \widehat{G}^S$$

of central extensions of N_{ad} by $Z(N)$. Note that $G^S \cong \widehat{G}^S/Z(N)$ acts on $\gamma^{-1}(N_{\text{ad}})$ because $Z(N)$ acts trivially on this group.

Proof. — Since the extension \widehat{G} corresponds to $[S]$, it is equivalent to an extension of the type $N \times_{(S,\omega)} G$ with $q(n, g) = g$, where (S, ω) is a smooth factor system (Proposition 2.8). This means that

$$(4.1) \quad C_N(n, g) = C_N(n)S(g) = C_N(n)S(q(n, g)),$$

so that $\gamma(n, g) = (C_N(n), g)$. Now the explicit formulas for the multiplications in $N \times_{(S,\omega)} G$ and G^S imply that γ is a surjective morphism of Lie groups. Its kernel is $Z(N)$, so that the assumption that this is a split Lie subgroup implies that γ defines a Lie group extension of G^S by $Z(N)$.

(1) If $\varphi: \widehat{G}_1 \rightarrow \widehat{G}_2$ is an equivalence of N -extensions of G , then the representations C_N^j , $j = 1, 2$, of \widehat{G}_j on N satisfy $C_N^2 \circ \varphi = C_N^1$ because $\varphi|_N = \text{id}_N$. Therefore the quotient maps

$$\gamma_j = (C_N^j \cdot (S \circ q_j)^{-1}, q_j): \widehat{G}_j \rightarrow G^S$$

satisfy $\gamma_2 \circ \varphi = ((C_N^2 \circ \varphi) \cdot (S \circ q_2 \circ \varphi)^{-1}, q_2 \circ \varphi) = (C_N^1 \cdot (S \circ q_1)^{-1}, q_1) = \gamma_1$. This means that $\varphi: \widehat{G}_1 \rightarrow \widehat{G}_2$ also is an equivalence of extensions of G^S by $Z(N)$.

(2) Suppose first that the extension $\gamma: \widehat{G}^S \rightarrow G^S$ by $Z(N)$ comes from the N -extension $q: \widehat{G} \rightarrow G$ corresponding to $[S]$ in the sense that $\widehat{G} = \widehat{G}^S$ with $\widehat{G}/Z(N) \cong G^S$. We may assume that $\widehat{G} = G_{(S,\omega)}$ (Proposition 2.8). Then (4.1) shows that $(n, g) \in \widehat{G}$ acts on N by $C_N(n)S(g) = \rho(C_N(n), g)$. Therefore the inclusion $N \hookrightarrow \widehat{G}$ onto the subset $\gamma^{-1}(N_{\text{ad}})$ is equivariant with respect to the action of G^S , and therefore in particular for the action of N_{ad} , so that it is an equivalence of central extensions of N_{ad} by $Z(N)$.

Suppose, conversely, that $\gamma: \widehat{G}^S \rightarrow G^S$ is an extension of G^S by $Z(N)$ for which there exists a G^S -equivariant equivalence $\alpha: N \rightarrow \gamma^{-1}(N_{\text{ad}})$ of central extensions of N_{ad} by $Z(N)$. Then

$$\widehat{G}^S/\alpha(N) = \widehat{G}^S/\gamma^{-1}(N_{\text{ad}}) \cong G^S/N_{\text{ad}} \cong G,$$

so that we obtain an extension of G by N by the quotient map $q = q_S \circ \gamma: \widehat{G}^S \rightarrow G$ with kernel $\gamma^{-1}(N_{\text{ad}})$. The action of $G^S \cong \widehat{G}^S/Z(N)$ on N induced by the conjugation action of \widehat{G}^S on N coincides with the given action

$$\rho: G^S \rightarrow \text{Aut}(N), \quad (C_N(n), g) \mapsto C_N(n)S(g)$$

of G^S on N because α is G^S -equivariant. Therefore the G -kernel of the extension $q: \widehat{G}^S \rightarrow G$ is $[S]$.

For the conjugation action C_N of \widehat{G}^S on N we have $C_N = \rho \circ \gamma$ and $q = q_S \circ \gamma$, so that the corresponding map $\widehat{G}^S \rightarrow G^S$ coincides with γ . This

means that $\gamma: \widehat{G}^S \rightarrow G^S$ is associated to the extension $q: \widehat{G}^S \rightarrow G$ by the process described above. \square

Remark 4.5. — In Lemma 4.1 we have seen that the map $\alpha: N \rightarrow G^S$ defines a smooth crossed module with smooth outer action $S: G \rightarrow \text{Aut}(N)$. According to Theorem 3.8, the central extension

$$Z(N) \hookrightarrow N \xrightarrow{\alpha} N_{\text{ad}}$$

can be embedded in a G^S -equivariant way into an abelian extension

$$Z(N) \hookrightarrow \widehat{G}^S \xrightarrow{\alpha} G^S$$

if and only if $\chi_{ss}(S) \in H_{ss}^3(G, Z(N))_S$ vanishes. If G is connected, we have the simpler criterion $\chi_s(S) = \mathbf{1}$ in $H_s^3(G, Z(N))_S$. Comparing with Lemma 4.3, we see that this happens if and only if $\text{Ext}(G, N)_{[S]} \neq \emptyset$ (cf. also Theorem 2.17).

Appendix A. Automorphisms of Lie group extensions

In this appendix we collect several useful results on automorphisms of group extensions.

Let $q: \widehat{G} \rightarrow G$ be a Lie group extension of G by N corresponding to the G -kernel $[S]$. In this appendix we discuss the group

$$\text{Aut}(\widehat{G}, N) := \{\varphi \in \text{Aut}(\widehat{G}) : \varphi(N) = N\}$$

of all Lie group automorphisms of \widehat{G} preserving the split normal Lie subgroup N . We have a group homomorphism

$$\Phi: \text{Aut}(\widehat{G}, N) \rightarrow \text{Aut}(N) \times \text{Aut}(G), \quad \varphi \mapsto (\varphi_N, \varphi_G) := (\Phi_N(\varphi), \Phi_G(\varphi))$$

with

$$\varphi_N := \varphi|_N \text{ and } \varphi_G \circ q = q \circ \varphi$$

(cf. Lemmas 1.7 and 2.3). The kernel of Φ consists of those automorphisms of \widehat{G} inducing the identity on the subgroup N and the quotient group G . One of the main results of this appendix is an exact sequence

$$\mathbf{0} \rightarrow Z_s^1(G, Z(N))_S \xrightarrow{\Psi} \text{Aut}(\widehat{G}, N) \xrightarrow{\Phi} (\text{Aut}(N) \times \text{Aut}(G))_{[S]} \xrightarrow{I} H_{ss}^2(G, Z(N))_S,$$

where I is a 1-cocycle for the action of the stabilizer $(\text{Aut}(N) \times \text{Aut}(G))_{[S]}$ of $[S]$ on the group $H_{ss}^2(G, Z(N))_S$. In the context of abstract groups, a similar result has been obtained by Wells in [28] (cf. also [22]).

Automorphisms of Lie group extensions

First we take a closer look at the action of $\text{Aut}(N) \times \text{Aut}(G)$ on factor systems and extensions. In the following we shall often refer to the natural action of $\text{Aut}(N) \times \text{Aut}(G)$ on the sets $C_s^p(G, N)$ by

$$((\varphi, \psi).f)(g_1, \dots, g_p) := \varphi(f(\psi^{-1}(g_1), \dots, \psi^{-1}(g_p))).$$

LEMMA A.1. — *The group $\text{Aut}(N) \times \text{Aut}(G)$ acts naturally on $C_s^1(G, \text{Aut}(N)) \times C_{ss}^2(G, N)$ by*

$$(\varphi, \psi).(S, \omega) := (c_\varphi \circ S \circ \psi^{-1}, \varphi \circ \omega \circ (\psi^{-1} \times \psi^{-1})).$$

For $(\varphi, \psi).(S, \omega) = (S', \omega')$ the following assertions hold:

- (1) If $\delta_S = C_N \circ \omega$, then $\delta_{S'} = C_N \circ \omega'$.
- (2) $d_{S'}\omega' = \varphi \circ (d_S\omega) \circ (\psi \times \psi \times \psi)^{-1}$. In particular, the subset $Z_{ss}^2(G, N)$ is invariant under the action of $\text{Aut}(N) \times \text{Aut}(G)$.
- (3) For $h \in C_s^1(G, N)$ and $(\varphi, \psi) \in \text{Aut}(N) \times \text{Aut}(G)$ we have

$$(\varphi, \psi).(h.(S, \omega)) = ((\varphi, \psi).h).((\varphi, \psi).(S, \omega)).$$

In particular, we obtain an action of the semidirect product group

$$C_s^1(G, N) \rtimes (\text{Aut}(N) \times \text{Aut}(G))$$

and hence an action of $\text{Aut}(N) \times \text{Aut}(G)$ on the set $H_{ss}^2(G, N)$ of $C_s^1(G, N)$ -orbits.

(4) Let $\mathcal{E}: N \xrightarrow{\iota} \widehat{G} \xrightarrow{q} G$ be a short exact sequence describing a Lie group extension of G by N and consider the exact sequence

$$\mathcal{E}' := (\varphi, \psi).\mathcal{E}: \quad \mathbf{1} \rightarrow N \xrightarrow{\iota \circ \varphi^{-1}} \widehat{G} \xrightarrow{\psi \circ q} G \rightarrow \mathbf{1}.$$

For $\widehat{G} = N \times_{(S, \omega)} G$ the sequence \mathcal{E}' describes an extension equivalent to $N \times_{(\varphi, \psi).(S, \omega)} G$ and the map

$$\mu_{(\varphi, \psi)}: N \times_{(S, \omega)} G \rightarrow N \times_{(\varphi, \psi).(S, \omega)} G, \quad (n, g) \mapsto (\varphi(n), \psi(g))$$

is an isomorphism of Lie groups.

Proof.

(1) For $S' := (\varphi, \psi).S$ and $\delta_S = C_N \circ \omega$ we have

$$\begin{aligned} \delta_{S'} &= c_\varphi \circ \delta_S \circ (\psi \times \psi)^{-1} = c_\varphi \circ C_N \circ \omega \circ (\psi \times \psi)^{-1} \\ &= c_\varphi \circ C_N \circ \varphi^{-1} \circ (\varphi, \psi).\omega = C_N \circ (\varphi, \psi).\omega. \end{aligned}$$

(2) follows from

$$\begin{aligned} (d_{S'}\omega')(g, g', g'') &= S'(g)(\omega'(g', g''))\omega'(g, g'g'')\omega'(gg', g'')^{-1}\omega'(g, g')^{-1} \\ &= \varphi\left(S(g)(\omega(\psi^{-1}g', \psi^{-1}g'')) \cdots \omega(\psi^{-1}g, \psi^{-1}g')^{-1}\right) \\ &= \varphi \circ d_S\omega \circ (\psi \times \psi \times \psi)^{-1}(g, g', g''). \end{aligned}$$

In view of (1), this formula shows that the action of $\text{Aut}(N) \times \text{Aut}(G)$ on the product set $C_s^1(G, \text{Aut}(N)) \times C_{ss}^2(G, N)$ preserves the subset $Z_{ss}^2(G, N)$ of smooth factor systems.

(3) follows from an easy calculation. For the identification of $\text{Ext}(G, N)$ with the set $H_{ss}^2(G, N)$ of $C_s^1(G, N)$ -orbits in $Z_{ss}^2(G, N)$, we refer to Proposition 2.10.

(4) Suppose that $\widehat{G} = N \times_{(S, \omega)} G$ holds for a smooth factor system (S, ω) (Proposition 2.8). If $\sigma: G \rightarrow N \times_{(S, \omega)} G, g \mapsto (\mathbf{1}, g)$ is the canonical section, then $\sigma' := \sigma \circ \psi^{-1} \in C_s^1(G, \widehat{G})$ satisfies

$$(\psi \circ q) \circ \sigma' = \psi \circ q \circ \sigma \circ \psi^{-1} = \text{id}_G,$$

so that we can interpret σ' as a section for the extension \mathcal{E}' . We now have

$$\delta_{\sigma'} = \delta_\sigma \circ \psi^{-1} = \iota \circ \omega \circ \psi^{-1},$$

and the action of \widehat{G} on N corresponding to the extension \mathcal{E}' is given by

$$\begin{aligned} C'_N(x)(g) &:= \iota'^{-1}(x\iota'(g)x^{-1}) = \varphi \circ \iota^{-1}(x\iota(\varphi^{-1}(g))x^{-1}) \\ &= \varphi \circ C_N(x)(\varphi^{-1}(g)) = (c_\varphi \circ C_N(x))(g). \end{aligned}$$

Therefore

$$S' := C'_N \circ \sigma' = c_\varphi \circ C_N \circ \sigma \circ \psi^{-1} = c_\varphi \circ S \circ \psi^{-1} = (\varphi, \psi).S.$$

Moreover, the relation $\iota \circ \omega = \delta_\sigma$ leads to

$$\begin{aligned} \delta_{\sigma'} &= \delta_\sigma \circ (\psi \circ \psi)^{-1} = \iota \circ \omega \circ (\psi \circ \psi)^{-1} \\ &= \iota \circ \varphi^{-1} \circ \varphi \circ \omega \circ (\psi \circ \psi)^{-1} = \iota' \circ (\varphi, \psi).\omega. \end{aligned}$$

This implies that $\omega' = (\varphi, \psi).\omega$, so that the sequence \mathcal{E}' is represented by the extension $N \times_{(S', \omega')} G$ with $(S', \omega') = (\varphi, \psi).(S, \omega)$. Furthermore, the

map $\mu_{(\varphi, \psi)}$ is a group homomorphism:

$$\begin{aligned} \mu_{(\varphi, \psi)}((n, g)(n', g')) &= \mu_{(\varphi, \psi)}(nS(g)(n')\omega(g, g'), gg') \\ &= (\varphi(n)(\varphi \circ S(g)(n'))(\varphi \circ \omega(g, g')), \psi(gg')) \\ &= (\varphi(n)(c_\varphi \cdot S(g))(\varphi(n'))\omega'(\psi(g), \psi(g')), \psi(gg')) \\ &= (\varphi(n)S'(g)(\varphi(n'))\omega'(\psi(g), \psi(g')), \psi(gg')) \\ &= (\varphi(n), \psi(g))(\varphi(n'), \psi(g')) = \mu_{(\varphi, \psi)}(n, g)\mu_{(\varphi, \psi)}(n', g'). \end{aligned}$$

That it is an isomorphism of Lie groups follows from the fact that it is a local diffeomorphism. \square

The following two lemmas provide descriptions of kernel and range of Φ .

LEMMA A.2. — On $Z(N)$ we consider the action S_Z of G induced by the conjugation action of \widehat{G} on N . For each $f \in Z_s^1(G, Z(N))_{S_Z}$ we obtain an element $\varphi_f := (f \circ q) \cdot \text{id}_{\widehat{G}} \in \ker \Phi$, and the map

$$\Psi: Z_s^1(G, Z(N))_{S_Z} \rightarrow \ker \Phi, \quad f \mapsto \varphi_f$$

is a bijective group homomorphism.

Proof. — Let $f \in Z_s^1(G, Z(N))$. Clearly $\varphi_f: \widehat{G} \rightarrow \widehat{G}$ is a diffeomorphism of \widehat{G} with $q \circ \varphi_f = q$ whose inverse is given by $\varphi_{f^{-1}}$. That it is a group homomorphism follows for $\gamma = q(\widehat{\gamma})$ and $\gamma' = q(\widehat{\gamma}')$ from

$$\begin{aligned} \text{(A.1)} \quad \varphi_f(\widehat{\gamma}\widehat{\gamma}') &= f(\gamma\gamma')\widehat{\gamma}\widehat{\gamma}' = f(\gamma)\gamma(f(\gamma'))\widehat{\gamma}\widehat{\gamma}' \\ &= f(\gamma)\widehat{\gamma}f(\gamma')\widehat{\gamma}^{-1}\widehat{\gamma}\widehat{\gamma}' = \varphi_f(\gamma)\varphi_f(\gamma'). \end{aligned}$$

From $q \circ \varphi_f = q$ we immediately obtain that $\varphi_f \circ \varphi_{f'} = \varphi_{ff'}$ for $f, f' \in Z_s^1(G, Z(N))$, showing that Ψ is a group homomorphism. This homomorphism is obviously injective. To see that it is also surjective, let $\varphi \in \ker \Phi$. Then $\varphi = \widehat{f} \cdot \text{id}_{\widehat{G}}$ with a smooth function $\widehat{f}: \widehat{G} \rightarrow N$. The fact that φ is a homomorphism implies that $\widehat{f} \in Z_s^1(\widehat{G}, N)$ with respect to the conjugation action of \widehat{G} on N (cf. (A.1)). Moreover, $\widehat{f}(N) = \{1\}$ implies that \widehat{f} is constant on the N -cosets:

$$\widehat{f}(gn) = \widehat{f}(g)c_g(\widehat{f}(n)) = \widehat{f}(g), \quad g \in \widehat{G}, n \in N.$$

Hence \widehat{f} can be written as $f \circ q$ for some smooth function $f: G \rightarrow N$. Since N is normal in \widehat{G} , we also have for each $g \in \widehat{G}$ and $n \in N$ the relation

$$\widehat{f}(g) = \widehat{f}(g \cdot (g^{-1}ng)) = \widehat{f}(ng) = \widehat{f}(n)c_n(\widehat{f}(g)) = c_n(\widehat{f}(g)),$$

and hence that $\text{im}(f) \subseteq Z(N)$. This proves that $f \in Z_s^1(G, Z(N))$, so that Ψ is surjective. \square

PROPOSITION A.3. — Let $(S, \omega) \in Z_{ss}^2(G, N)$, $\widehat{G} = N \times_{(S, \omega)} G$ be the corresponding Lie group extension of G by N and $(\varphi, \psi) \in \text{Aut}(N) \times \text{Aut}(G)$.

(a) We write the extension \widehat{G} as the exact sequence

$$\mathcal{E}: \mathbf{1} \rightarrow N \xrightarrow{\iota} \widehat{G} \xrightarrow{q} G \rightarrow \mathbf{1}.$$

Then $(\varphi, \psi) \in \text{im}(\Phi)$ if and only if $(\varphi, \psi).\mathcal{E} \sim \mathcal{E}$ holds for the extension

$$(\varphi, \psi).\mathcal{E}: \mathbf{1} \rightarrow N \xrightarrow{\iota \circ \varphi^{-1}} \widehat{G} \xrightarrow{\psi \circ q} G \rightarrow \mathbf{1}.$$

(b) An automorphism $\nu \in \text{Aut}(\widehat{G}, N)$ satisfies $\Phi(\nu) = (\varphi, \psi)$ if and only if it is of the form

$$(A.2) \quad \nu(n, g) = (\varphi(n)h(\beta(g)), \beta(g))$$

with $h \in C_s^1(G, N)$ satisfying $(\varphi, \psi).(S, \omega) = h.(S, \omega)$. In particular, $(\varphi, \psi) \in \text{im}(\Phi)$ if and only if we have in $\text{Ext}(G, N) \cong Z_{ss}^2(G, N)/C_s^1(G, N)$ the relation

$$(\varphi, \psi).[(S, \omega)] = [(S, \omega)].$$

Proof.

(a) For $\lambda \in \text{Aut}(\widehat{G}, N)$ we consider the extension $\mathcal{E}' := (\lambda_N, \lambda_G).\mathcal{E}$ and put $\iota' := \iota \circ \lambda_N^{-1}$ and $q' := \lambda_G \circ q$. Then the map $\lambda: \widehat{G} \rightarrow \widehat{G}$ yields an equivalence of extensions

$$\begin{array}{ccccc} N & \xrightarrow{\iota \circ \lambda_N^{-1}} & \widehat{G} & \xrightarrow{\lambda_G \circ q} & G \\ \downarrow \text{id}_N & & \downarrow \lambda & & \downarrow \text{id}_G \\ N & \xrightarrow{\iota} & \widehat{G} & \xrightarrow{q} & G. \end{array}$$

Therefore $\Phi(\lambda).\mathcal{E} \sim \mathcal{E}$. If, conversely, $(\varphi, \psi).\mathcal{E} \sim \mathcal{E}$, there exists an equivalence of extensions

$$\begin{array}{ccccc} N & \xrightarrow{\iota \circ \varphi^{-1}} & \widehat{G} & \xrightarrow{\psi \circ q} & G \\ \downarrow \text{id}_N & & \downarrow \lambda & & \downarrow \text{id}_G \\ N & \xrightarrow{\iota} & \widehat{G} & \xrightarrow{q} & G. \end{array}$$

This means that $\lambda_N = \varphi$ and $\lambda_G = \psi$.

(b) Let $\nu \in \text{Aut}(\widehat{G}, N)$ with $\Phi(\nu) = (\varphi, \psi)$. In the product coordinates of $\widehat{G} = N \times_{(S, \omega)} G$ we then have

$$\nu(n, g) = (\varphi(n)h(\psi(g)), \psi(g))$$

for some $h \in C_s^1(G, N)$. Let $(S', \omega') := (\varphi, \psi).(S, \omega)$ and consider the isomorphism

$$\mu: N \times_{(S, \omega)} G \rightarrow N \times_{(S', \omega')} G, \quad (n, g) \mapsto (\varphi(n), \psi(g))$$

(Lemma A.1(4)). Then we obtain an isomorphism

$$\lambda := \nu \circ \mu^{-1}: N \times_{(S', \omega')} G \rightarrow N \times_{(S, \omega)} G, \quad (n, g) \mapsto (nh(g), g)$$

which is an equivalence of extensions, so that Proposition 2.10 implies that $(S', \omega') = h.(S, \omega)$.

If, conversely, $(S', \omega') = h.(S, \omega)$, then λ is an equivalence of extensions and $\nu := \lambda \circ \mu$ an automorphism of Lie groups. □

COROLLARY A.4. — *Suppose that N is a smooth abelian G -module, where the action of G on N is given by $S: G \rightarrow \text{Aut}(N)$, $\omega \in Z_{ss}^2(G, N)$, and that $\widehat{G} = N \times_{\omega} G$ is the corresponding abelian extension of G by N . Further let $(\varphi, \psi) \in \text{Aut}(N) \times \text{Aut}(G)$. Then an automorphisms $\nu \in \text{Aut}(\widehat{G}, N)$ satisfies $\Phi(\nu) = (\varphi, \psi)$ if and only if it is of the form*

$$(A.3) \quad \nu(n, g) := (\varphi(n)h(\beta(g)), \beta(g))$$

with $h \in C_s^1(G, N)$ satisfying

$$(\varphi, \psi).S = S \text{ and } (\varphi, \psi).\omega - \omega = d_S h.$$

The preceding observations provide the exact sequence

$$\mathbf{1} \rightarrow Z_s^1(G, Z(N)) \xrightarrow{\Psi} \text{Aut}(\widehat{G}, N) \xrightarrow{\Phi} (\text{Aut}(N) \times \text{Aut}(G))_{[\mathcal{E}]} \rightarrow \mathbf{1}.$$

The following lemma expresses that the actions of \widehat{G} on N and G preserve the extension class \mathcal{E} .

LEMMA A.5. — *Let N be a G -Lie group and $q_G: \widehat{G} \rightarrow G$ be an N -extension of G for which the corresponding outer action is contained in the class $[S]$. Then for each $g \in G$ there is an automorphism $\varphi \in \text{Aut}(\widehat{G}, N)$ with $\varphi_N = S(g)$ and $\varphi_G = c_g$.*

Proof. — Let $\widehat{g} \in \widehat{G}$ with $q(\widehat{g}) = g$. Then $\psi := C_{\widehat{G}}(\widehat{g}) \in \text{Aut}(\widehat{G}, N)$ satisfies $\psi_G = c_g$ and ψ_N is of the form $c_n \circ S(g)$ for some $n \in N$ because $[\widehat{G}] \in \text{Ext}(G, N)_{[S]}$. On the other hand $C_{\widehat{G}}(n)$ satisfies $C_{\widehat{G}}(n)_G = \text{id}_G$ and $C_{\widehat{G}}(n)_N = c_n$, so that $\varphi := C_{\widehat{G}}(n)^{-1} \circ \psi$ meets all requirements. □

Remark A.6. — The group N also acts on the set $Z_{ss}^2(G, N)$ of factor systems by the homomorphism $(C_N, \mathbf{1}): N \rightarrow \text{Aut}(N) \times \text{Aut}(G)$. This means that

$$(n.S)(g) = C_N(n) \cdot S(g) \cdot C_N(n)^{-1} = C_N(nS(g)(n^{-1}))S(g)$$

and

$$n.\omega = C_N(n) \circ \omega.$$

We consider the function $h \in C_s^1(G, N)$ given by $h(g) := nS(g)(n^{-1}) = d_S(n)(g)$ and claim that

$$h *_S \omega = n.\omega,$$

which implies that $n.(S, \omega) = h.(S, \omega)$. In fact, we have

$$\begin{aligned} (h *_S \omega)(g, g') &= h(g)S(g)(h(g'))\omega(g, g')h(gg')^{-1} \\ &= nS(g)(n)^{-1}S(g)(nS(g')(n^{-1}))\omega(g, g')S(gg')(n)n^{-1} \\ &= nS(g)S(g')(n^{-1})\omega(g, g')S(gg')(n)n^{-1} \\ &= n(C_N(\omega(g, g'))S(gg')(n^{-1}))\omega(g, g')S(gg')(n)n^{-1} \\ &= n\omega(g, g')S(gg')(n^{-1})\omega(g, g')^{-1}\omega(g, g')S(gg')(n)n^{-1} \\ &= n\omega(g, g')n^{-1} = C_N(n)(\omega(g, g')). \end{aligned}$$

Remark A.7. — For an extension $q: \widehat{G} \rightarrow G$ it is also interesting to consider its *gauge automorphism group*

$$\text{Gau}(\widehat{G}) := \{\varphi \in \text{Aut}(\widehat{G}, N) : q \circ \varphi = q\}$$

of all automorphisms of \widehat{G} inducing the identity on G . It is the kernel of the natural homomorphism

$$\Phi_G : \text{Aut}(\widehat{G}, N) \rightarrow \text{Aut}(G)$$

and it obviously contains $\ker \Phi \cong Z_s^1(G, Z(N))$ (Proposition A.3).

Each automorphism of \widehat{G} preserving N that induces the identity on G can be written in the form $\varphi = \varphi_f := f \cdot \text{id}_{\widehat{G}}$, where $f: \widehat{G} \rightarrow N$ is a smooth function. We then have

$$\varphi_{f_1} \circ \varphi_{f_2} = \varphi_{(f_1 \circ \varphi_{f_2}) \cdot f_2}.$$

If $f|_N \cdot \text{id}_N : N \rightarrow N$ is a diffeomorphism, then the map φ_f is a diffeomorphism of \widehat{G} with $\varphi_f^{-1} = \varphi_{f^{-1} \circ \varphi_f^{-1}}$.

It is easy to see that φ_f is a group homomorphism if and only if $f \in Z_s^1(\widehat{G}, N)_{C_N}$ with respect to the conjugation action C_N of \widehat{G} on N . This means that

$$\text{Gau}(\widehat{G}) \cong (Z_s^1(\widehat{G}, N), *)^\times$$

is the unit group of the monoid $Z_s^1(\widehat{G}, N)$, where the monoid structure on $Z_s^1(\widehat{G}, N)_{C_N}$ is given by $\varphi_{f_1 * f_2} = \varphi_{f_1} \circ \varphi_{f_2}$ and

$$\begin{aligned} (f_1 * f_2)(\widehat{\gamma}) &= f_1(f_2(\widehat{\gamma}))f_2(\widehat{\gamma}) \\ &= f_1(f_2(\widehat{\gamma}))c_{f_2(\widehat{\gamma})}(f_1(\widehat{\gamma}))f_2(\widehat{\gamma}) \\ &= f_1(f_2(\widehat{\gamma}))f_2(\widehat{\gamma})f_1(\widehat{\gamma}). \end{aligned}$$

This formula shows in which way the pointwise product on the group $C_s^1(\widehat{G}, N)$ is twisted to obtain the monoid structure on $Z_s^1(\widehat{G}, N)$.

Instead of considering the subgroup $\text{im}(\Phi) = (\text{Aut}(N) \times \text{Aut}(G))_{[\mathcal{E}]}$, one may also consider the larger subgroup

$$\begin{aligned} \text{Comp}(S) &:= (\text{Aut}(N) \times \text{Aut}(G))_{[S]} \\ &= \{(\varphi, \psi) \in \text{Aut}(N) \times \text{Aut}(G) : (\varphi, \psi).S \sim S\} \end{aligned}$$

of *S-compatible pairs of automorphisms of N and G*, and then identify $\text{im}(\varphi)$ as a subgroup of $\text{Comp}(S)$. The group $\text{Comp}(S)$ acts on $\text{Ext}(G, N)_{[S]}$ because its action on $\text{Ext}(G, N)$ preserves the subset

$$\text{Ext}(G, N)_{[S]} \subseteq \text{Ext}(G, N).$$

We also have a simply transitive action of the abelian group $H_{ss}^2(G, Z(N))$ on $\text{Ext}(G, N)_{[S]}$ induced from the action of $Z_{ss}^2(G, Z(N))$ on $Z_{ss}^2(G, N)$ by

$$\beta.(S, \omega) := (S, \beta \cdot \omega)$$

(Corollary 2.13). For $(\varphi, \psi) \in \text{Comp}(S)$ we then have

$$(\varphi, \psi).(\beta.(S, \omega)) = (\varphi, \psi).(S, \beta \cdot \omega) = ((\varphi, \psi).S, ((\varphi, \psi).\beta) \cdot (\varphi, \psi).\omega).$$

This implies that the actions of $\text{Comp}(S)$ and $H_{ss}^2(G, Z(N))$ on $\text{Ext}(G, N)_{[S]}$ satisfy

$$(\varphi, \psi).([\beta].[\widehat{G}]) = ((\varphi, \psi).\beta).((\varphi, \psi).[\widehat{G}]).$$

PROPOSITION A.8. — For $(\varphi, \psi) \in \text{Comp}(S)$ let $I(\varphi, \psi) \in H_{ss}^2(G, Z(N))_S$ denote the unique cohomology class with $(\varphi, \psi).[\widehat{G}] = I(\varphi, \psi).[\widehat{G}]$. Then the following assertions hold:

- (1) The action of $\text{Comp}(S)$ on $\text{Ext}(G, N)_{[S]} = H_{ss}^2(G, Z(N))_S \cdot [\widehat{G}]$ has the form

$$(\varphi, \psi).([\beta].[\widehat{G}]) = ((\varphi, \psi).\beta) + I(\varphi, \psi).[\widehat{G}].$$

- (2) $I: \text{Comp}(S) \rightarrow H_{ss}^2(G, Z(N))_S$ is a 1-cocycle for the action of $\text{Comp}(S)$ on $H_{ss}^2(G, Z(N))_S$.
- (3) If $\widehat{G} \cong N \times_{(S, \omega)} G$, then there exists an $h_0 \in C_s^1(G, N)$ with $(\varphi, \psi).S = h_0.S$ and

$$I(\varphi, \psi) = [(\varphi, \psi).\omega - h_0 *_S \omega].$$

- (4) If $I(\varphi, \psi) = 0$, then there exists some $\gamma \in C_s^1(G, Z(N))$ with $d_S \omega = (\varphi, \psi).\omega - h_0 *_S \omega$, and for $h := h_0 \cdot \gamma$ the map

$$\nu: N \times_{(S, \omega)} G \rightarrow N \times_{(S, \omega)} G, \quad (n, g) \mapsto (\varphi(n)h(\psi(g)), \psi(g))$$

is an element of $\text{Aut}(\widehat{G}, N)$ with $\Phi(\nu) = (\varphi, \psi)$.

Proof.

(1) is clear.

(2) For $(\varphi, \psi), (\varphi', \psi') \in \text{Comp}(S)$ we have

$$\begin{aligned} I((\varphi, \psi)(\varphi', \psi')).[\widehat{G}] &= ((\varphi, \psi)(\varphi', \psi')).[\widehat{G}] = (\varphi, \psi).((\varphi', \psi')).[\widehat{G}] \\ &= (\varphi, \psi).(I(\varphi', \psi').[\widehat{G}]) = ((\varphi, \psi).I(\varphi', \psi')).((\varphi, \psi).[\widehat{G}]) \\ &= ((\varphi, \psi).I(\varphi', \psi')).(I(\varphi, \psi).[\widehat{G}]) \\ &= ((\varphi, \psi).I(\varphi', \psi') + I(\varphi, \psi)).[\widehat{G}]. \end{aligned}$$

(3) As $(\varphi, \psi) \in \text{Comp}(S)$, we have $S' := (\varphi, \psi).S \sim S$, so that there exists an $h_0 \in C_s^1(G, N)$ with $S' = h_0.S = (C_N \circ h_0).S$. Let $\omega' := (\varphi, \psi).\omega$. Then

$$C_N \circ \omega' = \delta_{S'} = C_N \circ (h_0 *_S \omega)$$

implies that $\beta := \omega' - h_0 *_S \omega \in Z_{ss}^2(G, Z(N))_S$, and we have

$$(\varphi, \psi).(S, \omega) = (S', \omega') = \beta.(S', h_0 *_S \omega) = \beta.(h_0.S, h_0 *_S \omega) \sim \beta.(S, \omega).$$

Therefore the cocycle I is given by

$$I(\varphi, \psi) = [\beta] = [\omega' - h_0 *_S \omega] = [(\varphi, \psi).\omega - h_0 *_S \omega] \in H_{ss}^2(G, Z(N))_S.$$

(4) If $I(\varphi, \psi) = \mathbf{1}$, then there exists some $\gamma \in C_s^1(G, Z(N))$ with $\beta = d_S \gamma$, so that we obtain for $h := h_0 \cdot \gamma$ the relations $S' = h.S$ and $\omega' = h_0 *_S \omega + d_S \gamma = h *_S \omega$, i.e., $(S', \omega') = h.(S, \omega)$. Then the map

$$\lambda_h : N \times_{(S', \omega')} G \rightarrow N \times_{(S, \omega)} G, \quad (n, g) \mapsto (nh(g), g)$$

is an equivalence of extensions (Proposition 2.10), and composition with the isomorphism

$$\mu_{(\varphi, \psi)} : N \times_{(S, \omega)} G \rightarrow N \times_{(S', \omega')} G, \quad (n, g) \mapsto (\varphi(n), \psi(g))$$

(Lemma A.1) leads to the automorphism ν . □

We now see that the exact sequence from above has a prolongation:

THEOREM A.9. — *The following sequence is exact*

$$\mathbf{1} \rightarrow Z_s^1(G, Z(N))_S \rightarrow \text{Aut}(\widehat{G}, N) \xrightarrow{\Phi} \text{Comp}(S) \xrightarrow{I} H_{ss}^2(G, Z(N))_S.$$

Proof. — This follows from Lemma A.2 and Propositions A.3 and A.8, which yields

$$\text{im}(\Phi) = \text{Comp}(S)_{[\widehat{G}]} = I^{-1}(0).$$

□

Automorphism groups of Lie group extensions

So far we dealt with pairs $(\varphi, \psi) \in \text{Aut}(N) \times \text{Aut}(G)$ coming from automorphisms of the extension $q: \widehat{G} \rightarrow G$ with kernel N , which lead to the exact sequence

$$(A.4) \quad \mathbf{1} \rightarrow Z_s^1(G, Z(N))_S \hookrightarrow \text{Aut}(\widehat{G}, N) \twoheadrightarrow (\text{Aut}(N) \times \text{Aut}(G))_{[\widehat{G}]} \rightarrow \mathbf{1}.$$

If we are given a group homomorphism

$$\psi: H \rightarrow (\text{Aut}(N) \times \text{Aut}(G))_{[\widehat{G}]} = (\text{Aut}(N) \times \text{Aut}(G))_{[S, \omega]} = \text{im}(\Phi),$$

then each element $\psi(h)$ comes from an automorphism $\widehat{\psi}(h) \in \text{Aut}(\widehat{G}, N)$, but this automorphism is not unique if $Z_s^1(G, Z(N))_S \neq \mathbf{1}$. To obtain a group homomorphism $\widehat{\psi}: H \rightarrow \text{Aut}(\widehat{G}, N)$ lifting ψ , a certain cohomology class in $H^2(H, Z_s^1(G, Z(N))_S)$ has to vanish, i.e., the abelian extension $\psi^* \text{Aut}(\widehat{G}, N)$ of H by $Z_s^1(G, Z(N))_S$ has to split. That this is not always the case is shown in the following example, where we describe a central extension $Z \hookrightarrow \widehat{G} \twoheadrightarrow G$ for which (A.4) does not split.

Example A.10.

(a) We construct an example of a central extension $Z \hookrightarrow \widehat{G} = Z \times_f G \twoheadrightarrow G$ of finite-dimensional real Lie groups, for which the sequence

$$Z_s^1(G, Z) \cong \text{Hom}(G, Z) \hookrightarrow \text{Aut}(\widehat{G}, Z) \twoheadrightarrow (\text{Aut}(Z) \times \text{Aut}(G))_{[f]}$$

is not split. As this is an exact sequence of finite-dimensional real Lie groups, it suffices to show that the corresponding sequence

$$Z^1(\mathfrak{g}, \mathfrak{z}) \cong \text{Hom}_{\text{Lie}}(\mathfrak{g}, \mathfrak{z}) \hookrightarrow \text{der}(\widehat{\mathfrak{g}}, \mathfrak{z}) \twoheadrightarrow (\text{der}(\mathfrak{z}) \times \text{der}(\mathfrak{g}))_{[\omega]}$$

of Lie algebras is not split if $\omega \in Z^2(\mathfrak{g}, \mathfrak{z})$ satisfies $\widehat{\mathfrak{g}} \sim \mathfrak{z} \oplus_{\omega} \mathfrak{g}$.

Let G be the 3-dimensional real Heisenberg group and \mathfrak{g} is Lie algebra. Then \mathfrak{g} has a basis of the form p, q, z satisfying

$$[p, q] = z, \quad [p, z] = [q, z] = 0$$

and we identify G with \mathfrak{g} , endowed with the Campbell–Hausdorff product

$$x * y := x + y + \frac{1}{2}[x, y].$$

Note that all left multiplications $\lambda_x(y) := x * y$ are affine maps.

We consider the cocycle $\omega \in Z^2(\mathfrak{g}, \mathbb{R})$ with

$$\omega(p, z) = 1, \quad \omega(q, z) = \omega(q, z) = 0.$$

We then obtain a central extension $\widehat{\mathfrak{g}} := \mathbb{R} \oplus_{\omega} \mathfrak{g}$ of \mathfrak{g} by $\mathfrak{z} := \mathbb{R}$, and in [20, Ex. A.8] it is shown that the exact sequence

$$\mathrm{Hom}_{\mathrm{Lie}}(\widehat{\mathfrak{g}}, \mathfrak{z}) \cong \mathrm{Hom}_{\mathrm{Lie}}(\mathfrak{g}, \mathfrak{z}) \hookrightarrow \mathrm{der}(\widehat{\mathfrak{g}}, \mathfrak{z}) \twoheadrightarrow (\mathrm{der}(\mathfrak{g}) \times \mathrm{der}(\mathfrak{z}))_{[\omega]}$$

is not split. More concretely, the action of $\mathfrak{b} := \mathrm{Hom}_{\mathrm{Lie}}(\mathfrak{g}, \mathfrak{z}(\mathfrak{g})) \subseteq \mathrm{der}(\mathfrak{g})$ preserves the class $[\omega] \in H^2(\mathfrak{g}, \mathfrak{z})$, but the action of \mathfrak{b} on \mathfrak{g} does not lift to an action of the abelian Lie algebra \mathfrak{b} on $\widehat{\mathfrak{g}}$.

Let $\widehat{\mathfrak{b}} := \mathfrak{g} \rtimes \mathfrak{b}$ be the semidirect sum. Then $[\omega] \in H^2(\mathfrak{g}, \mathfrak{z})^{\mathfrak{b}}$, but there is no representation \widehat{S} of $\widehat{\mathfrak{b}}$ on $\widehat{\mathfrak{g}}$ lifting the representation on \mathfrak{g} .

LEMMA A.11. — *Suppose that $\widehat{G} = N \times_{(S, \omega)} G$. Further let $\theta: H \rightarrow C_s^1(G, N)$ be a map with*

$$h \cdot (S, \omega) = \theta(h)^{-1} \cdot (S, \omega) \text{ for all } h \in H$$

and define the automorphism $\widehat{\psi}(h) \in \mathrm{Aut}(\widehat{G}, N)$ by

$$\widehat{\psi}(h)(n, g) := (h \cdot n \cdot \theta(h)(h \cdot g)^{-1}, h \cdot g).$$

Then $\widehat{\psi}: H \rightarrow \mathrm{Aut}(\widehat{G}, N)$ is a homomorphism if and only if $\theta \in Z^1(H, C_s^1(G, N)_{\psi})$, where $C_s^1(G, N)_{\psi}$ stands for the group $C_s^1(G, N)$, endowed with the H -action given by $h \cdot \chi := \psi(h) \cdot \chi$.

Proof. — That the maps $\widehat{\psi}(h)$ define elements of $\mathrm{Aut}(\widehat{G}, N)$ follows from Proposition A.3(b). We have

$$\begin{aligned} \widehat{\psi}(h)\widehat{\psi}(h')(n, g) &= \widehat{\psi}(h)(h' \cdot n \cdot \theta(h')(h' \cdot g)^{-1}, h' \cdot g) \\ &= \left(h \cdot (h' \cdot n \cdot \theta(h')(h' \cdot g)^{-1}) \cdot \theta(h)(hh' \cdot g)^{-1}, hh' \cdot g \right) \\ &= \left(hh' \cdot n \cdot h \cdot (\theta(h')(h' \cdot g)^{-1}) \cdot \theta(h)(hh' \cdot g)^{-1}, hh' \cdot g \right) \\ &= \left(hh' \cdot n \cdot (\theta(h) \cdot h \cdot \theta(h'))(hh' \cdot g)^{-1}, hh' \cdot g \right). \end{aligned}$$

This implies that

$$\delta_{\widehat{\psi}}(h, h') = \widehat{\psi}(h)\widehat{\psi}(h')\widehat{\psi}(hh')^{-1} \in Z_s^1(G, Z(N)) = \ker \Phi \subseteq \mathrm{Aut}(\widehat{G}, N).$$

In view of

$$\begin{aligned} \widehat{\psi}(h)^{-1}(n, g) &= (h^{-1} \cdot n \cdot h^{-1} \cdot (\theta(h)(g)), h^{-1} \cdot g) \\ &= (h^{-1} \cdot n \cdot (h^{-1} \cdot \theta(h))(h^{-1} \cdot g), h^{-1} \cdot g), \end{aligned}$$

the cocycle $\delta_{\widehat{\psi}}$ is given by

$$\begin{aligned} \theta(h) \cdot h \cdot \theta(h') \cdot hh' \cdot ((hh')^{-1} \cdot \theta(hh'))^{-1} &= \theta(h) \cdot h \cdot \theta(h') \cdot \theta(hh')^{-1} \\ &= (d_H \theta)(h, h'). \end{aligned}$$

We conclude that $\widehat{\psi}$ is a homomorphism if and only if θ is a cocycle. □

Remark A.12. — For a given homomorphism

$$\psi: H \rightarrow (\text{Aut}(N) \times \text{Aut}(G))_{[(S,\omega)]}$$

we may choose a map $\theta: H \rightarrow C_s^1(G, N)$ with $h \cdot (S, \omega) = \theta(h)^{-1} \cdot (S, \omega)$ for each $h \in H$. We have seen in the preceding proof that

$$\text{im}(d_H\theta) \subseteq Z_s^1(G, Z(N))$$

and that for the corresponding map $\widehat{\psi}: H \rightarrow \text{Aut}(\widehat{G}, N)$ we have $\delta_{\widehat{\psi}} = d_H\theta$. This is the cocycle of the abelian extension $\psi^* \text{Aut}(\widehat{G}, N)$ of H by $Z_s^1(G, Z(N)) = \ker \Phi$ obtained by pulling back the abelian extension

$$Z_s^1(G, Z(N)) \hookrightarrow \text{Aut}(\widehat{G}, N) \rightarrow (\text{Aut}(N) \times \text{Aut}(G))_{[(S,\omega)]}.$$

We may thus assign to ψ the cohomology class

$$[d_H\theta] = [\psi^* \text{Aut}(\widehat{G}, N)] \in H^2(H, Z_s^1(G, Z(N))_\psi).$$

If the cohomology class $[d_H\theta]$ vanishes, then there exists a map $\eta: H \rightarrow Z_s^1(G, Z(N))$ with $d_H\theta = d_H\eta$, which implies that $\widetilde{\theta} := \theta \cdot \eta^{-1}$ satisfies $d_H\widetilde{\theta} = 0$ and, in addition, $h \cdot (S, \omega) = \widetilde{\theta}(h)^{-1} \cdot (S, \omega)$ for each $h \in H$. Therefore $\widetilde{\theta} \in Z^1(H, C_s^1(G, N)_\psi)$ defines a lift $\widehat{\psi}$ of ψ . Hence ψ lifts to a homomorphism $\widehat{\psi}: H \rightarrow \text{Aut}(\widehat{G}, N)$ if and only if the cohomology class $[d_H\theta]$ vanishes, which means that there exists a cocycle $\theta: H \rightarrow C_s^1(G, N)$ with $h \cdot (S, \omega) = \theta(h)^{-1} \cdot (S, \omega)$ for each $h \in H$.

In addition to the setting of the preceding subsection, we now assume that H is a Lie group and ask for smoothness properties of actions on \widehat{G} .

PROPOSITION A.13. — *Suppose we are given smooth actions of H on G and N by automorphisms and that we have an action of H on $\widehat{G} := N \times_{(S,\omega)} G$ given by $\theta \in Z^1(H, C_s^1(G, N))$ via*

$$h \cdot (n, g) = (h \cdot n \cdot \theta(h)(h \cdot g)^{-1}, h \cdot g).$$

Then the following assertions hold:

- (1) *Each $h \in H$ acts by a Lie group automorphism.*
- (2) *The action map $\sigma: H \times \widehat{G} \rightarrow \widehat{G}$ is smooth in an identity neighborhood of $H \times \widehat{G}$ if and only if the map*

$$\widetilde{\theta}: H \times G \rightarrow N, \quad (h, g) \mapsto \theta(h)(g)$$

is smooth in an identity neighborhood.

- (3) *If N is abelian, then the action has smooth orbit maps if and only if all maps*

$$\theta_g: H \rightarrow N, \quad h \mapsto \theta(h)(h \cdot g)^{-1} \cdot \omega(h \cdot g, g^{-1})$$

are smooth in an identity neighborhood of H .

- (4) The action σ of H on \widehat{G} is smooth if and only if σ is smooth in an identity neighborhood of $H \times \widehat{G}$ and all orbit maps are smooth. If G is connected, then the smoothness of the orbit maps follows from the local smoothness of the action.

Proof.

(1) Each $h \in H$ acts by a group automorphism which is smooth in an identity neighborhood, hence a Lie group automorphism.

(2) Suppose that the action is smooth in an identity neighborhood. Then the function $(h, g) \mapsto \theta(h)(h.g)$ has this property and the map

$$H \times G \rightarrow H \times G, \quad (h, g) \mapsto (h, h^{-1}.g)$$

is a diffeomorphism fixing $(\mathbf{1}, \mathbf{1})$. Therefore $\tilde{\theta}$ is smooth on an identity neighborhood. The converse is clear.

(3) Since H acts by Lie group automorphisms, the orbit maps are smooth if and only if they are smooth in an identity neighborhood of H .

It suffices to consider orbit maps of elements of the form $(\mathbf{1}, g)$ because H acts smoothly on N . For $g \in G$ the orbit map is smooth in an identity neighborhood of H if and only if the map

$$\begin{aligned} h \mapsto (h.(\mathbf{1}, g))(\mathbf{1}, g)^{-1} &= (\theta(h)(h.g)^{-1}, h.g)(\omega(g^{-1}, g)^{-1}, g^{-1}) \\ &= \left(\theta(h)(h.g)^{-1} S(h.g)(\omega(g^{-1}, g)^{-1}) \cdot \omega(h.g, g^{-1}), (h.g)g^{-1} \right) \end{aligned}$$

(cf. Lemma 2.7 for the inversion formula) is smooth in an identity neighborhood, which is equivalent to the smoothness of

$$h \mapsto \theta(h)(h.g)^{-1} \cdot S(h.g)(\omega(g^{-1}, g)^{-1}) \cdot \omega(h.g, g^{-1})$$

in an identity neighborhood.

If, in addition, N is abelian, then S defines a smooth action of G on N , so that $S(h.g)(\omega(g^{-1}, g)^{-1})$ is a smooth function of h . Hence it suffices that the map

$$h \mapsto \theta(h)(h.g)^{-1} \cdot \omega(h.g, g^{-1})$$

is smooth.

- (4) Since H acts by smooth automorphisms of \widehat{G} , the action of H is smooth if and only if all orbit maps are smooth and it is locally smooth in the sense of (2).

The set of all elements of \widehat{G} with smooth orbit maps is a subgroup containing N . If the action is locally smooth, then this subgroup also contains an identity neighborhood, hence the group $N \times_{(S_0, \omega_0)} G_0$, where G_0 is the identity component of G , $S_0 := S|_{G_0}$ and $\omega_0 := \omega|_{G_0 \times G_0}$. If G is connected,

this argument shows that the local smoothness of the action already implies the smoothness of all orbit maps. \square

Appendix B. Lie group structures on groups

THEOREM B.1. — *Let G be a group and $K = K^{-1}$ a symmetric subset. We further assume that K is a smooth manifold such that*

(L1) *there exists an open $\mathbf{1}$ -neighborhood $V \subseteq K$ with $V^2 = V \cdot V \subseteq K$ such that the group multiplication*

$$\mu_V: V \times V \rightarrow K \text{ is smooth,}$$

(L2) *the inversion map $\eta_K: K \rightarrow K, k \mapsto k^{-1}$ is smooth, and*

(L3) *for each $g \in G$ there exists an open $\mathbf{1}$ -neighborhood $K_g \subseteq K$ with $c_g(K_g) \subseteq K$ and such that the conjugation map*

$$c_g: K_g \rightarrow K, \quad x \mapsto gxg^{-1}$$

is smooth.

Then there exists a unique structure of a Lie group on G for which there exists an open $\mathbf{1}$ -neighborhood $K_1 \subseteq K$ such that the inclusion map $K_1 \rightarrow G$ induces a diffeomorphism onto an open subset of G .

If G is generated by each $\mathbf{1}$ -neighborhood $U \subseteq K$, then condition (L3) can be omitted.

Proof. — [19, Theorem A.4 and Remark A.5]. \square

BIBLIOGRAPHY

- [1] R. BAER, “Erweiterungen von Gruppen und ihren Isomorphismen”, *Math. Zeit.* **38** (1934), p. 375-416.
- [2] M. V. BOROVoi, “Abelianization of the second nonabelian Galois cohomology”, *Duke Math. J.* **72** (1993), no. 1, p. 217-239.
- [3] L. G. BROWN, “Extensions of topological groups”, *Pacific J. Math.* **39** (1971), p. 71-78.
- [4] L. CALABI, “Sur les extensions des groupes topologiques”, *Ann. Mat. Pura Appl.* (4) **32** (1951), p. 295-370.
- [5] M. CEDERWALL, G. FERRETTI, B. E. W. NILSSON & A. WESTERBERG, “Higher-dimensional loop algebras, non-abelian extensions and p -branes”, *Nuclear Phys. B.* **424** (1994), no. 1, p. 97-123.
- [6] S. EILENBERG & S. MACLANE, “Group extensions and homology”, *Ann. of Math.* (2) **43** (1942), p. 757-831.
- [7] ———, “Cohomology theory in abstract groups. II. Group extensions with a non-Abelian kernel”, *Ann. of Math.* (2) **48** (1947), p. 326-341.
- [8] H. GLÖCKNER, “Infinite-dimensional Lie groups without completeness condition”, in *Geometry and analysis on finite- and infinite-dimensional Lie groups*, A. Strasburger et al Eds., vol. 55, Banach Center Publications, Warszawa, 2002, p. 53-59.
- [9] M. GOTO, “On an arcwise connected subgroup of a Lie group”, *Proc. Amer. Math. Soc.* **20** (1969), p. 157-162.

- [10] J. HUEBSCHMANN, “Automorphisms of group extensions and differentials in the Lyndon-Hochschild-Serre spectral sequence”, *J. Algebra* **72** (1981), no. 2, p. 296-334.
- [11] A. KRIEGL & P. W. MICHOR, *The convenient setting of global analysis*, Mathematical Surveys and Monographs, vol. 53, American Mathematical Society, Providence, RI, 1997.
- [12] G. W. MACKEY, “Les ensembles boréliens et les extensions des groupes”, *J. Math. Pures Appl. (9)* **36** (1957), p. 171-178.
- [13] S. MACLANE, *Homological Algebra*, Springer-Verlag, 1963.
- [14] J. MILNOR, “On the existence of a connection with curvature zero”, *Comment. Math. Helv.* **32** (1958), p. 215-223.
- [15] ———, “Remarks on infinite-dimensional Lie groups”, in *Relativity, groups and topology, II (Les Houches, 1983)*, North-Holland, Amsterdam, 1984, p. 1007-1057.
- [16] C. C. MOORE, “Extensions and low dimensional cohomology theory of locally compact groups. I, II”, *Trans. Amer. Math. Soc.* **113** (1964), p. 40-63, 63-86.
- [17] K.-H. NEEB, “Exact sequences for Lie group cohomology with non-abelian coefficients”, in preparation.
- [18] ———, “Central extensions of infinite-dimensional Lie groups”, *Ann. Inst. Fourier (Grenoble)* **52** (2002), no. 5, p. 1365-1442.
- [19] ———, “Abelian extensions of infinite-dimensional Lie groups”, in *Travaux mathématiques. Fasc. XV, Trav. Math., XV, Univ. Luxemb., Luxembourg, 2004*, p. 69-194.
- [20] ———, “Non-abelian extensions of topological Lie algebras”, *Comm. Algebra* **34** (2006), no. 3, p. 991-1041.
- [21] I. RAEBURN, A. SIMS & D. P. WILLIAMS, “Twisted actions and obstructions in group cohomology”, in *C*-algebras (Münster, 1999)*, Springer, Berlin, 2000, p. 161-181.
- [22] D. J. S. ROBINSON, “Automorphisms of group extensions”, in *Algebra and its applications (New Delhi, 1981)*, Lecture Notes in Pure and Appl. Math., vol. 91, Dekker, New York, 1984, p. 163-167.
- [23] O. SCHREIER, “Über die Erweiterungen von Gruppen I”, *Monatshefte f. Math.* **34** (1926), p. 165-180.
- [24] ———, “Über die Erweiterungen von Gruppen II”, *Abhandlungen Hamburg* **4** (1926), p. 321-346.
- [25] A. M. TURING, “The extensions of a group”, *Compos. Math.* **5** (1938), p. 357-367.
- [26] V. S. VARADARAJAN, *Geometry of quantum theory*, second ed., Springer-Verlag, New York, 1985.
- [27] C. A. WEIBEL, *An introduction to homological algebra*, Cambridge Studies in Advanced Mathematics, vol. 38, Cambridge University Press, Cambridge, 1994.
- [28] C. WELLS, “Automorphisms of group extensions”, *Trans. Amer. Math. Soc.* **155** (1971), p. 189-194.

Manuscrit reçu le 19 avril 2005,
 accepté le 23 mars 2006.

Karl-Hermann NEEB
 Technische Universität Darmstadt
 Schlossgartenstrasse 7
 64289 Darmstadt(Deutschland)
 neeb@mathematik.tu-darmstadt.de