



ANNALES DE L'INSTITUT FOURIER

Bharathwaj PALVANNAN

Height one specializations of Selmer groups

Tome 69, n° 1 (2019), p. 303-334.

http://aif.centre-mersenne.org/item/AIF_2019__69_1_303_0

© Association des Annales de l'institut Fourier, 2019,

Certains droits réservés.



Cet article est mis à disposition selon les termes de la licence

CREATIVE COMMONS ATTRIBUTION – PAS DE MODIFICATION 3.0 FRANCE.

<http://creativecommons.org/licenses/by-nd/3.0/fr/>



Les Annales de l'institut Fourier sont membres du

Centre Mersenne pour l'édition scientifique ouverte

www.centre-mersenne.org

HEIGHT ONE SPECIALIZATIONS OF SELMER GROUPS

by Bharathwaj PALVANNAN

ABSTRACT. — We provide applications to studying the behavior of Selmer groups under specialization. We consider Selmer groups associated to 4-dimensional Galois representations coming from

- (i) the tensor product of two cuspidal Hida families F and G ,
- (ii) its cyclotomic deformation,
- (iii) the tensor product of a cusp form f and the Hida family G , where f is a classical specialization of F with weight $k \geq 2$.

We prove control theorems to relate

- (a) the Selmer group associated to the tensor product of Hida families F and G to the Selmer group associated to its cyclotomic deformation, and
- (b) the Selmer group associated to the tensor product of f and G to the Selmer group associated to the tensor product of F and G .

On the analytic side of the main conjectures, Hida has constructed one variable, two variable and three variable Rankin–Selberg p -adic L -functions. Our specialization results enable us to verify that Hida’s results relating

- (a) the two variable p -adic L -function to the three variable p -adic L -function, and
- (b) the one variable p -adic L -function to the two variable p -adic L -function,

and our control theorems for Selmer groups are completely consistent with the main conjectures.

RÉSUMÉ. — Nous donnons des applications de l’étude du comportement des groupes de Selmer sous la spécialisation. Nous considérons les groupes de Selmer associés à de représentations galoisiennes de dimension 4 provenant

- (i) du produit tensoriel de deux familles cuspidales de Hida F et G ,
- (ii) de la déformation cyclotomique du dernier,
- (iii) du produit tensoriel d’une forme cuspidale f par une famille de Hida G , où f est une spécialisation classique de F de poids $k \geq 2$.

Nous démontrons des théorèmes de contrôle qui relient

- (a) le groupe de Selmer associé au produit tensoriel des familles de Hida F et G au groupe de Selmer associé à sa déformation cyclotomique,
- (b) le groupe de Selmer associé au produit tensoriel de f par G au groupe de Selmer associé au produit tensoriel de F et G .

Du côté analytique des conjectures principales, Hida a construit des fonctions L p -adiques de Rankin–Selberg à une variable, à deux variables et à trois variables.

Nos résultats sur la spécialisation nous permettent de vérifier les résultats de Hida qui relie

- (a) la fonction L p -adique à deux variables à la fonction L p -adique à trois variables, et
- (b) la fonction L p -adique à une variables à la fonction L p -adique à deux variables,

et nos théorèmes de contrôle pour les groupes de Selmer sont complètement compatibles avec les conjectures principales.

1. Introduction

Studying the behavior of Selmer groups under specialization has been quite fruitful in the context of the main conjectures in Iwasawa theory. We cite three examples where the topic of specializations of Selmer groups has come up in the past; it has come up in a work of Greenberg [6] to prove a result involving classical Iwasawa modules corresponding to a factorization formula proved by Gross involving Katz's 2-variable p -adic L -function, in the work of Greenberg–Vatsal [11] while studying an elliptic curve with an isogeny of degree p and in an earlier work of ours [28] to prove a result involving Selmer groups corresponding to a factorization formula proved by Dasgupta involving Hida's Rankin–Selberg 3-variable p -adic L -function.

The main results of this paper will deal with Selmer groups to which the Rankin–Selberg p -adic L -functions, constructed by Hida in [16], are associated. The results of Lei–Loeffer–Zerbes [24] and Kings–Loeffer–Zerbes [23] suggest that the proofs of the main conjectures associated to these Rankin–Selberg p -adic L -functions are imminent; the results of this paper provide evidence to these conjectures. The results of this paper contrast well with the three examples cited earlier since in this paper we deal with “critical” specializations (the three examples cited earlier dealt with “non-critical” specializations). After the completion of this project, the author learnt about the paper [12] where the behavior of Selmer groups under certain critical specializations has also been studied. Hara and Ochiai (see Theorem D in [12]) deduce the cyclotomic Iwasawa main conjecture over a CM field E (formulated over a regular local ring with Krull dimension 2) for Hilbert cuspforms with complex multiplication from the main conjecture formulated (over a regular local ring with Krull dimension greater than or equal to $\frac{[E:\mathbb{Q}]}{2} + 2$) by Hida and Tilouine in [21]; the Hida–Tilouine main conjecture has been proved by Ming-Lun Hsieh [22] under certain hypotheses. To place our work involving specializations of Selmer groups in the context of main conjectures, this result of [12] might serve as a useful template for the reader to keep in mind.

To prove our results, we will use the techniques developed in an earlier work of ours [28] (which is briefly summarized in Section 5). The novelty in the techniques introduced in [28] is that the local domains, appearing in Hida theory and over which the main conjectures are formulated, are not always known to be regular or even a UFD. See Section 6 where we provide such examples. These examples are based on the circle of ideas first developed by Hida in [19] and later explored in works of Cho [2] and Cho–Vatsal [3]. In contrast, the main conjectures over CM fields are formulated over regular local rings. One obstacle from the perspective of commutative algebra that thus needs to be overcome is that the kernel of the specialization map (which in our case is a prime ideal of height one) is not known to be principal. See Remark 1.11

Let us introduce some notations. Let $p \geq 5$ be a fixed prime number. Let $F = \sum_{n=1}^{\infty} a_n q^n \in R_F[[q]]$ and $G = \sum_{n=1}^{\infty} b_n q^n \in R_G[[q]]$ be two ordinary cuspidal Hida families. Here, we let R_F and R_G denote the integral closures of the irreducible components of the ordinary primitive cuspidal Hecke algebras through F and G respectively. The rings R_F and R_G turn out to be finite integral extensions of $\mathbb{Z}_p[[x_F]]$ and $\mathbb{Z}_p[[x_G]]$ respectively, where x_F and x_G are “weight variables” for F and G respectively. We assume the following hypotheses on F and G :

IRR. — The residual representations associated to F and G are absolutely irreducible.

p -DIS-IN. — The restrictions, to the inertia subgroup I_p at p , of the residual representations associated to F and G have non-scalar semi-simplifications.

Let Σ be a finite set of primes in \mathbb{Q} containing the primes p, ∞ , a finite prime $l_0 \neq p$ and all the primes dividing the levels of F and G . Let $\Sigma_0 = \Sigma \setminus \{p\}$. Let O denote the ring of integers in a finite extension of \mathbb{Q}_p . Let \mathbb{Q}_Σ denote the maximal extension of \mathbb{Q} unramified outside Σ . Let G_Σ denote $\text{Gal}(\mathbb{Q}_\Sigma/\mathbb{Q})$. We have Galois representations $\rho_F : G_\Sigma \rightarrow \text{GL}_2(R_F)$ and $\rho_G : G_\Sigma \rightarrow \text{GL}_2(R_G)$ associated to F and G respectively (see [14]). We let L_F (and L_G) be the free R_F -module (and R_G -module) of rank 2 on which G_Σ acts to let us obtain the Galois representation ρ_F (and ρ_G respectively). Assume that the integral closures of \mathbb{Z}_p in R_F and R_G are equal (extending scalars if necessary)⁽¹⁾ to O .

⁽¹⁾The ring $R_{F,G}$ turns out to be a domain. This is because the completed tensor product $R_{F,G}$ can be shown to be isomorphic to the tensor product $R_F[[x_G]] \otimes_{O[[x_G]]} R_G$. The fraction field of R_G is a finite field extension of the fraction field of $O[[x_G]]$, while

1.1. Specializing from 3-variables to 2-variables

Let $R_{F,G}$ denote the completed tensor product $R_F \widehat{\otimes} R_G$. The completed tensor product is the co-product in the category of complete semi-local Noetherian O -algebras (where the morphisms are continuous O -algebra maps). We have the natural inclusions $i_F : R_F \hookrightarrow R_{F,G}$ and $i_G : R_G \hookrightarrow R_{F,G}$. One can construct a 4-dimensional Galois representation $\rho_{F,G} : G_\Sigma \rightarrow \mathrm{GL}_4(R_{F,G})$, that is given by the action of G_Σ on the following free $R_{F,G}$ -module of rank 4:

$$L_2 := \mathrm{Hom}_{R_{F,G}}(L_F \otimes_{R_F} R_{F,G}, L_G \otimes_{R_G} R_{F,G}).$$

Let $\rho_{F,G} \otimes \kappa^{-1} : G_\Sigma \rightarrow \mathrm{GL}_4(R_{F,G}[[\Gamma]])$ denote the Galois representation given by the action of G_Σ on the following free $R_{F,G}[[\Gamma]]$ -module of rank 4:

$$L_3 := L_2 \otimes_{R_{F,G}} R_{F,G}[[\Gamma]](\kappa^{-1}).$$

We let \mathbb{Q}_∞ denote the cyclotomic \mathbb{Z}_p -extension of \mathbb{Q} and we let Γ denote the Galois group $\mathrm{Gal}(\mathbb{Q}_\infty/\mathbb{Q})$. Here, $\kappa : G_\Sigma \twoheadrightarrow \Gamma \hookrightarrow \mathbb{Z}_p[[\Gamma]]^\times$ denotes the tautological character. To simplify notations, we will let ρ_2 denote $\rho_{F,G}$ and ρ_3 denote $\rho_{F,G} \otimes \kappa^{-1}$. Hida has constructed elements $\theta_2^{\Sigma_0}$ and $\theta_3^{\Sigma_0}$, in the fraction fields of $R_{F,G}$ and $R_{F,G}[[\Gamma]]$ respectively, as generators for non-primitive p -adic L -functions associated to ρ_2 and ρ_3 respectively. See [17, §7.4 and 10.4] for their constructions and the precise interpolation properties that they satisfy. The elements $\theta_2^{\Sigma_0}$ and $\theta_3^{\Sigma_0}$ generate a 2-variable p -adic L -function and 3-variable p -adic L -function respectively. For the 2-variable p -adic L -function, one can vary the weights of F and G . For the 3-variable p -adic L -function, in addition to the weights of F and G , one can also vary the cyclotomic variable. Notice that the subscripts in our notation for the Galois representations or the associated Galois lattices indicate the number of variables in the corresponding p -adic L -functions.

Let $\pi_{3,2} : R_{F,G}[[\Gamma]] \rightarrow R_{F,G}$ be the map defined by sending every element of Γ to 1. We have the Galois representation $\pi_{3,2} \circ \rho_3 : G_\Sigma \xrightarrow{\rho_3} \mathrm{GL}_4(R_{F,G}[[\Gamma]]) \xrightarrow{\pi_{3,2}} \mathrm{GL}_4(R_{F,G})$, along with the following isomorphism of Galois representations:

$$(1.1) \quad \pi_{3,2} \circ \rho_3 \cong \rho_2.$$

Hida has proved the following theorem relating $\theta_2^{\Sigma_0}$ and $\theta_3^{\Sigma_0}$. See [17, §10.5, Thm. 1] and [16, Thm. 5.1d’].

the fraction field of $R_F[[x_G]]$ is a purely transcendental extension of the fraction field of $O[[x_G]]$. Since the fraction fields of $R_F[[x_G]]$ and R_G are “linearly disjoint over the fraction field of $O[[x_G]]$ ”, the tensor product $R_F[[x_G]] \otimes_{O[[x_G]]} R_G$ (and hence $R_{F,G}$) turns out to be a domain.

THEOREM 1.1 (Hida). — $\pi_{3,2} \left(\theta_3^{\Sigma_0} \right) = \left(1 - \frac{i_F(a_p)}{i_G(b_p)} \right) \cdot \theta_2^{\Sigma_0}$.

Recall that the ordinarity assumption on F and G ensures us that the coefficients $i_F(a_p)$ and $i_G(b_p)$ would be units in the ring $R_{F,G}$. On the algebraic side, we can construct Selmer groups $\text{Sel}_{\rho_2}^{\Sigma_0}(\mathbb{Q})$ and $\text{Sel}_{\rho_3}^{\Sigma_0}(\mathbb{Q})$ associated to the Galois representations ρ_2 and ρ_3 respectively. We prove the following control theorem relating $\text{Sel}_{\rho_2}^{\Sigma_0}(\mathbb{Q})$ and $\text{Sel}_{\rho_3}^{\Sigma_0}(\mathbb{Q})$:

THEOREM 1.2. — *We have the following equality in the divisor group of $R_{F,G}$:*

$$\begin{aligned} \text{Div} \left(\text{Sel}_{\rho_3}^{\Sigma_0}(\mathbb{Q})^\vee \otimes_{R_{F,G}[[\Gamma]]} R_{F,G} \right) + \text{Div} \left(H^0(G_\Sigma, D_{\rho_3})^\vee [\ker(\pi_{3,2})] \right) \\ = \text{Div} \left(\text{Sel}_{\rho_2}^{\Sigma_0}(\mathbb{Q})^\vee \right) + \text{Div} \left(1 - \frac{i_F(a_p)}{i_G(b_p)} \right). \end{aligned}$$

Let us explain some of our notations. If A is a profinite ring, we let \hat{A} denote the Pontryagin dual of A . If M is a continuous A -module, we let M^\vee denote its Pontryagin dual. To a finitely generated torsion module over an integrally closed Noetherian local domain or a non-zero element of its fraction field, we shall associate an element of the divisor group following [7]. If M is an A -module and I is an ideal of A , the A/I -module $M[I]$ is defined to be the set $\{m \in M \mid i \cdot m = 0, \forall i \in I\}$. The discrete modules D_{ρ_3} , D_{ρ_2} and D_{ρ_1} will be precisely defined in Section 2. We will assume throughout this paper that the $R_{F,G}[[\Gamma]]$ -module $\text{Sel}_{\rho_3}^{\Sigma_0}(\mathbb{Q})^\vee$ and the $R_{F,G}$ -module $\text{Sel}_{\rho_2}^{\Sigma_0}(\mathbb{Q})^\vee$ are torsion.

One can formulate a main conjecture in Iwasawa theory for ρ_2 and ρ_3 following [7]. The main conjecture for ρ_3 predicts the following equality of divisors in $R_{F,G}[[\Gamma]]$:

$$\text{(MC-}\rho_3) \quad \text{Div} \left(\theta_3^{\Sigma_0} \right) \stackrel{?}{=} \text{Div} \left(\text{Sel}_{\rho_3}^{\Sigma_0}(\mathbb{Q})^\vee \right) - \text{Div} \left(H^0(G_\Sigma, D_{\rho_3})^\vee \right).$$

The main conjecture for ρ_2 predicts the following equality of divisors in $R_{F,G}$:

$$\text{(MC-}\rho_2) \quad \text{Div} \left(\theta_2^{\Sigma_0} \right) \stackrel{?}{=} \text{Div} \left(\text{Sel}_{\rho_2}^{\Sigma_0}(\mathbb{Q})^\vee \right) - \text{Div} \left(H^0(G_\Sigma, D_{\rho_2})^\vee \right).$$

We prove the following theorem to show that Theorem 1.1 and Theorem 1.2 are completely consistent with the main conjectures (MC- ρ_3) and (MC- ρ_2):

THEOREM 1.3. — *If the main conjecture (MC- ρ_3) holds, then the main conjecture (MC- ρ_2) also holds.*

Remark 1.4. — As we will show in equation (3.2), we have the following isomorphism of $R_{F,G}$ -modules:

$$(1.2) \quad \mathrm{Tor}_1^{R_{F,G}[\Gamma]}(R_{F,G}, H^0(G_\Sigma, D_{\rho_3})^\vee) \cong H^0(G_\Sigma, D_{\rho_3})^\vee[\ker(\pi_{3,2})].$$

Regarding Theorem 1.2, it will be useful to keep this isomorphism in mind. Since we invoke results from homological algebra, we will need to use the Tor functor frequently.

1.2. Specializing from 2-variables to 1-variable

Let f be a cuspidal p -stabilized eigenform obtained by specializing F at a classical height one prime ideal P_k of weight $k \geq 2$. Let $\rho_f : G_\Sigma \rightarrow \mathrm{GL}_2(O_f)$ denote the Galois representation associated to f (constructed by Deligne). Here, O_f denotes the integral closure of $\frac{R_F}{P_k}$. We also obtain a natural map $\pi_f : R_F \rightarrow O_f$. We let L_f denote the free O_f -module of rank 2 on which G_Σ acts to let us obtain the Galois representation ρ_f . Let $R_{f,G}$ be the completed tensor product $O_f \widehat{\otimes} R_G$. We have natural maps $j_F : R_F \rightarrow O_f \hookrightarrow R_{f,G}$ and $j_G : R_G \hookrightarrow R_{f,G}$. Consider the 4-dimensional Galois representation $\rho_1 : G_\Sigma \rightarrow \mathrm{GL}_4(R_{f,G})$ given by the action of G_Σ on the rank 4 free $R_{f,G}$ -module:

$$L_1 := \mathrm{Hom}_{R_{f,G}}(L_F \otimes_{R_F} R_{f,G}, L_G \otimes_{R_G} R_{f,G}).$$

Hida has also constructed an element $\theta_1^{\Sigma_0}$, in the fraction field of $R_{f,G}$, that generates a non-primitive 1-variable Rankin–Selberg p -adic L -function associated to ρ_1 . See [17, §7.4] for its construction and the precise interpolation properties that $\theta_1^{\Sigma_0}$ satisfies. For the 1-variable p -adic L -function, one can vary the weight of G . We have a natural map $\pi_{2,1} : R_{F,G} \rightarrow R_{f,G}$ obtained by the maps $j_F : R_F \rightarrow O_f \hookrightarrow R_{f,G}$ and $j_G : R_G \hookrightarrow R_{f,G}$. The map $\pi_{2,1} : R_{F,G} \rightarrow R_{f,G}$ lets us obtain the following isomorphism of Galois representations:

$$(1.3) \quad \pi_{2,1} \circ \rho_2 \cong \rho_1.$$

Hida has proved following theorem relating $\theta_2^{\Sigma_0}$ and $\theta_1^{\Sigma_0}$ (see the comment just before Theorem 2 in [17, §7.4]):

THEOREM 1.5 (Hida). — $\pi_{2,1}(\theta_2^{\Sigma_0}) = \theta_1^{\Sigma_0}$.

Associated to ρ_1 , we can construct the Selmer group $\mathrm{Sel}_{\rho_1}^{\Sigma_0}(\mathbb{Q})$. We will assume that $\mathrm{Sel}_{\rho_1}^{\Sigma_0}(\mathbb{Q})^\vee$ is a torsion $R_{f,G}$ -module. We prove the following control theorem relating $\mathrm{Sel}_{\rho_1}^{\Sigma_0}(\mathbb{Q})$ and $\mathrm{Sel}_{\rho_2}^{\Sigma_0}(\mathbb{Q})$:

THEOREM 1.6. — We have the following equality in the divisor group of $R_{f,G}$:

$$\begin{aligned} \text{Div} \left(\text{Sel}_{\rho_2}^{\Sigma_0}(\mathbb{Q})^\vee \otimes_{R_{f,G}} R_{f,G} \right) + \text{Div} \left(\text{Tor}_1^{R_{f,G}}(R_{f,G}, H^0(G_\Sigma, D_{\rho_2})^\vee) \right) \\ = \text{Div} \left(\text{Sel}_{\rho_1}^{\Sigma_0}(\mathbb{Q})^\vee \right). \end{aligned}$$

The main conjecture for ρ_1 predicts the following equality of divisors in $R_{f,G}$:

$$\text{(MC-}\rho_1) \quad \text{Div} \left(\theta^{\Sigma_0} \right) \stackrel{?}{=} \text{Div} \left(\text{Sel}_{\rho_1}^{\Sigma_0}(\mathbb{Q})^\vee \right) - \text{Div} \left(H^0(G_\Sigma, D_{\rho_1})^\vee \right).$$

We prove the following theorem to show that Theorem 1.5 and Theorem 1.6 are completely consistent with the main conjectures (MC- ρ_2) and (MC- ρ_1):

THEOREM 1.7. — If the main conjecture (MC- ρ_2) holds, then the main conjecture (MC- ρ_1) also holds.

Remark 1.8. — One can combine Proposition 5.3 and Lemma 5.4 to obtain the following equality of divisors in $R_{f,G}$:

$$\begin{aligned} (1.4) \quad \text{Div} \left(\text{Tor}_1^{R_{f,G}}(R_{f,G}, H^0(G_\Sigma, D_{\rho_2})^\vee) \right) \\ = \text{Div} \left(\left(H^0(G_\Sigma, D_{\rho_2})^\vee[\ker(\pi_{2,1})] \right) \otimes_{\frac{R_{f,G}}{\ker(\pi_{2,1})}} R_{f,G} \right), \end{aligned}$$

which is similar to the one obtained in equation (1.2). As mentioned in Remark 1.4, we prefer stating Theorem 1.6 using the term on the left hand side of equation (1.4) involving the Tor functor (instead of the term appearing on the right hand side).

Remark 1.9. — The main conjectures, formulated in [7], relate the primitive Selmer groups to the primitive p -adic L -functions. It is possible to relate the primitive Selmer groups to the non-primitive Selmer groups; it is also possible to relate the primitive p -adic L -functions to the non-primitive p -adic L -functions. As a result, it will be possible to formulate a conjecture, that is equivalent to the one formulated in [7], relating the non-primitive Selmer groups to the non-primitive p -adic L -functions. It is the latter formulation that we adopt in this paper. Since we are motivated by Hida’s results involving the non-primitive p -adic L -functions, we will only work with the non-primitive Selmer groups; and since Hida’s results simply serve as heuristics to motivate our theorems, we will not concern ourselves with proving that Greenberg’s formulation of the main conjectures for ρ_3 , ρ_2 and ρ_1 involving primitive Selmer groups and primitive p -adic L -functions in [7] are equivalent to (MC- ρ_3), (MC- ρ_2), (MC- ρ_1) respectively.

Remark 1.10. — We would like make the reader aware of a point that was conveyed to us by Haruzo Hida. The p -adic L -functions $\theta_1^{\Sigma_0}$, $\theta_2^{\Sigma_0}$ and $\theta_3^{\Sigma_0}$ are “genuine” in the sense of [18] and do not exactly equal the p -adic L -functions constructed in [17]. Recall that we have inclusions $R_G \hookrightarrow R_{f,G}$ and $R_G \hookrightarrow R_{F,G} \hookrightarrow R_{F,G}[[\Gamma]]$. There is an element $\Theta \in R_G$ that corresponds to the divisor of a one-variable p -adic L -function associated to the 3-dimensional adjoint representation $\text{Ad}^0(\rho_G)$ (see Conjecture 1.0.1 in [18]). The p -adic L -functions, that are constructed in [17] as elements of the fraction fields of $R_{f,G}$, $R_{F,G}$ and $R_{F,G}[[\Gamma]]$ respectively, are in fact equal to $\frac{\theta_1^{\Sigma_0}}{\Theta}$, $\frac{\theta_2^{\Sigma_0}}{\Theta}$ and $\frac{\theta_3^{\Sigma_0}}{\Theta}$ respectively. A discussion surrounding the need to introduce this modification, which is related to the choice of a certain period (Neron period versus a period involving the Peterson inner product), is carefully explained in [18] (see the introduction and Section 6 there). This point will not matter to us since the maps $\pi_{3,2}$ and $\pi_{2,1}$ are R_G -linear.

Remark 1.11. — The obstacle, presented by the fact the local domains appearing in Hida theory need not even be UFDs, manifests itself in the following way. Although one is interested in results, such as the one in Theorem 1.2 (and Theorem 1.6 respectively), in the divisor group of $R_{F,G}$ (and $R_{f,G}$ respectively), one is forced to consider the localizations of $\text{Sel}_{\rho_3}^{\Sigma_0}(\mathbb{Q})^\vee$ (and $\text{Sel}_{\rho_2}^{\Sigma_0}(\mathbb{Q})^\vee$ respectively) at certain height two prime ideals of $R_{F,G}[[\Gamma]]$ (and $R_{F,G}$ respectively).

Remark 1.12. — We always work with the hypothesis (labeled TOR in Section 2) that the $R_{F,G}[[\Gamma]]$ -module $\text{Sel}_{\rho_3}^{\Sigma_0}(\mathbb{Q})^\vee$, the $R_{F,G}$ -module $\text{Sel}_{\rho_2}^{\Sigma_0}(\mathbb{Q})^\vee$ and the $R_{f,G}$ -module $\text{Sel}_{\rho_1}^{\Sigma_0}(\mathbb{Q})^\vee$ are torsion. To establish the hypothesis TOR, one can utilize the techniques involving Euler systems developed by Lei–Loeffler–Zerbes [24] and Kings–Loeffler–Zerbes [23]. We refer the reader to [24] and [23] for the precise conditions when the Euler system machinery is available. Under these conditions, to establish TOR, one has to show that Hida’s Rankin–Selberg p -adic L -functions $\theta_3^{\Sigma_0}$, $\theta_2^{\Sigma_0}$ and $\theta_1^{\Sigma_0}$ are non-zero. The non-vanishing of these p -adic L -functions is a recent work in progress of Jeanine Van Order. See [26] and [27].

It will be possible to obtain results, consistent with the various main conjectures, without assuming TOR. We briefly indicate how to obtain such results, though we leave the precise details to the interested reader. Suppose the $R_{F,G}$ -module $\text{Sel}_{\rho_2}^{\Sigma_0}(\mathbb{Q})^\vee$ is not torsion. The Iwasawa main conjecture predicts that $\theta_2^{\Sigma_0}$ equals zero. On the analytic side, Hida’s theorem (Theorem 1.1) predicts that $\pi_{3,2}(\theta_3^{\Sigma_0})$ equals zero. On the algebraic side, one can

establish a control theorem to show that $\ker(\pi_{3,2})$ belongs to the support of the $R_{F,G}[[\Gamma]]$ -module $\text{Sel}_{\rho_3}^{\Sigma_0}(\mathbb{Q})^\vee$.

Suppose the $R_{f,G}$ -module $\text{Sel}_{\rho_1}^{\Sigma_0}(\mathbb{Q})^\vee$ is not torsion. The Iwasawa main conjecture predicts that $\theta_1^{\Sigma_0}$ equals zero. On the analytic side, Hida’s theorem (Theorem 1.5) predicts that $\pi_{2,1}(\theta_2^{\Sigma_0})$ equals zero. On the algebraic side, one can similarly establish a control theorem to show that $\ker(\pi_{2,1})$ belongs to the support of the $R_{F,G}$ -module $\text{Sel}_{\rho_2}^{\Sigma_0}(\mathbb{Q})^\vee$.

Acknowledgements. We are greatly indebted to Ralph Greenberg. This project was completed while the author was his student at University of Washington. We would like to thank Haruzo Hida for indicating to us how to compute the examples in Section 6. We are grateful to Antonio Lei, Jeanine Van Order and the referee for their helpful comments.

2. Selmer groups

The hypothesis p -DIS-IN lets us obtain following $\text{Gal}(\overline{\mathbb{Q}}_p/\mathbb{Q}_p)$ -equivariant short exact sequences of free R_F and R_G modules respectively:

$$(2.1) \quad \begin{aligned} 0 \rightarrow \underbrace{\text{Fil}^+ L_F}_{\text{Rank}=1} \rightarrow \underbrace{L_F}_{\text{Rank}=2} \rightarrow \underbrace{\frac{L_F}{\text{Fil}^+ L_F}}_{\text{Rank}=1} \rightarrow 0, \\ 0 \rightarrow \underbrace{\text{Fil}^+ L_G}_{\text{Rank}=1} \rightarrow \underbrace{L_G}_{\text{Rank}=2} \rightarrow \underbrace{\frac{L_G}{\text{Fil}^+ L_G}}_{\text{Rank}=1} \rightarrow 0. \end{aligned}$$

Here, the action of $\text{Gal}(\overline{\mathbb{Q}}_p/\mathbb{Q}_p)$ on the rank 1 modules $\text{Fil}^+ L_F$ and $\text{Fil}^+ L_G$ is given by ramified characters $\delta_F : \text{Gal}(\overline{\mathbb{Q}}_p/\mathbb{Q}_p) \rightarrow R_F^\times$ and $\delta_G : \text{Gal}(\overline{\mathbb{Q}}_p/\mathbb{Q}_p) \rightarrow R_G^\times$ respectively. The action of $\text{Gal}(\overline{\mathbb{Q}}_p/\mathbb{Q}_p)$ on the rank 1 modules $\frac{L_F}{\text{Fil}^+ L_F}$ and $\frac{L_G}{\text{Fil}^+ L_G}$ is given by unramified characters $\epsilon_F : \text{Gal}(\overline{\mathbb{Q}}_p/\mathbb{Q}_p) \rightarrow R_F^\times$ and $\epsilon_G : \text{Gal}(\overline{\mathbb{Q}}_p/\mathbb{Q}_p) \rightarrow R_G^\times$ respectively. Note that $\epsilon_F(\text{Frob}_p) = a_p$ and $\epsilon_G(\text{Frob}_p) = b_p$. Furthermore, using the residual characters $\bar{\delta}_F : \text{Gal}(\overline{\mathbb{Q}}_p/\mathbb{Q}_p) \rightarrow \overline{\mathbb{F}}_p^\times$ and $\bar{\delta}_G : \text{Gal}(\overline{\mathbb{Q}}_p/\mathbb{Q}_p) \rightarrow \overline{\mathbb{F}}_p^\times$ associated to δ_F and δ_G respectively, the hypothesis p -DIS-IN can be restated in the following way:

$$(p\text{-DIS-IN}) \quad \bar{\delta}_F|_{I_p} \neq \mathbf{1}, \quad \bar{\delta}_G|_{I_p} \neq \mathbf{1}.$$

Here, I_p denotes the inertia subgroup inside the decomposition group $\text{Gal}(\overline{\mathbb{Q}}_p/\mathbb{Q}_p)$. Analogous to the short exact sequences in (2.1), we can now

form the following $\text{Gal}(\overline{\mathbb{Q}}_p/\mathbb{Q}_p)$ -equivariant short exact sequence of free $R_{F,G}[[\Gamma]]$ -modules:

$$0 \rightarrow \underbrace{\text{Fil}^+ L_3}_{\text{Rank}=2} \rightarrow \underbrace{L_3}_{\text{Rank}=4} \rightarrow \underbrace{\frac{L_3}{\text{Fil}^+ L_3}}_{\text{Rank}=2} \rightarrow 0,$$

where

$$\text{Fil}^+ L_3 := \text{Hom}_{R_{F,G}}(L_F \otimes_{R_F} R_{F,G}, \text{Fil}^+ L_G \otimes_{R_G} R_{F,G}) \otimes_{R_{F,G}} R_{F,G}[[\Gamma]](\kappa^{-1}).$$

We would like to emphasize the following isomorphisms of $R_{F,G}$ and $R_{f,G}$ -modules respectively.

$$(2.2) \quad L_2 \cong L_3 \otimes_{R_{F,G}[[\Gamma]]} R_{F,G}, \quad L_1 \cong L_2 \otimes_{R_{F,G}} R_{f,G}.$$

Once again, analogous to the short exact sequences in (2.1), we can now form the following $\text{Gal}(\overline{\mathbb{Q}}_p/\mathbb{Q}_p)$ -equivariant short exact sequence of free $R_{F,G}$ and $R_{f,G}$ modules respectively:

$$\begin{aligned} 0 \rightarrow \underbrace{\text{Fil}^+ L_2}_{\text{Rank}=2} \rightarrow \underbrace{L_2}_{\text{Rank}=4} \rightarrow \underbrace{\frac{L_2}{\text{Fil}^+ L_2}}_{\text{Rank}=2} \rightarrow 0, \\ 0 \rightarrow \underbrace{\text{Fil}^+ L_1}_{\text{Rank}=2} \rightarrow \underbrace{L_1}_{\text{Rank}=4} \rightarrow \underbrace{\frac{L_1}{\text{Fil}^+ L_1}}_{\text{Rank}=2} \rightarrow 0, \end{aligned}$$

where

$$\text{Fil}^+ L_2 := \text{Fil}^+ L_3 \otimes_{R_{F,G}[[\Gamma]]} R_{F,G}, \quad \text{Fil}^+ L_1 := \text{Fil}^+ L_2 \otimes_{R_{F,G}} R_{f,G}.$$

To the Galois representations ρ_3 , ρ_2 and ρ_1 , we can now associate the following discrete modules:

$$\begin{aligned} D_{\rho_3} &= L_3 \otimes_{R_{F,G}[[\Gamma]]} \widehat{R_{F,G}[[\Gamma]]}, & \text{Fil}^+ D_{\rho_3} &= \text{Fil}^+ L_3 \otimes_{R_{F,G}[[\Gamma]]} \widehat{R_{F,G}[[\Gamma]]} \\ D_{\rho_2} &= L_2 \otimes_{R_{F,G}} \widehat{R_{F,G}}, & \text{Fil}^+ D_{\rho_2} &= \text{Fil}^+ L_2 \otimes_{R_{F,G}} \widehat{R_{F,G}} \\ D_{\rho_1} &= L_1 \otimes_{R_{f,G}} \widehat{R_{f,G}}, & \text{Fil}^+ D_{\rho_1} &= \text{Fil}^+ L_1 \otimes_{R_{f,G}} \widehat{R_{f,G}}. \end{aligned}$$

We now define the (discrete) non-primitive Selmer groups associated to the Galois representations ρ_3 , ρ_2 and ρ_1 respectively. Since we do not

discuss primitive Selmer groups in this paper, we will refer to the non-primitive Selmer groups simply as Selmer groups (without the prefix “non-primitive”). Let $i \in \{1, 2, 3\}$.

$$\text{Sel}_{\rho_i}^{\Sigma_0}(\mathbb{Q}) := \ker \left(H^1(G_\Sigma, D_{\rho_i}) \xrightarrow{\phi_{\rho_i}^{\Sigma_0}} H^1 \left(I_p, \frac{D_{\rho_i}}{\text{Fil}^+ D_{\rho_i}} \right)^{\Gamma_p} \right).$$

Here, $\Gamma_p = \frac{\text{Gal}(\overline{\mathbb{Q}_p}/\mathbb{Q}_p)}{I_p}$. Throughout this paper, we will suppose that the following hypothesis holds.

$$\begin{aligned} \text{TOR. —} \quad & \text{Rank}_{R_{F,G}[\Gamma]} \left(\text{Sel}_{\rho_3}^{\Sigma_0}(\mathbb{Q})^\vee \right) = 0, \\ & \text{Rank}_{R_{F,G}} \left(\text{Sel}_{\rho_2}^{\Sigma_0}(\mathbb{Q})^\vee \right) = 0, \\ & \text{Rank}_{R_{f,G}} \left(\text{Sel}_{\rho_1}^{\Sigma_0}(\mathbb{Q})^\vee \right) = 0. \end{aligned}$$

Let us now discuss the proofs of Theorems 1.2 and 1.3 along with the proofs of Theorems 1.6 and 1.7. These proofs will utilize the control theorems and specialization results developed in our earlier work [28]. The control theorems are developed in [28, §6] (see [28, Prop. 6.2]). The specialization results are developed [28, Section 5] (see [28, Prop. 5.2]). We will treat those results as black boxes. We summarize the results of [28] in Section 5.

3. Proofs of Theorem 1.2 and 1.3

One can use Proposition 6.2 in [28, §6] to establish a control theorem relating $\text{Sel}_{\rho_3}^{\Sigma_0}(\mathbb{Q})$ to $\text{Sel}_{\rho_2}^{\Sigma_0}(\mathbb{Q})$ (see Section 5). We have the following equality of divisors in $R_{F,G}$:

$$\begin{aligned} (3.1) \quad & \text{Div} \left(\text{Sel}_{\rho_3}^{\Sigma_0}(\mathbb{Q})^\vee \otimes_{R_{F,G}[\Gamma]} R_{F,G} \right) \\ & + \text{Div} \left(\text{Tor}_1^{R_{F,G}[\Gamma]} \left(R_{F,G}, H^0(G_{\Sigma, D_{\rho_3}})^\vee \right) \right) \\ & = \text{Div} \left(\text{Sel}_{\rho_2}^{\Sigma_0}(\mathbb{Q})^\vee \right) \\ & + \text{Div} \left(\text{Tor}_1^{R_{F,G}[\Gamma]} \left(R_{F,G}, H^0 \left(I_p, \frac{D_{\rho_3}}{\text{Fil}^+ D_{\rho_3}} \right)^{\vee}_{\Gamma_p} \right) \right). \end{aligned}$$

We will view $R_{F,G}$ as an $R_{F,G}[\Gamma]$ -module via the map $\pi_{3,2} : R_{F,G}[\Gamma] \rightarrow R_{F,G}$. As a topological group, $\Gamma \cong \mathbb{Z}_p$. Let’s choose a topological generator γ_0 of Γ . The prime ideal $\ker(\pi_{3,2})$ is generated by $\gamma_0 - 1$ as an

$R_{F,G}[[\Gamma]]$ -module. Consider the short exact sequence $0 \rightarrow R_{F,G}[[\Gamma]] \xrightarrow{\gamma_0^{-1}} R_{F,G}[[\Gamma]] \rightarrow R_{F,G} \rightarrow 0$ of $R_{F,G}[[\Gamma]]$ -modules. Taking the tensor product with the $R_{F,G}[[\Gamma]]$ -module $H^0(G_\Sigma, D_{\rho_3})^\vee$ over the ring $R_{F,G}[[\Gamma]]$ gives us following isomorphism:

$$(3.2) \quad \text{Tor}_1^{R_{F,G}[[\Gamma]]} (R_{F,G}, H^0(G_\Sigma, D_{\rho_3})^\vee) \cong H^0(G_\Sigma, D_{\rho_3})^\vee [\ker(\pi_{3,2})].$$

Lemma 5.8 establishes the following isomorphism of $R_{F,G}$ -modules:

$$(3.3) \quad \text{Tor}_1^{R_{F,G}[[\Gamma]]} \left(R_{F,G}, H^0 \left(I_p, \frac{D_{\rho_3}}{\text{Fil}^+ D_{\rho_3}} \right)_{\Gamma_p}^\vee \right) \cong \frac{R_{F,G}}{1 - \frac{i_F(a_p)}{i_G(b_p)}}.$$

Combining equations (3.1), (3.2), and (3.3) completes the proof of Theorem 1.2.

Now suppose that the main conjecture (MC- ρ_3) holds. That is, we have the following equality in the divisor group of $R_{F,G}[[\Gamma]]$:

$$(MC-\rho_3) \quad \text{Div} \left(\theta_3^{\Sigma_0} \right) = \text{Div} \left(\text{Sel}_{\rho_3}^{\Sigma_0}(\mathbb{Q})^\vee \right) - \text{Div} \left(H^0(G_\Sigma, D_{\rho_3})^\vee \right).$$

We will later outline in Section 5 how the specialization result ([28, §5, Prop. 5.2]) lets us obtain the following equality of divisors in $R_{F,G}$:

$$(3.4) \quad \begin{aligned} \text{Div} \left(\pi_{3,2} \left(\theta_3^{\Sigma_0} \right) \right) &= \text{Div} \left(\text{Sel}_{\rho_3}^{\Sigma_0}(\mathbb{Q})^\vee \otimes_{R_{F,G}[[\Gamma]]} R_{F,G} \right) \\ &\quad + \text{Div} \left(\text{Tor}_1^{R_{F,G}[[\Gamma]]} (R_{F,G}, H^0(G_\Sigma, D_{\rho_3})^\vee) \right) \\ &\quad - \text{Div} \left(H^0(G_\Sigma, D_{\rho_2})^\vee \right). \end{aligned}$$

Theorem 1.1 establishes the following equality in the divisor group of $R_{F,G}$:

$$(3.5) \quad \text{Div} \left(\pi_{3,2} \left(\theta_3^{\Sigma_0} \right) \right) = \text{Div} \left(\theta_{\rho_2}^{\Sigma_0} \right) + \text{Div} \left(1 - \frac{i_F(a_p)}{i_G(b_p)} \right).$$

Theorem 1.2 along with equation (3.2) establishes the following equality in the divisor group of $R_{F,G}$:

$$(3.6) \quad \begin{aligned} \text{Div} \left(\text{Sel}_{\rho_3}^{\Sigma_0}(\mathbb{Q})^\vee \otimes_{R_{F,G}[[\Gamma]]} R_{F,G} \right) \\ + \text{Div} \left(\text{Tor}_1^{R_{F,G}[[\Gamma]]} (R_{F,G}, H^0(G_\Sigma, D_{\rho_3})^\vee) \right) \\ = \text{Div} \left(\text{Sel}_{\rho_2}^{\Sigma_0}(\mathbb{Q})^\vee \right) + \text{Div} \left(1 - \frac{i_F(a_p)}{i_G(b_p)} \right). \end{aligned}$$

Now combining equations (3.4), (3.5) and (3.6), we have the following equality of divisors in $R_{F,G}$:

$$(MC-\rho_2) \quad \text{Div} \left(\theta_2^{\Sigma_0} \right) = \text{Div} \left(\text{Sel}_{\rho_2}^{\Sigma_0}(\mathbb{Q})^\vee \right) - \text{Div} \left(H^0(G_\Sigma, D_{\rho_2})^\vee \right).$$

This completes the proof of Theorem 1.3.

4. Proofs of Theorem 1.6 and 1.7

The proofs of Theorem 1.6 and 1.7 will be similar to the proofs of Theorem 1.2 and 1.3. Proposition 6.2 in [28, §6] will allow us to establish a control theorem relating $\text{Sel}_{\rho_2}^{\Sigma_0}(\mathbb{Q})^\vee \otimes_{R_{F,G}} R_{f,G}$ to $\text{Sel}_{\rho_1}^{\Sigma_0}(\mathbb{Q})^\vee$. We have the following equality of divisors in $R_{f,G}$:

$$\begin{aligned}
 (4.1) \quad & \text{Div} \left(\text{Sel}_{\rho_2}^{\Sigma_0}(\mathbb{Q})^\vee \otimes_{R_{F,G}} R_{f,G} \right) \\
 & + \text{Div} \left(\text{Tor}_1^{R_{F,G}} \left(R_{f,G}, H^0(G_{\Sigma, D_{\rho_2}})^\vee \right) \right) \\
 & = \text{Div} \left(\text{Sel}_{\rho_1}^{\Sigma_0}(\mathbb{Q})^\vee \right) \\
 & + \text{Div} \left(\text{Tor}_1^{R_{F,G}} \left(R_{f,G}, H^0 \left(I_p, \frac{D_{\rho_2}}{\text{Fil}^+ D_{\rho_2}} \right)^\vee \right)_{\Gamma_p} \right).
 \end{aligned}$$

Lemma 5.9 establishes the following equality:

$$(4.2) \quad \text{Tor}_1^{R_{F,G}} \left(R_{f,G}, H^0 \left(I_p, \frac{D_{\rho_2}}{\text{Fil}^+ D_{\rho_2}} \right)^\vee \right) = 0.$$

Combining equations (4.1) and (4.2) completes the proof of Theorem 1.6.

Now suppose that the main conjecture (MC- ρ_2) holds. That is, we have the following equality in the divisor group of $R_{F,G}$:

$$(\text{MC-}\rho_2) \quad \text{Div} \left(\theta_2^{\Sigma_0} \right) = \text{Div} \left(\text{Sel}_{\rho_2}^{\Sigma_0}(\mathbb{Q})^\vee \right) - \text{Div} \left(H^0(G_{\Sigma, D_{\rho_2}})^\vee \right).$$

As before, we will later outline in Section 5 how the specialization result ([28, §5, Prop. 5.2]) lets us obtain the following equality of divisors in $R_{f,G}$:

$$\begin{aligned}
 (4.3) \quad & \text{Div} \left(\pi_{2,1} \left(\theta_2^{\Sigma_0} \right) \right) = \text{Div} \left(\text{Sel}_{\rho_2}^{\Sigma_0}(\mathbb{Q})^\vee \otimes_{R_{F,G}} R_{f,G} \right) \\
 & + \text{Div} \left(\text{Tor}_1^{R_{F,G}} \left(R_{f,G}, H^0(G_{\Sigma, D_{\rho_2}})^\vee \right) \right) \\
 & - \text{Div} \left(H^0(G_{\Sigma, D_{\rho_1}})^\vee \right).
 \end{aligned}$$

Theorem 1.5 establishes the following equality in the divisor group of $R_{f,G}$:

$$(4.4) \quad \text{Div} \left(\pi_{2,1} \left(\theta_2^{\Sigma_0} \right) \right) = \text{Div} \left(\theta_{\rho_1}^{\Sigma_0} \right).$$

Theorem 1.6 establishes the following equality in the divisor group of $R_{f,G}$:

$$(4.5) \quad \text{Div} \left(\text{Sel}_{\rho_2}^{\Sigma_0}(\mathbb{Q})^\vee \otimes_{R_{F,G}} R_{f,G} \right) + \text{Div} \left(\text{Tor}_1^{R_{F,G}} \left(R_{F,G}, H^0(G_\Sigma, D_{\rho_2})^\vee \right) \right) = \text{Div} \left(\text{Sel}_{\rho_1}^{\Sigma_0}(\mathbb{Q})^\vee \right).$$

Now combining equations (4.3), (4.4) and (4.5), we have the following equality of divisors in $R_{f,G}$:

$$(MC\text{-}\rho_1) \quad \text{Div} \left(\theta_1^{\Sigma_0} \right) = \text{Div} \left(\text{Sel}_{\rho_1}^{\Sigma_0}(\mathbb{Q})^\vee \right) - \text{Div} \left(H^0(G_\Sigma, D_{\rho_1})^\vee \right).$$

This completes the proof of Theorem 1.7.

5. Towards the control theorems and specialization results

The hypothesis TOR plays a crucial role in establishing many structural properties of the Galois cohomology groups and Selmer groups. The control theorems and specialization results rely heavily on Greenberg’s foundational works ([8], [9] and [10]). In these works, the Weak Leopoldt conjecture often comes into play. Let $i \in \{1, 2, 3\}$. We consider the following group:

$$\text{III}^2(D_{\rho_i}) := \ker \left(H^2(G_\Sigma, D_{\rho_i}) \rightarrow \prod_{\nu \in \Sigma} H^2(\text{Gal}(\overline{\mathbb{Q}}_\nu/\mathbb{Q}_\nu), D_{\rho_i}) \right).$$

The validity of TOR ensures us that the following statement Weak Leo holds:

Weak Leo. — $\text{III}^2(D_{\rho_3})^\vee$, $\text{III}^2(D_{\rho_2})^\vee$ and $\text{III}^2(D_{\rho_1})^\vee$ are torsion $R_{F,G}[[\Gamma]]$, $R_{F,G}$ and $R_{f,G}$ modules respectively.

For each $i \in \{1, 2, 3\}$ and each $\nu \in \Sigma_0$, the validity of TOR also lets us deduce the following equalities:

$$H^2(\text{Gal}(\overline{\mathbb{Q}}_\nu/\mathbb{Q}_\nu), D_{\rho_i}) = 0, \forall \nu \in \Sigma_0; \quad H^2 \left(\text{Gal}(\overline{\mathbb{Q}}_p/\mathbb{Q}_p), \frac{D_{\rho_i}}{\text{Fil}^+ D_{\rho_i}} \right) = 0.$$

The Galois representation ρ_3 is related to the cyclotomic deformation of ρ_2 (see [7, §3] for a description of the cyclotomic deformation). Equation (5.1) can be deduced from the arguments given in [9, §5]. The arguments involve combining local duality, Proposition 3.10 in [8] along with results from [7, §3]. These arguments are also described in [28, §4].

$$(5.1) \quad H^2(\text{Gal}(\overline{\mathbb{Q}}_p/\mathbb{Q}_p), D_{\rho_3}) = H^2(\text{Gal}(\overline{\mathbb{Q}}_p/\mathbb{Q}_p), \text{Fil}^+ D_{\rho_3}) = 0.$$

We will use local duality to establish a similar result for ρ_2 and ρ_1 . See Lemma 5.10.

$$\begin{aligned} H^2(\text{Gal}(\overline{\mathbb{Q}}_p/\mathbb{Q}_p), D_{\rho_2}) &= H^2(\text{Gal}(\overline{\mathbb{Q}}_p/\mathbb{Q}_p), \text{Fil}^+ D_{\rho_2}) = 0, \\ H^2(\text{Gal}(\overline{\mathbb{Q}}_p/\mathbb{Q}_p), D_{\rho_1}) &= H^2(\text{Gal}(\overline{\mathbb{Q}}_p/\mathbb{Q}_p), \text{Fil}^+ D_{\rho_1}) = 0. \end{aligned}$$

These observations, along with Corollary 5.11, establish the following statement:

Van-H2. — Let $i \in \{1, 2, 3\}$. Let $\nu \in \Sigma_0$. We have

$$\begin{aligned} H^2(G_\Sigma, D_{\rho_i}) &= H^2(\text{Gal}(\overline{\mathbb{Q}}_\nu/\mathbb{Q}_\nu), D_{\rho_i}) = H^2(\text{Gal}(\overline{\mathbb{Q}}_p/\mathbb{Q}_p), D_{\rho_i}) \\ &= H^2(\text{Gal}(\overline{\mathbb{Q}}_p/\mathbb{Q}_p), \text{Fil}^+ D_{\rho_i}) \\ &= H^2\left(\text{Gal}(\overline{\mathbb{Q}}_p/\mathbb{Q}_p), \frac{D_{\rho_i}}{\text{Fil}^+ D_{\rho_i}}\right) = 0. \end{aligned}$$

We will now summarize the arguments involved in deducing the specialization results given in equations (3.4) and (4.3). Let $X_3 = \text{Sel}_{\rho_3}^{\Sigma_0}(\mathbb{Q})^\vee$ and $X_2 = \text{Sel}_{\rho_2}^{\Sigma_0}(\mathbb{Q})^\vee$. Given the divisor $\text{Div}(X_3)$ in $R_{F,G}[[\Gamma]]$ (and $\text{Div}(X_2)$ in $R_{F,G}$ respectively), we would like to find the divisor $\text{Div}(X_3 \otimes_{R_{F,G}[[\Gamma]]} R_{F,G})$ in $R_{F,G}$ (and $\text{Div}(X_2 \otimes_{R_{F,G}} R_{f,G})$ in $R_{f,G}$ respectively). For this purpose, we will use Proposition 5.2 in [28, §5]. One of the hypotheses involved in the Proposition there concerns the following statement:

Fin-Proj. — For every height two prime ideal $Q_{3,2}$ in $R_{F,G}[[\Gamma]]$ containing $\ker(\pi_{3,2})$, the projective dimension of $(\text{Sel}_{\rho_3}^{\Sigma_0}(\mathbb{Q})^\vee)_{Q_{3,2}}$ is finite.

For every height two prime ideal $Q_{2,1}$ in $R_{F,G}$ containing $\ker(\pi_{2,1})$, the projective dimension of $(\text{Sel}_{\rho_2}^{\Sigma_0}(\mathbb{Q})^\vee)_{Q_{2,1}}$ is finite.

Let us suppose that Fin-Proj holds (we will briefly discuss later why Fin-Proj holds). Another hypothesis involved in Proposition 5.2 in [28, §5] requires us to show that the maximal $R_{F,G}[[\Gamma]]$ (and $R_{F,G}$ respectively) pseudo-null submodule of X_3 (and X_2 respectively) is trivial. For this purpose, we will first define the strict (non-primitive) Selmer group, for each $i \in \{1, 2, 3\}$, as follows:

$$\text{Sel}_{\rho_i}^{\Sigma_0, str}(\mathbb{Q}) := \ker \left(H^1(G_\Sigma, D_{\rho_i}) \xrightarrow{\phi_{\rho_i}^{\Sigma_0, str}} H^1\left(\text{Gal}(\overline{\mathbb{Q}}_p/\mathbb{Q}_p), \frac{D_{\rho_i}}{\text{Fil}^+ D_{\rho_i}}\right) \right).$$

Combining Proposition 4.2.1 and Proposition 4.3.2 from [10] with the statements TOR and Van-H2, we obtain that the $R_{F,G}[[\Gamma]]$ -module $\text{Sel}_{\rho_3}^{\Sigma_0, str}(\mathbb{Q})^\vee$ and the $R_{F,G}$ -module $\text{Sel}_{\rho_2}^{\Sigma_0, str}(\mathbb{Q})^\vee$ have no non-trivial pseudo-null submodules. Also for each $i \in \{1, 2, 3\}$, one has the following short exact

sequence relating $\text{Sel}_{\rho_i}^{\Sigma_0, str}(\mathbb{Q})$ to $\text{Sel}_{\rho_i}^{\Sigma_0}(\mathbb{Q})$:

$$0 \rightarrow \text{Sel}_{\rho_i}^{\Sigma_0, str}(\mathbb{Q}) \xrightarrow{\alpha_i} \text{Sel}_{\rho_i}^{\Sigma_0}(\mathbb{Q}) \rightarrow H^1\left(\Gamma_p, H^0\left(I_p, \frac{D_{\rho_i}}{\text{Fil}^+ D_{\rho_i}}\right)\right) \rightarrow 0.$$

In the cases we are interested in, Lemma 5.7 tells us that

$$H^1\left(\Gamma_p, H^0\left(I_p, \frac{D_{\rho_i}}{\text{Fil}^+ D_{\rho_i}}\right)\right) = 0,$$

for each $i \in \{1, 2, 3\}$. This lets us conclude that the map α_i is an isomorphism for each $i \in \{1, 2, 3\}$. We have the following statement:

No-PN. — The $R_{F,G}[[\Gamma]]$ -module $\text{Sel}_{\rho_3}^{\Sigma_0}(\mathbb{Q})^\vee$ has no non-trivial pseudo-null submodules.

The $R_{F,G}$ -module $\text{Sel}_{\rho_2}^{\Sigma_0}(\mathbb{Q})^\vee$ has no non-trivial pseudo-null submodules.

Let $Q_{3,2}$ (and $Q_{2,1}$ respectively) be a height two prime ideal in $R_{F,G}[[\Gamma]]$ (and $R_{F,G}$ respectively) containing $\ker(\pi_{3,2})$ (and $\ker(\pi_{2,1})$ respectively). No-PN allows us to conclude that

$$(5.2) \quad \text{depth}_{(R_{F,G}[[\Gamma]])_{Q_{3,2}}}(X_3)_{Q_{3,2}} \geq 1, \text{depth}_{(R_{F,G})_{Q_{2,1}}}(X_2)_{Q_{2,1}} \geq 1.$$

The hypotheses Fin-Proj allows us to use the Auslander–Buchsbaum formula. Combining equation (5.2) along with the Auslander–Buchsbaum formula over the two-dimensional local rings $(R_{F,G}[[\Gamma]])_{Q_{3,2}}$ and $(R_{F,G})_{Q_{2,1}}$ allows us to make the following conclusion involving projective dimensions:

$$(5.3) \quad \text{pd}_{(R_{F,G}[[\Gamma]])_{Q_{3,2}}}(X_3)_{Q_{3,2}} \leq 1, \quad \text{pd}_{(R_{F,G})_{Q_{2,1}}}(X_2)_{Q_{2,1}} \leq 1.$$

This allows us to write down projective resolutions for $(X_3)_{Q_{3,2}}$ (and $(X_2)_{Q_{2,1}}$ respectively) over $(R_{F,G}[[\Gamma]])_{Q_{3,2}}$ (and $(R_{F,G})_{Q_{2,1}}$ respectively). Let $A_3 = (R_{F,G}[[\Gamma]])_{Q_{3,2}}$ and $A_2 = (R_{F,G})_{Q_{2,1}}$. Both the rings A_3 and A_2 have Krull dimension two. We have the following short exact sequences:

$$0 \rightarrow A_3^{n_3} \xrightarrow{\omega_3} A_3^{n_3} \rightarrow (X_3)_{\mathfrak{q}_3} \rightarrow 0, \quad 0 \rightarrow A_2^{n_2} \xrightarrow{\omega_2} A_2^{n_2} \rightarrow (X_2)_{\mathfrak{q}_2} \rightarrow 0.$$

Here, we have matrices $\omega_3 \in M_{n_3}(A_3)$ and $\omega_2 \in M_{n_2}(A_2)$. Consider the rings $B_2 := A_3 \otimes_{R_{F,G}[[\Gamma]]} R_{F,G}$ and $B_1 := A_2 \otimes_{R_{F,G}} R_{f,G}$, both of which have Krull dimension one. Taking the tensor product with the $R_{F,G}[[\Gamma]]$ -module $R_{F,G}$ (and $R_{F,G}$ -module $R_{f,G}$ respectively), we obtain the following short exact sequences:

$$\begin{aligned} 0 \rightarrow B_2^{n_3} &\xrightarrow{\pi_{3,2}(\omega_3)} B_2^{n_3} \rightarrow X_3 \otimes_{R_{F,G}[[\Gamma]]} B_2 \rightarrow 0, \\ 0 \rightarrow B_1^{n_2} &\xrightarrow{\pi_{2,1}(\omega_2)} B_1^{n_2} \rightarrow X_2 \otimes_{R_{F,G}} B_1 \rightarrow 0. \end{aligned}$$

If we let $\text{len}_A(M)$ denote the length of an A -module M , we have

$$\begin{aligned} \text{len}_{B_2} \left(\frac{B_2}{(\pi_{3,2}(\det(\omega_3)))} \right) &= \text{len}_{B_2} (X_3 \otimes_{R_{F,G}[\Gamma]} B_2), \\ \text{len}_{B_1} \left(\frac{B_1}{(\pi_{2,1}(\det(\omega_2)))} \right) &= \text{len}_{B_1} (X_2 \otimes_{R_{F,G}} B_1). \end{aligned}$$

It is this observation that we will maneuver to obtain information about the divisors $\text{Div} (X_3 \otimes_{R_{F,G}[\Gamma]} R_{F,G})$ and $\text{Div} (X_2 \otimes_{R_{F,G}} R_{f,G})$ in $R_{F,G}$ and $R_{f,G}$ respectively.

In what follows, we will keep the isomorphism $\text{Sel}_{\rho_i}^{\Sigma_0}(\mathbb{Q}_\infty) \cong \text{Sel}_{\rho_i}^{\Sigma_0, \text{str}}(\mathbb{Q}_\infty)$ in mind to establish Fin-Proj. We can combine Corollary 3.2.3 in [9] with TOR and Van-H2 to establish the following statement:

Surj. — The global-to-local maps $\phi_{\rho_3}^{\Sigma_0}$, $\phi_{\rho_2}^{\Sigma_0}$ and $\phi_{\rho_1}^{\Sigma_0}$ defining the Selmer groups $\text{Sel}_{\rho_3}^{\Sigma_0}(\mathbb{Q})$, $\text{Sel}_{\rho_2}^{\Sigma_0}(\mathbb{Q})$ and $\text{Sel}_{\rho_1}^{\Sigma_0}(\mathbb{Q})$ respectively are surjective.

Let $Q_{3,2}$ (and $Q_{2,1}$ respectively) be a height two prime ideal in $R_{F,G}[\Gamma]$ (and $R_{F,G}$ respectively) containing $\ker(\pi_{3,2})$ (and $\ker(\pi_{2,1})$ respectively). To establish Fin-Proj, it will suffice to prove that the projective dimensions of the $(R_{F,G}[\Gamma])_{Q_{3,2}}$ -modules

$$\begin{aligned} &H^1(G_\Sigma, D_{\rho_3})^\vee \otimes_{R_{F,G}[\Gamma]} R_{F,G}[\Gamma]_{Q_{3,2}}, \\ &H^1 \left(\text{Gal}(\overline{\mathbb{Q}}_p/\mathbb{Q}_p), \frac{D_{\rho_3}}{\text{Fil}^+ D_{\rho_3}} \right)^\vee \otimes_{R_{F,G}[\Gamma]} (R_{F,G}[\Gamma])_{Q_{3,2}}, \end{aligned}$$

and the projective dimensions of the $(R_{F,G})_{Q_{2,1}}$ -modules

$$\begin{aligned} &H^1(G_\Sigma, D_{\rho_2})^\vee \otimes_{R_{F,G}} (R_{F,G})_{Q_{2,1}}, \\ &H^1 \left(\text{Gal}(\overline{\mathbb{Q}}_p/\mathbb{Q}_p), \frac{D_{\rho_2}}{\text{Fil}^+ D_{\rho_2}} \right)^\vee \otimes_{R_{F,G}} (R_{F,G})_{Q_{2,1}}, \end{aligned}$$

respectively are finite. The hypothesis Van-H2 allows us to apply Proposition 5.5 in [28]. It will now suffice to prove the following statements Reg- ρ_3 and Reg- ρ_2 :

Reg- ρ_3 . — For every height two prime ideal $Q_{3,2}$ containing $\ker(\pi_{3,2})$ and in the support of $H^0(G_\Sigma, D_{\rho_3})^\vee$ or $H^0 \left(\text{Gal}(\overline{\mathbb{Q}}_p/\mathbb{Q}_p), \frac{D_{\rho_3}}{\text{Fil}^+ D_{\rho_3}} \right)^\vee$, the 2-dimensional local ring $(R_{F,G}[\Gamma])_{Q_{3,2}}$ is regular.

Reg- ρ_2 . — For every height two prime ideal $Q_{2,1}$ containing $\ker(\pi_{2,1})$ and in the support of $H^0(G_\Sigma, D_{\rho_2})^\vee$,


- the 2-dimensional local ring $(R_{F,G})_{Q_{2,1}}$ is regular;

- the 1-dimensional ring $\left(\frac{R_{F,G}}{\ker(\pi_{2,1})}\right)_{Q_{2,1}}$ is integrally closed (and hence regular too).

The projective dimension of the $R_{F,G}$ -module $H^0\left(\text{Gal}(\overline{\mathbb{Q}}_p/\mathbb{Q}_p), \frac{D_{\rho_2}}{\text{Fil}^+ D_{\rho_2}}\right)^\vee$ equals one.

Proposition 5.3, Lemma 5.4 and Lemma 5.8 establish $\text{Reg-}\rho_2$. Lemma 4.16 and Proposition 4.17 in [28] establish $\text{Reg-}\rho_3$.

The control theorems that relate $\text{Sel}_{\rho_3}^{\Sigma_0}(\mathbb{Q})^\vee \otimes_{R_{F,G}[[\Gamma]]} R_{F,G}$ with $\text{Sel}_{\rho_2}^{\Sigma_0}(\mathbb{Q})^\vee$, given in equation (3.1), and $\text{Sel}_{\rho_2}^{\Sigma_0}(\mathbb{Q})^\vee \otimes_{R_{F,G}} R_{f,G}$ with $\text{Sel}_{\rho_1}^{\Sigma_0}(\mathbb{Q})^\vee$, given in equation (4.1), use Proposition 6.2 in [28]. The following flowchart summarizes the logical flow involved in those arguments:



5.1. Calculations involving global Galois cohomology groups

LEMMA 5.1. — *Let Q be a height two prime ideal in $R_{F,G}$ satisfying the following properties:*

- $p \notin Q$.
- $i_F(\mathfrak{P}_F) \subset Q$, for some classical height one prime ideal \mathfrak{P}_F of R_F with weight ≥ 2 .

Then, $(R_{F,G})_Q$ is a regular local ring with Krull dimension 2.

Proof. — We have the following natural inclusions:

$$i_1 : O[[x_F]] \hookrightarrow R_F, \quad i_F : R_F \hookrightarrow R_{F,G}, \quad i_{F,G} : R_G[[x_F]] \hookrightarrow R_{F,G}.$$

Let us define the following prime ideals:

$$\underbrace{\mathfrak{p}_0}_{\text{height}=1} := i_1^{-1}(\mathfrak{P}_F), \quad \underbrace{Q_0}_{\text{height}=2} := i_{F,G}^{-1}(Q).$$

Since $p \notin Q$ and the Krull dimension of R_F equals two, we have the equality $i_F^{-1}(Q) = \mathfrak{P}_F$. Furthermore, since \mathfrak{P}_F is a classical height one prime ideal in R_F , the extension

$$(O[[x_F]])_{\mathfrak{p}_0} \hookrightarrow (R_F)_{\mathfrak{P}_F}$$

of discrete valuation rings is finite and étale (see [14, Cor. 1.4]). Let

$$\mathcal{R} = (R_G[[x_F]])_{Q_0} \otimes_{(O[[x_F]])_{\mathfrak{p}_0}} (R_F)_{\mathfrak{P}_F}.$$

We have the following commutative diagrams:

$$\begin{array}{ccc} O[[x_F]] & \longrightarrow & R_G[[x_F]] & & (O[[x_F]])_{\mathfrak{p}_0} & \longrightarrow & (R_G[[x_F]])_{Q_0} \\ \downarrow i_1 & & \downarrow i_{F,G} & & \downarrow & & \downarrow \\ R_F & \xrightarrow{i_F} & R_{F,G} & & (R_F)_{\mathfrak{P}_F} & \longrightarrow & \mathcal{R} \end{array}$$

Étale morphisms are stable under base change ([25, Chap. 4, Prop. 3.22]). So, the morphism $(R_G[[x_F]])_{Q_0} \rightarrow \mathcal{R}$ is also étale. Since \mathfrak{P}_F is a classical height one prime ideal in R_F , there exists integers n_1 and n_2 such that $(1 + x_F)^{n_1} - (1 + p)^{n_2}$ belongs to \mathfrak{P}_F (and hence belongs to Q_0 as well). The polynomial $(1 + x_F)^{n_1} - (1 + p)^{n_2}$ is a non-constant monic polynomial. Lemma 4.16 in [28] now forces $(R_G[[x_F]])_{Q_0}$ to be a regular local ring with Krull dimension 2. Since the morphism $(R_G[[x_F]])_{Q_0} \rightarrow \mathcal{R}$ is étale, the ring \mathcal{R} is regular too ([25, Chap. 4, Cor. 3.24]).

The natural $(O[[x_F]])_{\mathfrak{p}_0}$ -algebra maps

$$(R_G[[x_F]])_{Q_0} \hookrightarrow (R_{F,G})_Q, \quad (R_F)_{\mathfrak{P}_F} \hookrightarrow (R_{F,G})_Q$$

give us a natural $(O[[x_F]])_{\mathfrak{p}_0}$ -algebra map $\mathcal{R} \rightarrow (R_{F,G})_Q$. If we let Q_1 be the prime ideal in \mathcal{R} that Q pullbacks to, we get a natural map $\beta : \mathcal{R}_{Q_1} \rightarrow (R_{F,G})_Q$ of local rings. Using the fact that the completed tensor product $R_{F,G}$ satisfies the universal property of being a co-product in the category of complete semi-local Noetherian O -algebras, we can deduce the following natural isomorphism:

$$(5.4) \quad R_G[[x_F]] \otimes_{O[[x_F]]} R_F \cong R_{F,G}.$$

Equation (5.4) gives us a natural map $R_{F,G} \rightarrow \mathcal{R}$. We obtain the two following sequence of maps:

$$R_{F,G} \rightarrow \mathcal{R}_{Q_1} \xrightarrow{\beta} (R_{F,G})_Q, \quad (R_{F,G})_Q \xrightarrow{\beta'} \mathcal{R}_{Q_1} \xrightarrow{\beta} (R_{F,G})_Q$$

The second sequence of maps in the equation above is obtained by localizing the first sequence of maps with respect to the prime ideal (corresponding to) Q . The composition $\beta \circ \beta'$ is the identity map. Thus, the map β must be surjective. In fact, β turns out to be an isomorphism. To see

this, it suffices to show that $\ker(\beta)$ equals 0. Since the Krull dimensions of both \mathcal{R}_{Q_1} and $(R_{f,G})_Q$ equal two, the height of the prime ideal $\ker(\beta)$ must equal 0. Since \mathcal{R}_{Q_1} is a domain (it is in fact a regular local ring), $\ker(\beta)$ must equal 0. This completes the proof since \mathcal{R}_{Q_1} is a regular local ring. \square

LEMMA 5.2. — *The $R_{f,G}$ -module $H^0(G_\Sigma, D_{\rho_1})^\vee$ is not supported at any height 1 prime ideal of $R_{f,G}$ containing the prime number p . Consequently, the $R_{f,G}$ -module $H^0(G_\Sigma, D_{\rho_1})^\vee$ is torsion.*

Proof. — Let \mathfrak{q} be a height one prime ideal in $R_{f,G}$ containing p . Let K denote the fraction field of $\frac{R_{f,G}}{\mathfrak{q}}$. Note that the characteristic of K is equal to p . We have the following G_Σ -equivariant isomorphisms:

$$\begin{aligned} D_{\rho_1}^\vee &\cong \text{Hom}_{R_{f,G}}(L_G \otimes_{R_G} R_{f,G}, L_f \otimes_{O_f} R_{f,G}), \\ &\implies D_{\rho_1}^\vee \otimes_{R_{f,G}} K \cong \text{Hom}_K(V_G, V_f). \end{aligned}$$

Here, V_f and V_G are K -vector spaces with a G_Σ -action given by following representations σ_f and σ_G respectively:

$$\begin{aligned} \sigma_f : G_\Sigma &\xrightarrow{\rho_f} \text{GL}_2(O_f) \hookrightarrow \text{GL}_2(R_{f,G}) \rightarrow \text{GL}_2(K), \\ \sigma_G : G_\Sigma &\xrightarrow{\rho_G} \text{GL}_2(R_G) \hookrightarrow \text{GL}_2(R_{f,G}) \rightarrow \text{GL}_2(K). \end{aligned}$$

To show that $H^0(G_\Sigma, D_{\rho_1})^\vee$ is not supported at \mathfrak{p} , it will be sufficient to show that $H^0(G_\Sigma, \text{Hom}_K(V_G, V_f))$ is equal to 0. Both σ_f and σ_G are Galois representations over K , a field of characteristic p . In fact as a G_Σ -representation, σ_f is isomorphic to the residual representation $\bar{\rho}_F$, after extending scalars to K . The residual representation $\bar{\rho}_F$ is absolutely irreducible (due to the hypothesis IRR). Consequently, the Galois representation σ_f is irreducible. Thus, to show that $H^0(G_\Sigma, \text{Hom}_K(V_G, V_f))$ is equal to 0, it will be sufficient to show that σ_f and σ_G are not isomorphic as G_Σ -representations over K . For this purpose, it will be sufficient to show that $\det(\sigma_f)$ and $\det(\sigma_G)$ are not isomorphic as G_Σ -representations. The image of $\det(\sigma_f)$ lies inside $\overline{\mathbb{F}}_p^\times$. The pullback $j_G^{-1}(\mathfrak{q})$ of the height one prime ideal \mathfrak{q} under the map $j_G : R_G \hookrightarrow R_{f,G}$ is a height one prime ideal (and not the maximal ideal) of R_G . Consequently, the image of $\det(\sigma_G)$ lies inside $\overline{\mathbb{F}}_p[[x_G]]^\times$ but does not lie inside $\overline{\mathbb{F}}_p^\times$. Consequently, $\det(\sigma_f)$ and $\det(\sigma_G)$ are not isomorphic as G_Σ -representations. The lemma follows. \square

Note that $\frac{R_{F,G}}{\ker(\pi_{2,1})} \hookrightarrow R_{f,G}$ is an integral extension. We have the following isomorphism of $R_{f,G}$ -modules:

$$(5.5) \quad \left(H^0(G_\Sigma, D_{\rho_2})^\vee \otimes_{R_{F,G}} \frac{R_{F,G}}{\ker(\pi_{2,1})} \right) \otimes_{\frac{R_{F,G}}{\ker(\pi_{2,1})}} R_{f,G} \cong H^0(G_\Sigma, D_{\rho_1})^\vee.$$

Let $Q_{2,1}$ be a height two prime ideal in $R_{F,G}$ containing $\ker(\pi_{2,1})$. It uniquely corresponds to a height one prime ideal \mathfrak{p} in $\frac{R_{F,G}}{\ker(\pi_{2,1})}$; And there exist finitely many height one prime ideals $\mathfrak{q}_1, \dots, \mathfrak{q}_n$ in $R_{f,G}$ lying above \mathfrak{p} . A simple application of Nakayama’s lemma along with equation (5.5) lets us make the following observation: A height two prime ideal $Q_{2,1}$ in $R_{F,G}$ containing $\ker(\pi_{2,1})$ belongs to the support of $H^0(G_\Sigma, D_{\rho_2})^\vee$ if and only if one of the height one prime ideals \mathfrak{q}_i (for $1 \leq i \leq n$) in $R_{f,G}$ belongs to the support $H^0(G_\Sigma, D_{\rho_1})^\vee$. These observations along with Lemma 5.1 and Lemma 5.2 immediately give us the following proposition:

PROPOSITION 5.3. — *Let $Q_{2,1}$ be a height two prime ideal containing $\ker(\pi_{2,1})$.*

- *If $p \in Q_{2,1}$, then $(H^0(G_\Sigma, D_{\rho_2})^\vee)_{Q_{2,1}} = 0$.*
- *If $p \notin Q_{2,1}$, then $(R_{F,G})_{Q_{2,1}}$ is a regular local ring of dimension 2.*

Let P_k be the classical height one prime ideal corresponding to f . Here, P_k is defined to be the kernel of the natural map $\pi_f : R_F \rightarrow O_f$. The ring O_f , defined to be the integral closure of $\frac{R_F}{P_k}$, is the ring of integers in a finite extension of \mathbb{Q}_p . As a result, the index $[O_f : \frac{R_F}{\ker(\pi_f)}]$ is finite and equal to a power of p . The ring $R_{f,G}$ is the completed tensor product $O_f \widehat{\otimes} R_G$ (over the ring O). The map $\pi_{2,1} : R_{F,G} \rightarrow R_{f,G}$ is obtained naturally via the maps

$$\pi_f : R_F \rightarrow O_f, \quad \text{id} : R_G \xrightarrow{\cong} R_G.$$

So, the cokernel of the natural inclusion $\frac{R_{F,G}}{\ker(\pi_{2,1})} \hookrightarrow R_{f,G}$ is also annihilated by a power of p . We obtain the following lemma:

LEMMA 5.4. — *Let \mathfrak{q} be a height one prime ideal in $R_{f,G}$ not containing p . Let $Q_{2,1}$ denote the height two prime ideal in $R_{F,G}$, containing $\ker(\pi_{2,1})$, given by $\pi_{2,1}^{-1}(\mathfrak{q})$. Then, the following natural inclusion is an equality:*

$$\left(\frac{R_{F,G}}{\ker(\pi_{2,1})} \right)_{Q_{2,1}} \xrightarrow{\cong} (R_{f,G})_{\mathfrak{q}}.$$

Lemma 5.2 also lets us conclude that $H^0(G_\Sigma, D_{\rho_1})^\vee$ is a torsion $R_{f,G}$ -module. Combining equation (5.5) with Nakayama’s lemma, we can then conclude that $(H^0(G_\Sigma, D_{\rho_2})^\vee)_{\ker(\pi_{2,1})} = 0$.

What we also obtain is that $H^0(G_\Sigma, D_{\rho_2})^\vee$ is a torsion $R_{F,G}$ -module. Considering $R_{F,G}$ as an $R_{F,G}[[\Gamma]]$ -module via the map $\pi_{3,2} : R_{F,G}[[\Gamma]] \rightarrow R_{F,G}$, we also have the following isomorphism of $R_{F,G}$ -modules:

$$(5.6) \quad H^0(G_\Sigma, D_{\rho_3})^\vee \otimes_{R_{F,G}[[\Gamma]]} R_{F,G} \cong H^0(G_\Sigma, D_{\rho_2})^\vee.$$

Nakayama’s lemma lets us conclude that $(H^0(G_\Sigma, D_{\rho_3})^\vee)_{\ker(\pi_{3,2})} = 0$. We have shown that the following equality holds:

$$(H^0(G_\Sigma, D_{\rho_3})^\vee)_{\ker(\pi_{3,2})} = (H^0(G_\Sigma, D_{\rho_2})^\vee)_{\ker(\pi_{2,1})} = 0.$$

Since the Tor functor commutes with localization, we immediately get the following lemma:

LEMMA 5.5. — *For all $i \geq 0$, the following statements hold:*

- *The $R_{F,G}$ -module $\text{Tor}_i^{R_{F,G}[[\Gamma]]}(R_{F,G}, H^0(G_\Sigma, D_{\rho_3})^\vee)$ is torsion,*
- *The $R_{f,G}$ -module $\text{Tor}_i^{R_{F,G}}(R_{f,G}, H^0(G_\Sigma, D_{\rho_2})^\vee)$ is torsion.*

5.2. Calculations involving local Galois cohomology groups

As for the local Galois cohomology groups, we will begin with the following exact sequence of $R_{F,G}[[\Gamma]]$ -modules that is $\text{Gal}(\overline{\mathbb{Q}}_p/\mathbb{Q}_p)$ -equivariant:

$$(5.7) \quad 0 \rightarrow \underbrace{W}_{\text{corank}=1} \rightarrow \underbrace{\frac{D_{\rho_3}}{\text{Fil}^+ D_{\rho_3}}}_{\text{corank}=2} \rightarrow \underbrace{W'}_{\text{corank}=1} \rightarrow 0, \text{ where}$$

$$W := \text{Hom}_{R_{F,G}} \left(\frac{L_F}{\text{Fil}^+ L_F} \otimes_{R_F} R_{F,G}, \frac{L_G}{\text{Fil}^+ L_G} \otimes_{R_G} R_{F,G} \right) \otimes_{R_{F,G}} \widehat{R_{F,G}[[\Gamma]]}(\kappa^{-1}),$$

$$W' := \text{Hom}_{R_{F,G}} \left(\text{Fil}^+ L_F \otimes_{R_F} R_{F,G}, \frac{L_G}{\text{Fil}^+ L_G} \otimes_{R_G} R_{F,G} \right) \otimes_{R_{F,G}} \widehat{R_{F,G}[[\Gamma]]}(\kappa^{-1}).$$

Note that we also have the following $\text{Gal}(\overline{\mathbb{Q}}_p/\mathbb{Q}_p)$ -equivariant isomorphisms of $R_{F,G}[[\Gamma]]$ -modules:

$$W \cong R_{F,G} \left((i_F \circ \epsilon_F^{-1}) \cdot (i_G \circ \epsilon_G) \right) \otimes_{R_{F,G}} \widehat{R_{F,G}[[\Gamma]]}(\kappa^{-1}),$$

$$W' \cong R_{F,G} \left((i_F \circ \delta_F^{-1}) \cdot (i_G \circ \epsilon_G) \right) \otimes_{R_{F,G}} \widehat{R_{F,G}[[\Gamma]]}(\kappa^{-1}).$$

The hypotheses p -DIS-IN ensures us that the residual representation, defined over $\overline{\mathbb{F}}_p$, associated to $(i_F \circ \delta_F^{-1}) \cdot (i_G \circ \epsilon_G)$ is non-trivial. Consequently, $H^0(I_p, W') = 0$. So, we get the following $\frac{\text{Gal}(\overline{\mathbb{Q}}_p/\mathbb{Q}_p)}{I_p}$ -equivariant isomorphism of $R_{F,G}[[\Gamma]]$ -modules:

$$(5.8) \quad H^0 \left(I_p, \frac{D_{\rho_3}}{\text{Fil}^+ D_{\rho_3}} \right) \cong H^0(I_p, W).$$

Let η_p denote the unique prime in \mathbb{Q}_∞ lying above p and let I_{η_p} denote the corresponding inertia subgroup at η_p . The Galois representation ρ_3 is related to the cyclotomic deformation of ρ_2 . We have denoted the quotient $\frac{\text{Gal}(\overline{\mathbb{Q}}_p/\mathbb{Q}_p)}{I_p}$ by Γ_p . The arguments in [7, §3] combined with the fact that both the characters ϵ_F and ϵ_G are unramified at p give us the following isomorphism that is Γ_p -equivariant:

$$\begin{aligned} H^0(I_p, W) &\cong H^0 \left(I_{\eta_p}, \widehat{R_{F,G}} \left((i_F \circ \epsilon_F^{-1}) \cdot (i_G \circ \epsilon_G) \right) \right) \\ &\cong \widehat{R_{F,G}} \left((i_F \circ \epsilon_F^{-1}) \cdot (i_G \circ \epsilon_G) \right) \end{aligned}$$

Taking the Pontryagin duals of the modules in the above equation gives us the following lemma:

LEMMA 5.6. — *We have the following isomorphism of $R_{F,G}[[\Gamma]]$ -modules that is Γ_p -equivariant.*

$$H^0 \left(I_p, \frac{D_{\rho_3}}{\text{Fil}^+ D_{\rho_3}} \right)^\vee = R_{F,G} \left((i_F \circ \epsilon_F) \cdot (i_G \circ \epsilon_G^{-1}) \right).$$

Here, we consider $R_{F,G}$ as an $R_{F,G}[[\Gamma]]$ -module via the map $\pi_{3,2} : R_{F,G}[[\Gamma]] \rightarrow R_{F,G}$.

Note that a_p and b_p are elements of R_F and R_G respectively but not elements of O . This is because the values that a_p and b_p take at classical specializations of F and G respectively vary as one varies the weight (see [13, Lem. 3.2]). Consequently, the ratio $\frac{i_F(a_p)}{i_G(b_p)}$ is not an element of O (in particular, it is not equal to 1). Thus, the following map of free $R_{F,G}$ -modules (of rank 1), which is given by multiplication by $\frac{i_F(a_p)}{i_G(b_p)} - 1$, is injective:

$$H^0 \left(I_p, \frac{D_{\rho_3}}{\text{Fil}^+ D_{\rho_3}} \right)^\vee \xrightarrow{\text{Frob}_p - 1} H^0 \left(I_p, \frac{D_{\rho_3}}{\text{Fil}^+ D_{\rho_3}} \right)^\vee.$$

The Frobenius at p , denoted by Frob_p , is a topological generator for Γ_p . This gives us the following equality:

$$H^1 \left(\Gamma_p, H^0 \left(I_p, \frac{D_{\rho_3}}{\text{Fil}^+ D_{\rho_3}} \right) \right) = 0.$$

Arguing similarly, we get the following results for the modules related to ρ_2 and ρ_1 :

$$(5.9) \quad \begin{aligned} H^0 \left(I_p, \frac{D_{\rho_2}}{\text{Fil}^+ D_{\rho_2}} \right) &\cong \widehat{R_{F,G}} \left((i_F \circ \epsilon_F^{-1}) \cdot (i_G \circ \epsilon_G) \right), \\ H^1 \left(\Gamma_p, H^0 \left(I_p, \frac{D_{\rho_2}}{\text{Fil}^+ D_{\rho_2}} \right) \right) &= 0. \\ H^0 \left(I_p, \frac{D_{\rho_1}}{\text{Fil}^+ D_{\rho_1}} \right) &\cong \widehat{R_{f,G}} \left((j_F \circ \epsilon_F^{-1}) \cdot (i_G \circ \epsilon_G) \right), \\ H^1 \left(\Gamma_p, H^0 \left(I_p, \frac{D_{\rho_1}}{\text{Fil}^+ D_{\rho_1}} \right) \right) &= 0. \end{aligned}$$

Combining these observations, we get the following lemma:

LEMMA 5.7. — *Let $i \in \{1, 2, 3\}$. Then, $H^1 \left(\Gamma_p, H^0 \left(I_p, \frac{D_{\rho_i}}{\text{Fil}^+ D_{\rho_i}} \right) \right) = 0$.*

Consider the short exact sequence $0 \rightarrow R_{F,G}[\Gamma] \xrightarrow{\gamma_0^{-1}} R_{F,G}[\Gamma] \rightarrow R_{F,G} \rightarrow 0$ of $R_{F,G}[\Gamma]$ -modules. Here, γ_0 is a topological generator for the pro-cyclic group Γ . Taking the tensor product with the $R_{F,G}[\Gamma]$ -module $H^0 \left(I_p, \frac{D_{\rho_3}}{\text{Fil}^+ D_{\rho_3}} \right)^\vee$ over the ring $R_{F,G}[\Gamma]$, we get the following exact sequence:

$$\begin{aligned} 0 \rightarrow \text{Tor}_1^{R_{F,G}[\Gamma]} \left(R_{F,G}, H^0 \left(I_p, \frac{D_{\rho_3}}{\text{Fil}^+ D_{\rho_3}} \right)^\vee \right) \\ \rightarrow H^0 \left(I_p, \frac{D_{\rho_3}}{\text{Fil}^+ D_{\rho_3}} \right)^\vee \xrightarrow{\gamma_0^{-1}} H^0 \left(I_p, \frac{D_{\rho_3}}{\text{Fil}^+ D_{\rho_3}} \right)^\vee \\ \rightarrow H^0 \left(I_p, \frac{D_{\rho_3}}{\text{Fil}^+ D_{\rho_3}} \right)^\vee \otimes_{R_{F,G}[\Gamma]} R_{F,G} \rightarrow 0. \end{aligned}$$

By Nakayama’s lemma, the natural surjection

$$H^0 \left(I_p, \frac{D_{\rho_3}}{\text{Fil}^+ D_{\rho_3}} \right)^\vee \rightarrow H^0 \left(I_p, \frac{D_{\rho_3}}{\text{Fil}^+ D_{\rho_3}} \right)^\vee \otimes_{R_{F,G}[\Gamma]} R_{F,G}$$

must be an isomorphism since it is a map between free $R_{F,G}$ -modules of rank 1 (see Lemma 5.6 and equation (5.9)). This tells us that the map $H^0 \left(I_p, \frac{D_{\rho_3}}{\text{Fil}^+ D_{\rho_3}} \right)^\vee \xrightarrow{\gamma_0^{-1}} H^0 \left(I_p, \frac{D_{\rho_3}}{\text{Fil}^+ D_{\rho_3}} \right)^\vee$, given by multiplication by the

element $\gamma_0 - 1$, is the zero map. Using Lemma 5.6, we get the following isomorphism of $R_{F,G}$ -modules that is Γ_p -equivariant:

$$\begin{aligned} \text{Tor}_1^{R_{F,G}[[\Gamma]]} \left(R_{F,G}, H^0 \left(I_p, \frac{D_{\rho_3}}{\text{Fil}^+ D_{\rho_3}} \right)^\vee \right) &\cong H^0 \left(I_p, \frac{D_{\rho_3}}{\text{Fil}^+ D_{\rho_3}} \right)^\vee \\ &\cong R_{F,G} \left((i_F \circ \epsilon_F) \cdot (i_G \circ \epsilon_G^{-1}) \right). \end{aligned}$$

This observation, along with equation (5.9), immediately gives us the following lemma:

LEMMA 5.8. — We have the following isomorphism of torsion $R_{F,G}$ -modules:

$$\begin{aligned} \text{Tor}_1^{R_{F,G}[[\Gamma]]} \left(R_{F,G}, H^0 \left(I_p, \frac{D_{\rho_3}}{\text{Fil}^+ D_{\rho_3}} \right)^\vee \right)_{\Gamma_p} &\cong \frac{R_{F,G}}{\left(1 - \frac{i_F(a_p)}{i_G(b_p)} \right)}, \\ H^0 \left(\text{Gal}(\overline{\mathbb{Q}}_p/\mathbb{Q}_p), \frac{D_{\rho_2}}{\text{Fil}^+ D_{\rho_2}} \right)^\vee &\cong \frac{R_{F,G}}{\left(1 - \frac{i_F(a_p)}{i_G(b_p)} \right)}. \end{aligned}$$

LEMMA 5.9.

- For all $i \geq 2$, we have $\text{Tor}_i^{R_{F,G}[[\Gamma]]} \left(R_{F,G}, H^0 \left(I_p, \frac{D_{\rho_3}}{\text{Fil}^+ D_{\rho_3}} \right)^\vee \right) = 0$.
- For all $i \geq 1$, we have $\text{Tor}_i^{R_{F,G}} \left(R_{f,G}, H^0 \left(I_p, \frac{D_{\rho_2}}{\text{Fil}^+ D_{\rho_2}} \right)^\vee \right) = 0$.

Proof. — The first statement simply follows from the fact that as an $R_{F,G}[[\Gamma]]$ -module, the projective dimension of $R_{F,G}$ (which is isomorphic to $\frac{R_{F,G}[[\Gamma]]}{(\gamma_0 - 1)}$ as an $R_{F,G}[[\Gamma]]$ -module) is equal to 1. The second statement follows from equation (5.9), since $H^0 \left(I_p, \frac{D_{\rho_2}}{\text{Fil}^+ D_{\rho_2}} \right)^\vee$ is a free $R_{F,G}$ -module. \square

LEMMA 5.10.

$$\begin{aligned} H^2 \left(\text{Gal}(\overline{\mathbb{Q}}_p/\mathbb{Q}_p), D_{\rho_1} \right) &= H^2 \left(\text{Gal}(\overline{\mathbb{Q}}_p/\mathbb{Q}_p), \text{Fil}^+ D_{\rho_1} \right) \\ &= H^2 \left(\text{Gal}(\overline{\mathbb{Q}}_p/\mathbb{Q}_p), D_{\rho_2} \right) = H^2 \left(\text{Gal}(\overline{\mathbb{Q}}_p/\mathbb{Q}_p), \text{Fil}^+ D_{\rho_2} \right) = 0. \end{aligned}$$

Proof. — We will show that $H^2 \left(\text{Gal}(\overline{\mathbb{Q}}_p/\mathbb{Q}_p), D_{\rho_2} \right) = 0$. The rest of the lemma follows using similar arguments. By local duality, we have

$$\begin{aligned} H^2 \left(\text{Gal}(\overline{\mathbb{Q}}_p/\mathbb{Q}_p), D_{\rho_2} \right) &\cong H^0 \left(\text{Gal}(\overline{\mathbb{Q}}_p/\mathbb{Q}_p), L_2^* \right), \\ \text{where } L_2^* &\cong \text{Hom}_{R_{F,G}} \left(L_G \otimes_{R_G} R_{F,G}, L_F \otimes_{R_F} R_{F,G}(\chi_p) \right). \end{aligned}$$

Here, $\chi_p : G_\Sigma \rightarrow \mathbb{Z}_p^\times$ is the p -adic cyclotomic character given by the action of G_Σ on μ_{p^∞} , the p -power roots of unity. Let E denote the fraction field

of $R_{F,G}$. We define the following vector spaces over E that have an action of $\text{Gal}(\overline{\mathbb{Q}}_p/\mathbb{Q}_p)$:

$$V_F(\chi_p) := L_F \otimes_{R_F} E(\chi_p), \quad V_G := L_G \otimes_{R_G} E.$$

It will be sufficient to show that

$$(5.10) \quad \text{Hom}_E(V_G, V_F(\chi_p))^{I_p} \stackrel{?}{=} 0.$$

Both $V_F(\chi_p)$ and V_G are reducible representations for the action of $\text{Gal}(\overline{\mathbb{Q}}_p/\mathbb{Q}_p)$. Let $V_F(\chi_p)^{ss}$ and V_G^{ss} denote the semi-simplifications of $V_F(\chi_p)$ and V_G respectively for the action of the inertia subgroup I_p . We have the following I_p -equivariant isomorphisms:

$$V_F(\chi_p)^{ss} \cong E(\chi_p(i_F \circ \delta_F)) \oplus E(\chi_p), \quad V_G^{ss} \cong E(i_G \circ \delta_G) \oplus E.$$

In this paragraph, we will consider characters over the inertia group I_p . The values of the character $\chi_p(i_F \circ \delta_F)$ lie inside the ring R_F but not inside the ring O . Similarly, the values of the character $i_G \circ \delta_G$ lie inside the ring R_G but not inside the ring O . As a result, the character $\chi_p(i_F \circ \delta_F)$ cannot equal $i_G \circ \delta_G$ or the trivial character. Also, the p -adic cyclotomic character χ_p (which takes values in \mathbb{Z}_p^\times) cannot equal the character $i_G \circ \delta_G$ or the trivial character. Equation (5.10) now follows from these observations; and hence so does the lemma. \square

The validity of the Weak Leopoldt conjecture Weak Leo along with Proposition 6.1 in [8] and Proposition 5.2.4 in [9] gives us the following corollary to Lemma 5.10.

COROLLARY 5.11.

$$H^2(G_\Sigma, D_{\rho_1}) = H^2(G_\Sigma, D_{\rho_2}) = H^2(G_\Sigma, D_{\rho_3}) = 0.$$

6. Examples when R_F is not regular

To end, we will provide some examples when R_F , the normalization of a primitive component of the ordinary Hecke algebra, is not a UFD (and hence not regular). These examples will be based on the circle of ideas developed in [19], [2] and [3]. We have used SAGE [29] to perform the computations. Let $E = \mathbb{Q}(\sqrt{D})$ be a real quadratic field with discriminant $D > 0$, and where the prime p splits; we have $(p) = \mathfrak{p}_1\mathfrak{p}_2$ for two distinct prime ideals \mathfrak{p}_1 and \mathfrak{p}_2 in the ring of integers of the number field E . Let k be an even integer strictly greater than 2. Let $\chi_E : \text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q}) \rightarrow \{\pm 1\}$ be

the quadratic character associated to E . Let ϵ be a fundamental unit for E . Let us suppose that the following condition holds:

$$(6.1) \quad p \text{ divides } \text{Norm}_{E/\mathbb{Q}}(\epsilon^{k-1} - 1).$$

We will assume as a result, without loss of generality, that $\epsilon^{k-1} - 1 \in \mathfrak{p}_1$. Equation (6.1) allows us to apply Proposition 2.1 in [19]. We have an ordinary p -stabilized newform $f = \sum_{n \geq 1} a_n(f)q^n$ in $S_k^{\text{ord}}(Dp, \chi_E)$ such that the residual representation $\bar{\rho}_f$ (defined over a finite field \mathbb{F} with characteristic p) associated to f satisfies the following properties:

$$\begin{aligned} \bar{\rho}_f \text{ is absolutely irreducible, } \quad \bar{\rho}_f \otimes \chi_E &\cong \bar{\rho}_f, \\ \bar{\rho}_f &\text{ is } p\text{-ordinary (in the sense of [1]).} \end{aligned}$$

Let O denote the ring of integers in a finite extension of \mathbb{Q}_p containing the unique unramified quadratic extension of \mathbb{Q}_p , the ring of Witt vectors $W(\mathbb{F})$ and the p^r -th roots of unity (where p^r denotes the exponent of the p -Sylow subgroup of the ideal class group of E). Let $h^{\text{ord}}(Dp^\infty, \chi_E, O)$ denote the universal p -ordinary Hecke algebra, with coefficients in O , with tame level D and whose prime-to- p part of the Nebentypus equals the character χ_E . Let h_m be the unique local ring of $h^{\text{ord}}(Dp^\infty, \chi_E, O)$ through which the following ring homomorphism factors:

$$\begin{aligned} \lambda_f : h^{\text{ord}}(Dp^\infty, \chi_E, O) &\rightarrow \bar{\mathbb{F}}_p, \\ \text{given by } \begin{cases} \lambda_f(T_l) = \overline{a_l(f)}, & \forall \text{ primes } l \text{ not dividing } Dp, \\ \lambda_f(U_l) = \overline{a_l(f)}, & \forall \text{ primes } l \text{ dividing } Dp. \end{cases} \end{aligned}$$

See [14] for a definition of the Hecke operators T_l and U_l . Also note that h_m is the “full” Hecke algebra generated by the Hecke operators T_l along with the diamond operators $\langle l \rangle$ for primes l not dividing Dp and the U_l operators for primes l dividing Dp . Let $\rho_m : G_{\mathbb{Q}} \rightarrow \text{GL}_2(h_m)$ denote the Galois representation ([4, §2.4]) such that $\text{Trace}(\rho_m(\text{Frob}_l)) = T_l$, for all primes $l \nmid Dp$.

Let CNL_O denote the category of complete, Noetherian, local O -algebras whose residue field equals the residue field of O . Let SETS denote the category of sets. Consider the deformation functor $\mathfrak{F} : \text{CNL}_O \rightarrow \text{SETS}$ that, for every $B \in \text{CNL}_O$, is given by strict equivalence classes of deformations $\varrho : G_{\mathbb{Q}} \rightarrow \text{GL}_2(B)$ that are of type \mathfrak{D} and such that the residual representation associated to the deformation ϱ is isomorphic to $\bar{\rho}_f$. A Galois representation $\varrho : G_{\mathbb{Q}} \rightarrow \text{GL}_2(B)$ is said to be of type \mathfrak{D} if it satisfies the following properties:

- ϱ is unramified at all primes l not dividing Dp .

- The restriction $\varrho \mid_{\text{Gal}(\overline{\mathbb{Q}}_p/\mathbb{Q}_p)}$ to the decomposition group at p is p -ordinary.
- If l is a prime dividing D , we have $\rho \mid_{\text{Gal}(\overline{\mathbb{Q}}_l/\mathbb{Q}_l)} \sim \begin{pmatrix} 1 & 0 \\ 0 & \chi_E \end{pmatrix}$.

Suppose also that the following condition holds:

$$(6.2) \quad 2(k - 1) \not\equiv 0 \pmod{p - 1}.$$

By Theorem 3.9 in [1], the ring h_m along with the Galois representation $\rho_m : G_{\mathbb{Q}} \rightarrow \text{GL}_2(h_m)$ represents the deformation functor \mathfrak{F} . The isomorphism $\bar{\rho}_f \cong \bar{\rho}_f \otimes \chi_E$ induces an involution $\tau : h_m \rightarrow h_m$ that is $O[[x_F]]$ -equivariant. Let h_m^+ denote the subring of h_m fixed by the involution τ . We have the following natural inclusions of $O[[x_F]]$ -algebras:

$$O[[x_F]] \hookrightarrow h_m^+ \hookrightarrow h_m.$$

Theorems 1.1 and 1.2 in [14] give us a natural surjection $h_m \twoheadrightarrow h_{k,m}$. Here, the ring $h_{k,m}$ denotes the local factor of $h_k^{\text{ord}}(\Gamma_1(Dp), \chi_E, O)$ (the “full” Hecke algebra acting on the space of ordinary cuspforms of weight k with Nebentypus χ_E and conductor Dp generated by the operators $T_l^{(k)}$ for primes $l \nmid Dp$ and $U_l^{(k)}$ for primes $l \mid Dp$) through which the following ring homomorphism factors:

$$\beta_f : h_k^{\text{ord}}(\Gamma_1(Dp), \chi_E, O) \rightarrow \overline{\mathbb{F}}_p, \begin{cases} \beta_f(T_l^{(k)}) = \overline{a_l(f)}, & \forall l \nmid Dp, \\ \beta_f(U_l^{(k)}) = \overline{a_l(f)}, & \forall l \mid Dp. \end{cases}$$

Let us now suppose that the following condition holds:

$$(6.3) \quad \text{Rank}_O h_{k,m} = 2.$$

Nakayama’s lemma and the fact h_m is a free $O[[x_F]]$ -algebra ([15, Thm. 3.1]) then let us conclude that $\text{Rank}_{O[[x_F]]} h_m = 2$. Note that by Corollary 3.12 in [3], the natural inclusion $h_m^+ \xrightarrow{\cong} h_m$ is not an isomorphism. So, the natural inclusion $O[[x_F]] \xrightarrow{\cong} h_m^+$ must be an isomorphism. By Theorem 1.1 in [2], we have the following isomorphism of $O[[x_F]]$ -algebras:

$$(6.4) \quad h_m \cong h_m^+[\vartheta] \cong O[[x_F]][\vartheta], \quad \text{where } \vartheta^2 = \sqrt{(1 + x_F)^{\frac{\log(\epsilon)}{\log(1+p)}} - 1}.$$

The function \log denotes the p -adic logarithm on E that is determined by the inclusion $E \hookrightarrow E_{\mathfrak{p}_1}$, where the field $E_{\mathfrak{p}_1}$ denotes the completion of the field E with respect to the valuation $\text{Val}_{\mathfrak{p}_1}$ given by the prime ideal \mathfrak{p}_1 . Note that $E_{\mathfrak{p}_1} \cong \mathbb{Q}_p$. Now, let us further suppose that ϵ is a p -th power in $E_{\mathfrak{p}_1}$. This condition is satisfied precisely when the following inequality holds:

$$(6.5) \quad \text{Val}_{\mathfrak{p}_1}(\epsilon^{p-1} - 1) \geq 2.$$

In this case, the element $(1 + x_F)^{\frac{\log(\epsilon)}{\log(1+p)}} - 1$ is a square-free polynomial in $O[[x_F]]$ and a product of at least 2 distinct irreducible elements of $O[[x_F]]$. As a result, the local ring h_m is an integrally closed domain that is not a UFD. Finally, we let F equal the unique Hida family passing through f .

Remark 6.1. — The Galois representation ρ_m , as constructed in [4, §2.4], takes values in the “reduced” ordinary Hecke algebra h_m^{red} (generated by the Hecke operators T_l and the diamond operators $\langle l \rangle$, for primes l not dividing Dp). Note that h_m^{red} is a subring of h_m . Theorem 3.9 in [1], in fact, identifies the universal deformation ring for the functor \mathfrak{F} with h_m^{red} . It is Proposition 2.4.2 in [4] that provides us an isomorphism $h_m^{\text{red}} \cong h_m$. See [5, Thm. 12] that also addresses this fact.

Table 6.1. $R_F = h_m = O[[x_F]] \left[\sqrt{(1 + x_F)^{\frac{\log(\epsilon)}{\log(1+p)}} - 1} \right]$

$\mathbb{Q}(\sqrt{D})$	(k, p)	ϵ	$\text{Norm}_{E/\mathbb{Q}}(\epsilon^{k-1} - 1)$	$\text{Val}_{p_1}(\epsilon^{p-1} - 1)$	$[K_f : \mathbb{Q}]$
$\mathbb{Q}(\sqrt{10*4})$	(20, 191)	$\sqrt{10} + 3$	$-1 * 2 * 3 * 191^2 * 4523 * 1021973$	2	112
$\mathbb{Q}(\sqrt{33})$	(10, 37)	$4\sqrt{33} + 23$	$-1 * 2^2 * 11 * 37^4 * 47^2 * 71^2$	2	32
$\mathbb{Q}(\sqrt{89})$	(4, 5)	$53\sqrt{89} + 500$	$-1 * 2^3 * 5^3 * 1000003$	3	22
$\mathbb{Q}(\sqrt{89})$	(6, 5)	$53\sqrt{89} + 500$	$-1 * 2^3 * 5^4 * 11 * 18181909091$	3	36
$\mathbb{Q}(\sqrt{629})$	(4, 5)	$\frac{\sqrt{629+25}}{2}$	$-1 * 2^2 * 5^2 * 157$	2	168

Remark 6.2. — We will now explain how to ascertain that $\text{Rank}_O h_{k,m} = 2$. Let us define the following set of Hecke eigenforms of conductor Dp , weight k and Nebentypus χ_E with coefficients in $\overline{\mathbb{Q}}_p$:

$$\mathcal{B}_{k,m,st} = \left\{ g \mid \begin{array}{l} g \text{ is a } p\text{-stabilized Hecke eigenform in } S_k^{\text{ord}}(Dp, \chi_E), \\ a_1(g) = 1, a_l(f) \equiv a_l(g) \pmod{l}, \forall l \nmid Dp \end{array} \right\}.$$

By Corollary 3.7 in [14], it suffices to show that $|\mathcal{B}_{k,m,st}| = 2$. Note that since f and its twist $f \otimes \chi_E$ belong to the set $\mathcal{B}_{k,m,st}$, we have $|\mathcal{B}_{k,m,st}| \geq 2$. For each $g \in \mathcal{B}_{k,m,st}$, let g_0 be the unique newform associated to it ([20, Thm. 3.22]). Throughout this remark, it will be useful to keep in mind that since the weight k is strictly greater than 2 and that the conductor of χ_E is equal to D (and hence of order prime to p), the conductor of g_0 is also equal to D ([13, Prop. 3.1]). We will now use the following code in SAGE to

construct a set \mathcal{B}_{full} , whose $\text{Gal}(E/\mathbb{Q})$ -orbit equals the set of all normalized (new) Hecke eigenforms of weight k , conductor D and Nebentypus χ_E :

```
%sage B_full = Newforms(chi_E, k, names='alpha');
```

For each $g \in \mathcal{B}_{full}$, let K_g denote the number field generated by the Hecke eigenvalues of g . Note that if the p -stabilization of g belongs to $\mathcal{B}_{k,m,st}$ and if l is a prime that remains inert in the extension E/\mathbb{Q} , then l divides $\text{Norm}_{\mathbb{Q}}^{K_g}(a_l(g))$. This is because the residual Galois representation $\bar{\rho}_g$ associated to g (which is isomorphic to $\bar{\rho}_f$) would be isomorphic to $\bar{\rho}_g \otimes \chi_E$ (since $\bar{\rho}_f \cong \bar{\rho}_f \otimes \chi_E$). Define a subset $\mathcal{B}'_{k,m,new}$ of \mathcal{B}_{full} which contains those newforms g such that l divides $\text{Norm}_{\mathbb{Q}}^{K_g}(a_l(g))$, for all primes l that remain inert in the extension E/\mathbb{Q} and such that $3 \leq l \leq 100$. In the examples listed in Table 6.1, $|\mathcal{B}'_{k,m,new}|$ turns out to be 1. And hence in those examples, the $\text{Gal}(E/\mathbb{Q})$ -orbit of $\mathcal{B}'_{k,m,new}$ is in bijection with the set $\mathcal{B}_{k,m,st}$ (the bijection is obtained through the process of p -stabilization) leading us to conclude that $|\mathcal{B}_{k,m,st}| = 2 \times |\mathcal{B}'_{k,m,new}| = 2$. This will ascertain that $\text{Rank}_O h_{k,m} = 2$.

Remark 6.3. — Fixing a quadratic field $\mathbb{Q}(\sqrt{D})$ and letting the primes p vary, it seems computationally difficult to find a large number of primes p such that equation (6.5) holds. William Stein informed us that the complexity of the code in Remark 6.2 is $\mathcal{O}((Dk)^3)$. This explains the paucity of our examples.

BIBLIOGRAPHY

- [1] G. BÖCKLE, “On the density of modular points in universal deformation spaces”, *Am. J. Math.* **123** (2001), no. 5, p. 985-1007.
- [2] S. CHO, “Deformation rings for induced Galois representations”, PhD Thesis, University of California, Los Angeles (USA), 1999.
- [3] S. CHO & V. VATSAL, “Deformations of induced Galois representations”, *J. Reine Angew. Math.* **556** (2003), p. 79-98.
- [4] M. EMERTON, R. POLLACK & T. WESTON, “Variation of Iwasawa invariants in Hida families”, *Invent. Math.* **163** (2006), no. 3, p. 523-580.
- [5] F. Q. GOUVÊA, “Deforming Galois representations: controlling the conductor”, *J. Number Theory* **34** (1990), no. 1, p. 95-113.
- [6] R. GREENBERG, “Iwasawa’s theory and p -adic L -functions for imaginary quadratic fields”, in *Number theory related to Fermat’s last theorem*, Progress in Math., vol. 26, Birkhäuser, 1982, p. 275-285.
- [7] ———, “Iwasawa theory and p -adic deformations of motives”, in *Motives (Seattle, WA, 1991)*, Proceedings of Symposia in Pure Mathematics, vol. 55, American Mathematical Society, 1994, p. 193-223.
- [8] ———, “On the structure of certain Galois cohomology groups”, *Doc. Math.* (2006), p. 335-391.

- [9] ———, “Surjectivity of the global-to-local map defining a Selmer group”, *Kyoto J. Math.* **50** (2010), no. 4, p. 853-888.
- [10] ———, “On the structure of Selmer groups”, in *Elliptic curves, modular forms and Iwasawa theory*, Springer Proc. Math. Stat., vol. 188, Springer, 2016, p. 225-252.
- [11] R. GREENBERG & V. VATSAL, “On the Iwasawa invariants of elliptic curves”, *Invent. Math.* **142** (2000), no. 1, p. 17-63.
- [12] T. HARA & T. OCHIAI, “The cyclotomic Iwasawa main conjecture for Hilbert cuspforms with complex multiplication”, *Kyoto J. Math.* **58** (2018), no. 1, p. 1-100.
- [13] H. HIDA, “A p -adic measure attached to the zeta functions associated with two elliptic modular forms. I”, *Invent. Math.* **79** (1985), no. 1, p. 159-195.
- [14] ———, “Galois representations into $GL_2(\mathbf{Z}_p[[X]])$ attached to ordinary cusp forms”, *Invent. Math.* **85** (1986), no. 3, p. 545-613.
- [15] ———, “Iwasawa modules attached to congruences of cusp forms”, *Ann. Sci. Éc. Norm. Supér.* **19** (1986), no. 2, p. 231-273.
- [16] ———, “A p -adic measure attached to the zeta functions associated with two elliptic modular forms. II”, *Ann. Inst. Fourier* **38** (1988), no. 3, p. 1-83.
- [17] ———, *Elementary theory of L -functions and Eisenstein series*, London Mathematical Society Student Texts, vol. 26, Cambridge University Press, 1993, xii+386 pages.
- [18] ———, “On the search of genuine p -adic modular L -functions for $GL(n)$ ”, *Mém. Soc. Math. Fr., Nouv. Sér.* (1996), no. 67, p. 1-110, With a correction to: “On p -adic L -functions of $GL(2) \times GL(2)$ over totally real fields”, *Ann. Inst. Fourier* **41** (1991), no. 2, p. 311-391.
- [19] ———, “Global quadratic units and Hecke algebras”, *Doc. Math.* **3** (1998), p. 273-284.
- [20] ———, *Modular forms and Galois cohomology*, Cambridge Studies in Advanced Mathematics, vol. 69, Cambridge University Press, 2000, x+343 pages.
- [21] H. HIDA & J. TILOUINE, “On the anticyclotomic main conjecture for CM fields”, *Invent. Math.* **117** (1994), no. 1, p. 89-147.
- [22] M.-L. HSIEH, “Eisenstein congruence on unitary groups and Iwasawa main conjectures for CM fields”, *J. Am. Math. Soc.* **27** (2014), no. 3, p. 753-862.
- [23] G. KINGS, D. LOEFFLER & S. L. ZERBES, “Rankin-Eisenstein classes and explicit reciprocity laws”, *Camb. J. Math.* **5** (2017), no. 1, p. 1-122.
- [24] A. LEI, D. LOEFFLER & S. L. ZERBES, “Euler systems for Rankin-Selberg convolutions of modular forms”, *Ann. Math.* **180** (2014), no. 2, p. 653-771.
- [25] Q. LIU, *Algebraic geometry and arithmetic curves*, Oxford Graduate Texts in Mathematics, vol. 6, Oxford University Press, 2002, Translated from the French by Reinie Ern e, xvi+576 pages.
- [26] J. VAN ORDER, “Rankin-Selberg L -functions in cyclotomic towers, I”, <https://arxiv.org/abs/1207.1672>, 2012.
- [27] ———, “Rankin-Selberg L -functions in cyclotomic towers, II”, <https://arxiv.org/abs/1207.1673>, 2012.
- [28] B. PALVANNAN, “On Selmer groups and factoring p -adic L -functions”, *Int. Math. Res. Not.* (2017), published online, to appear in print.
- [29] THE SAGE DEVELOPERS, “SageMath, the Sage Mathematics Software System (Version 7.2)”, <http://www.sagemath.org>.

Manuscrit reçu le 22 août 2016,
révisé le 19 septembre 2017,
accepté le 2 février 2018.

Bharathwaj PALVANNAN
Department of Mathematics
University of Pennsylvania
209 South 33rd Street, 4N53 DRL
Philadelphia 19104-6395 (USA)
pbharath@math.upenn.edu