



ANNALES DE L'INSTITUT FOURIER

Stéphanie NIVOCHÉ

Transcendental versions in \mathbb{C}^n of the Nagata conjecture

Tome 71, n° 1 (2021), p. 27-52.

<http://aif.centre-mersenne.org/item/AIF_2021__71_1_27_0>

© Association des Annales de l'institut Fourier, 2021,

Certains droits réservés.



Cet article est mis à disposition selon les termes de la licence

CREATIVE COMMONS ATTRIBUTION – PAS DE MODIFICATION 3.0 FRANCE.

<http://creativecommons.org/licenses/by-nd/3.0/fr/>



Les Annales de l'institut Fourier sont membres du

Centre Mersenne pour l'édition scientifique ouverte

www.centre-mersenne.org

TRANSCENDENTAL VERSIONS IN \mathbb{C}^n OF THE NAGATA CONJECTURE

by Stéphanie NIVOCHÉ (*)

ABSTRACT. — The Nagata Conjecture is one of the most intriguing open problems in the area of curves in the plane. It is easily stated. Namely, it predicts that the smallest degree d of a plane curve passing through $r \geq 10$ general points in the projective plane \mathbb{P}^2 with multiplicities at least l at every point, satisfies the inequality $d > \sqrt{r} \cdot l$. This conjecture has been proven by M. Nagata in 1959, if r is a perfect square greater than 9. Up to now, it remains open for every non-square $r \geq 10$, after more than a half century of attention by many researchers.

In this paper, we formulate new transcendental versions of this conjecture coming from pluripotential theory and which are equivalent to a version in \mathbb{C}^n of the Nagata Conjecture.

RÉSUMÉ. — La conjecture de Nagata est l'un des problèmes ouverts les plus intrigants dans le domaine des courbes du plan complexe. Elle s'énonce simplement. En effet, elle affirme que le plus petit degré d d'une courbe plane passant par $r \geq 10$ points généraux dans le plan projectif \mathbb{P}^2 avec des multiplicités au moins l en chaque point, satisfait l'inégalité $d > \sqrt{r} \cdot l$. Cette conjecture a été vérifiée par M. Nagata en 1959, si r est un carré parfait strictement supérieur à 9. Jusqu'à présent, elle est restée ouverte pour tout entier $r \geq 10$ non carré, après plus d'un demi-siècle d'attention de la part de nombreux chercheurs.

Dans cet article, nous formulons de nouvelles versions transcendentales de cette conjecture issues de la théorie du pluripotentiel, et qui sont équivalentes à une version dans \mathbb{C}^n de la conjecture Nagata.

1. Introduction

1.1. History and known results

In 1900, at the international Congress of Mathematicians in Paris, D. Hilbert posed twenty-three problems. The fourteenth one may be formulated as follows:

Keywords: Nagata conjecture, pluripotential theory, pluricomplex Green function.

2020 Mathematics Subject Classification: 14H50, 32U10, 32U25, 32U35, 32U40, 32W20.

(*) Research of S. Nivoche was supported by ANR grant “ANR-14-CE34-0002-01” and the FWF grant I1776 for the international cooperation project “Dynamics and CR geometry”.

Let \mathbb{K} be a field and x_1, \dots, x_n algebraically independent elements over \mathbb{K} . Let L be a subfield of $\mathbb{K}(x_1, \dots, x_n)$ containing \mathbb{K} . Is the ring $\mathbb{K}[x_1, \dots, x_n] \cap L$ finitely generated over \mathbb{K} ?

Hilbert conjectured that all such algebras are finitely generated over \mathbb{K} . Contributions to the fourteenth problem are obtained confirming Hilbert's conjecture in special cases and for certain classes of rings. In 1953, a significant contribution was made by O. Zariski, who generalized the fourteenth problem in the following way:

PROBLEM OF ZARISKI. — Let \mathbb{K} be a field and $\mathbb{K}[a_1, \dots, a_n]$ an affine normal domain (i.e. a finitely generated integrally closed domain over \mathbb{K}). Let L be a subfield of $\mathbb{K}(a_1, \dots, a_n)$ containing \mathbb{K} . Is the ring $\mathbb{K}[a_1, \dots, a_n] \cap L$ finitely generated over \mathbb{K} ?

He answered the question in the affirmative when $\text{trans.deg}_{\mathbb{K}} L \leq 2$. Later, in 1957, D. Rees gave a counter example to the problem of Zariski when $\text{trans.deg}_{\mathbb{K}} L = 3$. Finally Masayoshi Nagata in [24] gave a counter example to the original fourteenth problem itself. This counter example is in the case of $\text{trans.deg}_{\mathbb{K}} L = 13$.

In 1959, M. Nagata [22] gave another counter example (a suitably constructed ring of invariants for the action of a linear algebraic group) in the case of $\text{trans.deg}_{\mathbb{K}} L = 4$. In this work, he finally formulated a conjecture, which governs the minimal degree required for a plane algebraic curve to pass through a collection of general points with prescribed multiplicities:

Suppose p_1, \dots, p_r are r general points in \mathbb{P}^2 and that m_1, \dots, m_r are given positive integers. Then for $r > 9$, any curve C in \mathbb{P}^2 that passes through each of the points p_i with multiplicity m_i must satisfy $\deg C > \frac{1}{\sqrt{r}} \cdot \sum_{i=1}^r m_i$.

One says that a property \mathcal{P} holds for r general points in \mathbb{P}^2 if there is a Zariski-open subset W of $(\mathbb{P}^2)^r$ such that \mathcal{P} holds for every set (p_1, \dots, p_r) of r points in W . As Nagata pointed out, it is enough to consider the uniform case. Thus this conjecture is equivalent to the following one, which it usually called:

THE NAGATA CONJECTURE. — Suppose p_1, \dots, p_r are r general points in \mathbb{P}^2 and that m is a given positive integer. Then for $r > 9$, any curve C in \mathbb{P}^2 that passes through each of the points p_i with multiplicity at least m must satisfy $\deg C > \sqrt{r} \cdot m$.

The only case when this is known to hold true is when $r > 9$ is a perfect square. This was proved by Nagata with technics of specializations of the

points in the plane. More recently, without any extra condition connecting the degree, m and r , G. Xu [30] proved that $\deg C \geq \frac{\sqrt{r-1}}{r} \cdot \sum_{i=1}^r m_i$ and H. Tutaj-Gasinska [27] proved that $\deg C \geq \frac{1}{\sqrt{r+(1/12)}} \cdot \sum_{i=1}^r m_i$ (see also B. Harbourne [13] and B. Harbourne and J. Roé [14]).

A more modern formulation of this conjecture is often given in terms of Seshadri constants, introduced by J.-P. Demailly ([8] and [10]) in the course of his work on Fujita’s conjecture. The Nagata Conjecture is generalized to other surfaces under the name of the Nagata–Biran conjecture (let X be a smooth algebraic surface and L be an ample line bundle on X of degree d . The Nagata–Biran conjecture states that for sufficiently large r the Seshadri constant satisfies $\epsilon(p_1, \dots, p_r; X, L) = \frac{d}{\sqrt{r}}$). Nagata has also remarked that the condition $r > 9$ is necessary. The cases $r > 9$ and $r \leq 9$ are distinguished by whether or not the anti-canonical bundle on the blowup of \mathbb{P}^2 at a collection of r points is nef.

Fix r general points p_1, \dots, p_r in \mathbb{P}^2 and a nonnegative integer m . Define $\delta(r, m)$ to be the least integer d such that there is a curve of degree d vanishing at each point p_i with multiplicity at least m . For $r \leq 9$, applying methods of [23], Harbourne [13] shows that $\delta(r, m) = \lceil c_r m \rceil$, where $c_r = 1; 1; 3/2; 2; 2; 12/5; 21/8; 48/17$ and 3 for $r = 1, \dots, 9$ respectively (for any real number c , $\lceil c \rceil$ is the smallest integer larger than or equal to c). When $r = 1, 4$ and 9 , we can remark that the Nagata conjecture holds with a weak inequality instead of a strict one.

Iarrobino [16] generalized Nagata Conjecture in any dimension $n \geq 2$. Based on a conjecture of Fröberg, Iarrobino predicted that:

An hypersurface in \mathbb{P}^n passing through r generic points with multiplicity m has a degree $> r^{1/n} \cdot m$, except for an explicit finite list of (r, n) .

L. Evain [12] proved this conjecture when the number of points in \mathbb{P}^n is of the form s^n (in this case the list is $(4, 2)$, $(9, 2)$ and $(8, 3)$).

An affirmative answer to the Nagata conjecture would provide important applications in the theory of linear systems in the projective plane. It would also provide information for the study of the singular degrees (works of Bombieri, Skoda, Waldschmidt, Chudnovsky) with a lot of applications in number theory (arithmetic nature of values of Abelian functions of several variables), in symplectic geometry (symplectic packings in the unit ball) and in algebraic geometry (study of multiple-point Seshadri constants for generic rational surfaces).

During the last three/four decades, the use of pluripotential theory in analytic/algebraic geometry has been very fruitful since it allows a lot

of flexibility while keeping track of the analytic features. This point of view was proven to be very efficient by J.-P. Demailly, Y.-T. Siu and their schools, to name a few. Our idea in this paper is to develop transcendental techniques, to overcome the intrinsic rigidity of polynomials and to obtain a new approach to this problem of algebraic geometry. Instead of considering complex polynomials, we work with plurisubharmonic (written psh for simplicity) functions, having logarithmic poles at prescribed points. These last functions are much more flexible than the first ones (Lelong). See Section 2.1 for some recalls about pluripotential theory. Thus two points of view are possible. A global one and a local one. We can consider psh functions in all \mathbb{C}^n , with a logarithmic behavior at infinity. This class of plurisubharmonic functions contains in particular the logarithm of modulus of polynomials, with prescribed properties in the Nagata conjecture. The subclass of plurisubharmonic functions which are maximal (for the complex Monge–Ampère operator) outside prescribed points is of particular interest. They inevitably have to satisfy certain conditions of growth at infinity. We look for the minimal growth that they can have. In [5], D. Coman and S. Nivoche have obtained some preliminary results about this subject and have established a link between a quantity of the same nature such as the singular degrees of M. Waldschmidt [28], [29], and a quantity coming directly from pluripotential theory. We can also consider psh functions in a bounded domain in \mathbb{C}^n , with logarithmic poles at prescribed points and with zero value at the boundary. We study in particular the subclass of pluricomplex Green functions in this domain with logarithmic poles of weight 1 at fixed points. Especially we look at what happens when the poles collide to a single point in the domain. The nature of the logarithmic singularity of the limit function is in connection with the algebraic properties of the set of fixed points and its singular degree of M. Waldschmidt.

In this paper, we establish a link between some well known algebraic quantities of the same nature as the singular degree of M. Waldschmidt, and some others quantities coming directly from pluripotential theory. We outline new conjectures in term of pluripotential theory, and we prove that they are all equivalent to the Nagata conjecture.

1.2. Results

1.2.1. A first conjecture of pluripotential theory

We are going to study the convergence of multipole pluricomplex Green functions for a bounded hyperconvex domain in \mathbb{C}^n , in the case where

poles contract to one single point. First we start with the case where the domain is the unit ball centered at the origin. Let S be a finite set of distinct points in \mathbb{C}^n and denote $|S|$ its cardinality. Let R be a positive real number sufficiently large such that $S \subset B(O, R)$. Denote by $g_R(S, \cdot)$ the pluricomplex Green function in ball $B(O, R)$ with logarithmic poles in S , of weight one. $g_R(S, z) = g_1(S/R, z/R)$ for any $z \in B(O, R)$. Thus its is natural to study g_∞ a negative psh function defined in the unit ball $B(O, 1)$ by the following upper semi-continuous regularization

$$g_\infty(z) = \left(\limsup_{t \in \mathbb{C}^* \rightarrow 0} g_1(tS, z) \right)^* .$$

Clearly, if we replace the set S by λS , where $\lambda \in \mathbb{C}^*$, then we obtain the same limit function g_∞ . $\|\cdot\|$ is the Euclidean norm in \mathbb{C}^n and $B(z_0, R)$ is the hermitian ball centered at z_0 with radius R . g_∞ always satisfies in $B(O, 1)$ the following inequalities

$$|S| \cdot \ln \|z\| \leq g_\infty(z) \leq g_{B(O,1)}(O, z) = \ln \|z\| .$$

It tends to 0 on the boundary of the unit ball and it has an unique logarithmic singularity at the origin. Let us also consider another family $(\tilde{g}_1(tS, \cdot))_{t \in \mathbb{C}^*}$ of continuous and psh functions, defined in $B(O, 1)$ by $\tilde{g}_1(tS, z) = \sup\{g_1(tS, w) : \|w\| = \|z\|\}$ and the following continuous and psh function in $B(O, 1)$

$$\tilde{g}_\infty(z) = \sup\{g_\infty(w) : \|w\| = \|z\|\} .$$

No upper regularization is needed in this definition, since it is a convex increasing (hence continuous) function of $\log \|z\|$. We want to study precisely the nature of the logarithmic singularity of these functions g_∞ and \tilde{g}_∞ at the origin. In particular we want to understand what their Lelong numbers at the origin are.

We recall that if u is a psh function, then the *classical Lelong number* $\nu(u, z)$ of u at a point z is (P. Lelong, 1969) the $(2n - 2)$ -dimensional density of the measure $dd^c u$ at z :

$$\nu(u, z) := \lim_{r \rightarrow 0} \frac{1}{(\pi r^2)^{n-1}} \int_{|w-z| < r} dd^c u \wedge (dd^c |w - z|^2)^{n-1} .$$

We can also compute this number as follows (V. Avannissian, C. Kiselman):

$$\nu(u, z) = \lim_{y \rightarrow -\infty} \frac{\sup_{|w|=1} u(z + we^y)}{y} = \lim_{y \rightarrow -\infty} \frac{1}{y} \int_{|w|=1} u(z + we^y) d\tilde{\lambda}(w) ,$$

where $d\tilde{\lambda}$ is the normalized surface measure on the unit sphere.

There already exist some results about the convergence of multipole Green functions in [20], [26] and [11]. In this paper, we establish a direct connection between the nature of the logarithmic singularity of these psh functions g_∞ and \tilde{g}_∞ and the algebraic properties of the set S . More precisely, by using a Schwarz' Lemma for finite sets ([21], [29]) and a generalization of a result in [25], we prove in Section 2.3 the following theorem which describes properties of these two psh functions, in connection with $\Omega(S)$, the singular degree of S introduced by Waldschmidt ([28], [3]) and which is an affine invariant in connection with the Nagata Conjecture. For any polynomial $P \in \mathbb{C}[z] = \mathbb{C}[z_1, \dots, z_n]$, $\deg P$ is its degree and $\text{ord}(P, p)$ denotes the vanishing order of P at any point p . If l is a positive integer we define

$$\Omega(S, l) = \min\{\deg P : P \in \mathbb{C}[z], \text{ord}(P, p) \geq l, \forall p \in S\}.$$

The limit

$$\Omega(S) := \lim_{l \rightarrow +\infty} \Omega(S, l)/l = \inf_{l \geq 1} \Omega(S, l)/l$$

exists and is called the singular degree of S . On the other hand, according to Theorem 1.1 (and Example 5.8) in [26], we already know that the family $(g_1(tS, \cdot))_{t \in \mathbb{C}^*}$ converges locally uniformly outside the origin in $B(O, 1)$ to g_∞ .

THEOREM 1.1. — *Let S be a finite set of points in \mathbb{C}^n . The two psh functions g_∞ and \tilde{g}_∞ satisfy several properties:*

- (i) $\nu(g_\infty, O) = \Omega(S)$ and $(\text{dd}^c g_\infty)^n = 0$ in $B(O, 1) \setminus \{O\}$. $\Omega(S)^n \leq (\text{dd}^c g_\infty)^n(\{O\}) \leq \int_{B(O, 1)} (\text{dd}^c g_\infty)^n \leq |S|$ and we have

$$g_\infty(z) \leq \Omega(S) \ln \|z\|, \text{ in } B(O, 1).$$

- (ii) *The family $(\tilde{g}_1(tS, \cdot))_{t \in \mathbb{C}^*}$ converges locally uniformly outside the origin in $B(O, 1)$ to \tilde{g}_∞ which is equal to $\Omega(S) \ln \|z\|$ in $\bar{B}(O, 1)$.*
 (iii) *If $\Omega(S) = |S|^{1/n}$ then $(\text{dd}^c g_\infty)^n(\{O\}) = \Omega(S)^n = |S|$ and*

$$g_\infty(z) = |S|^{1/n} \ln \|z\|, \text{ in } B(O, 1).$$

- (iv) *Conversely if g_∞ is equal to $\Omega(S) \ln \|\cdot\|$ in $\bar{B}(O, 1)$, then $\Omega(S) = |S|^{1/n}$.*

We will deduce several applications from this theorem, in particular an equivalence between the following conjecture of pluripotential theory in \mathbb{C}^n and a weak version of the Nagata Conjecture in \mathbb{P}^n .

CONJECTURE (\mathcal{P}_1). — In \mathbb{C}^n , except for a finite number of integer values r , for any general set $S = \{p_1, \dots, p_r\}$ of r points, the family of pluricomplex Green functions $(g_{B(O,1)}(tS, \cdot))_{t \in \mathbb{C}^*}$ converges locally uniformly outside the origin of $B(O, 1)$ to $r^{1/n}g_{B(O,1)}(O, \cdot)$, when t tends to 0.

CONJECTURE (\mathcal{A}_1) (Weak Version of the Nagata Conjecture in \mathbb{P}^n). — In \mathbb{P}^n , except for a finite number of integer values r , for any general set $S = \{p_1, \dots, p_r\}$ of r points, $\Omega(S) = r^{1/n}$.

With the homogeneous coordinates $[z_0 : z_1 : \dots : z_n]$ in \mathbb{P}^n , we know that \mathbb{P}^n is a complex manifold of dimension n , which is covered by n copies $U_i = \{[z_0 : z_1 : \dots : z_n] \in \mathbb{P}^n \mid z_i \neq 0\}$ of \mathbb{C}^n . If $S = \{p_1, \dots, p_r\}$ is a set of r distinct points in \mathbb{P}^n , in the previous definition of $\Omega(S)$, we have just to replace polynomials $P \in \mathbb{C}[z] = \mathbb{C}[z_1, \dots, z_n]$ by homogeneous polynomials in $\mathbb{C}[z_0, z_1, \dots, z_n]$. We can remark that if $n = 2$, this Weak version of the Nagata Conjecture is identical to the original one when the number of points is not a perfect square. More generally in \mathbb{P}^n , a stronger version of the Weak version of the Nagata Conjecture is satisfied for a number of points of the form s^n , according to [12]. When the number of points is not of the form s^n , this weak version corresponds to Iarrobino’s conjecture in \mathbb{P}^n .

In the previous construction of this function g_∞ , we can replace the unit ball $B(O, 1)$ by any bounded hyperconvex domain D in \mathbb{C}^n (see the definition in Section 2.1) and the origin by any point z_o in D . In this case the function g_∞ is defined in D by

$$\left(\limsup_{t \in \mathbb{C}^* \rightarrow 0} g_D(z_o + tS, \cdot) \right)^*$$

where $g_D(z_o + tS, \cdot)$ is the pluricomplex Green function in D with logarithmic poles of weight one at any points of the set $z_o + tS$. g_∞ has similar properties as in Theorem 1.1, where we replace the pluricomplex Green function in the unit ball with logarithmic pole at the origin $\ln \|\cdot\| = g_{B(O,1)}(O, \cdot)$ by the pluricomplex Green function in D with a logarithmic pole at z_o , $g_D(z_o, \cdot)$. First, according to Theorem 1.1 (and Example 5.8) in [26], the family $(g_D(z_o + tS, \cdot))_{t \in \mathbb{C}^*}$ converges locally uniformly outside z_o in D to g_∞ . In addition, this function satisfies the following theorem.

THEOREM 1.2. — Let S be a finite set of points in \mathbb{C}^n . Let D be a bounded hyperconvex domain in \mathbb{C}^n . Fix z_o in D . The psh function g_∞ satisfies several properties:

- (i) $\nu(g_\infty, z_o) = \Omega(S)$ and $(dd^c g_\infty)^n = 0$ in $D \setminus \{z_o\}$. Also $\Omega(S)^n \leq (\ddot{d}^c g_\infty)^n(\{z_o\}) \leq \int_D (\ddot{d}^c g_\infty)^n \leq |S|$ and we have

$$g_\infty(z) \leq \Omega(S)g_D(z_o, z), \text{ in } D.$$

- (ii) If $\Omega(S) = |S|^{1/n}$ then $(\ddot{d}^c g_\infty)^n(\{z_o\}) = \Omega(S)^n = |S|$ and

$$g_\infty(z) = |S|^{1/n}g_D(z_o, z), \text{ in } D.$$

- (iii) Conversely if g_∞ is equal to $\Omega(S)g_D(z_o, \cdot)$ in D , then $\Omega(S) = |S|^{1/n}$.

We shall not develop the proof of this theorem, because it is very similar to the proof of Theorem 1.1. Indeed, if D is a bounded hyperconvex domain in \mathbb{C}^n and z_o is any point in D , there exists two positive real numbers R_1 and R_2 such that $B(z_o, R_1) \subset D \subset B(z_o, R_2)$,

$$g_{B(z_o, R_2)}(z_o + tS, z) \leq g_D(z_o + tS, z) \leq g_{B(z_o, R_1)}(z_o + tS, z) \text{ in } B(z_o, R_1)$$

and

$$g_{B(z_o, R_i)}(z_o + tS, z) = g_{B(O, 1)}\left(\frac{tS}{R_i}, \frac{z - z_o}{R_i}\right) \text{ in } B(z_o, R_i), \text{ for } i = 1, 2.$$

Thus we have

$$g_{B(O, 1)}\left(\frac{tS}{R_2}, \frac{z - z_o}{R_2}\right) \leq g_D(z_o + tS, z) \leq g_{B(O, 1)}\left(\frac{tS}{R_1}, \frac{z - z_o}{R_1}\right), \text{ in } B(z_o, R_1).$$

1.2.2. New affine invariants and others conjectures of pluripotential theory

Instead of considering psh functions in bounded domains in \mathbb{C}^n , here we study a class of entire psh functions in \mathbb{C}^n , with logarithmic poles in a finite set of points S and with a logarithmic growth at infinity. In particular we are interested in the subclass of such functions which are also locally bounded outside of S . If u is a psh function in \mathbb{C}^n , let us denote γ_u the following upper limit:

$$\gamma_u := \limsup_{|z| \rightarrow \infty} \frac{u(z)}{\log \|z\|} \in [0, +\infty].$$

If $S = \{p_1, \dots, p_r\} \subset \mathbb{C}^n$ is a finite set of distinct points, we associate to any psh function u in \mathbb{C}^n , a number $\omega(S, u)$ defined by

$$\omega(S, u) := \frac{\sum_{j=1}^k \nu(u, p_j)}{\gamma_u}.$$

For instance, if the psh function u is of the form $\ln|Q|$ where Q is a polynomial, then $\nu(u, z)$ is the vanishing order of Q at z and γ_u is the degree of Q . We associate to S an affine invariant $\omega_{\text{psh}}(S)$ defined by

$$\omega_{\text{psh}}(S) := \sup\{\omega(S, u) : u \in \text{PSH}(\mathbb{C}^n)\}.$$

We also consider the class of psh functions in \mathbb{C}^n which are locally bounded in $\mathbb{C}^n \setminus S$. We set

$$\omega_{\text{psh}}^+(S) := \sup\{\omega(S, u) : u \in \text{PSH}(\mathbb{C}^n) \cap L_{\text{loc}}^\infty(\mathbb{C}^n \setminus S)\}.$$

$\omega_{\text{psh}}^+(S)$ is also an affine invariant. Then naturally, a second conjecture of pluripotential theory can be formulated as follow:

CONJECTURE (\mathcal{P}_2). — *In \mathbb{C}^n , except for a finite number of integer values r , for any general set $S = \{p_1, \dots, p_r\}$ of r points, $\omega_{\text{psh}}(S) = \omega_{\text{psh}}^+(S)$.*

This problem is non-trivial and it is related to the algebraic geometric properties of S . There exists another well known invariant of algebraic geometry defined by

$$\omega(S) := \sup \left\{ \frac{\sum_{j=1}^r \text{ord}(P, p_j)}{\text{deg } P} : P \in \mathbb{C}[z] \right\}.$$

$\omega(S)$ is an affine invariant, related to the singular degree of S . G.V. Chudnovsky defined in [3] the very singular degree of S , $|S|/\omega(S)$. Since for any positive integer l , $\omega(S) \geq \frac{|S|l}{\Omega(S,l)}$, the following relation between $\omega(S)$ and $\Omega(S)$ is always satisfied:

$$\omega(S) \geq \frac{|S|}{\Omega(S)}.$$

In Section 1.1, we have stated a general version of the Nagata conjecture (see also Harbourne in [13]): *if $r > 9$ then for generic set $S \subset \mathbb{P}^2$ with $|S| = r$ one has $\sum_{j=1}^r \text{ord}(P, p_j) < \sqrt{r} \text{deg } P$, for every polynomial $P \in \mathbb{C}[z]$.* As Nagata has pointed out, this version is equivalent to the uniform one “the usual Nagata conjecture”. More generally in \mathbb{P}^n , we state the following conjecture of algebraic geometry.

CONJECTURE (\mathcal{A}_2). — *In \mathbb{P}^n , except for a finite number of integer values r , for any general set $S = \{p_1, \dots, p_r\}$ of r points, $\omega(S) = |S|^{1-\frac{1}{n}}$.*

It is well known that Conjectures (\mathcal{A}_1) and (\mathcal{A}_2) are equivalent. Finally we can state a last conjecture of pluripotential theory, which can be seen as the dual version of the first one (\mathcal{P}_1).

CONJECTURE (\mathcal{P}_3). — In \mathbb{C}^n , except for a finite number of integer values r , for any general set S of r points, we have: for any $\epsilon > 0$, there exists an entire continuous psh function v in $L_{\text{loc}}^\infty(\mathbb{C}^n \setminus S)$, such that $\nu(v, p) \geq 1$ for any $p \in S$ and $\gamma_v \leq (1 + \epsilon)|S|^{1/n}$.

THEOREM 1.3. — Each conjecture (\mathcal{P}_1), (\mathcal{P}_2) and (\mathcal{P}_3) is equivalent to (\mathcal{A}_1) and (\mathcal{A}_2).

We will prove Theorem 1.3 in Section 3.3.

Acknowledgments

I would like to thank Professors André Hirschowitz and Joaquim Roé for helpful discussions.

2. Conjecture (\mathcal{P}_1) in connection with pluricomplex Green functions in domains in \mathbb{C}^n

2.1. Some recalls of pluripotential theory

For a bounded domain $D \subset \mathbb{C}^n$, the pluricomplex Green function in D with logarithmic poles in a finite subset S of D , generalizes the one-variable Green functions (for the Laplacian). It is defined by

$$g_D(S, z) = \sup \left\{ u(z) : \begin{array}{l} u \text{ psh on } D, u \leq 0, \\ u(z) \leq \ln \|z - p\| + O(1) \text{ for any point } p \text{ in } S \end{array} \right\}.$$

([6], [7], [17], [18], [19]). If D is hyperconvex (i.e. there exists an exhaustive continuous psh function $D \rightarrow (-\infty, 0)$) then we have an alternative description of the pluricomplex Green functions in terms of the complex Monge–Ampère operator, namely $g_D(S, \cdot)$ is the unique solution to the following Dirichlet problem:

$$\begin{cases} u \text{ is plurisubharmonic and negative on } D, \text{ continuous on } \bar{D} \setminus S, \\ (dd^c u)^n = 0 \text{ on } D \setminus S, \\ u(z) = \ln \|z - p\| + O(1) \text{ as } z \rightarrow p, \forall p \in S, \\ u(z) \rightarrow 0 \text{ as } z \rightarrow \partial D. \end{cases}$$

In this case, $(dd^c u)^n = \sum_{p \in S} \delta_p$.

The complex Monge–Ampère operator is a good candidate to replace Laplacian in one variable. But a important difference between cases of one

variable and several variables is that this complex Monge–Ampère operator is not linear anymore. Here $d = \partial + \bar{\partial} = \sum_{j=1}^n \frac{\partial}{\partial z_j} + \sum_{j=1}^n \frac{\partial}{\partial \bar{z}_j}$ and $d^c = \frac{i}{2\pi}(\bar{\partial} - \partial)$. $dd^c = \frac{i}{\pi}\partial\bar{\partial}$. The normalization of the d^c operator is chosen such that we have precisely $(dd^c \ln\|z - p\|)^n = \delta_p$, the Dirac measure at point p . The n th exterior power of $dd^c = \frac{i}{\pi}\partial\bar{\partial}$, i.e. $(dd^c)^n = dd^c \wedge \dots \wedge dd^c$ (n times), defines the complex Monge–Ampère operator in \mathbb{C}^n . If $u \in \mathcal{C}^2(D)$, then $(dd^c u)^n = \frac{2^n n!}{\pi^n} \det[\frac{\partial^2 u}{\partial z_j \partial \bar{z}_k}] dV$, where $dV = (\frac{i}{2})^n dz_1 \wedge d\bar{z}_1 \wedge \dots \wedge dz_n \wedge d\bar{z}_n$ is the usual volume form in \mathbb{C}^n . It is a positive measure, defined inductively for locally bounded psh functions according to the definition of Bedford–Taylor [1], [2], and it can also be extended to psh functions with isolated or compactly supported poles [9].

2.2. The singular degree of a finite set

The singular degree of S introduced by Waldschmidt [29] (see also [3]) is an affine invariant in connection with the Nagata Conjecture. If l is a positive integer we define

$$\Omega(S, l) = \min\{\deg P : P \in \mathbb{C}[z], \text{ord}(P, p_j) \geq l, 1 \leq j \leq r\}.$$

$\Omega(S, 1)$ is sometimes called the degree of S . We clearly have $\Omega(S, l_1 + l_2) \leq \Omega(S, l_1) + \Omega(S, l_2)$, and in particular $\Omega(S, l) \leq \Omega(S, 1)l$. The limit

$$\Omega(S) := \lim_{l \rightarrow +\infty} \frac{\Omega(S, l)}{l}$$

exists and is called the singular degree of S . We have for each $l \geq 1$

$$\frac{\Omega(S, 1)}{n} \leq \Omega(S) = \inf_l \frac{\Omega(S, l)}{l} \leq \frac{\Omega(S, l)}{l} \leq \Omega(S, 1).$$

The second, the third and the fourth inequalities are trivial, while the proof of the first one uses complex analysis. By using Hormander–Bombieri–Skoda theorem, M. Waldschmidt proved more generally that for any positive integer l_1 and l_2

$$\frac{\Omega(S, l_1)}{l_1 + n - 1} \leq \Omega(S) \leq \frac{\Omega(S, l_2)}{l_2}.$$

Upper bound for the numbers $\Omega(S, l)$ is well known. It is a result of Waldschmidt ([29, Lemma 1.3.13]):

$$\Omega(S, l) \leq (l + n - 1)|S|^{1/n} - (n - 1).$$

And a consequence is $\Omega(S) \leq |S|^{1/n}$. An important and difficult problem is to find a lower bound for $\Omega(S)$. The Nagata Conjecture can be stated again in term of the invariants $\Omega(S, l)$:

In \mathbb{C}^2 , if $r > 9$, then $\Omega(S, l) > l\sqrt{r} \forall l \geq 1$ holds for a set S of r points in general position.

This statement doesn't hold for $r \leq 9$.

2.3. Proof of Theorem 1.1

Let S be a finite set of distinct points in \mathbb{C}^n . We use the same notation as in Section 1.2.1. For any $t \in \mathbb{C}^*$ sufficiently small, the function $g_1(tS, \cdot)$ always satisfies the following inequalities

$$\sum_{p \in S} g_1(tp, z) \leq g_1(tS, z) \leq \inf_{p \in S} g_1(tp, z),$$

where $g_1(tp, \cdot)$ is the pluricomplex Green function with an unique logarithmic pole at tp with weight one. Consequently, the psh function g_∞ satisfies in $B(O, 1)$

$$|S| \cdot \ln \|z\| \leq g_\infty(z) \leq g_1(O, z) = \ln \|z\|.$$

In addition, this function tends to 0 on the boundary of the unit ball and it has an unique singularity at the origin. The functions $g_1(tS, \cdot)$ are continuous in $\overline{B}(O, 1)$ valued in $[-\infty, 0]$ and psh in $B(O, 1)$ with a finite set of singularities tS .

The following very simple example, helps us to understand what different situations can occur in Theorem 1.1.

Example 2.1. — Let $S = \{(1/2, 0), (-1/2, 0)\}$ in \mathbb{C}^2 . We know that $\Omega(S) = 1$. To simplify computation, we use the sup norm $|z| = \max\{|z_1|, |z_2|\}$ instead of the Euclidean norm and we are in the unit polydisc instead of the unit ball. For any $t \in \mathbb{C}^*$ sufficiently small,

$$g_1(tS, z) = \max \left\{ \ln \left| \frac{(z_1 - t/2)(z_1 + t/2)}{(1 - \bar{t}z_1/2)(1 + \bar{t}z_1/2)} \right|, \ln |z_2| \right\}$$

is the pluricomplex Green function in the unit polydisc $P(O, 1)$ with logarithmic poles in tS with weight 1. $(g_1(tS, \cdot))_{t \in \mathbb{C}^*}$ converges uniformly on any compact set of the form $\overline{P}(O, 1) \setminus P(O, \varrho)$ (where $0 < \varrho < 1$) to g_∞ which is explicitly equal to $g_\infty(z) = \max\{2 \ln |z_1|, \ln |z_2|\}$ in $P(O, 1)$. It is also easy to verify that $\tilde{g}_\infty(z) = \max\{\ln |z_1|, \ln |z_2|\} = \ln |z|$ in $P(O, 1)$.

Proof of Theorem 1.1. — In the first three steps and step 7, we prove item (ii). In steps 4 and 5, we prove item (i). In step 6, we prove item (iii) and in step 8, we prove item (iv).

Step 1. — Let Q be a holomorphic polynomial and ϱ be a positive real number. It is well known that

$$\frac{1}{\deg(Q)} \ln \left(\frac{|Q(z)|}{\|Q\|_\varrho} \right) \leq \ln^+ \left(\frac{\|z\|}{\varrho} \right), \text{ in } \mathbb{C}^n.$$

$\|f\|_\varrho = \sup_{z \in B(O, \varrho)} |f(z)|$ is the sup-norm in $B(O, \varrho)$. The function on the right hand side, is called the pluricomplex Green function with logarithmic pole at infinity for the compact set $\bar{B}(O, \varrho)$ (Zahariuta, Siciak). \ln^+ is the function $\max\{\ln, 0\}$. Consequently, we obtain for any positive real number R such that $0 < \varrho \leq R$,

$$(2.1) \quad \ln\|Q\|_\varrho - \ln\|Q\|_R \geq \deg(Q) \cdot \ln \left(\frac{\varrho}{R} \right).$$

On the other hand, according to a Schwarz' Lemma (Moreau [21], Waldschmidt [29, p. 146]): for any $\epsilon > 0$, there exists a real number $r_2(S, \epsilon)$ such that for any $r_2 < \varrho < R$ with $2e^n \cdot \varrho < R$ and for any polynomial Q such that $\text{ord}(Q, p) \geq l$ for all $p \in S$, we have

$$(2.2) \quad \begin{aligned} \ln\|Q\|_\varrho - \ln\|Q\|_R &\leq (\Omega(S, l) - l\epsilon) \cdot \ln \left(\frac{2e^n \varrho}{R} \right) \\ &\leq -l(\Omega(S) - \epsilon) \ln \left(\frac{R}{2e^n \varrho} \right). \end{aligned}$$

We can suppose in addition that $r_2(S, \epsilon) \geq \sup_{p \in S} \|p\| := \|S\|$.

Step 2. — We can generalize Theorem 1.1 of [25] (for a pluricomplex Green function with one logarithmic pole) to the case of a pluricomplex Green function with a finite number of logarithmic poles:

$$g_R(S, \cdot) = \sup_{l \geq 1} H_{S, R, l} = \lim_{l \rightarrow \infty} H_{S, R, l},$$

where each psh function $H_{S, R, l}$ (l is a positive integer) is defined by

$$H_{S, R, l} = \sup \left\{ \frac{1}{l} \ln|f| : f \in \mathcal{O}(B(O, R)), \|f\|_R \leq 1, \right. \\ \left. \text{ord}(f, s) \geq l \text{ for any } s \in S \right\}.$$

Here $\mathcal{O}(D)$ is the Frechet space of holomorphic functions in an open set D . We can replace in the previous definition of $H_{S, R, l}$, holomorphic functions by polynomials satisfying the same properties. Indeed, if l and d are two positive fixed integers, then we have a continuous linear map φ from $\mathbb{C}_d[z]$ to \mathbb{C}^{m_l} ($m_l = |S|^{\binom{l-1+n}{n}}$) as follow

$$\varphi : P \mapsto (P^{(\alpha)}(s), \forall |\alpha| \leq l - 1, \forall s \in S).$$

$\mathbb{C}_d[z]$ is the linear space of polynomials of degree less or equal to d of dimension $\binom{d+n}{n}$. If d is sufficiently large, then this map φ is surjective (we can prove it by using Cartan–Serre's Theorem). $\text{Ker } \varphi$ is the kernel of

φ and it has $\dim(\text{Ker } \varphi) = \binom{d+n}{n} - |S| \binom{l-1+n}{n}$. We choose E a subspace of $\mathbb{C}_d[z]$, such that $\text{Ker } \varphi \oplus E = \mathbb{C}_d[z]$. Then there exists a map ψ from \mathbb{C}^{m_l} to E , which is the inverse of the restriction of φ to E . ψ is a bijective continuous linear map from \mathbb{C}^{m_l} to E with norm $\|\psi\|$ (which depends on l and d). According to the fact that the family $(H_{S,R',l})_{R' > R}$ converges uniformly on $B(O, R)$ to $H_{S,R,l}$ when R' goes to R and because $B(O, R)$ is a Runge domain in \mathbb{C}^n ; for any f with $\|f\|_R = 1$ which appears in the definition of $H_{S,R,l}$, we can find a sequence $(P_j)_j$ of polynomials such that $1 - 1/j \leq \|P_j\|_R \leq 1 + 1/j$ and $\|f - P_j\|_R \leq 1/j$. There exists $\delta > 0$ such that $P(s, \delta) \subset B(O, R)$, for all $s \in S$. $P(s, \delta)$ is the polydisc centered in s with multiradius (δ, \dots, δ) . According to Cauchy's inequalities applied in each polydisc $P(s, \delta)$ and since $\|f - P_j\|_{P(s, \delta)} \leq 1/j$, we obtain that $|(f - P_j)^{(\alpha)}(s)| = |(P_j)^{(\alpha)}(s)| \leq \alpha!/(j r^{|\alpha|})$, for all $|\alpha| \leq l - 1$ and for all $s \in S$. Denote by Q_j the polynomial in $\mathbb{C}_d[z]$, equal to $\psi((P_j)^{(\alpha)}(s), \forall |\alpha| \leq l - 1, \forall s \in S)$. There exists a positive constant $c_{l,d}$ such that $\|Q_j\|_R \leq c_{l,d}/j$. Then if we replace P_j by $R_j := (P_j - Q_j)/\|P_j - Q_j\|_R$ we obtain that $\|f - R_j\|_R \leq O(1/j)$, with $\text{ord}(R_j, p) \geq l$, for all $p \in S$.

Step 3. — Since for any positive integer l , there exists a polynomial Q such that $\text{ord}(Q, p) \geq l$ for any $p \in S$, $\deg(Q) = \Omega(S, l)$ and $\|Q\|_R = 1$, according to inequalities (2.1) and (2.2), we obtain

$$\frac{\Omega(S, l)}{l} \ln \left(\frac{\varrho}{R} \right) \leq \sup_{z \in B(O, \varrho)} H_{S, R, l}(z) \leq \left(\frac{\Omega(S, l)}{l} - \epsilon \right) \ln \left(\frac{2\varrho e^n}{R} \right).$$

Since $H_{S, 1/|t|, l}(z) = H_{tS, 1, l}(tz)$ when $|z| \leq 1/|t|$ and $t \in \mathbb{C}^*$, we have in particular

$$\frac{\Omega(S, l)}{l} \ln(|t|\varrho) \leq \sup_{z \in B(O, |t|\varrho)} H_{tS, 1, l}(z) \leq \left(\frac{\Omega(S, l)}{l} - \epsilon \right) \ln(2\varrho|t|e^n),$$

or which is equivalent: for any $t \in \mathbb{C}^*$ and $\varrho' > 0$ such that $|t|r_2(S, \epsilon) \leq \varrho' \leq 1$

$$\frac{\Omega(S, l)}{l} \ln \varrho' \leq \sup_{z \in B(O, \varrho')} H_{tS, 1, l}(z) \leq \left(\frac{\Omega(S, l)}{l} - \epsilon \right) \ln(2\varrho' e^n).$$

We can suppose in addition that t is sufficiently small such that $tS \subset B(O, \varrho')$. We know that the sequence $(H_{tS, 1, l})_l$ converges uniformly on $\overline{B}(O, 1) \setminus B(O, \varrho')$ to $g_1(tS, \cdot)$ when l tends to infinity ([25]). From the previous estimates, we deduce

$$\Omega(S) \ln \varrho' \leq \sup_{z \in B(O, \varrho')} g_1(tS, z) \leq (\Omega(S) - \epsilon) \ln(2\varrho' e^n),$$

for any $\varrho' > 0$ and $t \in \mathbb{C}^*$ satisfying $|t|r_2(S, \epsilon) \leq \varrho' \leq 1$ and $tS \subset B(O, \varrho')$. If $z \neq O$, $|t|r_2(S, \epsilon) \leq \|z\| \leq 1$ and $tS \subset B(O, \|z\|)$,

$$(2.3) \quad \Omega(S) \ln \|z\| \leq \tilde{g}_1(tS, z) \leq (\Omega(S) - \epsilon) \ln(2\|z\|e^n).$$

In particular if $|t|r_2(S, \epsilon) = \|z\|$ (we have $r_2(S, \epsilon) \geq \|S\|$), we have

$$\begin{aligned} \tilde{g}_1(tS, z) &\leq (\Omega(S) - \epsilon) \ln(2e^n |t|r_2(S, \epsilon)) \\ &= \left((\Omega(S) - \epsilon) \frac{\ln(2e^n)}{\ln(|t|r_2)} + (\Omega(S) - \epsilon) \right) \ln \|z\|. \end{aligned}$$

For t sufficiently small, $(\Omega(S) - \epsilon) \frac{\ln(2e^n)}{\ln(|t|r_2)} + (\Omega(S) - \epsilon) > 0$, the psh function $\left((\Omega(S) - \epsilon) \frac{\ln(2e^n)}{\ln(|t|r_2)} + (\Omega(S) - \epsilon) \right) \ln \|\cdot\|$ is maximal in $B(O, 1) \setminus \{O\}$ and equal to 0 on $\partial B(O, 1)$. Thus we obtain in $B(O, 1) \setminus B(O, |t|r_2)$,

$$\Omega(S) \ln \|z\| \leq \tilde{g}_1(tS, z) \leq \left((\Omega(S) - \epsilon) \frac{\ln(2e^n)}{\ln(|t|r_2)} + (\Omega(S) - \epsilon) \right) \ln \|z\|.$$

Consequently, for any $0 < \varrho < 1$, $(\tilde{g}_1(tS, \cdot))_t$ converges uniformly in $\bar{B}(O, 1) \setminus B(O, \varrho)$ to $\Omega(S) \ln \|\cdot\|$ when t converges to 0.

Step 4. — Let $0 < \varrho < 1$ be fixed. Let ϵ_1 be a positive real number such that $\epsilon_1 r_2(S, \epsilon) \leq \varrho \leq 1$. Since $\sup_{|t| \leq \epsilon_1} \sup_{z \in B(O, \varrho)} g_1(tS, z) = \sup_{z \in B(O, \varrho)} \sup_{|t| \leq \epsilon_1} g_1(tS, z)$, we obtain according to the previous estimates (2.3)

$$(2.4) \quad \Omega(S) \ln \varrho \leq \sup_{z \in B(O, \varrho)} \left(\sup_{|t| \leq \epsilon_1} g_1(tS, z) \right)^* \leq (\Omega(S) - \epsilon) \ln(2\varrho e^n).$$

Since $g_\infty = \lim_{\epsilon_1 \rightarrow 0} (\sup_{|t| \leq \epsilon_1} g_1(tS, \cdot))^*$, where this limit decreases when ϵ_1 decreases, and according to the fact that the family $(g_1(tS, \cdot))_{t \in \mathbb{C}^*}$ converges locally uniformly outside the origin in $B(O, 1)$ to g_∞ ([26, Theorem 1.1]), we can deduce that

$$\Omega(S) \ln \varrho \leq \sup_{z \in B(O, \varrho)} g_\infty(z) \leq (\Omega(S) - \epsilon) \ln(2\varrho e^n).$$

Consequently, $\lim_{\varrho \rightarrow 0} \frac{\sup_{z \in B(O, \varrho)} g_\infty(z)}{\ln \varrho} := \nu(g_\infty, O) = \Omega(S)$. In particular, according to the maximality of the psh function $\ln \|z\|$ in $B(O, 1) \setminus \{O\}$, we obtain in $B(O, 1)$ that

$$g_\infty(z) \leq \Omega(S) \ln \|z\|.$$

And $(\text{dd}^c g_\infty)^n(\{O\}) \geq \Omega(S)^n$.

Step 5. — By the comparison principle for the complex Monge–Ampère operator [2], we have for any $\epsilon > 0$

$$\int_{\{u < v\}} (\mathrm{dd}^c v)^n \leq \int_{\{u < v\}} (\mathrm{dd}^c u)^n,$$

where u and v are two psh functions in $B(O, 1)$ defined by $u = (1 + \epsilon) \times g_1(tS, \cdot)$ and $v = (\sup_{|t| \leq \epsilon_1} g_1(tS, \cdot))^*$. u has isolated logarithmic singularities and v is bounded. Since u and v tend to 0 on the boundary of the unit ball, we deduce that

$$\int_{B(O, 1)} (\mathrm{dd}^c (\sup_{|t| \leq \epsilon_1} g_1(tS, \cdot))^*)^n \leq \int_{B(O, 1)} (\mathrm{dd}^c (1 + \epsilon) g_1(tS, \cdot))^n = (1 + \epsilon)^n |S|.$$

According to the monotonicity of the family $((\sup_{|t| \leq \epsilon_1} g_1(tS, \cdot))^*)_{\epsilon_1}$ when ϵ_1 decreases, and by making tend ϵ to 0, we deduce that

$$\int_{B(O, 1)} (\mathrm{dd}^c g_\infty)^n \leq |S|.$$

Since the family $(g_1(tS, \cdot))_{t \in \mathbb{C}^*}$ converges locally uniformly outside the origin in $B(O, 1)$ to g_∞ , then the complex Monge–Ampère measure $(\mathrm{dd}^c g_1(tS, \cdot))^n$ converges to $(\mathrm{dd}^c g_\infty)^n$ in $B(O, 1) \setminus \{O\}$. Since $(\mathrm{dd}^c g_1(tS, \cdot))^n = \sum_{p \in S} \delta_{tp}$, we have $(\mathrm{dd}^c g_\infty)^n = 0$ in $B(O, 1) \setminus \{O\}$.

Step 6. — If $\Omega(S) = |S|^{1/n}$ then all inequalities in item (i) are equalities. $\nu(g_\infty, O) = \Omega(S)$, $(\mathrm{dd}^c g_\infty)^n = 0$ in $B(O, 1) \setminus \{O\}$ and $(\mathrm{dd}^c g_\infty)^n(\{O\}) = \Omega(S)^n = |S|$. We conclude this case by proving that $g_\infty(z) = |S|^{1/n} \ln \|z\|$ in $B(O, 1)$. Assume that there exists a point $a \in B(O, 1) \setminus \{O\}$ such that $g_\infty(a) < |S|^{1/n} \ln \|a\|$. Let γ be a $\mathcal{C}^\infty(\overline{B(O, 1)})$ and strictly psh function such that $\gamma \leq -1$ in $\overline{B(O, 1)}$ (classical argument, used in the proof of Theorem 4.3 in [7] for instance. We can choose $\gamma(z) = \|z\|^2 - 2$). Let $\epsilon > 0$ be sufficiently small such that the following psh function v defined by

$$v(z) = \max\{\Omega(S) \ln \|z\| + \epsilon \gamma(z), g_\infty(z)\}$$

satisfies

- (a) $v(z) = \Omega(S) \ln \|z\| + O(1)$ when $z \rightarrow O$,
- (b) $v(z) = g_\infty(z)$ in a neighborhood of ∂B ,
- (c) $v(z) = \Omega(S) \ln \|z\| + \epsilon \gamma(z)$ in a neighborhood of a .

According to (a) and (b), we deduce that $\mathbf{1}_{\{O\}}(\mathrm{dd}^c v)^n = \Omega(S)^n \delta_O = |S| \delta_O$ and $\int_{B(O, 1)} (\mathrm{dd}^c v)^n = \int_{B(O, 1)} (\mathrm{dd}^c g_\infty)^n = |S|$ (according to the comparison principle for MA operator). Hence $(\mathrm{dd}^c v)^n = 0$ in $B(O, 1) \setminus \{O\}$. This is in contradiction with the third above property. Consequently $g_\infty \equiv |S|^{1/n} \ln \|\cdot\|$ in $B(O, 1)$.

Step 7. — According to the third section of this proof, $g_\infty(z) \leq \Omega(S) \ln \|z\|$ in $\bar{B}(O, 1)$. Then \tilde{g}_∞ satisfies the same estimate in $\bar{B}(O, 1)$:

$$\tilde{g}_\infty(z) \leq \Omega(S) \ln \|z\|.$$

Let us prove that $\tilde{g}_\infty(z) = \Omega(S) \ln \|z\|$ in $\bar{B}(O, 1)$. Suppose that there exists $O \neq z \in B(O, 1)$, such that $-\infty < \tilde{g}_\infty(z) = c \ln \|z\| < \Omega(S) \ln \|z\|$. Then $\tilde{g}_\infty(w)$ satisfies the same estimates for any w such that $\|w\| = \|z\|$ and we deduce that

$$g_\infty(w) \leq c \ln \|w\| < \Omega(S) \ln \|w\|, \quad \forall w \in \partial B(O, \|z\|).$$

By definition we have $\limsup_{t \rightarrow 0} g_1(tS, w) \leq (\limsup_{t \rightarrow 0} g_1(tS, w))^* = g_\infty(w)$. According to Hartog’s Lemma, we obtain for the positive constant $d = (\Omega(S) - c) \ln \|z\|$, that there exists ϵ_0 such that for any $|t| \leq \epsilon_0$

$$g_1(tS, w) \leq c \ln \|w\| + d/2 < \Omega(S) \ln \|w\|, \quad \forall w \in \partial B(O, \|z\|).$$

In particular, $\sup_{w \in \partial B(O, \|z\|)} g_1(tS, w) \leq c \ln \|z\| + d/2 < \Omega(S) \ln \|z\|$. This contradicts what it has been proved in Step 3 and consequently, $\tilde{g}_\infty \equiv \Omega(S) \ln \|\cdot\|$ in $\bar{B}(O, 1)$.

Step 8. — Let us prove the last item (iv). We suppose that $(g_1(tS, \cdot))_{t \in \mathbb{C}^*}$ converges locally uniformly outside the origin in $B(O, 1)$ to $g_\infty = \Omega(S) \ln \|\cdot\|$ in $\bar{B}(O, 1)$. Then for any $\epsilon > 0$ and for any $0 < \varrho < 1$, there exists $\epsilon_0 > 0$ such that for any $|t| \leq \epsilon_0$, we have $tS \subset B(O, \varrho)$ and

$$(1 + \epsilon)\Omega(S) \ln \|z\| \leq g_1(tS, z) \leq (1 - \epsilon)\Omega(S) \ln \|z\|, \quad \forall z \in \bar{B}(O, 1) \setminus B(O, \varrho).$$

Let us construct a continuous and psh entire function v such that

$$v(z) = \begin{cases} g_1(tS, z), & z \in B(O, \varrho), \\ \max\{g_1(tS, z), (1 + \epsilon)\Omega(S) \ln \|z\|\}, & z \in B(O, 1) \setminus B(O, \varrho), \\ (1 + \epsilon)\Omega(S) \ln \|z\|, & z \in \mathbb{C}^n \setminus B(O, 1). \end{cases}$$

We obtain that $|S| = \int_{B(O, \varrho)} (\text{dd}^c v)^n \leq \int_{\mathbb{C}^n} (\text{dd}^c v)^n = (1 + \epsilon)^n \Omega(S)^n$. We already know that $\Omega(S) \leq |S|^{1/n}$. Consequently, $\Omega(S) = |S|^{1/n}$. This achieves the proof of Theorem 1.1. □

2.4. A Schwarz’ Lemma

With the first item of the previous theorem, we obtain a slightly different version of Schwarz’ Lemma from the previous ones [21].

COROLLARY 2.2. — For any positive real numbers $\varrho < 1$ and ϵ , there exists a positive real number $R_{\epsilon, \varrho}$ such that: if R and r satisfy $R \geq R_{\epsilon, \varrho}$ and $\varrho R \leq r \leq R$ we have:

$$\ln \|f\|_r \leq \ln \|f\|_R - l(\Omega(S) - \epsilon) \ln(R/r),$$

for any integer $l \geq 1$ and for any entire function f with $\text{ord}(f, p) \geq l$ for all $p \in S$.

Proof. — According to Theorem 1.1, $g_\infty(z) \leq \Omega(S) \ln \|z\|$ in $B(O, 1)$. If we apply Hartogs lemma to the compact set $\partial B(O, \varrho)$, we obtain that: there exists a positive real number $R_{\epsilon, \varrho}$ such that for any $R \geq R_{\epsilon, \varrho}$, we have $S/R \subset B(O, \varrho)$ and

$$g_1(S/R, z) \leq (\Omega(S) - \epsilon) \ln \|z\|, \quad \text{on } \partial B(O, \varrho).$$

$(\Omega(S) - \epsilon) \ln \|\cdot\|$ is maximal in $B(O, 1) \setminus \{O\}$ and these two functions are equal to 0 on the boundary of $B(O, 1)$. Consequently, the same inequality is satisfied in $B(O, 1) \setminus B(O, \varrho)$ and

$$g_R(S, z) \leq (\Omega(S) - \epsilon) \ln \|z/R\|, \quad \text{on } B(O, R) \setminus B(O, \varrho R).$$

Finally, if l is a positive integer and f is an entire function with $\text{ord}(f, p) \geq l$ for any $p \in S$ ($\ln(|f|/\|f\|_R) \leq l.g_R(S, \cdot)$ in $B(O, R)$), then for any $\varrho R \leq r \leq R$ we obtain

$$\ln \|f\|_r \leq \ln \|f\|_R - l(\Omega(S) - \epsilon) \ln(R/r). \quad \square$$

2.5. Pluricomplex Green functions with a finite set of logarithmic poles with different homogeneous weights

Let D be a bounded hyperconvex domain in \mathbb{C}^n . Let $S = \{p_1, \dots, p_r\}$ be a finite set of r distinct points in D and $\mathcal{M} = \{m_1, \dots, m_r\}$ be a set of r positive integers. Denote by $g_D((S, \mathcal{M}), \cdot)$, the pluricomplex Green function in the domain D , with logarithmic poles at any points p_j of S and with homogeneous weight m_j respectively. According to [12], to any point $p_j \in S$ we can associate a generic set S_j of m_j^n distinct points in \mathbb{C}^n such that $\Omega(S_j) = m_j$. Denote by $\Sigma(t_1, \dots, t_r)$ the set $\{p_j + t_j S_j : 1 \leq j \leq r\}$ of $\sum_{j=1}^r m_j^n$ distinct points. For sufficiently small $(t_1, \dots, t_r) \in (\mathbb{C}^*)^r$ the set $\Sigma(t_1, \dots, t_r)$ is contained in D . Denote by $g_D(\Sigma(t_1, \dots, t_r), \cdot)$ the pluricomplex Green function in D , with logarithmic poles at any points of $\Sigma(t_1, \dots, t_r)$ and with weight 1. These functions satisfy in D

$$\sum_{j=1}^r g_D(p_j + t_j S_j, z) \leq g_D(\Sigma(t_1, \dots, t_r), z) \leq \inf_{1 \leq j \leq r} g_D(p_j + t_j S_j, z).$$

According to Theorem 1.1, for any $1 \leq j \leq r$, the family $(g_D(p_j + t_j S_j, \cdot))_{t_j \in \mathbb{C}^*}$ converges locally uniformly outside the point p_j in D to $m_j g_D(p_j, \cdot)$. By using the same technics as in the proof of Theorem 1.1, we deduce the following theorem.

THEOREM 2.3. — *The family of pluricomplex Green functions $(g_D(\Sigma(t_1, \dots, t_r), \cdot))_{(t_1, \dots, t_r) \in (\mathbb{C}^*)^r}$ converges locally uniformly outside the set S in D to $g_D((S, \mathcal{M}), \cdot)$, when (t_1, \dots, t_r) tends to O .*

3. New affine invariants and others conjectures of pluripotential theory

3.1. Affine invariants associated to finite sets of points in \mathbb{C}^n

Fix a finite set $S = \{p_1, \dots, p_r\} \subset \mathbb{C}^n$ of r distinct points, $|S| = r$ is its cardinality. In Section 1.2.2 we have defined two affine invariants $\omega_{\text{psh}}(S)$ and $\omega_{\text{psh}}^+(S)$.

$$\omega_{\text{psh}}(S) := \sup\{\omega(S, u) : u \in \text{PSH}(\mathbb{C}^n)\}$$

and

$$\omega_{\text{psh}}^+(S) := \sup\{\omega(S, u) : u \in \text{PSH}(\mathbb{C}^n) \cap L_{\text{loc}}^\infty(\mathbb{C}^n \setminus S)\},$$

where for any psh function u in \mathbb{C}^n ,

$$\gamma_u := \limsup_{|z| \rightarrow \infty} \frac{u(z)}{\log |z|} \quad \text{and} \quad \omega(S, u) := \frac{\sum_{j=1}^k \nu(u, p_j)}{\gamma_u}.$$

$\nu(u, z)$ is the Lelong number of the psh function u at the point z . Let us remark that for any psh function u in \mathbb{C}^n , we have always $\gamma_u \in [0, +\infty]$. Indeed, if u is psh in \mathbb{C}^n and $u(z) \leq o(\log |z|)$ as $|z| \rightarrow \infty$, then u is constant (it is a generalization of Liouville's Theorem, see for example [15]). Conversely, if $u(z) = |z|^2$ then $\gamma_u = +\infty$. $\omega_{\text{psh}}(S)$ and $\omega_{\text{psh}}^+(S)$ are related to the algebraic geometric properties of S and in particular to the affine invariant $\omega(S)$ of S , that they generalize. Here are some simple properties of these invariants.

LEMMA 3.1.

- (i) $\omega(S) \leq \omega_{\text{psh}}(S)$ and $\omega_{\text{psh}}^+(S) \leq \omega_{\text{psh}}(S)$.
 - (ii) The sets $\{\omega(S, u) : u \in \text{PSH}(\mathbb{C}^n)\}$ and $\{\omega(S, u) : u \in \text{PSH}(\mathbb{C}^n) \cap L_{\text{loc}}^\infty(\mathbb{C}^n \setminus S)\}$ are connected.
- $$(0, \omega_{\text{psh}}(S)) \subseteq \{\omega(S, u) : u \in \text{PSH}(\mathbb{C}^n)\} \subseteq (0, \omega_{\text{psh}}(S)),$$
- $$(0, \omega_{\text{psh}}^+(S)) \subseteq \{\omega(S, u) : u \in \text{PSH}(\mathbb{C}^n) \cap L_{\text{loc}}^\infty(\mathbb{C}^n \setminus S)\} \subseteq (0, \omega_{\text{psh}}^+(S)).$$

Proof. — For $a > 1$ let $f_a : \mathbb{R} \rightarrow \mathbb{R}$ be the convex increasing function defined by $f_a(t) = t$ if $t < 0$, $f_a(t) = at$ if $t \geq 0$. If $u \in \text{PSH}(\mathbb{C}^n)$ then $f_a \circ u \in \text{PSH}(\mathbb{C}^n)$, $\gamma_{f_a \circ u} = a\gamma_u$, $\nu(f_a \circ u, z) = \nu(u, z)$ for any point z and finally $\omega(S, f_a \circ u) = \omega(S, u)/a$. \square

For technical reasons, we need to introduce some other constants. Let l be a positive real, we define

$$\Omega_{\text{psh}}(S, l) := \inf\{\gamma_u : u \in \text{PSH}(\mathbb{C}^n), \nu(u, p) \geq l \text{ for any } p \in S\}.$$

It is easy to see that $\Omega_{\text{psh}}(S, l)/l$ doesn't depend on l . Then we just introduce the common value $\Omega_{\text{psh}}(S, 1)$, that we denote by $\Omega_{\text{psh}}(S)$. In the same way we have

$$\Omega_{\text{psh}}^+(S) := \inf\{\gamma_u : u \in \text{PSH}(\mathbb{C}^n) \cap L_{\text{loc}}^\infty(\mathbb{C}^n \setminus S), \nu(u, p) \geq 1 \text{ for any } p \in S\}.$$

This invariant was already introduced in [5]. All these invariants can be compared.

LEMMA 3.2.

- (i) *The sets $\{\gamma_u : u \in \text{PSH}(\mathbb{C}^n), \nu(u, p) \geq 1 \text{ for any } p \in S\}$ and $\{\gamma_u : u \in \text{PSH}(\mathbb{C}^n) \cap L_{\text{loc}}^\infty(\mathbb{C}^n \setminus S), \nu(u, p) \geq 1 \text{ for any } p \in S\}$ are connected and since the function $u(z) = \sum_{p \in S} \ln|z - p| + |z|$ satisfies $\gamma_u = +\infty$, we have*

$$\begin{aligned} (\Omega_{\text{psh}}(S), +\infty) &\subset \{\gamma_u : u \in \text{PSH}(\mathbb{C}^n), \nu(u, p) \geq 1, \forall p \in S\} \\ &\subset [\Omega_{\text{psh}}(S), +\infty] \end{aligned}$$

and

$$\begin{aligned} (\Omega_{\text{psh}}^+(S), +\infty) &\subset \{\gamma_u : u \in \text{PSH}(\mathbb{C}^n) \cap L_{\text{loc}}^\infty(\mathbb{C}^n \setminus S), \nu(u, p) \geq 1, \forall p \in S\} \\ &\subset [\Omega_{\text{psh}}^+(S), +\infty]. \end{aligned}$$

- (ii) *We have $\Omega_{\text{psh}}(S) \leq \Omega(S)$ and $\Omega_{\text{psh}}(S) \leq \Omega_{\text{psh}}^+(S)$.*

- (iii) *We have $\omega(S) \geq \frac{|S|}{\Omega(S)}$, $\omega_{\text{psh}}(S) \geq \frac{|S|}{\Omega_{\text{psh}}(S)}$ and $\omega_{\text{psh}}^+(S) \geq \frac{|S|}{\Omega_{\text{psh}}^+(S)}$.*

- (iv) *If $S' \subseteq S \subset \mathbb{C}^n$ then*

$$\omega_{\text{psh}}^+(S') \leq \omega_{\text{psh}}^+(S) \leq \omega_{\text{psh}}^+(S') + \omega_{\text{psh}}^+(S \setminus S').$$

The proof of this lemma is similar to the proof of Proposition 3.1 in [5]. In what follows, we will prove that $\omega_{\text{psh}}^+(S) \leq \Omega_{\text{psh}}(S) \leq \Omega(S)$, in the 2-dimensional case. So we are interested in finding lower bounds for $\omega_{\text{psh}}^+(S)$, and then for $\Omega(S)$. It is relevant for Conjecture (\mathcal{A}_1) . On the other hand we are interested in finding upper bounds for $\omega_{\text{psh}}(S)$, and then for $\omega(S)$. It is relevant for Conjecture (\mathcal{A}_2) , which is equivalent to Conjecture (\mathcal{A}_1) .

3.2. Comparisons of these invariants

3.2.1. A very simple situation in one complex variable

The situation is very simple for subharmonic functions in \mathbb{C} . Let v be an entire subharmonic (sh) function such that $\nu(v, p_j) = \alpha_j$ (positive real number) for $1 \leq j \leq r$. Let u be the sh function defined by $u(z) = \sum_{j=1}^r \alpha_j \ln|z - p_j|$. Then $w := v - u$ is a sh function in \mathbb{C} .

$$0 \leq \gamma_w = \limsup_{|z| \rightarrow +\infty} \frac{v(z) - u(z)}{\ln|z|} = \limsup_{|z| \rightarrow +\infty} \frac{v(z)}{\ln|z|} - \lim_{|z| \rightarrow +\infty} \frac{u(z)}{\ln|z|} = \gamma_v - \gamma_u,$$

then $\gamma_v \geq \gamma_u = \sum_j \alpha_j$. $\omega(S, u) = 1$ and $\omega(S, v) \leq 1$. Consequently, $\omega_{sh}(S) = \omega_{sh}^+(S) = 1$. For polynomials, it is also very simple. Indeed, if P is a polynomial such that $\text{ord}(P, p_j) = \alpha_j$ (positive integer) for $1 \leq j \leq r$, then P can be divided by $\prod_{j=1}^r (z - p_j)^{\alpha_j}$. There exists a polynomial Q such that $P(z) = \prod_{j=1}^r (z - p_j)^{\alpha_j} Q(z)$ and $\deg(P) \geq \deg(Q)$. Consequently, $\omega(S) = 1$. It is easy to see that we also have in this situation, $\Omega(S) = \Omega_{sh}(S) = \Omega_{sh}^+(S) = |S|$.

3.2.2. In the multivariable case

Here is a comparison principle which relates Lelong numbers at the points of S of two psh functions with their logarithmic growth at infinity. The proof of this result is simple and is similar to the proof of Theorem 3.4 in [5] (or Proposition 2.1 in [4]).

THEOREM 3.3. — *Let $S \subset \mathbb{C}^n$ be a finite set. Let v be a psh function in \mathbb{C}^n . Let $u \in \text{PSH}(\mathbb{C}^n) \cap L_{\text{loc}}^\infty(\mathbb{C}^n \setminus S)$. Then*

$$\sum_{p \in S} \nu(u, p)^{n-1} \nu(v, p) \leq \gamma_u^{n-1} \gamma_v.$$

Proof. — Let $u \in \text{PSH}(\mathbb{C}^n) \cap L_{\text{loc}}^\infty(\mathbb{C}^n \setminus S)$ be such that $\gamma_u > 0$. Let us suppose that u has singularities at any point of S . It follows from a comparison theorem for Lelong numbers with weights due to Demailly [9] that for any $p \in S$

$$\lim_{r \rightarrow 0} \int_{B(p,r)} (\text{dd}^c u)^{n-1} \wedge \text{dd}^c v = \nu_u(\text{dd}^c v, p) \geq \nu(u, p)^{n-1} \nu(v, p)$$

and

$$\int_{\mathbb{C}^n} (\text{dd}^c u)^{n-1} \wedge \text{dd}^c v \geq \sum_{p \in S} \nu(u, p)^{n-1} \nu(v, p).$$

In addition,

$$\int_{\mathbb{C}^n} (\mathrm{dd}^c u)^{n-1} \wedge \mathrm{dd}^c v \leq \gamma_u^{n-1} \int_{\mathbb{C}^n} (\mathrm{dd}^c \log |z|)^{n-1} \wedge \mathrm{dd}^c v \leq \gamma_u^{n-1} \gamma_v.$$

and the proof is completed. \square

COROLLARY 3.4. — *For any finite set $S \subset \mathbb{C}^n$, we have*

$$(3.1) \quad \omega_{\mathrm{psh}}^+(S)^{n-1} \leq \Omega_{\mathrm{psh}}(S)|S|^{n-2} \leq \Omega(S)|S|^{n-2} \leq |S|^{(n-1)^2/n},$$

$$(3.2) \quad \frac{|S|}{\Omega(S)} \leq \omega(S) \leq \omega_{\mathrm{psh}}(S) \leq \Omega_{\mathrm{psh}}^+(S)^{n-1} \leq |S|^{n-1}.$$

In particular, for $n = 2$, $\omega_{\mathrm{psh}}^+(S) \leq \Omega_{\mathrm{psh}}(S)$ and $\omega_{\mathrm{psh}}(S) \leq \Omega_{\mathrm{psh}}^+(S)$.

The inequalities $\omega(S) \leq \Omega_{\mathrm{psh}}^+(S)^{n-1} \leq |S|^{n-1}$ are already proved in [5].

Proof.

Step 1. — Let v be a psh function in \mathbb{C}^n such that $\nu(v, p) \geq 1$ for any $p \in S$. Let u be a function in $\mathrm{PSH}(\mathbb{C}^n) \cap L_{\mathrm{loc}}^\infty(\mathbb{C}^n \setminus S)$. According to Theorem 3.3, we have $\frac{1}{\gamma_u^{n-1}} \sum_{p \in S} \nu(u, p)^{n-1} \leq \gamma_v$. If we take the infimum in the right hand side of this inequality for any such v , then we obtain $\frac{1}{\gamma_u^{n-1}} \sum_{p \in S} \nu(u, p)^{n-1} \leq \Omega_{\mathrm{psh}}(S)$. By using Hölder inequality, we deduce $\frac{1}{\gamma_u} \sum_{p \in S} \nu(u, p) \leq \Omega_{\mathrm{psh}}(S)^{1/(n-1)} |S|^{(n-2)/(n-1)}$. Finally by taking the supremum in the left hand side of this inequality for any $u \in \mathrm{PSH}(\mathbb{C}^n) \cap L_{\mathrm{loc}}^\infty(\mathbb{C}^n \setminus S)$, we obtain

$$\omega_{\mathrm{psh}}^+(S)^{n-1} \leq \Omega_{\mathrm{psh}}(S) |S|^{n-2}.$$

Step 2. — Let v be a psh function in \mathbb{C}^n and u be a function in $\mathrm{PSH}(\mathbb{C}^n) \cap L_{\mathrm{loc}}^\infty(\mathbb{C}^n \setminus S)$ such that $\nu(u, p) \geq 1$ for any $p \in S$. According to Theorem 3.3, we have $\sum_{p \in S} \nu(v, p) / \gamma_v \leq \gamma_u^{n-1}$. We take the supremum in the left side of this inequality for any such v and the infimum in the right side of this inequality for any such previous u and we obtain

$$\omega_{\mathrm{psh}}(S) \leq \Omega_{\mathrm{psh}}^+(S)^{n-1}.$$

Step 3. — By definition of $\Omega_{\mathrm{psh}}^+(S)$, with the psh function $u(z) = \sum_{p \in S} \log |z - p|$ we obtain

$$\Omega_{\mathrm{psh}}^+(S) \leq |S|.$$

The others inequalities are obvious. This achieves the proof. \square

3.3. Proof of Theorem 1.3

We already know that Conjectures (\mathcal{P}_1) and (\mathcal{A}_1) are equivalent, according to Theorem 1.1. Indeed, with item (iii), we have: (\mathcal{A}_1) implies (\mathcal{P}_1) and with item (i), we have: (\mathcal{P}_1) implies (\mathcal{A}_1) . In addition it is well know that Conjectures (\mathcal{A}_1) and (\mathcal{A}_2) are equivalent ([22]).

Step 1. — From Conjecture (\mathcal{P}_3) we deduce Conjectures (\mathcal{A}_1) , (\mathcal{P}_2) and (\mathcal{A}_2) . Let us suppose that Conjecture (\mathcal{P}_3) is satisfied: for any $\epsilon > 0$, there exists an entire psh function v in $L^\infty_{\text{loc}}(\mathbb{C}^n \setminus S)$, such that $\nu(v, p) \geq 1$ for any $p \in S$ and such that $\gamma_v \leq (1 + \epsilon)|S|^{1/n}$. Then $\omega(S, v) \geq \frac{|S|}{(1+\epsilon)|S|^{1/n}} = \frac{|S|^{1-1/n}}{1+\epsilon}$. Consequently,

$$(3.3) \quad \omega_{\text{psh}}(S) \geq \omega_{\text{psh}}^+(S) \geq |S|^{1-1/n}$$

and

$$(3.4) \quad \Omega_{\text{psh}}(S) \leq \Omega_{\text{psh}}^+(S) \leq |S|^{1/n}.$$

On the other hand, we have the chain (3.1) of inequalities:

$$\omega_{\text{psh}}^+(S)^{n-1} \leq \Omega_{\text{psh}}(S)|S|^{n-2} \leq \Omega(S)|S|^{n-2} \leq |S|^{(n-1)^2/n}.$$

Then we deduce that

$$\begin{aligned} \omega_{\text{psh}}^+(S)^{n-1} &= \Omega_{\text{psh}}(S)|S|^{n-2} = \Omega(S)|S|^{n-2} = |S|^{(n-1)^2/n}, \\ \omega_{\text{psh}}^+(S) &= |S|^{(n-1)/n} \text{ and } \Omega_{\text{psh}}(S) = \Omega(S) = |S|^{1/n}. \end{aligned}$$

Conjecture (\mathcal{A}_1) is solved. According to (3.4), we deduce that $\Omega_{\text{psh}}^+(S) = |S|^{1/n}$. According to (3.2), we obtain $\omega_{\text{psh}}(S) \leq \Omega_{\text{psh}}^+(S)^{n-1} = |S|^{1-1/n}$. And according to inequalities (3.3), we deduce that

$$\omega_{\text{psh}}(S) = |S|^{1-1/n}.$$

Conjecture (\mathcal{P}_2) is solved. Since $\omega(S) \leq \omega_{\text{psh}}(S) = |S|^{1-1/n}$ and $\omega(S) \geq \frac{|S|}{\Omega(S)}$, conjecture (\mathcal{A}_2) is solved.

Step 2. — From conjecture (\mathcal{P}_2) we deduce conjectures (\mathcal{A}_1) and (\mathcal{A}_2) . Let us suppose that conjecture (\mathcal{P}_2) is satisfied: $\omega_{\text{psh}}(S) = \omega_{\text{psh}}^+(S)$. According to the two chains of inequalities (3.1) and (3.2), we obtain some identities.

$$\begin{aligned} \frac{|S|^{n-1}}{\Omega(S)^{n-1}} &\leq \omega(S)^{n-1} \leq \omega_{\text{psh}}(S)^{n-1} = \omega_{\text{psh}}^+(S)^{n-1} \\ &\leq \Omega_{\text{psh}}(S)|S|^{n-2} \leq \Omega(S)|S|^{n-2} \leq |S|^{(n-1)^2/n}. \end{aligned}$$

We deduce in particular that $\frac{|S|^{n-1}}{\Omega(S)^{n-1}} \leq \Omega(S)|S|^{n-2}$, which is equivalent to $\Omega(S) \geq |S|^{1/n}$. And since we always have $\Omega(S) \leq |S|^{1/n}$, Conjecture (\mathcal{A}_1) is finally solved and all previous inequalities are equalities:

$$|S|^{(n-1)^2/n} = \omega(S)^{n-1} = \omega_{\text{psh}}^{n-1}(S) = \omega_{\text{psh}}^+(S)^{n-1} = \Omega_{\text{psh}}(S)|S|^{n-2}.$$

We deduce that $\omega(S) = \omega_{\text{psh}}(S) = \omega_{\text{psh}}^+(S) = |S|^{(n-1)/n}$ and $\Omega_{\text{psh}}(S) = |S|^{1/n}$. In particular conjecture (\mathcal{A}_2) is solved.

Step 3. — Now let us prove that conjecture (\mathcal{A}_1) implies conjecture (\mathcal{P}_3) . Since conjectures (\mathcal{A}_1) and (\mathcal{P}_1) are equivalent, $(g_1(tS, \cdot))_{t \in \mathbb{C}^*}$ converges locally uniformly in $\overline{B(O, 1)} \setminus \{O\}$ to $|S|^{1/n} \ln \|\cdot\|$: for any $\epsilon > 0$ and any $0 < \varrho < 1$, there exists $\epsilon_0 > 0$ such that for any $|t| \leq \epsilon_0$, we have

$$(1 + \epsilon)|S|^{1/n} \ln \|z\| \leq g_1(tS, z) \leq (1 - \epsilon)|S|^{1/n} \ln \|z\| \quad \text{in } \overline{B(O, 1)} \setminus B(O, \varrho).$$

Consequently, the following psh and continuous function v is well defined in \mathbb{C}^n

$$v(z) = \begin{cases} g_1(tS, z), & z \in B(O, \varrho) \\ \max\{g_1(tS, z), (1 + \epsilon)|S|^{1/n} \ln \|z\|\}, & z \in \overline{B(O, 1)} \setminus B(O, \varrho) \\ (1 + \epsilon)|S|^{1/n} \ln \|z\|, & z \in \mathbb{C}^n \setminus B(O, 1) \end{cases}$$

Let us denote by w the entire psh function defined by: $w(z) = v(tz)$. w is locally bounded in $\mathbb{C}^n \setminus S$. $\nu(w, p) = 1$ for any $p \in S$ and $\gamma_w = (1 + \epsilon)|S|^{1/n}$, as required in conjecture (\mathcal{P}_3) . The proof is complete. \square

BIBLIOGRAPHY

- [1] E. BEDFORD & B. A. TAYLOR, “The Dirichlet problem for a complex Monge–Ampère equation”, *Invent. Math.* **37** (1976), no. 1, p. 1-44.
- [2] ———, “A new capacity for plurisubharmonic functions”, *Acta Math.* **149** (1982), no. 1-2, p. 1-40.
- [3] G. V. CHUDNOVSKY, “Singular points on complex hypersurfaces and multidimensional Schwarz lemma”, in *Seminar on Number Theory, Paris 1979/80*, Progress in Mathematics, vol. 12, Birkhäuser, 1981, p. 29-69.
- [4] D. COMAN, “Entire pluricomplex Green functions and Lelong numbers of projective currents”, *Proc. Am. Math. Soc.* **134** (2006), no. 7, p. 1927-1935.
- [5] D. COMAN & S. NIVOICHE, “Plurisubharmonic functions with singularities and affine invariants for finite sets in \mathbb{C}^n ”, *Math. Ann.* **322** (2002), no. 2, p. 317-332.
- [6] J.-P. DEMAILLY, “Mesures de Monge–Ampère et caractérisation géométrique des variétés algébriques affines”, *Mém. Soc. Math. Fr., Nouv. Sér.* **19** (1985), p. 1-125.
- [7] ———, “Mesures de Monge–Ampère et mesures pluriharmoniques”, *Math. Z.* **194** (1987), no. 4, p. 519-564.
- [8] ———, “Singular Hermitian metrics on positive line bundles”, in *Complex algebraic varieties (Bayreuth, 1990)*, Lecture Notes in Mathematics, vol. 1507, Springer, 1992, p. 87-104.

- [9] ———, “Monge–Ampère operators, Lelong numbers and intersection theory”, in *Complex analysis and geometry*, The University Series in Mathematics, Plenum Press, 1993, p. 115-193.
- [10] ———, “Multiplier ideal sheaves and analytic methods in algebraic geometry”, in *School on Vanishing Theorems and Effective Results in Algebraic Geometry (Trieste, 2000)*, ICTP Lecture Notes, vol. 6, Abdus Salam International Centre for Theoretical Physics, 2001, p. 1-148.
- [11] N. Q. DIEU & P. J. THOMAS, “Convergence of multipole Green functions”, *Indiana Univ. Math. J.* **65** (2016), no. 1, p. 223-241.
- [12] L. EVAIN, “On the postulation of s^d fat points in \mathbb{P}^d ”, *J. Algebra* **285** (2005), no. 2, p. 516-530.
- [13] B. HARBOURNE, “On Nagata’s Conjecture”, *J. Algebra* **236** (2001), no. 2, p. 692-702.
- [14] B. HARBOURNE & J. ROÉ, “Discrete behavior of Seshadri constants on surfaces”, *J. Pure Appl. Algebra* **212** (2008), no. 3, p. 616-627.
- [15] L. HÖRMANDER, *Notions of Convexity*, Progress in Mathematics, vol. 127, Birkhäuser, 1994.
- [16] A. LARROBINO, “Inverse system of a symbolic power. III. Thin algebras and fat points”, *Compos. Math.* **108** (1997), no. 3, p. 319-356.
- [17] M. KLIMEK, “Extremal plurisubharmonic functions and invariant pseudodistances”, *Bull. Soc. Math. Fr.* **113** (1985), p. 231-240.
- [18] P. LELONG, “Notions capacitaires et fonctions de Green pluricomplexes dans les espaces de Banach”, *C. R. Math. Acad. Sci. Paris* **305** (1987), no. 3, p. 71-76.
- [19] ———, “Fonction de Green pluricomplexe et lemmes de Schwarz dans les espaces de Banach”, *J. Math. Pures Appl.* **68** (1989), p. 319-347.
- [20] J. I. MAGNÚSSON, A. RASHKOVSKII, R. SIGURDSSON & P. J. THOMAS, “Limits of multipole pluricomplex Green functions”, *Int. J. Math.* **23** (2012), no. 6, article no. 1250065 (38 pages).
- [21] J.-C. MOREAU, “Lemmes de Schwarz en plusieurs variables et applications arithmétiques”, in *Séminaire Pierre Lelong (Analyse) année 1975/76*, Lecture Notes in Mathematics, vol. 822, Springer, 1980, p. 174-190.
- [22] M. NAGATA, “On the 14-th problem of Hilbert”, *Am. J. Math.* **81** (1959), p. 766-772.
- [23] ———, “On rational surfaces, II”, *Mem. Coll. Sci., Univ. Kyoto, Ser. A* **33** (1960), p. 271-293.
- [24] ———, “On the fourteenth problem of Hilbert”, in *Proceedings of the International Congress of Mathematicians (Edinburgh, 1958)*, Cambridge University Press, 1960, p. 459-462.
- [25] S. NIVOCHÉ, “The pluricomplex Green function, capacitative notions, and approximation problems in \mathbb{C}^n ”, *Indiana Univ. Math. J.* **44** (1995), no. 2, p. 489-510.
- [26] A. RASHKOVSKII & P. J. THOMAS, “Powers of ideals and convergence of Green functions with colliding poles”, *Int. Math. Res. Not.* **2014** (2014), no. 5, p. 1253-1272.
- [27] H. TUTAJ-GASIŃSKA, “A bound for Seshadri constants on \mathbb{P}^2 ”, *Math. Nachr.* **257** (2003), p. 108-116.
- [28] M. WALDSCHMIDT, “Propriétés arithmétiques de fonctions de plusieurs variables”, in *Séminaire Pierre Lelong (Analyse) année 1975/76*, Lecture Notes in Mathematics, vol. 578, Springer, 1977, p. 108-135.
- [29] ———, *Nombres transcendants et groupes algébriques*, second ed., Astérisque, Société Mathématique de France, 1987, 218 pages.
- [30] G. XU, “Curves in \mathbb{P}^2 and symplectic packings”, *Math. Ann.* **299** (1994), no. 4, p. 609-613.

Manuscrit reçu le 13 avril 2018,
révisé le 8 avril 2019,
accepté le 13 mai 2020.

Stéphanie NIVOCHÉ
Université Côte d'Azur
Laboratoire J.A. Dieudonné, CNRS UMR 7351
Université de Nice Sophia Antipolis
Parc Valrose
06108 Nice cedex 2 (France)
Stephanie.nivoche@univ-cotedazur.fr