



INSTITUT DE FRANCE
Académie des sciences

Comptes Rendus

Mathématique

Mohamed Saïdi

On sections of arithmetic fundamental groups of open p -adic annuli

Volume 360 (2022), p. 291-296

Published online: 31 March 2022

<https://doi.org/10.5802/crmath.303>



This article is licensed under the
CREATIVE COMMONS ATTRIBUTION 4.0 INTERNATIONAL LICENSE.
<http://creativecommons.org/licenses/by/4.0/>



Les Comptes Rendus. Mathématique sont membres du
Centre Mersenne pour l'édition scientifique ouverte
www.centre-mersenne.org
e-ISSN : 1778-3569



Number theory / *Théorie des nombres*

On sections of arithmetic fundamental groups of open p -adic annuli

Mohamed Saïdi^a

^a College of Engineering, Mathematics, and Physical Sciences, University of Exeter, Harrison Building, North Park Road, EXETER EX4 4QF, United Kingdom
E-mail: M.Saidi@exeter.ac.uk

Abstract. We show the non-existence of sections of arithmetic fundamental groups of open p -adic annuli of small radii. This implies the non-existence of sections of arithmetic fundamental groups of formal boundaries of formal germs of p -adic curves.

Manuscript received 20th March 2021, accepted 25th November 2021.

1. Introduction/Statement of the Main Result

Let $p \geq 2$ be a prime integer, k a p -adic local field (i.e., k/\mathbb{Q}_p is a finite extension), with ring of integers \mathcal{O}_k , uniformiser π , and residue field F . Thus, F is a finite field of characteristic p .

Let $X \rightarrow \text{Spec } \mathcal{O}_k$ be a flat, proper, relative \mathcal{O}_k -curve, with X normal, and $X_k \stackrel{\text{def}}{=} X \times_{\text{Spec } \mathcal{O}_k} \text{Spec } k$ geometrically connected. Assume $X(F) \neq \emptyset$. Let $x \in X^{\text{cl}}(F)$ be a closed point, $\mathcal{O}_{X,x}$ the local ring at x , $\widehat{\mathcal{O}}_{X,x}$ its completion, and $E \stackrel{\text{def}}{=} \widehat{\mathcal{O}}_{X,x} \otimes_{\mathcal{O}_k} k = \widehat{\mathcal{O}}_{X,x}[\frac{1}{\pi}]$. Write $\mathcal{X} \stackrel{\text{def}}{=} \text{Spec } E$, which we assume to be geometrically connected. We shall refer to \mathcal{X} as the *formal germ of X at x* .

Let η be a geometric point of \mathcal{X} with values in its generic point. Thus, η determines an algebraic closure \bar{k} of k , and a geometric point $\bar{\eta}$ of $\mathcal{X}_{\bar{k}} \stackrel{\text{def}}{=} \mathcal{X} \times_{\text{Spec } k} \text{Spec } \bar{k}$. There exists a canonical exact sequence of profinite groups (cf. [3, Exposé IX, Théorème 6.1])

$$1 \longrightarrow \pi_1(\mathcal{X}_{\bar{k}}, \bar{\eta}) \longrightarrow \pi_1(\mathcal{X}, \eta) \longrightarrow G_k \longrightarrow 1. \quad (1)$$

Here, $\pi_1(\mathcal{X}, \eta)$ denotes the arithmetic étale fundamental group of \mathcal{X} with base point η , $\pi_1(\mathcal{X}_{\bar{k}}, \bar{\eta})$ the étale fundamental group of $\mathcal{X}_{\bar{k}}$ with base point $\bar{\eta}$, and $G_k \stackrel{\text{def}}{=} \text{Gal}(\bar{k}/k)$ the absolute Galois group of k .

The sequence (1) splits if $\mathcal{X}(k) \neq \emptyset$. This is for example the case if the morphism $X \rightarrow \text{Spec } \mathcal{O}_k$ is smooth at x . If $\mathcal{X}(k) = \emptyset$; for instance if X is stable and regular, and x is an ordinary F -rational double point of $X_F \stackrel{\text{def}}{=} X \times_{\text{Spec } \mathcal{O}_k} F$, the existence of sections $s: G_k \rightarrow \pi_1(\mathcal{X}, \eta)$ of the projection $\pi_1(\mathcal{X}, \eta) \twoheadrightarrow G_k$ would provide examples of sections of the projection $\pi_1(X_k, \eta) \twoheadrightarrow G_k$ which are *non-geometric* (η induces a geometric point of X_k , denoted also η , via the morphism $\mathcal{X}_k \rightarrow X_k$), i.e., which do not arise from rational points. These in turn will provide *counter-examples to*

the p -adic version of the Grothendieck anabelian section conjecture. This prompts the following question.

Question 1. *With the above notations, assume $\mathcal{X}(k) = \emptyset$. Does the exact sequence (1) split?*

In this note we investigate the case where \mathcal{X} is a p -adic open annulus. Let

$$A \stackrel{\text{def}}{=} \mathcal{O}_k[[S]], \quad B \stackrel{\text{def}}{=} A \otimes_{\mathcal{O}_k} k = A \left[\frac{1}{\pi} \right],$$

$D \stackrel{\text{def}}{=} \text{Spf} A$ is the formal standard open disc, and $\mathcal{D} \stackrel{\text{def}}{=} D_k = \text{Spec} B$ its “generic fibre” which is the standard open disc centred at the point “ $S = 0$ ”. Let $\mathbb{A}_k^1 = \text{Spec} k[S]$, $Y_k = \mathbb{P}_k^1$ its smooth compactification with function field $k(S)$, and $Y = \mathbb{P}_{\mathcal{O}_k}^1$ the smooth compactification of $\mathbb{A}_{\mathcal{O}_k}^1 = \text{Spec} \mathcal{O}_k[S]$. We shall identify A with the completion of the local ring of Y at the closed point “ $S = 0$ ”. We have a natural morphism $\mathcal{D} \rightarrow \mathbb{P}_k^1$, which induces an identification between the set of closed points of \mathcal{D} and the set

$$\{x \in \mathbb{P}_k^1 : |S(x)| < 1\}.$$

For an integer $n \geq 1$, let

$$A_n \stackrel{\text{def}}{=} \frac{\mathcal{O}_k[[S, T]]}{(S^n T - \pi)}, \quad B_n \stackrel{\text{def}}{=} A_n \otimes_{\mathcal{O}_k} k, \quad \text{and} \quad \mathcal{C}_n \stackrel{\text{def}}{=} \text{Spec} B_n.$$

The natural embedding $\mathcal{C}_n \hookrightarrow \mathcal{D}$ induces an identification between the set of closed points of \mathcal{C}_n and the open annulus

$$\{x \in \mathcal{D} : |\pi|^{\frac{1}{n}} < |S(x)| < 1\}.$$

Further, let $P \stackrel{\text{def}}{=} A_{(\pi)}$ be the localisation of A at the ideal (π) , and \widehat{P} the completion of P , which is a complete discrete valuation ring isomorphic to

$$\mathcal{O}_k[[S]]\{S^{-1}\} \stackrel{\text{def}}{=} \left\{ \sum_{i=-\infty}^{\infty} a_i S^i : a_i \in \mathcal{O}_k, \lim_{i \rightarrow -\infty} |a_i| = 0 \right\},$$

where $|\cdot|$ is a normalised absolute value of \mathcal{O}_k (cf. [2, §2, 5]). Let $L \stackrel{\text{def}}{=} \text{Fr}(\widehat{P})$ be the fraction field of \widehat{P} , and $\mathcal{C}_\infty \stackrel{\text{def}}{=} \text{Spec} L$. We shall refer to \mathcal{C}_∞ as a *formal boundary* of the formal germs \mathcal{D} , and \mathcal{C}_i for $i \geq 1$. We have natural scheme morphisms

$$\mathcal{C}_\infty \longrightarrow \cdots \longrightarrow \mathcal{C}_{n+1} \longrightarrow \mathcal{C}_n \longrightarrow \cdots \longrightarrow \mathcal{C}_1 \longrightarrow \mathcal{D} \longrightarrow \mathbb{P}_k^1.$$

Let η be a geometric point of \mathcal{C}_∞ , which induces a geometric point (denoted also η) of \mathcal{C}_n for $n \geq 1$. For $i \in \mathbb{N} \cup \{\infty\}$, we have an exact sequence of arithmetic fundamental groups

$$1 \longrightarrow \pi_1(\mathcal{C}_{i, \bar{k}}, \bar{\eta}) \longrightarrow \pi_1(\mathcal{C}_i, \eta) \longrightarrow G_k \longrightarrow 1, \tag{2}$$

where $\pi_1(\mathcal{C}_i, \eta)$ denotes the arithmetic étale fundamental group of \mathcal{C}_i with base point η , $\pi_1(\mathcal{C}_{i, \bar{k}}, \bar{\eta})$ the étale fundamental group of $\mathcal{C}_{i, \bar{k}} \stackrel{\text{def}}{=} C_i \times_{\text{Spec} k} \text{Spec} \bar{k}$ with base point $\bar{\eta}$; which is induced by η , and $G_k \stackrel{\text{def}}{=} \text{Gal}(\bar{k}/k)$ the absolute Galois group of k . Here \bar{k} is the algebraic closure of k determined by η .

Our main result in this note is the following.

Theorem 2. *We use notations as above. There exists an integer $N \geq 1$, such that for every integer $n \geq N$, the projection $\pi_1(\mathcal{C}_n, \eta) \rightarrow G_k$ doesn't split.*

The author ignores, for the time being, if the projection $\pi_1(\mathcal{C}_1, \eta) \rightarrow G_k$ splits or not.

As a corollary of Theorem 2, we obtain the following.

Theorem 3. *The projection $\pi_1(\mathcal{C}_\infty, \eta) \rightarrow G_k$ doesn't split.*

One of the consequences of Theorems 2 and 3 is that one can not produce examples of sections of hyperbolic curves over p -adic local fields, which arise from sections of arithmetic fundamental groups of boundaries of formal fibres, or open annuli with small radii. Those sections would be non-geometric, hence would provide counter-examples to the p -adic version of the Grothendieck anabelian section conjecture.

Finally we observe the following. For $i \in \mathbb{N} \cup \{\infty\}$, let $\pi_1(\mathcal{C}_{i,\bar{k}}, \bar{\eta})^{\text{ab}}$ be the maximal abelian quotient of $\pi_1(\mathcal{C}_{i,\bar{k}}, \bar{\eta})$, and consider the push-out diagram

$$\begin{array}{ccccccc}
 1 & \longrightarrow & \pi_1(\mathcal{C}_{i,\bar{k}}, \bar{\eta}) & \longrightarrow & \pi_1(\mathcal{C}_i, \eta) & \longrightarrow & G_k \longrightarrow 1 \\
 & & \downarrow & & \downarrow & & \parallel \\
 1 & \longrightarrow & \pi_1(\mathcal{C}_{i,\bar{k}}, \bar{\eta})^{\text{ab}} & \longrightarrow & \pi_1(\mathcal{C}_i, \eta)^{\text{(ab)}} & \longrightarrow & G_k \longrightarrow 1
 \end{array} \tag{3}$$

Thus, $\pi_1(\mathcal{C}_i, \eta)^{\text{(ab)}}$ is the geometrically abelian quotient of $\pi_1(\mathcal{C}_i, \eta)$.

Proposition 4. *The projection $\pi_1(\mathcal{C}_i, \eta)^{\text{(ab)}} \rightarrow G_k$ splits, $\forall i \in \mathbb{N}$.*

The author ignores, for the time being, if the projection $\pi_1(\mathcal{C}_\infty, \eta)^{\text{(ab)}} \rightarrow G_k$ splits or not.

2. Proof of Theorem 2

In this section we shall prove Theorem 2. We use the notations used in Section 1. We argue by contradiction, and assume that the projection $\pi_1(\mathcal{C}_n, \eta) \rightarrow G_k$ splits, $\forall n \geq 1$.

Proposition 5. *There exists a relative curve $X \rightarrow \text{Spec } \mathcal{O}_k$ with the following properties.*

- (i) *The morphism $X \rightarrow \text{Spec } \mathcal{O}_k$ is flat, proper, stable, and $X_k \stackrel{\text{def}}{=} X \times_{\text{Spec } \mathcal{O}_k} \text{Spec } k$ is geometrically connected.*
- (ii) *X is regular.*
- (iii) *The set of singular points X_F^{sing} of the special fibre $X_F \stackrel{\text{def}}{=} X \times_{\text{Spec } \mathcal{O}_k} \text{Spec } F$ of X consists of F -rational ordinary double points, $U \stackrel{\text{def}}{=} X_F \setminus X_F^{\text{sing}}$ is F -smooth, and $U(F) = \emptyset$ holds.*
- (iv) *$X(k) = \emptyset$ holds.*

Proof. First, assume $p \neq 2$. Let $\tilde{C} \stackrel{\text{def}}{=} \mathbb{P}_F^1$ with function field $k(\tilde{C})$. Thus, $\text{Card}(\tilde{C}(F)) = \text{Card } F + 1$ is even. Arrange the set $\tilde{C}(F)$ in pairs of F -rational points: $\tilde{C}(F) = \{(x_i, y_i)\}_{1 \leq i \leq \frac{\text{Card } F + 1}{2}}$. One can identify in \tilde{C} the points x_i and y_i ; $1 \leq i \leq \frac{\text{Card } F + 1}{2}$, to construct a stable proper F -curve C which is geometrically connected and geometrically reduced, with normalisation $\tilde{C} \rightarrow C$. Moreover, the set of singular points $C^{\text{sing}} = \{c_i\}_{1 \leq i \leq \frac{\text{Card } F + 1}{2}}$ consists of F -rational ordinary double points, and the pre-image of c_i in \tilde{C} consists of the two F -rational points $\{x_i, y_i\}$. In particular, $C(F) = C^{\text{sing}} = \{c_i\}_{1 \leq i \leq \frac{\text{Card } F + 1}{2}}$. More precisely, for $1 \leq i \leq \frac{\text{Card } F + 1}{2}$, let $\tilde{\mathcal{O}}_i \stackrel{\text{def}}{=} \mathcal{O}_{\tilde{C}, x_i} \cap \mathcal{O}_{\tilde{C}, y_i} \subset k(\tilde{C})$, $\mathcal{N}_{x_i} \stackrel{\text{def}}{=} \mathfrak{m}_{x_i} \cap \tilde{\mathcal{O}}_i$, and $\mathcal{N}_{y_i} \stackrel{\text{def}}{=} \mathfrak{m}_{y_i} \cap \tilde{\mathcal{O}}_i$, where \mathfrak{m}_{x_i} (resp. \mathfrak{m}_{y_i}) is the maximal ideal of $\mathcal{O}_{\tilde{C}, x_i}$ (resp. $\mathcal{O}_{\tilde{C}, y_i}$). Define $\mathcal{O}_{c_i} \stackrel{\text{def}}{=} F + \mathcal{N}_{x_i} \mathcal{N}_{y_i} \subset \tilde{\mathcal{O}}_i$. Then \mathcal{O}_{c_i} is a local ring (with maximal ideal $\mathcal{N}_{x_i} \mathcal{N}_{y_i}$, and residue field F) whose integral closure is $\mathcal{O}_{\tilde{C}, c_i}$ (cf. [1, Proposition 3.1, Theorem 3.4, and the references therein] for the properties of \mathcal{O}_{c_i} , as well as the existence of C with the required properties).

In case $p = 2$. Consider the affine F -curve $\text{Spec}(\frac{F[s,t]}{(st)})$, and \tilde{C} its smooth compactification. Thus, \tilde{C} consists of two F -smooth irreducible components $\tilde{C}_1 = \mathbb{P}_F^1$, and $\tilde{C}_2 = \mathbb{P}_F^1$, which intersect at the F -rational ordinary double point $c = (s, t) \in \text{Spec}(\frac{F[s,t]}{(st)})$. On each irreducible component \tilde{C}_i of \tilde{C} ; $1 \leq i \leq 2$, the set of F -rational points of $\tilde{C}_i \setminus \{c\}$ is non-empty and comes into pairs of rational points $\{(x_{i,j}, y_{i,j})\}_{1 \leq j \leq \frac{\text{Card } F}{2}}$. As above we can identify each of those pairs of F -rational points $(x_{i,j}, y_{i,j})$ into an F -rational ordinary double point $c_{i,j}$ to construct a reducible and geometrically connected stable curve F -curve C such that the set of singular points C^{sing} consists of F -rational

ordinary double points, a double point $c_{i,j}$ lies on a unique irreducible component of C , and $C^{\text{sing}} = C(F)$ (the local ring at c_i is defined as above; the case $p \neq 2$. See [7, §4], for a discussion of this procedure and the existence of such a curve C in the case of reducible curves).

Now the stable F -curve C can be deformed to a semi-stable \mathcal{O}_k -curve $X \rightarrow \text{Spec } \mathcal{O}_k$ with special fibre $X_F = C$ satisfying (i) and (ii) (cf. [9, Proposition 7.10, Corollary 7.11 and its proof]). By our construction (iii) holds also. If $x \in X(k)$, then X specialises in a point $\bar{x} \in C(F)$ which is a regular point of C and lies on a unique irreducible component of C (cf. [5, Corollary 9.1.32]). Thus, (iv) follows from (iii). \square

Let $X \rightarrow \text{Spec } \mathcal{O}_K$ be a regular, proper, flat, and stable \mathcal{O}_k -curve as in Proposition 5. Let $y \in X_F(F)$ be an F -rational point, which is an ordinary double point and a regular point of X (cf. Proposition 5 (ii) and (iii)). We fix an isomorphism $\rho : \widehat{\mathcal{O}}_{X,y} \xrightarrow{\sim} R[[S, T]]/(ST - \pi)$, and identify $\mathcal{X} \stackrel{\text{def}}{=} \text{Spec}(\widehat{\mathcal{O}}_{X,y} \otimes_{\mathcal{O}_k} k)$ with \mathcal{C}_1 via the isomorphism $\rho_k : \widehat{\mathcal{O}}_{X,y} \otimes_{\mathcal{O}_k} k \xrightarrow{\sim} \frac{R[[S, T]]}{(ST - \pi)} \otimes_{\mathcal{O}_k} k$ induced by ρ . Thus, we have scheme morphisms

$$\mathcal{C}_\infty \longrightarrow \cdots \longrightarrow \mathcal{C}_{n+1} \longrightarrow \mathcal{C}_n \longrightarrow \cdots \longrightarrow \mathcal{C}_1 \longrightarrow X_k.$$

For $n \geq 1$, write

$$\pi_1(X_k \setminus \mathcal{S}_n, \eta) \stackrel{\text{def}}{=} \varprojlim_{S_n \subset X_k \setminus \mathcal{C}_n} \pi_1(X_k \setminus S_n, \eta),$$

where the projective limit is over all finite sets of closed points $S_n \subset X_k \setminus \mathcal{C}_n$, and $\pi_1(X_k \setminus S_n, \eta)$ is the arithmetic fundamental group of the affine curve $X_k \setminus S_n$ with base point η . (Here, we identify the set of closed points of \mathcal{C}_n ; $n \geq 1$, with a subset of the set of closed points of X_k which specialise in y .) There is a natural projection $\pi_1(X_k \setminus \mathcal{S}_n, \eta) \twoheadrightarrow G_k$, and we have a commutative diagram

$$\begin{array}{ccc} \pi_1(\mathcal{C}_n, \eta) & \longrightarrow & G_k \\ \downarrow & & \parallel \\ \pi_1(X_k \setminus \mathcal{S}_n, \eta) & \longrightarrow & G_k \end{array} \tag{4}$$

where the left vertical map is induced by the morphism $\mathcal{C}_n \rightarrow X_k$.

Further, we have a natural map

$$\varprojlim_{n \geq 1} \pi_1(\mathcal{C}_n, \eta) \longrightarrow \varprojlim_{n \geq 1} \pi_1(X_k \setminus \mathcal{S}_n, \eta),$$

and $\varprojlim_{n \geq 1} \pi_1(X_k \setminus \mathcal{S}_n, \eta)$ is naturally identified with the absolute Galois group $G_{k(X)} \stackrel{\text{def}}{=} \text{Gal}(k(\overline{X})^{\text{sep}}/k(X))$, where $k(X)^{\text{sep}}$ is the separable closure of the function field $k(X)$ of X determined by the geometric point η .

Lemma 6. *The projection $G_{k(X)} \twoheadrightarrow G_k$ splits.*

Proof. First, our assumption that the projection $\pi_1(\mathcal{C}_n, \eta) \twoheadrightarrow G_k$ splits, implies that the projection $\pi_1(X_k \setminus \mathcal{S}_n, \eta) \twoheadrightarrow G_k$ splits, $\forall n \geq 1$ (cf. diagram (4)).

Let $(H_i)_{i \in I}$ be a projective system of quotients $G_{k(X)} \twoheadrightarrow H_i$, where H_i sits in an exact sequence $1 \rightarrow F_i \rightarrow H_i \rightarrow G_k \rightarrow 1$ with F_i finite, and $G_{k(X)} = \varprojlim_{i \in I} H_i$. [More precisely, write $G_{k(X)}$ as a projective limit of finite groups $\{\tilde{H}_i\}_{i \in I}$. Then \tilde{H}_i fits in an exact sequence $1 \rightarrow F_i \rightarrow \tilde{H}_i \rightarrow G_i \rightarrow 1$, where G_i is a quotient of G_k , and F_i a quotient of $\text{Gal}(k(X)^{\text{sep}}/k(X)\bar{k})$. Let $1 \rightarrow F_i \rightarrow H_i \rightarrow G_k \rightarrow 1$ be the pull-back of the group extension $1 \rightarrow F_i \rightarrow \tilde{H}_i \rightarrow G_i \rightarrow 1$ by $G_k \twoheadrightarrow G_i$. Then $G_{k(X)} = \varprojlim_{i \in I} H_i$]. The set $\text{Sect}(G_k, G_{k(X)})$ of group-theoretic sections of the projection $G_{k(X)} \twoheadrightarrow G_k$ is naturally identified with the projective limit $\varprojlim_{i \in I} \text{Sect}(G_k, H_i)$ of the sets $\text{Sect}(G_k, H_i)$ of group-theoretic sections of the projection $H_i \twoheadrightarrow G_k$. For each $i \in I$, the set $\text{Sect}(G_k, H_i)$ is non-empty. Indeed, H_i (being a quotient of $G_{k(X)}$) is a quotient of $\pi_1(X_k \setminus \mathcal{S}_n, \eta)$ for some $n \geq 1$, this quotient $\pi_1(X_k \setminus \mathcal{S}_n, \eta) \twoheadrightarrow H_i$ commutes with the projections onto G_k , and we know the projection

$\pi_1(X_k \setminus \mathcal{S}_n, \eta) \rightarrow G_k$ splits. Hence the projection $H_i \rightarrow G_k$ splits. Moreover, the set $\text{Sect}(G_k, H_i)$ is, up to conjugation by the elements of F_i , a torsor under the group $H^1(G_k, F_i)$ which is finite since k is a p -adic local field (cf. [6, (7.1.8) Theorem (iii)]). Thus, $\text{Sect}(G_k, H_i)$ is a nonempty finite set. Hence the set $\text{Sect}(G_k, G_{k(X)})$ is nonempty being the projective limit of nonempty finite sets. This finishes the proof of Lemma 6. (See also [8, the proof of Proposition 1.5] for similar arguments in a slightly different context.) \square

Let $s : G_k \rightarrow G_{k(X)}$ be a section of the projection $G_{k(X)} \twoheadrightarrow G_k$ (cf. Lemma 6).

Lemma 7. *The section s is geometric, i.e., $s(G_k) \subset D_x$, where $D_x \subset G_{k(X)}$ is a decomposition group associated to a (unique) rational point $x \in X(k)$. In particular, $X(k) \neq \emptyset$.*

Proof. This follows from [4, Proposition 2.4 (2)]. \square

Now the conclusion of Lemma 7 that $X(k) \neq \emptyset$ contradicts the assertion (iv) in Proposition 5 that $X(k) = \emptyset$. This is a contradiction. Thus, our assumption that the projection $\pi_1(\mathcal{C}_n, \eta) \twoheadrightarrow G_k$ splits, $\forall n \geq 1$, can not hold. Let $N \geq 1$ be such that the projection $\pi_1(\mathcal{C}_N, \eta) \twoheadrightarrow G_k$ doesn't splits. Then the projection $\pi_1(\mathcal{C}_n, \eta) \twoheadrightarrow G_k$ doesn't splits, $\forall n \geq N$ as required. Indeed, this follows from the fact that for $n \geq N$ we have a natural homomorphism $\pi_1(\mathcal{C}_n, \eta) \rightarrow \pi_1(\mathcal{C}_N, \eta)$ which commutes with the projections onto G_k . Hence if the projection $\pi_1(\mathcal{C}_n, \eta) \twoheadrightarrow G_k$ splits then the projection $\pi_1(\mathcal{C}_N, \eta) \twoheadrightarrow G_k$.

This finishes the proof of Theorem 2. \square

3. Proof of Theorem 3

Next, we explain how Theorem 3 can be derived from Theorem 2. We have, $\forall n \geq 1$, a commutative diagram

$$\begin{array}{ccc} \pi_1(\mathcal{C}_\infty, \eta) & \longrightarrow & G_k \\ \downarrow & & \parallel \\ \pi_1(\mathcal{C}_n, \eta) & \longrightarrow & G_k \end{array}$$

where the horizontal maps are the natural projections, and the left vertical map is induced by the morphism $\mathcal{C}_\infty \rightarrow \mathcal{C}_n$.

Now assume that the projection $\pi_1(\mathcal{C}_\infty, \eta) \twoheadrightarrow G_k$ splits. Then the projection $\pi_1(\mathcal{C}_n, \eta) \twoheadrightarrow G_k$ splits, $\forall n \geq 1$, by the above diagram. But this contradicts Theorem 2.

This finishes the proof of Theorem 3. \square

4. Proof of Proposition 4

Let $n \geq 1$ be an integer, and ℓ_1, ℓ_2 , distinct prime integers such that $\ell_1 \geq 2n$, and $\ell_2 \geq 2n$. Let \mathcal{O}_1 , and \mathcal{O}_2 , be totally ramified extensions of \mathcal{O}_k of degree ℓ_1 , and ℓ_2 , with fraction fields $L_1 = \text{Fr}(\mathcal{O}_1)$, and $L_2 = \text{Fr}(\mathcal{O}_2)$; respectively. Thus, the extensions L_1/k and L_2/k , are disjoint and $\mathcal{C}_n(L_i) \neq \emptyset$, for $i \in \{1, 2\}$. A restriction-corestriction argument shows that the class $[\pi_1(\mathcal{C}_n, \eta)^{\text{(ab)}}]$ of the group extension $\pi_1(\mathcal{C}_n, \eta)^{\text{(ab)}}$ in $H^2(G_k, \pi_1(\mathcal{C}_{n, \bar{k}}, \bar{\eta})^{\text{(ab)}})$ is trivial. Thus the group extension $\pi_1(\mathcal{C}_n, \eta)^{\text{(ab)}}$ splits.

This finishes the proof of Proposition 4. \square

References

- [1] Y. Aubry, A. Iezzi, “On the Maximum Number of Rational Points on Singular Curves over Finite Fields”, *Mosc. Math. J.* **15** (2015), no. 4, p. 615-627.
- [2] N. Bourbaki, *Éléments de mathématique. Algèbre commutative. Chapitre 9: Anneaux locaux noethériens complets*, Masson, 1983.
- [3] A. Grothendieck (ed.), *Revêtements étales et groupe fondamental*, Lecture Notes in Mathematics, vol. 224, Springer, 1971.
- [4] J. Koenigsmann, “On the ‘Section Conjecture’ in anabelian geometry”, *J. Reine Angew. Math.* **588** (2005), p. 221-235.
- [5] Q. Liu, *Algebraic geometry and arithmetic curves*, Oxford Graduate Texts in Mathematics, vol. 6, Oxford University Press, 2002.
- [6] J. Neukirch, A. Schmidt, K. Wingberg, “Cohomology of number fields”, 2000.
- [7] M. Rosenlicht, “Equivalence relations on algebraic curves”, *Ann. Math.* **56** (1952), no. 1, p. 169-191.
- [8] M. Saïdi, “On the existence of non-geometric sections of arithmetic fundamental groups”, *Math. Z.* **277** (2014), no. 1-2, p. 361-372.
- [9] M. Talpo, A. Vistoli, “Deformation theory from the point of view of fibered categories”, in *Handbook of moduli. Volume III*, Advanced Lectures in Mathematics (ALM), vol. 26, International Press, 2013, p. 281-397.