

MANUEL DE LEÓN

PAULO R. RODRIGUES

**Dynamical connections and non-autonomous
lagrangian systems**

Annales de la faculté des sciences de Toulouse 5^e série, tome 9, n° 2
(1988), p. 171-181

<http://www.numdam.org/item?id=AFST_1988_5_9_2_171_0>

© Université Paul Sabatier, 1988, tous droits réservés.

L'accès aux archives de la revue « Annales de la faculté des sciences de Toulouse » (<http://picard.ups-tlse.fr/~annales/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>

Dynamical connections and non-autonomous Lagrangian systems⁽¹⁾

MANUEL DE LEON⁽²⁾ and PAULO R. RODRIGUES⁽³⁾

RÉSUMÉ. — On montre que si ξ est une équation différentielle du deuxième ordre (semigerbe) sur le fibré des jets $J^1(\mathbf{R}, M)$ telle que les courbes intégrales sont des solutions de l'équation de Lagrange non-autonome alors il existe une connexion Γ sur $J^1(\mathbf{R}, M)$ dont les courbes intégrales sont aussi des solutions de la même équation. En plus, Γ est une connexion ayant comme semigerbe ξ . L'étude est une extension à la dynamique Lagrangienne non-autonome de quelques résultats de Grifone pour le cas autonome.

ABSTRACT. — We show that if ξ is a second-order differential equation (semispray) on the jet bundle $J^1(\mathbf{R}, M)$ whose paths are solutions of the non-autonomous Lagrange equations then there is a connection Γ on $J^1(\mathbf{R}, M)$ whose paths are also solutions of the same equations. Moreover, Γ is a connection whose associated semispray is precisely ξ . This is an extension to non-autonomous Lagrangian dynamics of a previous result due to Grifone for autonomous Lagrangians.

1. Introduction

The geometrical description of autonomous Lagrangian systems, started with GALLISOT [G], was elucidated by KLEIN [K1], [K2] (see also GODBILLON [GB]). He showed that the differential geometry of Lagrangian dynamics is intrinsically related to a (1.1) tensor field J , called *almost tangent structure*, defined on the tangent bundle of a manifold.

⁽¹⁾ Partially supported by Xunta de Galicia, Spain and CNPq-Brazil, Proc. 301115/79.

⁽²⁾ CECIME Consejo Superior de Investigaciones Científicas, Serrano 123, 28006 Madrid - Spain

⁽³⁾ Departamento de Geometria Instituto de Matematica, Universidade Federal Fluminense, 24.000 Niteroi, RJ - Brazil

Since the early works of KLEIN some articles have been published showing that almost tangent geometry provides a natural framework in which interesting generalizations of autonomous Lagrangian systems may be developed (see for instance, CRAMPIN [C], CRAMPIN et al. [CSC], de LEON and RODRIGUES [DLR1], [DLA2], GOTAY and NESTER [GN], SARLET et al. [SCC]). In particular, an extensive study about the theory of connections on tangent bundles in terms of the almost tangent geometry, including some aspects of autonomous Lagrangians, was proposed by GRIFONE [GR] around 1972.

As far as we know the non-autonomous case has been practically unknown in the literature (an exception, for example, is the recent paper of CRAMPIN and co-workers [CPT]). It is the purpose of this paper to establish some intrinsic properties about almost tangent theory of connections and its relation with non-autonomous Lagrangian dynamics. We will see that the theory of connections on the jet manifold $J^1(\mathbf{R}, M)$ is surprisingly more simpler than the theory of connections on TM , where M is a given manifold (the reader is invited to compare our results with GRIFONE).

2. Preliminaries

Throughout the text we shall keep in mind all results, definitions and notations previously introduced in [DLR 1] (see also [DLR 2]). All structures, functions, etc, are assumed to be smooth (C^∞). Let M be a manifold of dimension m (called *configuration* manifold) and Γ a connection on the tangent bundle TM of M . We recall here that a connection Γ on TM generates two projectors $h : T(TM) \rightarrow Hor(TM)$, $v : T(TM) \rightarrow Ver(TM)$ such that $T(TM) = Hor(TM) \oplus Ver(TM)$, where $Hor(TM)$ (resp. $Ver(TM)$) is the horizontal (resp. vertical) bundle over TM . If $\bar{\xi}$ is an arbitrary semispray (second-order differential equation) on TM then $\xi = h(\bar{\xi})$ is a semispray on TM which does not depend on the choice of $\bar{\xi}$. We call ξ the *associated* semispray of Γ . A connection Γ and its associated semispray have same paths.

If ξ is a semispray on TM then it can be shown that $\Gamma = -\mathcal{L}_\xi J$ is a connection on TM (here \mathcal{L}_ξ is the Lie derivative and $\mathcal{L}_\xi J$ is defined by

$$(\mathcal{L}_\xi J)(Y) = ([\xi, JY] - J[\xi, Y]).$$

When ξ is a spray (homogeneous second-order differential equation) then $\Gamma = -\mathcal{L}_\xi J$ is a connection on M such that its associated semispray is precisely ξ . For a semispray ξ there is a family of connections $\Gamma = -\mathcal{L}_\xi J + T$,

where T is a semibasic tensor field of type (1.1) on TM in equilibrium with ξ (in fact T is the strong torsion of Γ) (see [GR]). In the non-autonomous situation the relation between connections and semisprays becomes much more simpler, as we will show below.

The jet manifold $J^1(\mathbf{R}, M)$ is fibred over $\mathbf{R} \times M$, \mathbf{R} and M with projection maps π , π_1 , and π_2 . We notice that $J^1(\mathbf{R}, M)$ can be identified with $\mathbf{R} \times TM$ in a very natural way. Therefore we transport the geometric structures defined on TM to $J^1(\mathbf{R}, M)$ like the almost tangent structure J and the Liouville vector field C on TM . We may define a new tensor field \tilde{J} of type (1.1) on $J^1(\mathbf{R}, M)$ by

$$\tilde{J} = J - C \otimes dt, \quad (1)$$

which is locally characterized by

$$\tilde{J}(\partial/\partial t) = -C; \quad \tilde{J}(\partial/\partial x^i) = \partial/\partial x^i; \quad \tilde{J}(\partial/\partial y^i) = 0 \quad (2)$$

where (t, x, y) are local coordinates for $J^1(\mathbf{R}, M)$.

Hence \tilde{J} has rank m and satisfies $(\tilde{J})^2 = 0$. We define the adjoint of \tilde{J} , \tilde{J}^* , as the endomorphism of the exterior algebra $\Lambda(J^1(\mathbf{R}, M))$ of $J^1(\mathbf{R}, M)$ locally given by

$$\tilde{J}^*(dt) = 0, \quad \tilde{J}^*(dx^i) = 0, \quad \tilde{J}^*(dy^i) = dx^i - y^i dt. \quad (3)$$

Like in the autonomous situation we associate to \tilde{J} operators $i_{\tilde{J}}$ and $d_{\tilde{J}}$ on the algebra $\Lambda(J^1(\mathbf{R}, M))$ by

$$\left. \begin{aligned} i_{\tilde{J}}\omega(X_1, \dots, X_r) &= \sum_{\ell=1}^r \omega(X_1, \dots, \tilde{J}X_\ell, \dots, X_r), \\ d_{\tilde{J}} &= i_{\tilde{J}}d - di_{\tilde{J}}, \end{aligned} \right\} \quad (4)$$

and so we have

$$\left. \begin{aligned} i_{\tilde{J}}(df) &= \tilde{J}^*(df), \text{ for all } f \text{ on } J^1(\mathbf{R}, M) \\ i_{\tilde{J}}(dt) &= i_{\tilde{J}}(dx^i) = 0; \quad i_{\tilde{J}}(dy^i) = dx^i - y^i dt \end{aligned} \right\} \quad (5)$$

$$\left. \begin{aligned} d_{\tilde{J}}f &= \frac{\partial f}{\partial y^i} (dx^i - y^i dt) \\ d_{\tilde{J}}(dt) &= d_{\tilde{J}}(dx^i) = 0; \quad d_{\tilde{J}}(dy^i) = -d(dx^i - y^i dt) = dy^i \wedge dt. \end{aligned} \right\} \quad (6)$$

In the following we will set

$$\theta^i = dx^i - y^i dt, \quad 1 \leq i \leq m. \quad (7)$$

Also, it is not hard to see that a vector field ξ on $J^1(\mathbf{R}, M)$ is a semispray iff $\theta^i(\xi) = 0$ and $dt(\xi) = 1$, $1 \leq i \leq m$. In such a case ξ is locally given by

$$\xi = \partial/\partial t + y^i \partial/\partial x^i + \xi^i \partial/\partial y^i. \quad (8)$$

Furthermore, a vector field ξ on $J^1(\mathbf{R}, M)$ is a semispray iff $J\xi = C$ and $\tilde{J}\xi = 0$.

Let ξ be a semispray on $J^1(\mathbf{R}, M)$. A curve s in M is called a *path* of ξ if its canonical prolongation is an integral curve of ξ .

Let s be a curve in M locally given by $(x^i(t))$. Then $\tilde{s}^1(t) = (t, x^i(t), \dot{x}^i(t))$ and so s is a path of ξ if and only if satisfies the following non-autonomous system of differential equations

$$\frac{d^2 x^i}{dt^2} = \xi^i \left(t, x, \frac{dx}{dt} \right), \quad 1 \leq i \leq m$$

where ξ is given by (8).

3. Semisprays and dynamical connections

The tensor fields J and \tilde{J} on $J^1(\mathbf{R}, M)$ permit us to give a characterization of a kind of connections for the fibration $\pi : J^1(\mathbf{R}, M) \rightarrow \mathbf{R} \times M$.

DEFINITION (1).— *By a dynamical connection on $J^1(\mathbf{R}, M)$ we mean a tensor field Γ of type (1.1) on $J^1(\mathbf{R}, M)$ satisfying*

$$J\Gamma = \tilde{J}\Gamma = \tilde{J}, \quad \Gamma\tilde{J} = -\tilde{J}, \quad \Gamma J = -J. \quad (9)$$

By a straightforward computation from (9) we deduce that the local expressions of Γ are

$$\left. \begin{aligned} \Gamma(\partial/\partial t) &= -y^i \partial/\partial x^i + \Gamma^i \partial/\partial y^i, \\ \Gamma(\partial/\partial x^i) &= \partial/\partial x^i + \Gamma_i^j \partial/\partial y^j, \\ \Gamma(\partial/\partial y^i) &= -\partial/\partial y^i. \end{aligned} \right\} \quad (10)$$

The functions $\Gamma^i = \Gamma^i(t, x, y)$, $\Gamma_i^j = \Gamma_i^j(t, x, y)$ will be called the *components* of the connection Γ . From (10) we easily deduce that

$$\Gamma^3 - \Gamma = 0 \text{ and } \text{rank}(\Gamma) = 2m.$$

This type of polynomial structure is called $f(3, -1)$ -structure in the literature (see [YI]). Now, we can associate to Γ two canonical operators $\underline{\ell}$ and \underline{m} given by

$$\underline{\ell} = \Gamma^2, \quad \underline{m} = -\Gamma^2 + I.$$

Then we have

$$\underline{\ell}^2 = \underline{\ell}, \quad \underline{m}^2 = \underline{m}, \quad \underline{\ell}\underline{m} = \underline{m}\underline{\ell} = 0, \quad \underline{\ell} + \underline{m} = I, \quad (11)$$

and $\underline{\ell}$ and \underline{m} are complementary projectors. From (11) we deduce that $\underline{\ell}$ and \underline{m} are locally given by

$$\begin{aligned} \underline{\ell}(\partial/\partial t) &= -y^i \partial/\partial x^i - (\Gamma^i + y^j \Gamma_j^i) \partial/\partial y^i; \quad \underline{\ell}(\partial/\partial x^i) = \\ &= \partial/\partial x^i; \quad \underline{\ell}(\partial/\partial y^i) = \partial/\partial y^i; \quad \underline{m}(\partial/\partial t) = \\ &= \partial/\partial t + y^i \partial/\partial x^i + (\Gamma^i + y^j \Gamma_j^i) \partial/\partial y^i; \quad \underline{m}(\partial/\partial x^i) = \\ &= \underline{m}(\partial/\partial y^i) = 0. \end{aligned} \quad (12)$$

If we put $\mathcal{L} = I\mathbf{m}\underline{\ell}$, $\mathcal{M} = I\mathbf{m}\underline{m}$, then we have that \mathcal{L} and \mathcal{M} are complementary distributions on $J^1(\mathbf{R}, M)$, that is,

$$T(J^1(\mathbf{R}, M)) = \mathcal{M} \oplus \mathcal{L}.$$

From (12) we deduce that \mathcal{L} is $2m$ -dimensional and is locally spanned by $\{\partial/\partial x^i, \partial/\partial y^i\}$. \mathcal{M} is one-dimensional, globally spanned by the vector field $\xi = \underline{m}(\partial/\partial t)$. Taking into account the local expression of ξ , we deduce that ξ is a semispray which will be called the *canonical semispray associated to the dynamical connection* Γ .

Furthermore, we have $\Gamma^2 \underline{\ell} = \underline{\ell}$ and $\Gamma \underline{m} = 0$. Thus Γ acts on \mathcal{L} as an *almost product structure* and trivially on \mathcal{M} . Since $\mathcal{M} = \ker \Gamma$, Γ is said to be an $f(3, -1)$ -structure on $J^1(\mathbf{R}, M)$ of *rank* $2m$ and *parallelizable kernel*. Moreover, Γ/\mathcal{L} has eigenvalues $+1$ and -1 . From (10) the eigenspaces corresponding to the eigenvalue -1 are the vertical subspaces V_z , $z \in J^1(\mathbf{R}, M)$. Recall that for each $z \in J^1(\mathbf{R}, M)$, V_z is the set of all tangent vectors to $J^1(\mathbf{R}, M)$ at z which are projected to 0 by $T\pi$. Thus V is a distribution given by $z \mapsto V_z$. The eigenspace at $z \in J^1(\mathbf{R}, M)$ corresponding to the eigenvalue $+1$ will be denoted by H_z and called the *strong-horizontal subspace* at z . We have a canonical decomposition

$$T_z(J^1(\mathbf{R}, M)) = \mathcal{M}_z \oplus H_z \oplus V_z,$$

and obviously,

$$T(J^1(\mathbf{R}, M)) = \mathcal{M} \oplus H \oplus V, \quad (13)$$

where H is the distribution $z \mapsto H_z$.

Let us put $H'_z = \mathcal{M}_z \oplus H_z$; H'_z will be called the *weak-horizontal subspace at z* . Then we have the following decompositions

$$T_z(J^1(\mathbf{R}, M)) = H'_z \oplus V_z, \quad z \in J^1(\mathbf{R}, M)$$

and

$$T(J_1(\mathbf{R}, M)) = H' \oplus V, \quad (14)$$

where $H' : x \rightarrow H'_x$ is the corresponding distribution.

We notice that $\mathcal{L}, \mathcal{M}, H$ and H' may be considered as vector bundles over $J^1(\mathbf{R}, M)$; the bundles H and H' will be called *strong* and *weak-horizontal bundles*, respectively. Thus, from (14) Γ defines a connection on the fibration $\pi : J^1(\mathbf{R}, M) \rightarrow \mathbf{R} \times M$ with horizontal bundle H' (see ROUX [R] and de LEON & RODRIGUES [DLR 1]). But *not every connection on the fibration $\pi : J^1(\mathbf{R}, N) \rightarrow \mathbf{R} \times M$ arises in this way*.

A vector field X on $J^1(\mathbf{R}, M)$ which belongs to H (resp. H') will be called a *strong* (resp. *weak*) horizontal vector field. From (14), we have that the canonical projection $\pi : J^1(\mathbf{R}, M) \rightarrow \mathbf{R} \times M$ induces an isomorphism

$$\pi_* : H'_z \rightarrow T_{\pi(z)}(\mathbf{R} \times M), \quad z \in J^1(\mathbf{R}, M).$$

Then, if X is a vector field on $\mathbf{R} \times M$, there exists a unique vector field $X^{H'}$ on $J^1(\mathbf{R}, M)$ which is weak-horizontal and projects to X . The projection of $X^{H'}$ to H will be denoted by X^H .

From (10) and by a straightforward computation, we obtain

$$\begin{aligned} (\partial/\partial t)^{H'} &= \partial/\partial t + (\Gamma^j + \frac{1}{2} y^i \Gamma^j_i) \partial/\partial y^j \\ (\partial/\partial x^i)^{H'} &= \partial/\partial x^i + \frac{1}{2} \Gamma^j_i \partial/\partial y^j. \end{aligned} \quad (15)$$

Then, if we put $H_i = (\partial/\partial x^i)^{H'}$ and $V_i = \partial/\partial y^i$, one deduces that $\{\xi, H_i, V_i\}$ is a local basis of vector fields on $J^1(\mathbf{R}, M)$. In fact, $\mathcal{M} = \langle \xi \rangle$, $H = \langle H_i \rangle$, and $V = \langle V_i \rangle$; $\{\xi, H_i, V_i\}$ is called an *adapted basis* to the $f(3, -1)$ -structure Γ . In terms of $\{\xi, H_i, V_i\}$ (15) becomes

$$(\partial/\partial t)^{H'} = \xi - y^i H_i, \quad (\partial/\partial x^i)^{H'} = H_i.$$

Therefore, we obtain

$$(\partial/\partial t)^H = -y^i H_i, \quad (\partial/\partial x^i)^H = H_i.$$

If $X = \tau \partial/\partial t + X^i \partial/\partial x^i$ is a vector field on $R \times M$, we have

$$X^H = (X^i - \tau y^i) H_i \quad (16)$$

(compare with CRAMPIN, PRINCE and THOMPSON [CPT]). Finally, we notice that the dual local basis of 1-forms of the adapted basis $\{\xi, H_i, V_i\}$ is given by $\{dt, \theta^i, \psi^i\}$, where $\theta^i = dx^i - y^i dt$, and $\psi^i = -(\Gamma^i + \frac{1}{2} y^j \Gamma_j^i) dt - \frac{1}{2} \Gamma_j^i dx^j + dy^i$. This fact can be shown by a straightforward computation.

Let ξ be a semispray on $J^1(\mathbf{R}, M)$ and suppose that ξ is locally expressed by

$$\xi = \partial/\partial t + y^i \partial/\partial x^i + \xi^i \partial/\partial y^i. \quad (17)$$

Then a simple computation in local coordinates shows that

$$\left. \begin{aligned} [\xi, \partial/\partial t] &= -\frac{\partial \xi^j}{\partial t} \partial/\partial y^j, \\ [\xi, \partial/\partial x^i] &= -\frac{\partial \xi^j}{\partial x^i} \partial/\partial y^j, \\ [\xi, \partial/\partial y^i] &= -\frac{\partial}{\partial x^i} - \frac{\partial \xi^j}{\partial y^i} \partial/\partial y^j. \end{aligned} \right\} \quad (18)$$

PROPOSITION (1).— Let $\Gamma = -\mathcal{L}_\xi \tilde{J}$. Then Γ is a dynamical connection on $J^1(\mathbf{R}, M)$ whose associated semispray is, precisely, ξ .

Proof.— In fact from (18) we have

$$\left. \begin{aligned} \Gamma(\partial/\partial t) &= -y^i \partial/\partial x^i - \left(y^j \frac{\partial \xi^i}{\partial y^j} - \xi^i \right) \partial/\partial y^i, \\ \Gamma(\partial/\partial x^i) &= \partial/\partial x^i + \frac{\partial \xi^j}{\partial y^i} \partial/\partial y^j, \\ \Gamma(\partial/\partial y^i) &= -\partial/\partial y^i. \end{aligned} \right\} \quad (19)$$

Now, from (19) we easily deduce that Γ is a dynamical connection on $J^1(\mathbf{R}, M)$. Furthermore, taking into account (12), we have that the associated semispray to Γ is, precisely, ξ .

From Proposition (1) we may observe that the theory of dynamical connections on $J^1(\mathbf{R}, M)$ is more simpler than the theory of connections on TM .

Let Γ be a dynamical connection on $J^1(\mathbf{R}, M)$.

DEFINITION (2).— *A curve $u : \mathbf{R} \rightarrow M$ is called a path of Γ if the canonical prolongation $j^1 u$ of u to $J^1(\mathbf{R}, M)$ is a weak-horizontal curve.*

Now, we shall find the differential equations for the paths of Γ (the dots meaning time derivatives).

If $u : \mathbf{R} \rightarrow M$ is locally given by $t \mapsto (x^i(t))$, then we have $j^1 u(t) = (t, x^i(t), \dot{x}^i(t))$. Hence,

$$\dot{j}^1 u(t) = \frac{\partial}{\partial t} + \frac{dx^i}{dt} \frac{\partial}{\partial x^i} + \frac{d^2 x^i}{dt^2} \frac{\partial}{\partial y^i}.$$

Therefore, u is a path of Γ if and only if $\psi^i(\dot{j}^1 u(t)) = 0$, $1 \leq i \leq m$, that is u satisfies the following system of differential equations :

$$\frac{d^2 x^i}{dt^2} = \Gamma^i \left(t, x, \frac{dx}{dt} \right) + \Gamma_j^i \left(t, x, \frac{dx}{dt} \right) \frac{dx^j}{dt}. \quad (20)$$

Let ξ be the associated semispray of Γ . Then ξ is locally given by

$$\xi = \partial/\partial t + y^i \partial/\partial x^i + \xi^i \partial/\partial y^i,$$

where $\xi^i = \Gamma^i + y^j \Gamma_j^i$, $1 \leq i \leq m$.

From (20) it is clear that the paths of Γ and ξ satisfy the same system of differential equations. Then we have

PROPOSITION (2).— *A dynamical connection and its associated semispray on $J^1(\mathbf{R}, M)$ have the same paths.*

4. Dynamical connections and non-autonomous regular Lagrangian equations

Suppose that a non-autonomous regular Lagrangian L is given, that is, L is a non-degenerate real function on $J^1(\mathbf{R}, M) = \mathbf{R} \times TM$. Then it is

well-known that an extremal for L is a curve $s : \mathbf{R} \rightarrow M$ (or a section of $(\mathbf{R} \times M, p, \mathbf{R})$) such that

$$(\tilde{s})^*(i_X dL \wedge dt) = 0 \quad (21)$$

for all vertical vector fields on $\mathbf{R} \times TM$. Also, it is known that (21) is equivalent to

$$(\tilde{s}^2)^*(i_X d\Omega_L) = 0, \quad (22)$$

for all π_1 -vertical vector fields on $J^1(\mathbf{R}, M)$. In (22) Ω_L is the POINCARÉ-CARTAN canonical form on $J^1(\mathbf{R}, M)$ locally given by

$$\Omega_L = L(t, x, y)dt + \frac{\partial L}{\partial y^i} \theta^i,$$

where θ^i is defined in (7) of section 2.

In terms of the tensor field \tilde{J} and J and the Liouville vector field C on $J^1(\mathbf{R}, M)$, the POINCARÉ-CARTAN form takes the following expression :

$$\Omega_L = L dt + \frac{\partial L}{\partial y^i} \theta^i = L dt + d\tilde{J}L,$$

or equivalently,

$$\begin{aligned} \Omega_L &= L dt + \frac{\partial L}{\partial y^i} dx^i - y^i \frac{\partial L}{\partial y^i} dt = \left(L - y^i \frac{\partial L}{\partial y^i} \right) dt + d_J L \\ &= (L - CL)dt + d_J L = d_J L - E_L dt; E_L = CL - L. \end{aligned}$$

Thus

$$\Theta_L = d\Omega_L = dd\tilde{J}L + dL \wedge dt$$

or

$$\Theta_L = dd_J L - dE_L \wedge dt.$$

A straightforward computation in local coordinates shows that

$$\Theta_L \wedge \cdots \wedge \Theta_L = \pm \det \left(\frac{\partial^2 L}{\partial y^j \partial y^i} \right) dx^1 \wedge \cdots \wedge dx^m \wedge dy^1 \wedge \cdots \wedge dy^m$$

and if L is a non-autonomous regular Lagrangian we deduce that Θ_L is a contact form on $J^1(\mathbf{R}, M)$. Consequently, the *characteristic* bundle of Θ_L

$$R_{\Theta_L} = \{v \in T(J^1(\mathbf{R}, M)); i_v \Theta_L = 0\}$$

has one-dimensional fibers, that is, they are a line-bundle over $J^1(\mathbf{R}, M)$. Let us recall here that a vector field X on $J^1(\mathbf{R}, M)$ is *characteristic* if X is a section of R_{Θ_L} , that is, $i_X \Theta_L = 0$. The following result can be compared with the corresponding one for autonomous Lagrangian [see [DLR 1]].

PROPOSITION (3).— *Let L be a non-autonomous regular Lagrangian on $J^1(\mathbf{R}, M)$ and ξ a characteristic vector field which satisfies $i_\xi dt = 1$. Then ξ is a semispray on $J^1(\mathbf{R}, M)$ whose paths are the solutions of the Lagrange equations*

$$\frac{d}{dt} \left(\frac{\partial L}{\partial y^i} \right) - \frac{\partial L}{\partial x^i} = 0, \quad 1 \leq i \leq m.$$

We call ξ the Lagrange vector field for L .

THEOREM (1).— *Let L be a non-autonomous regular Lagrangian on $J^1(\mathbf{R}, M)$ and let ξ be a Lagrange vector field for L . Then there exists a dynamical connection Γ on $J^1(\mathbf{R}, M)$ whose paths are the solutions of the Lagrange equations. This connection is given by $\Gamma = -\mathcal{L}_\xi \tilde{J}$.*

Proof.— From Proposition (1) we deduce that $\Gamma = -\mathcal{L}_\xi \tilde{J}$ is a dynamical connection whose associated semispray is precisely ξ . Thus the theorem follows directly from Proposition (2) and (3).

Finally, let us remark that the results of CRAMPIN, PRINCE and THOMPSON [CPT] can be re-obtained in terms of Γ . In fact, with the notation of Section 3 we have a local basis of vector fields on $J^1(\mathbf{R}, M)$ given by $\{\xi, H_i, V_i\}$ where H_i is given by

$$H_i = \frac{\partial}{\partial x^i} + \frac{1}{2} \frac{\partial \xi^j}{\partial y^i} \frac{\partial}{\partial y^j}.$$

Thus the corresponding dual basis is $\{dt, \theta^i, \psi^i\}$, where

$$\psi^i = - \left(\xi^i - \frac{1}{2} y^j \frac{\partial \xi^i}{\partial y^j} \right) dt - \frac{1}{2} \frac{\partial \xi^i}{\partial y^j} dx^j + dy^i.$$

The significance of this dual basis is that the form Θ_L can be re-written as follows

$$\Theta_L = \frac{\partial^2 L}{\partial y^i \partial y^j} \theta^i \wedge \psi^j$$

and so the semispray ξ is uniquely determined by the equations

$$i_\xi \theta^i = i_\xi \psi^i = 0, \quad i_\xi dt = 1.$$

References

- [C] CRAMPIN (M.). — Tangent bundle geometry for Lagrangian dynamics, *J. Phys. A : Math. Gen.*, t. **16**, 1983, p. 3755-3772.
- [CPT] CRAMPIN (M.), PRINCE (G.), THOMPSON (G.). — A geometrical version of the Helmholtz conditions in time-dependent Lagrangian dynamics, *J. Phys. A : Math. Gen.*, t. **17**, 1984, p. 1437-1447.
- [DLR 1] DE LEON (M.), RODRIGUES (P.R.). — *Generalized Classical Mechanics and Field Theory*. — North-Holland Math. Studies, t. **112**, 1985.
- [DLR 2] DE LEON (M.), RODRIGUES (P.R.). — Formalisme hamiltonien symplectique sur les fibrés tangents d'ordre supérieur, *C.R. Acad. Sc. Paris*, t. **301**, ser. **II**, **7**, 1985, p. 455- 458.
- [G] GALLISSOT (T.). — Les formes extérieures en Mécanique, *Ann. Inst. Fourier, Grenoble*, t. **4**, 1952, p. 145-297.
- [GN] GOTAY (M.), NESTER (J.). — Presymplectic Lagrangian Systems, I/II. *Ann. Inst. Henri Poincaré* **30**,(2), 1979, p. 129-142; *ibidem*, **32**,(1), 1980, p. 1-13.
- [GR] GRIFONE (J.). — Structure presque-tangente et connexions, I/II. *Ann. Inst. Fourier (Grenoble)*, **22**,(3), 1972, p. 287-334; *ibidem*, **22**,(4), p. 291-338.
- [K 1] KLEIN (J.). — Espaces variationnels et Mécanique, *Ann. Inst. Fourier (Grenoble)*, t. **12**, 1962, p. 1-124.
- [K 2] KLEIN (J.). — Opérateurs différentiels sur les variétés presque- tangentes, *C.R. Acad. Sc. Paris*, t. **257 A**, 1963, p. 2392-2394.
- [R] ROUX (A.). — Jets et connexions, *Publ. Math. Univ. Lyon*, t. **7**, 1970, p. 1-42.
- [SCC] SARLET (W.), CANTRIJN (F.), CRAMPIN (M.). — A new look at second-order equations and Lagrangian mechanics, *J. Phys. A: Math. Gen.*, t. **17**, 1984, p. 1999-2009.
- [YI] YANO (K.), ISHIHARA (S.). — *Tangent and Cotangent bundles*. — Pure and Appl. Math. Ser. **16**, Marcel Dekker, N.Y., 1973.

(Manuscrit reçu le 27 janvier 1987)