

ANNALES DE L'I. H. P., SECTION B

MARIUSZ LEMAŃCZYK

Toeplitz Z_2 -extensions

Annales de l'I. H. P., section B, tome 24, n° 1 (1988), p. 1-43

http://www.numdam.org/item?id=AIHPB_1988__24_1_1_0

© Gauthier-Villars, 1988, tous droits réservés.

L'accès aux archives de la revue « Annales de l'I. H. P., section B » (<http://www.elsevier.com/locate/anihpb>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

Toeplitz Z_2 -extensions

by

Mariusz LEMAŃCZYK

Institute of Mathematics, Nicholas Copernicus University,
ul. Chopina 12/18, 87-100 Torun, Poland

RÉSUMÉ. — Nous allons nous occuper d'une sous-classe spéciale de la classe des Z_2 -extensions d'automorphismes ayant un spectre rationnel discret.

Quelques connections avec des points strictement transitifs dans l'espace $\{0,1\}^Z$ (particulièrement avec la théorie d'automates, la théorie de suites de Morse et de Toeplitz) sont établies.

On construit quelques exemples d'extension de cette sorte ayant $2k$ -spectres lebesguiens ($k \geq 1$) (sur l'orthocomplémentaire de l'espace des fonctions propres).

Une nouvelle classe de points strictement transitifs dans $\{0,1\}^Z$, nommée k -Morse-suites ($k \geq 1$) est introduite. Nous développons leur théorie métrique en démontrant qu'il y a continuum non isomorphe 2-Morse suites qui possède un 2-spectre lebesguien (sur l'orthocomplémentaire de l'espace des fonctions propres).

ABSTRACT. — We shall concern ourselves with some special subclass of ergodic Z_2 -extensions of automorphisms with rational discrete spectra. Some connections with strictly transitive points in the space $\{0,1\}^Z$ (in particular with automata theory, Morse sequences theory, Toeplitz sequences theory) are exhibited. Examples of such extensions with $2k$ -fold, $k \geq 1$, Lebesgue spectrum (in the orthocomplement of the space of eigenfunctions) are constructed. A new class of strictly transitive points in $\{0,1\}^Z$ called k -Morse sequences, $k \geq 1$, is introduced. We develop their metric theory proving there are continuum of 2-Morse sequences pairwise nonisomorphic having 2-fold Lebesgue spectrum (in the orthocomplement of the space of eigenfunctions).

Key words : Lebesgue spectrum, Toeplitz sequence, ergodic theory, automata theory.

INTRODUCTION

The theory presented in the paper combines methods and results from a few papers [1], [5], [8], [10], [11], [12], [14], [16] and it is mainly connected with the possibility of the construction of ergodic automorphisms with nonsingular spectra and finite spectral multiplicity.

So far, there are two extremely interesting examples of such automorphisms. The first arises from automata theory. In [6] there is a proof that Rudin-Shapiro automata has a Lebesgue component in its spectrum. Queffelec in her dissertation developed spectral theory of dynamical systems generated by automata. She calculated that Rudin-Shapiro automata has twofold Lebesgue spectrum (in the orthocomplement of the eigenfunctions space). The second example (in fact continuum examples) was constructed in [14]. The authors considered some Z_2 -extensions over an automorphism with discrete spectrum and proved that such examples had twofold Lebesgue spectrum in the orthocomplement of the eigenfunctions space.

Let us consider another two examples. In 1968 Keane [8] introduced to ergodic theory a class of strictly transitive points called generalized Morse sequences. However, in 1980 in [1] a new class of 0-1- and strictly transitive sequences was provided. The authors called them also generalized Morse sequences.

Are there some connections between those four examples? First of all, we see that the dynamical systems arising from the strictly transitive points above are always Z_2 -extensions of automorphisms with rational pure point spectra. But their main feature in common is that for such an $x \in \{0, 1\}^{\mathbb{N}}$ the sequence $\hat{x} \in \{0, 1\}^{\mathbb{N}}$ is a regular Toeplitz sequence ([5], [18]) ($\hat{x}[n] = x[n] + x[n+1] \pmod{2}$). On the other hand Mathew-Nadkarni's construction also leads to a class of regular Toeplitz sequences. In particular the results of this paper show that the theory of Rudin-Shapiro automata strictly includes in the Mathew-Nadkarni's theory.

Below, we intend to clarify in some details our approach to the problems arising.

Let (X, \mathcal{B}, μ) be a Lebesgue space and $T : (X, \mathcal{B}, \mu) \rightarrow (X, \mathcal{B}, \mu)$ be its automorphism. Suppose T to be ergodic and to have a rational discrete spectrum, i. e. there is a sequence $\{n_t\}$, $n_t | n_{t+1}$, $t \geq 0$ such that

$$\text{Sp}(T) = G \{ n_t : t \geq 0 \},$$

the group generated by $\{ \exp 2\pi i/n_t : i=0, \dots, n_t-1 \}$, $t \geq 0$ [here and in what follows $\text{Sp}(T)$ denotes always the point spectrum of an automorphism

T]. Consider now a measurable map $\Theta : X \rightarrow Z_2 = \{0, 1\}$. We can define T_Θ , a Z_2 -extension of T by

$$\left. \begin{aligned} T_\Theta : (X \times Z_2, \tilde{\mu}) \ni \\ T_\Theta(x, i) = (Tx, \Theta(x) + i), \end{aligned} \right\} \quad (1)$$

where $\tilde{\mu}$ is the product measure $\mu \times \nu$ ($\nu(i) = 1/2, i = 1, 0$) and $+$ is always meant in Z_2 (unless it states otherwise). We will require Θ to satisfy some additional properties. Our concept lies in the following. It is well-known that if T is ergodic then T has a rational pure point spectrum with $Sp(T) = G\{n_t : t \geq 0\}$ iff there is a sequence $\{D^{n_t}\}$ of n_t -T-stacks such that $D^{n_t} \nearrow \varepsilon$ (the point partition), where $D^{n_t} = \{D_0^{n_t}, \dots, D_{n_t-1}^{n_t}\}$, $T^i D_0^{n_t} = D_i^{n_t} \pmod{n_t}$ and D^{n_t} is a partition of X. Hence, to define a measurable $\Theta : X \rightarrow Z_2$ [i. e. two measurable sets $\Theta^{-1}(0), \Theta^{-1}(1)$] it is sufficient to assume

$$\Theta^{-1}(i) \subseteq \{D^{n_t}\}, \quad i = 0, 1. \quad (2)$$

This condition simply means that $\Theta^{-1}(i)$ consists of some levels of D^{n_t} , $t \geq 0$. In other words there exists a partition of the set $\{0, 1, \dots, n_t - 1\}$ into A^t, B^t, C^t such that

$$\begin{aligned} i \in A^t & \text{ iff } \Theta|_{D_i^{n_t}} = 0, \\ i \in B^t & \text{ iff } \Theta|_{D_i^{n_t}} = 1, \\ i \in C^t & \text{ iff } \Theta|_{D_i^{n_t}} \text{ is not constant.} \end{aligned} \quad (3)$$

Let us pass to the stack $D^{n_{t+1}}$. There is again a partition $A^{t+1}, B^{t+1}, C^{t+1}$ of $\{0, 1, \dots, n_{t+1} - 1\}$ satisfying (3). Denote

$$\lambda_{t+1} = n_{t+1}/n_t$$

Then

$$\begin{aligned} A^t, A^t + n_t, \dots, A^t(\lambda_{t+1} - 1)n_t \subset A^{t+1}, \\ B^t, B^t + n_t, \dots, B^t + (\lambda_{t+1} - 1)n_t \subset B^{t+1}. \end{aligned} \quad (4)$$

Further, given A^t, B^t, C^t we can define a finite sequence $\Theta^{(t)}$ of 0's and 1's and some "holes", $\Theta^{(t)} = (\Theta_0^{(t)}, \dots, \Theta_{n_t-1}^{(t)})$, where


$$\Theta_i^{(t)} = \begin{cases} 0 & \text{if } i \in A^t \\ 1 & \text{if } i \in B^t \\ \infty & \text{if } i \in C^t \end{cases}$$

Due to (4) $\Theta^{(t+1)}$ arises from $\Theta^{(t)}$ by repeating $\Theta^{(t)}$ λ_{t+1} times and by filling up some “holes”. This is just a method of constructing of Toeplitz sequences ([5], [18]). It is obvious that the method presented of constructing Θ leads always to a measurable function defined a. e. However, in order to assure T_Θ to have some “good” properties we claim Θ to be a regular Toeplitz sequence [18]. Now, we will consider some special examples of such Z_2 -extensions.

Let $\Theta \in \{0, 1\}^{\mathbb{Z}}$ be a regular Toeplitz sequence. Put $X = X(\Theta)$ (where $X(u) = \{\tau^i u : i \in \mathbb{Z}\}$, $u \in \{0, 1\}^{\mathbb{Z}}$, τ is the shift). Thus (X, τ) is a strictly ergodic dynamical system ([5], [18]), so there is a unique (ergodic) τ -invariant measure μ on X . Let $\{n_t\}$, $n_t | n_{t+1} t \geq 0$, define a period structure of Θ (see [18]). Then there is a sequence $D^{n_t} = \{D_0^{n_t}, \dots, D_{n_t-1}^{n_t}\}$, $t \geq 0$, and $D_i^{n_t}$ are simultaneously open and closed, $\tau^i D_0^{n_t} = D_i^{n_t} \pmod{n_t}$ and $D^{n_t} \nearrow \varepsilon$ [18]. Moreover,

$$D_i^{n_t} = \{y \in X(\Theta) : y \text{ has the same } n_t\text{-skeleton as } \tau^i \Theta\} \quad [18] \quad (5)$$

(here Θ is meant as two-sided sequence). In other words if n_t -skeleton of Θ is equal to, say, A , then y is of the form (Picture 1)



Picture 1

where the “holes” of A are filled up and y need not be a Toeplitz sequence at all.

It is well-known that $(X(\Theta), \tau, \mu)$ has a rational pure point spectrum with $\text{Sp}(\tau) = G\{n_t : t \geq 0\}$ where $\{n_t\}$ defines a period structure of Θ . Consider, now, the function $\Theta : X(\Theta) \rightarrow Z_2$ given by

$$\Theta(y) = y[0].$$

This function is continuous and taking into considerations D^{n_t} and n_t -skeleton of Θ we observe that $\Theta|_{D_i^{n_t}}$ is constant and equal to $\Theta[i]$ whenever $\Theta[i]$ is not the “hole” in the n_t -skeleton of Θ . Such extensions we will call *Toeplitz Z_2 -extensions*.

So, we will consider regular Toeplitz sequences as maps $\Theta : X \rightarrow Z_2$ and we will examine properties of τ_Θ . This is our first point of view.

To present our second approach (combinatorial) we have to introduce some operations on 0-1-sequences and blocks.

Let $B=(b_0, \dots, b_{r-1})$ be a block, $b_i \in \{0, 1\}$, $|B|=r$ is called the length of B . Then, by \hat{B} we mean the block $\hat{B}=(\hat{b}_0, \dots, \hat{b}_{r-2})$, $\hat{b}_i=b_i+b_{i+1}$, $i=0, \dots, r-2$. The inverse operation \check{B} yields two blocks C and \check{C} , where \check{C} is got from C by interchanging all of the 0's and 1's (\check{C} is the mirror image of C). These notations are easily adapted to the cases one-sided or two-sided sequences. In the sequel we treat $\check{}$ rather as a function. To avoid some difficulties, from now on, we will assume that if Θ is a regular Toeplitz sequence then it has the following property: *a block B appears on Θ iff the block \check{B} appears on it.* In the case of strict ergodicity of Θ this condition is equivalent to the one that $X(\Theta)$ is mirror-invariant. It will turn out (Section I) that there are some connections between τ on $X(\Theta)$ and τ_Θ . Moreover, from metric theory point of view these two automorphisms are indistinguishable.

Now, we briefly describe further results of the paper.

In Section II we present spectral theory of Z_2 -extensions. As a corollary we will get that in order to obtain a Z_2 -extension with Lebesgue maximal spectral type (in the orthocomplement of the eigenfunctions space) it is sufficient to construct a regular Toeplitz sequence $\Theta \in \{0,1\}^{\mathbb{N}}$ such that $\sum_{i \geq 1} |a_i^\Theta|^2 < \infty$, where

$$a_i^\Theta = \mu \left(\bigcup_{\substack{|B|=i \\ B \text{ contains} \\ \text{an even} \\ \text{number} \\ \text{of 1's}}} B \right) - \mu \left(\bigcup_{\substack{|B|=i \\ B \text{ contains} \\ \text{an odd} \\ \text{number} \\ \text{of 1's}}} B \right)$$

and μ is the measure defined by the average frequencies of blocks on Θ .

We also reach an estimation of spectral multiplicity of τ_Θ . This allows to provide for every $k \geq 1$ a class of Z_2 -extensions with 2^k -fold Lebesgue spectrum (in the orthocomplement of the space of eigenfunctions) (see III. 2).

In [1] the authors proposed to study properties of generalized Morse sequences. We do so for the rest of the paper. First of all we introduce a new class of strictly transitive points called k -Morse sequences, $k \geq 1$. Generalized Morse sequences introduced by Keane [8] are included in this theory for $k=1$, those introduced in [1] for $k=2^s$, $s \geq 1$. We develop metric theory of dynamical systems arising from such sequences. The tools used resemble those presented by the author in [10], [11], [12] and by M. K.

Mentzen and the author in [13]. In particular, we prove that Mathew-Nadkarni's construction contains continuum pairwise nonisomorphic dynamical systems.

The author would like to thank J. Kwiatkowski and M. K. Mentzen for helpful discussions.

I. ERGODIC TOEPLITZ Z_2 -EXTENSIONS AND SHIFT DYNAMICAL SYSTEMS

Let $\Theta \in \{0, 1\}^{\mathbb{N}}$ be a regular Toeplitz sequence. We recall here that τ_{Θ} is ergodic with respect to $\tilde{\mu}$ iff τ_{Θ} is uniquely ergodic [17]. Then for any partition $Q = (Q_0, Q_1)$ of $X(\Theta) \times Z_2$ and any $(x, i) \in X(\Theta) \times Z_2$ there exist the average frequencies of all blocks on Q - ∞ -name of (x, i) i. e. let

$$b_j = \begin{cases} 0 & \text{if } \tau_{\Theta}^j(x, i) \in Q_0 \\ 1 & \text{if } \tau_{\Theta}^j(x, i) \in Q_1 \end{cases}$$

[the one-sided sequence b is then called Q - ∞ -name of (x, i)], then

$$\lim_n \frac{1}{n} \text{fr}(\mathbf{B}, b[0, n-1]),$$

exists, where

$$\begin{aligned} \text{fr}(\mathbf{B}, b[0, n-1]) &= \text{card} \{ i : b[i, i + |\mathbf{B}| - 1] = \mathbf{B} \}, \\ \mathbf{B} &\in \{0, 1\}^k \end{aligned}$$

(in fact $\lim_n \frac{1}{n} \text{fr}(\mathbf{B}, b[i, i + n - 1])$ exists uniformly with respect to i).

THEOREM 1. — *Let τ_{Θ} be an ergodic Toeplitz Z_2 -extension. Then Θ is strictly transitive.*

Proof. — Let us take $Q_i = X(\Theta) \times \{i\}$, $i = 0, 1$. Then Q is a partition of $X(\Theta)$ into two open sets. Now, we will examine Q - n -names of points from Q_0 . We have $D_{\mathcal{G}}^n \times \{0\} \subset Q_0$, $t \geq 0$ and fix $(x, 0) \in D_{\mathcal{G}}^n \times \{0\}$. Let $\Theta^{(t)}$ be n_t -skeleton of Θ and suppose m_t to be the first place of appearing of a "hole", i. e.

$$\Theta^{(t)} = \Theta[0], \dots, \Theta[m_t - 1], \infty, \dots$$

and $\Theta [i] \in \{ 0, 1 \}$, $0 \leq i \leq m_t - 1$.

Then

$$m_t \leq m_{t+1} \quad \text{and} \quad m_t \xrightarrow{t} \infty \tag{7}$$

Let us observe that

$$\left. \begin{aligned} \tau_{\Theta}^k(x, 0) = (\tau^k x, \Theta(\tau^{k-1} x) + \dots + \Theta(\tau x) + \Theta(x)), \\ k = 0, 1, \dots \end{aligned} \right\} \tag{8}$$

From (8) it follows that

$$\begin{aligned} Q-n\text{-name of } (x, 0) = (0, \Theta(x), \Theta(\tau x) + \Theta(x), \\ \dots, \Theta(\tau^{n-1} x) + \dots + \Theta(\tau x) + \Theta(x)) \end{aligned}$$

This implies immediately that

$$(Q - m_t\text{-name of } (x, 0))^\wedge = \Theta[0, m_t - 1]. \tag{9}$$

Now, take into considerations the sets

$$X_k = \{ (x, 0) \in Q_0 : Q - m_k\text{-name of } (x, 0) = \check{\Theta}[0, m_k - 1], \check{\Theta}[0] = 0 \}$$

Then X_k is closed since τ_{Θ} is continuous (Θ is continuous) and from (9) $X_k \neq \emptyset$. Further

$$X_k \supset X_{k+1} \supset \dots$$

in view of (7). Since $X(\Theta) \times Z_2$ is compact, $\bigcap_{i \geq 1} X_i \neq \emptyset$ and therefore there is $(x, 0) \in Q_0$ such that $Q - \infty\text{-name of } (x, 0) = \check{\Theta}$. Hence $\check{\Theta}$ has the average frequencies of all blocks.

Remark. — If $\tilde{\mu}$ is the only ergodic measure then τ_{Θ} is in addition minimal since $\tilde{\mu}$ has positive measure on all open nonempty sets. Therefore in this case $\check{\Theta}$ is almost periodic (Q_i are open, $i=0, 1$).

LEMMA 1. — Let Θ be a Toeplitz sequence and $\check{\Theta}$ be almost periodic. Then $X(\check{\Theta}) = X(\Theta)^\check{=} = \{ \check{y} : y \in X(\Theta) \}$.

Proof. — If $y \in X(\Theta)$, then there must exist $\{i_s\}$ such that $\tau^{i_s} \Theta \rightarrow y$, this implies $\tau^{i_s} \check{\Theta} \rightarrow \check{y}$. Therefore $X(\Theta)^\check{=} \subset X(\check{\Theta})$. To see the inverse inclusion it is sufficient to prove $X(\Theta)^\check{=}$ is closed. So, let us assume $\check{y}^n \rightarrow z$,

$z \in X(\check{\Theta})$. We have, $\{\check{y}^n\}$ is a Cauchy sequence, so the sequence $\{y^n\}$ is also. Hence $y^n \rightarrow y \in X(\Theta)$ and $\check{y}^n \rightarrow \check{y}$.

Now, we are able to reverse Theorem 1.

THEOREM 2. — *Let Θ be a regular Toeplitz sequence and $\check{\Theta}$ be almost periodic. If $\check{\Theta}$ is also strictly transitive, then τ_Θ is (uniquely) ergodic.*

Proof. — Suppose, on the contrary, τ_Θ not to be ergodic. Then there is a measurable, nonconstant function $f: X(\Theta) \rightarrow S^1$ such that

$$f(\tau y) = (-1)^{\Theta(y)} f(y) \text{ a. e. } y \in X(\Theta) \quad [17]. \quad (10)$$

Let us take $g: X(\check{\Theta}) \rightarrow S^1$, $g(z) = (-1)^{z[0]} f(\hat{z})$. Then g is well defined from Lemma 1, measurable and it is not constant. We wish to show g is τ -invariant. To this aim we calculate

$$g(\tau z) = (-1)^{\tau z[0]} f(\tau z) = (-1)^{z[1]} f(\tau \hat{z}) \stackrel{(10)}{=} (-1)^{z[1] + \Theta(z)} f(\hat{z}).$$

To prove g is τ -invariant it remains to show $z[0] = z[1] + \Theta(\hat{z})$. However, from (6) $\Theta(\hat{z}) = \hat{z}[0]$ and the proof is complete.

As we have seen a regular Toeplitz sequence induces an ergodic Z_2 -extension iff it induces some strictly ergodic dynamical system. This does not specifically exclude the possibility that these two dynamical systems are distinct. The following theorem clarifies this situation.

THEOREM 3. — *Let Θ be a regular Toeplitz sequence and τ_Θ be ergodic. Then τ_Θ and τ on $X(\check{\Theta})$ are metrically isomorphic.*

Proof. — From the proof of Theorem 1 it follows that $(X(\check{\Theta}), \tau)$ is isomorphic to the factor of τ_Θ generated by the partition Q . All we have to show is that Q is a generator of τ_Θ .

Let \mathcal{B}_Q be a σ -algebra generated by $\bigvee_{-\infty}^{\infty} \tau_\Theta^i Q$ and η_Q be the corresponding measurable partition. Denote by η the partition of $X(\Theta) \times Z_2$ with the atoms of the form

$$\{(z, 0), (z, 1)\}, \quad z \in X(\Theta).$$

Then $\eta_Q \geq \eta$ since $\tau: X(\check{\Theta}) \curvearrowright$ has $\tau: X(\Theta) \curvearrowright$ as a factor and η is a canonical partition [15]. Moreover $\eta_Q \neq \eta$ since $Q_i \in \mathcal{B}_Q$ and $Q_i \notin \mathcal{B}(\eta)$ (the σ -algebra of η -measurable sets). This forces η_Q to be equal to ε because η_Q must have the same number of points in a. e. atom.

II. SOME SPECTRAL INVESTIGATIONS

II.1. Maximal spectral type

Let $T : (X, \mathcal{B}, \mu) \curvearrowright$ be an ergodic automorphism with discrete spectrum and $\text{Sp}(T) = G \{n_i : i \geq 0\}$. Let $\varphi : X \rightarrow Z_2$ be measurable. Then

$$L^2(X \times Z_2, \tilde{\mu}) = \mathcal{D} \oplus \mathcal{E}$$

where

$$\mathcal{D} = \{f \in L^2(X \times Z_2, \tilde{\mu}) : f \sigma = f\}, \quad \mathcal{E} = \{f \in L^2(X \times Z_2, \tilde{\mu}) : f \sigma = -f\},$$

where

$$\sigma : (X \times Z_2, \tilde{\mu}) \curvearrowright, \quad \sigma(x, i) = (x, i + 1).$$

Then $U_{T_\varphi}|_{\mathcal{D}}$ is spectrally isomorphic to $U_T : L^2(X, \mu)$.

Let us assume T_φ is ergodic. It is known [12] that if there is any point spectrum on \mathcal{E} then T_φ has again pure point spectrum.

Define also $V^\varphi : L^2(X, \mu) \curvearrowright$, by

$$V^\varphi(f)(x) = (-1)^{\varphi(x)} f(Tx).$$

Then V^φ is unitary and moreover

$$\left. \begin{aligned} U_{T_\varphi}|_{\mathcal{E}} \text{ is spectrally isomorphic to } V^\varphi \\ ([3], [14]). \end{aligned} \right\} \quad (11)$$

It turns out that when dealing with spectral properties of T_φ , the base role is played by the sequence

$$((V^\varphi)^n 1, 1) = \int_X (V^\varphi)^n(1) d\mu.$$

It is clear that

$$\int_X (V^\varphi)^n 1 d\mu = \mu \{x : \varphi(x) + \dots + \varphi(T^{n-1}x) = 0\} - \mu \{x : \varphi(x) + \dots + \varphi(T^{n-1}x) = 1\}.$$

Now, let Θ be a regular Toeplitz sequence. Denote by

$$a_n^\Theta = \mu \left(\bigcup_{\substack{|\mathbf{B}|=n \\ \mathbf{B} \text{ has an} \\ \text{even number} \\ \text{of 1's}}} \mathbf{B} \right) - \mu \left(\bigcup_{\substack{|\mathbf{B}|=n \\ \mathbf{B} \text{ has an} \\ \text{odd number} \\ \text{of 1's}}} \mathbf{B} \right) \tag{12}$$

$n = 1, 2, \dots$

PROPOSITION 1. — *Let Θ be a regular Toeplitz sequence. Then*

$$((V^\Theta)^k 1, 1) = a_k^\Theta, \quad k = 1, 2, \dots \tag{13}$$

Proof. — The assertion follows from the following simple facts:

$$\frac{1}{n_t} (\text{number of the "holes" in } n_t\text{-skeleton of } \Theta) \rightarrow 0, \tag{14}$$

$$\mu(D_i^{n_t}) = 1/n_t, \tag{15}$$

there are the average frequencies of all blocks on Θ . (16)

PROPOSITION 2. — *Let Θ be a regular Toeplitz sequence such that Θ is strictly transitive. Then*

$$a_n^\Theta = \mu_{\check{\Theta}}(00^{(n)} \cup 11^{(n)}) - \mu_{\check{\Theta}}(01^{(n)} \cup 10^{(n)}),$$

where $\mu_{\check{\Theta}}$ is the measure given by the average frequencies on $\check{\Theta}$ and

$$\begin{aligned} \mu(ij^{(n)}) &= \mu \{ y \in X(\check{\Theta}) : y[0] = i, y[n-1] = j \} \\ &= \lim_N \frac{1}{N} \text{fr} \left(\underbrace{i \infty \dots \infty j}_{n-2}, \check{\Theta}[0, N-1] \right). \end{aligned}$$

Proof. — Let us observe that if $\Theta[i] = 1$, then

$$\check{\Theta}[i] + \check{\Theta}[i+1] = 1$$

(i. e. the pair $(\check{\Theta}[i], \check{\Theta}[i+1])$ is disagreed). So, if $(\Theta_i, \Theta_{i+1}, \dots, \Theta_j)$ contains an even (odd) number of 1's, then $(\check{\Theta}_i, \check{\Theta}_{i+1}, \dots, \check{\Theta}_{j+1})$ contains an even (odd) number of pairs disagreed.

The following result is well-known (for instance [16]).

PROPOSITION 3. — *The maximal spectral type of V^Θ is equal to*

$$\sigma_m = \sum_{k=1}^{\infty} 2^{-k} (\sigma * \delta_{\alpha_k})$$

where σ is the spectral measure of 1, i. e. $\sigma = \sigma_1$ and $\{\alpha_k\}$ is the sequence of all eigenvalues of T .

**II. 2. Ergodicity,
mixing and Lebesgue spectrum**

Suppose T_φ to be ergodic and take $f(x, i) = (-1)^i$, $x \in X$, $i \in Z_2$. Then using Birkhoff theorem we see that

$$\frac{1}{N} \sum_{k=0}^{N-1} [f(T_\varphi^k(x, i))] \xrightarrow{N} 0.$$

By Lebesgue integral theorem

$$\frac{1}{N} \sum_{k=0}^{N-1} \int_X [(-1)^{\varphi(T^{k-1}x) + \dots + \varphi(x)}] d\mu(x) \xrightarrow{N} 0.$$

Thus, we obtain

$$\frac{1}{N} \sum_{k=0}^{N-1} ((V^\varphi)^k 1, 1) \xrightarrow{N} 0 \tag{17}$$

as a necessary condition T_φ to have the ergodicity property.

Denote

$$((V^\varphi)^n 1, 1) = a_n^\varphi, \quad n = 0, 1, \dots$$

COROLLARY 1. — T_φ is ergodic iff

$$\frac{1}{N} \sum_{i=0}^{N-1} \alpha_k^i a_i^\varphi \rightarrow 0, \quad \text{for all } \alpha_k \in \text{Sp}(T). \tag{18}$$

Proof. — The condition (18) gives

$$\frac{1}{N} \sum_{s=0}^{N-1} (U_{T_\varphi}^s F_k, F_k) \rightarrow 0,$$

where $F_k(x, s) = (-1)^s g_k(x)$ and g_k is the eigenfunction corresponding to $\alpha_k \in \text{Sp}(T)$.

But $\{F_k\}$ is an orthonormal base on \mathcal{C} , so

$$\frac{1}{N} \sum_{s=0}^{N-1} (U_{T_\varphi}^s f, f) \rightarrow 0$$

holds on the whole space \mathcal{C} . On \mathcal{D} (19) is valid since T was ergodic.

COROLLARY 2. — T_φ is weakly mixing on \mathcal{C} iff

$$\frac{1}{N} \sum_{s=0}^{N-1} |a_s^\varphi| \rightarrow 0, \quad (20)$$

iff

$$a_n^\varphi \rightarrow 0, \\ n \notin M$$

where the density of M is equal to 0.

Proof. — Let us notice that σ_1 is continuous iff (20) holds and σ_1 is continuous iff σ_m is continuous.

COROLLARY 3. — T_φ is mixing on \mathcal{C} iff $a_n^\varphi \rightarrow 0$.

Proof. — Let g_k be as in the proof of Corollary 1. Hence

$$((V^\varphi)^n g_k, g_k) = \alpha_k^n a_n^\varphi, \quad \text{so } ((V^\varphi)^n g_k, g_k) \rightarrow 0.$$

As a consequence of Proposition 3 we get then

$$\int_{S^1} z^n d\sigma_m \rightarrow 0.$$

So, from the generalized Lebesgue-Riemann lemma $\int_{S^1} z^n dv \rightarrow 0$ whenever

$v \ll \sigma_m$.

COROLLARY 4. — (a) $\sigma_m \equiv \lambda$ iff $\sigma_1 \ll \lambda$ (λ is the Lebesgue measure);

(b) if $\sigma_m \ll \lambda$, then $\sigma_m \equiv \lambda$;

(c) if $\sum_{n \geq 1} |\sigma_1(n)|^2 < \infty$, then $\sigma_m \equiv \lambda$;

(d) σ_m is singular iff σ_1 is singular.

Proof. — For the proof of (a) we notice that the set of eigenvalues $\{\alpha_k\}$ of T gives a dense set on S^1 . The remain statements are clear.

**II. AN ESTIMATION
OF SPECTRAL MULTIPLICITY**

THEOREM 4. — *Let Θ be a regular Toeplitz sequence with the following property: any n_t -skeleton $\Theta^{(t)}$ has at most K of the “holes” and $\Theta_{n_t-1}^{(t)} = \infty$. Then the maximal spectral multiplicity of V^Θ is also at most K .*

Proof. — We have $\Theta^{(t)} = (\Theta_0^{(t)}, \dots, \Theta_{n_t-1}^{(t)})$. Let i_1, \dots, i_K be the all indices such that $\Theta_{i_s}^{(t)} = \infty$, $s = 1, \dots, K$. Next, we define functions $f_1^t, \dots, f_K^t \in L^2(X(\Theta), \mu)$ putting

$$f_s^t = \chi_{D_{i_s+1}^{n_t}}, \quad s = 2, \dots, K, \quad f_1^t = \chi_{D_0^{n_t}}. \tag{21}$$

Now, let $Z(f_1^t, \dots, f_K^t)$ be the smallest V^Θ -invariant subspace of $L^2(X(\Theta), \mu)$.

Since

$$(V^\Theta)^r f_s^t = (-1)^{\varphi + \varphi r + \dots + \varphi r^{r-1}} f_s^t(\tau^r),$$

if Θ is constant on $D_{i_s+1}^{n_t}, \dots, D_{i_s+r}^{n_t}$ i.e. $\Theta_{i_s+1}^{(t)}, \dots, \Theta_{i_s+r}^{(t)}$ are not the “holes”), then

$$(V^\Theta)^r f_s^t = \pm f_s^t(\tau^r) = \pm \chi_{D_{i_s+r}^{n_t}} \pmod{n_t}. \tag{22}$$

This leads to the following

$$L^2(D^{n_t}) \subset Z(f_1^t, \dots, f_K^t), \tag{23}$$

where $L^2(D^{n_t})$ is the space of all measurable functions constant on the elements of D^{n_t} . Further

$$Z(f_1^t, \dots, f_K^t) \subset Z(f_1^{t+1}, \dots, f_K^{t+1}). \tag{24}$$

Indeed, it is sufficient to show $f_s^t \in Z(f_1^{t+1}, \dots, f_K^{t+1})$ and even, due to (23), that $f_s^t \in L^2(D^{n_t+1})$. But the latter is obvious, since $D^{n_t} \subseteq D^{n_t+1}$. In virtue of $D^{n_t} \nearrow \varepsilon$ we get

$$\overline{\bigcup_t Z(f_1^t, \dots, f_K^t)} = L^2(X, \mu)$$

and (24) completes the proof.

III. EXAMPLES

III. 1. Generalized Morse sequences [8]

We refer the reader to the Morse shifts theory to [8] and [11]. Our notations are as in [11].

It is well-known that the dynamical systems arising from Morse sequences are Z_2 -extensions (see for instance [9]). We intend to determine the form of the function defining the Z_2 -extension.

To this end, let us introduce the so called \star -product (see [2]). Let $B \in \{0, 1\}^n$, $C \in \{0, 1\}^m$

$$B \star C = \begin{cases} B c_0 B c_1 \dots B c_{m-1} B & \text{if } B \text{ contains an even number of } 1\text{'s} \\ B \tilde{c}_0 B \tilde{c}_1 \dots B \tilde{c}_{m-1} B & \text{otherwise} \end{cases}$$

Then if d^0, d^1, \dots is a sequence of blocks with lengths at least one and starting with 1 then the formula

$$z = d^0 \star d^1 \star \dots \quad (25)$$

defines a Toeplitz sequence. The classic Toeplitz sequence is then equal to $1 \star 1 \star \dots$ [4].

Let us denote $|d^i| = \lambda_i - 1$, $\lambda_i \geq 2$. Then $n_i = \lambda_0 \dots \lambda_i$ -skeleton of z is equal to $z^{(i)} = d^0 \star \dots \star d^i \infty$, $|z^{(i)}| = n_i$.

In other words the formula (25) leads always to the so called 1-spaced Toeplitz sequences (n_i -skeleton must contain only one "hole" at the last place). The reader can check that any 1-spaced Toeplitz sequence is of the form (25).

Now, we determine \check{z} . To this end let us recall that

$$B \times C = B^{c_0} B^{c_1} \dots B^{c_{m-1}}, \quad B^0 = B, \quad B^1 = \tilde{B}.$$

The following formula holds

$$(B \times C)^\wedge = \hat{B} \star \hat{C}. \quad (26)$$

Therefore $\check{z} = \check{d}^0 \times \check{d}^1 \times \dots$

So, if $x = b^0 \times b^1 \times \dots$ is a Morse sequence, then $\hat{x} = \hat{b}^0 \star \hat{b}^1 \star \dots$. In addition n_i -skeleton of \hat{x} is equal to

$$\hat{b}^0 \star \hat{b}^1 \star \dots \star \hat{b}^i \infty = \hat{c}_i \infty \quad (c_i = b^0 \times \dots \times b^i).$$

Morse sequences play a peculiar role in general ergodic theory (see [12])

and especially in the theory of Z_2 -extensions over automorphisms with rational discrete spectra. In [12] there is a proof that Morse sequences compose a “typical” set of automorphisms (in the strong topology). In fact, Katok-Stepin theory of an odd approximation of Z_2 -extensions is included in the theory of Morse sequences (for the speed of approximation at least $o(1/n^2)$ [12]).

As an immediate consequence of Theorem 4 we get all Morse sequences have simple spectra.

**III. 2. A class of Toeplitz
 Z_2 -extensions
with nonsingular spectra**

Let $m = 2^k$, $k \geq 1$. We will construct a class of Toeplitz Z_2 -extensions such that V^Θ will have uniform Lebesgue spectrum with maximal spectral multiplicity equal to m .

We will consider the Toeplitz sequence Θ as a limit of some periodic sequences $\text{Per } \Theta^{(n)}$ (see [5]), where

$$\text{Per } \Theta^{(n)} = | a_1^{(n)} a_2^{(n)} \dots a_{k_n}^{(n)} | a_1^{(n)} a_2^{(n)} \dots a_{k_n}^{(n)} | \dots \in \{ 0, 1, \infty \}^N$$

and Θ is constructed from $\{ \text{Per } \Theta^{(n)} \}$ in the following way. We first write down $\text{Per } \Theta^{(1)}$, next we write down $\text{Per } \Theta^{(2)}$ but only in the “holes”, $\text{Per } \Theta^{(3)}$ is written down in the remains “holes” and so on. The periodic part of $\text{Per } \Theta^{(n)}$ will be said to be an n -base-skeleton of Θ and denoted by $\text{per } \Theta^{(n)}$.

Now, set

$$\begin{aligned} \text{per } \Theta^{(r)} &= a_1^{(r)} \infty a_2^{(r)} \infty \dots \infty a_{2^{k-1}}^{(r)} \infty \tilde{a}_1^{(r)} \infty \tilde{a}_2^{(r)} \infty \dots \infty \tilde{a}_{2^{k-1}}^{(r)} \infty, \\ a_i^{(r)} &\in \{ 0, 1 \}, \quad r = 1, 2, \dots \end{aligned}$$

So, the corresponding Toeplitz sequence arises in the following way:

$$\begin{array}{cccccccccccc} a_1^{(1)} \infty a_2^{(1)} \infty a_3^{(1)} \infty a_4^{(1)} \infty \dots & \infty & \tilde{a}_{2^{k-1}}^{(1)} \infty & | & a_1^{(1)} \infty a_2^{(1)} \infty a_3^{(1)} \infty a_4^{(1)} \infty \dots & \infty & \tilde{a}_{2^{k+1}}^{(1)} \infty & | & \dots \\ a_1^{(2)} & \infty & a_2^{(2)} & \infty \dots & a_{2^k}^{(2)} & \infty & \tilde{a}_1^{(2)} & \infty & \tilde{a}_2^{(2)} & \infty \dots & \tilde{a}_{2^k}^{(2)} & \infty \dots \\ a_1^{(3)} & & & \infty \dots & & & & & & & & \\ \dots & & & & & & & & & & & \end{array}$$

Any sequence Θ arising as the above has the following properties:

$$a_1^\Theta = 0. \tag{27}$$

since on each base-skeleton the frequencies of 0's and 1's are the same,

$$2^r\text{-skeleton of } \Theta = \Theta^{(r)} = A_1^{(r)} \infty A_2^{(r)} \infty \dots \infty A_m^{(r)} \infty, \tag{28}$$

where

$$|A_i^{(r)}| = 2^r - 1, \quad i = 1, \dots, m.$$

We wish to get

$$a_n^\Theta = 0 \quad \text{for } n \geq 2. \tag{29}$$

Let us fix $n \geq 2$. Then, find $r \geq 1$ such that

$$2^r \leq n \leq 2^{r+1} \tag{30}$$

and consider 2^r -skeleton of Θ (*Picture 2*).

$$\underbrace{A_1^{(r)} \infty \dots \infty A_m^{(r)} \infty}_{\mathcal{C}_1} \quad \underbrace{A_1^{(r)} \infty A_2^{(r)} \infty \dots \infty A_m^{(r)} \infty}_{\mathcal{C}_2} \quad \underbrace{A_1^{(r)} \infty A_2^{(r)} \infty \dots \infty A_m^{(r)} \infty}_{\mathcal{C}_3} \infty \dots$$

Picture 2

In each group \mathcal{C}_i we number the “holes”, in the same way (*Picture 3*)

$$\underbrace{A_1^{(r)1} \infty A_2^{(r)2} \infty \dots \infty A_2^{(r)2^k}}_{\mathcal{C}_i}$$

Picture 3

Now, we fix our attention on $r+1, r+2, \dots, r+k$ -base skeletons of Θ .

We get

1^0 ($r+1$)-base-skeleton fills up in each \mathcal{C}_i 2^{k-1} of the “holes” at the places of the form $2^0 + 2^1 s$ (in each group \mathcal{C}_i the same places!)

2^0 ($r+2$)-base-skeleton fills up in each \mathcal{C}_i 2^{k-2} of the “holes” at the places of the form $2^1 + 2^2 s$ (in each group \mathcal{C}_i the same places!)

...
 j^0 ($r+j$)-base-skeleton fills up in each \mathcal{C}_i 2^{k-j} of the "holes" at the places of the form $2^{j-1} + 2^j s$ (in each group \mathcal{C}_i the same places!)

...
 k^0 ($r+k$)-base-skeleton fills up in each \mathcal{C}_i 1 of the "holes" at the place 2^{k-1} (in each \mathcal{C}_i the same place!).

In addition, we notice that for any j , $1 \leq j \leq k$, $r+j$ -base-skeleton can be written down as follows:


$$(b_1^{(1)} \infty \dots \infty b_{2^{k-j}}^{(1)}) \infty (b_1^{(2)} \infty \dots \infty b_{2^{k-j}}^{(2)}) \infty \dots$$

$$\infty (b_1^{(2^{j-1})} \infty \dots \infty b_{2^{k-j}^{(2^{j-1})}}^{(2^{j-1})}) \infty (\tilde{b}_1^{(1)} \infty \dots \infty \tilde{b}_{2^{k-j}}^{(1)})$$

$$\infty (\tilde{b}_1^{(2)} \infty \dots \infty \tilde{b}_{2^{k-j}}^{(2)}) \infty \dots \infty (\tilde{b}_1^{(2^{j-1})} \infty \dots \infty \tilde{b}_{2^{k-j}^{(2^{j-1})}}^{(2^{j-1})}) \infty = \text{per } \Theta^{(r+j)}$$

Now, let $|B| = n$. Due to (28) and (30) if we consider blocks of such length on 2^r -skeleton of Θ , then B must contain at least one "holes" and at most two "holes".

We take into consideration the case B is a subblock of $A_i^{(r)} \infty A_{i+1}^{(r)}$, $i \neq 2^k$. Then, there is a unique j , $1 \leq j \leq k-1$ that the i -th "hole" is filled up in $r+j$ -base-skeleton. Now, the easy proof follows from Picture 4 below [$t_1, \dots, t_{2^{k-j}}$ are the numbers of "holes" that are filled up by $(r+j)$ -base-skeleton]



Picture 4

The frequencies of blocks of length n , which are subblocks of $A_i^{(r)} \infty A_{i+1}^{(r)}$, $i \neq 2^k$ simply cancel out.

We assert that for any n the frequencies of blocks of length n starting at places of the form (32) cancel out. Indeed, for $n \geq 2$ this is the reasoning we have dealt with before a moment. It remains the case $n=1$. For i even the result is obvious. For i odd the reasoning as in case B contains one "hole" is deciding.

Now, notice that

$$((V^\Theta)^n i, \chi_{D_i^{2^k}}) = \lim_N \frac{1}{N} \text{fr}(\mathbf{B} : |\mathbf{B}| = n,$$

\mathbf{B} starts at a place of the form (32)

and \mathbf{B} contains an even number of 1's, $\Theta [0, N-1]$)

$$-\lim_N \frac{1}{N} \text{fr}(\mathbf{B} : |\mathbf{B}| = n,$$

\mathbf{B} starts at a place of the form (32)

and \mathbf{B} contains an odd number of 1's, $\Theta [0, N-1]$). (33)

Combining the considerations above and (33) we get

$$((V^\Theta)^n 1, \chi_{D_i^{2^k}}) = 0 \quad \text{for } n \geq 1 \quad (i=0, \dots, 2^k-1). \quad (34)$$

The following formulae are also obvious

$$((V^\Theta)^n \chi_{D_i^{2^k}}, \chi_{D_i^{2^k}}) = \left\{ \begin{array}{ll} ((V^\Theta)^n 1, \chi_{D_i^{2^k}}) & \text{if } n = i \bmod 2^k \\ 0 & \text{otherwise} \end{array} \right\} \quad (35)$$

$$((V^\Theta)^n \chi_{D_i^{2^k}}, \chi_{D_j^{2^k}}) = \left\{ \begin{array}{ll} ((V^\Theta)^n 1, \chi_{D_j^{2^k}}) & \text{if } n+i-j = j \bmod 2^k \\ 0 & \text{otherwise} \end{array} \right\} \quad (36)$$

Combining (34), (35) and (36) we obtain that the spectral multiplicity is at least 2^k and we have proved the desired result.

III. 3. Generalized Morse sequences [1]

In [1] the authors proposed to examine properties of the following sequences.

Let $u = u_{l-1} \dots u_0 \in \{0, 1, \infty\}^l$, $u_{l-1} \neq \infty$.

Now, define an infinite sequence $x_u \in \{0, 1\}^{\mathbb{N}}$ putting

$$x_u[n] = \text{card} \{ i \geq 0 : (\forall j = 0, \dots, l-1) u_j \neq \infty \rightarrow u_j = e_{j+1}(n) \} \bmod 2,$$

where

$$n = \sum_{i=0}^{\infty} e_i(n) 2^n$$

$$[e_i(n) = 0 \text{ for } i > \log n / \log 2],$$

i. e. $x_u[n]$ is the number of appearances of u on n written in binary expansion.

In [1] there is a proof that if u is not trivial (i. e. $u_i = 1$ for some i , $0 \leq i \leq l-1$), then u is recognizable by a 2-substitution. Moreover, there is more precise description in terms of $(k, 2k)$ -substitutions. Namely, let u be nontrivial and let $E(u)$ be the set of all $v = (v_0, \dots, v_{l-1}) \in \{0, 1\}^l$ obtaining from u by filling up all the "holes". Put

$$\text{Sp}(u) = \left\{ \sum_{i=0}^{l-1} v_i 2^i : v \in E(u) \right\} \quad \text{and} \quad k = 2^{l-1}.$$

Then x_u is a fixed point of the $(k, 2k)$ -substitution

$$\sigma(y_0, \dots, y_{k-1}) = \Theta^{(\alpha_0)} y_0 \Theta^{(\alpha_1)} y_1 \dots \Theta^{(\alpha_{k-1})} y_{k-1}. \quad (37)$$

$y_i \in \{0, 1\}$ and

$$\begin{aligned} \alpha_j &= 0 & \text{if } 2j \notin \text{Sp}(u) \text{ and } 2j+1 \notin \text{Sp}(u) \\ \alpha_j &= 1 & \text{if } 2j \in \text{Sp}(u) \text{ and } 2j+1 \notin \text{Sp}(u) \\ \alpha_j &= 2 & \text{if } 2j \notin \text{Sp}(u) \text{ and } 2j+1 \in \text{Sp}(u) \\ \alpha_j &= 3 & \text{if } 2j \in \text{Sp}(u) \text{ and } 2j+1 \in \text{Sp}(u) \end{aligned}$$

and $\Theta^{(0)}, \Theta^{(1)}, \Theta^{(2)}, \Theta^{(3)}$ are all of the continuous substitutions on $\{0, 1\}$ of length two, i. e. $\Theta^{(i)} 0 = \Theta^{(i)}(1)$ [1].

Let us observe that the column number [16] of the substitution (37) is equal to 2. It suggests that from ergodic theory point of view they are rational Z_2 -extensions. For a precise proof of this fact it is sufficient to show that \hat{x}_u is always a regular Toeplitz sequence. However, we do not prove this at the stage since we will introduce a more general class of strictly transitive sequences including both generalized Morse sequences introduced by Keane and those from [1].

III. 4. k -Morse sequences

Let k be a natural number, $k \geq 1$. We will consider blocks the lengths of which are some multiple of k (we call them k -blocks).

Let $B \in \{0, 1\}^{rk}$, $C \in \{0, 1\}^{sk}$. Then

$$B = B_1 \dots B_k, \quad |B_i| = r,$$

$$C = (c_0, \dots, c_{k-1})(c_k, \dots, c_{2k-1}) \dots (c_{(s-1)k}, \dots, c_{sk-1}).$$

We define the k -product of k -blocks,

$$B \times_k C = B_1^{c_0} B_2^{c_1} \dots B_k^{c_{k-1}} B_1^{c_k} B_2^{c_{k+1}} \dots B_k^{c_{2k-1}} \dots B_1^{c_{(s-1)k}} B_2^{c_{(s-1)k+1}} \dots B_k^{c_{sk-1}} \quad (38)$$

where

$$B_i^0 = B_i, \quad B_i^1 = \tilde{B}_i, \quad i = 1, \dots, k.$$

We have

$$|B \times_k C| = krs. \quad (39)$$

It is also worthwhile to note that k -product is an associative operation.

Now, let b^0, b^1, b^2, \dots be k -blocks and

$$|b^i| = k \lambda_i, \quad \lambda_i \geq 2,$$

$$b^i = (b_0^{(i)}, \dots, b_{k-1}^{(i)})(b_k^{(i)}, \dots, b_{2k-1}^{(i)}) \dots (b_{(\lambda_i-1)k}^{(i)}, \dots, b_{\lambda_i k-1}^{(i)})$$

$$(b_0^{(i)}, \dots, b_{k-1}^{(i)}) = (0, \dots, 0).$$

Then, we can define a one-sided sequence

$$x = b^0 \times_k b^1 \times_k b^2 \times_k \dots \quad (40)$$

The sequences of the form (40) are said to be k -Morse sequences provided that x is almost periodic and strictly transitive. Of course, 1-Morse sequences are precisely those introduced in [8].

Now, we fix our attention on $(k, 2k)$ -substitutions of the form (37). Assume in addition

$$\Theta^{(\alpha_0)} 0 \dots \Theta^{(\alpha_{k-1})} 0 = \underbrace{0 \dots 0}_k b_{k+1} \dots b_{2k-1} = B_1 \dots B_k \quad (41)$$

and put $B = \Theta^{(\alpha_0)} 0 \dots \Theta^{(\alpha_{k-1})} 0$. Then $|B| = 2k$ and the k -Morse sequence

$$B \times_k B \times_k \dots$$

is a fixed point of σ . In particular

$x_1 = 01 \times_1 01 \times_1 \dots$ — classic Thue-Morse sequence.

$x_{11} = 0001 \times_2 0001 \times_2 \dots$ — Rudin-Shapiro sequence (a 2-Morse sequence).

Now, let $x = b^0 \times_k b^1 \times_k \dots$ be a k -Morse sequence. By c_t we mean the product $b^0 \times_k b^1 \times_k \dots \times_k b^t$,

$$|c_t| = kn_t, \quad n_t = \lambda_0 \dots \lambda_t, \quad x_t = b^t \times_k b^{t+1} \times_k \dots$$

THEOREM 5. — *Let $x = b^0 \times_k b^1 \times_k \dots$ be a k -Morse sequence. Then \hat{x} is a regular Toeplitz sequence and moreover the sequence $\{kn_t\}$ is its period structure. The kn_t -skeleton of x has exactly k "holes" for any $t \geq 0$.*

Proof. — Due to the associativity property we can write down

$$x = b^0 \times_k (b^1 \times_k b^2 \times_k \dots) = b^0 \times_k x_1.$$

The sequence x_1 we divide into sequence of blocks of length k , i. e.

$$x_1 = (e_0^{(1)}, \dots, e_{k-1}^{(1)}) (e_k^{(1)}, \dots, e_{2k-1}^{(1)}) \dots$$

Since $b^0 = b_1^{(0)} b_2^{(0)} \dots b_k^{(0)}$, directly from the definition of k -product it follows that x is a juxtaposition of the form

$$b_1^{(0)} \square b_2^{(0)} \square \dots b_k^{(0)} \square b_1^{(0)} \square b_2^{(0)} \square \dots b_k^{(0)} \square \dots$$

where \square is equal to 0 or 1 (i. e. we take the block or take its mirror image). However, $b_j^{(0)} = \hat{b}_j^{(0)}$, so kn_0 -skeleton is of the form

$$\hat{b}_1^{(0)} \infty \hat{b}_2^{(0)} \infty \dots \infty \hat{b}_k^{(0)} \infty | \hat{b}_1^{(0)} \infty \hat{b}_2^{(0)} \infty \dots \infty \hat{b}_k^{(0)} \infty | \dots$$

Now, $x = c_t \times_k x_{t+1}$ and deducing as before we get that kn_t -skeleton of x is of the form

$$\hat{b}_1^{(01 \dots t)} \infty \dots \infty \hat{b}_k^{(01 \dots t)} \infty | \hat{b}_1^{(01 \dots t)} \infty \dots \infty \hat{b}_k^{(01 \dots t)} \infty | \dots$$

which completes the proof.

At present, we turn back again to $(k, 2k)$ -substitutions of the form (41) to give more detailed description of the structure of the base-skeletons. We will assume that k is an even number.

To compute s -base-skeleton of σ we take into considerations

$$\underbrace{\mathbf{B} \times_k \dots \times_k \mathbf{B}}_{s-1} \times_k \mathbf{B} = \mathbf{C}_1^{b_0} \dots \mathbf{C}_k^{b_{k-1}} \mathbf{C}_1^{b_k} \dots \mathbf{C}_k^{b_{2k-1}},$$

where

$$\underbrace{\mathbf{B} \times_k \dots \times_k \mathbf{B}}_{s-1} = \mathbf{C}_1 \dots \mathbf{C}_k, \quad |C_i| = |C_j|, \quad i, j = 1, \dots, k.$$

But k is an even number so

$$\mathbf{B} \times_k \dots \times_k \mathbf{B} \times_k \mathbf{B} = (\mathbf{C}_1^{b_0} \mathbf{C}_2^{b_1}) \times \dots \times (\mathbf{C}_{k-1}^{b_{k-2}} \mathbf{C}_k^{b_{k-1}}) (\mathbf{C}_1^{b_k} \mathbf{C}_2^{b_{k+1}}) \dots (\mathbf{C}_{k-1}^{b_{2k-2}} \mathbf{C}_k^{b_{2k-1}}).$$

Thus, from the proof of Theorem 5 we have obtained that s -base skeleton is equal to

$$\begin{aligned} & (\mathbf{C}_1 [|C_1| - 1] + \mathbf{C}_2 [0] + b_0 + b_1) \\ & \quad \infty (\mathbf{C}_3 [|C_3| - 1] + \mathbf{C}_4 [0] + b_2 + b_3) \infty \dots \\ & \quad \infty (\mathbf{C}_{k-1} [|C_{k-1}| - 1] + \mathbf{C}_k [0] + b_{k-2} + b_{k-1}) \\ & \quad \quad \infty (\mathbf{C}_1 [|C_1| - 1] + \mathbf{C}_2 [0] + b_k + b_{k+1}) \\ & \quad \infty (\mathbf{C}_3 [|C_3| - 1] + \mathbf{C}_4 [0] + b_{k+2} + b_{k+3}) \infty \dots \\ & \quad \infty (\mathbf{C}_{k-1} [|C_{k-1}| - 1] + \mathbf{C}_k [0] + b_{2k-2} + b_{2k-1}) \end{aligned} \tag{42}$$

Combining (41) and (42) we obtain that s -base-skeleton is of the form

$$\begin{aligned} & u_1^{(s)} \infty u_2^{(s)} \infty \dots \infty u_{k/2}^{(s)} \infty u_1^{(s)} + b_k + b_{k+1} \\ & \quad \quad \quad \infty u_2^{(s)} + b_{k+2} + b_{k+3} \infty \dots \infty u_{k/2}^{(s)} + b_{2k-2} + b_{2k-1} \infty \end{aligned}$$

Due to III.2 we have proved the following

THEOREM 6. — For any substitution (41) of the form

$$\sigma: \underbrace{0 \dots 0}_{2^s} \mapsto 0000 \dots 00 (i_1, 1 + i_1) (i_2, 1 + i_2) \dots (i_{2^{s-1}}, 1 + i_{2^{s-1}})$$

the dynamical system arising from σ has Lebesgue spectrum in the orthocomplement of the eigenfunctions space with spectral multiplicity equal to 2^s . In particular x_{11}, x_{10} have Lebesgue spectrum with multiplicity 2, $x_{1 \infty 1}, x_{1 \infty 0}$ have Lebesgue spectrum with multiplicity 4 and in general $x_{1 \infty \dots \infty 1}, x_{1 \infty \dots \infty 0}$ have Lebesgue spectrum with multiplicity $2^{|u|-1}$.

So far, we have not mentioned whether any sequence defined by (40) is a k -Morse sequence or not (i. e. whether it is almost periodic and strictly transitive). We will deal with this problem to get a new class of strictly ergodic dynamical systems as well as to get a class of ergodic Toeplitz Z_2 -extensions.

THEOREM 7. — *Let $x = b^0 \times_k b^1 \times_k \dots$ be a sequence such that*

$$x_t = b^t \times_k b^{t+1} \times_k \dots = \mathbf{B}_1^{(t)} \mathbf{B}_2^{(t)} \dots,$$

where $|\mathbf{B}_i^{(t)}| = k$, has the property

$$\text{for any } C \in \{0, 1\}^k \text{ there is } j \text{ such that } C = \mathbf{B}_j^{(t)}. \quad (43)$$

Then x is almost periodic.

Proof. — Let B be any block. Then, there is t such that $B \subset c_t$ (i. e. B is a subblock of c_t). Hence, it is sufficient to prove that c_t appears on x with a bounded frequency.

We have the operations

$$c_t^{i_1 \dots i_k}, \quad i_s \in \{0, 1\}, \quad s = 1, \dots, k,$$

where $c_t = \mathbf{B}_1^{(t)} \dots \mathbf{B}_k^{(t)}$ and

$$c_t^{i_1 \dots i_k} = (\mathbf{B}_1^{(t)})^{i_1} \dots (\mathbf{B}_k^{(t)})^{i_k}. \quad (44)$$

Given i_1, \dots, i_k we seek such s_{i_1, \dots, i_k} that

$$c_t^{i_1 \dots i_k} \subset c_{s_{i_1, \dots, i_k}}$$

[the block $c_t^{i_1, \dots, i_k}$ appears on x in view of (43)]. Next, we select a common s to have

$$c_t^{i_1 \dots i_k} \subset c_s \quad \text{for any } i_1, \dots, i_k.$$

It is possible by the simple fact $c_T \subset c_{T'}$ for $T \leq T'$. Further, we seek T with the property

$$c_T = \mathbf{B}_1^{(T)} \dots \mathbf{B}_k^{(T)}, \quad |\mathbf{B}_i^{(T)}| > |c_s|.$$

This simply means that on $\mathbf{B}_i^{(T)}$ all the blocks (44) appear. Moreover $x = c_T \times_k x_{T+1}$, so x is a juxtaposition

$$\mathbf{B}_1^{(T)} \square \dots \mathbf{B}_k^{(T)} \square \mathbf{B}_1^{(T)} \square \dots \mathbf{B}_k^{(T)} \square \dots$$

So, to get our claim it is enough to notice that

$$c_i^{i_1 \dots i_k} \in B_1^{(T)} \quad \text{iff} \quad c_i^{i_1 \dots i_k} \in \tilde{B}_1^{(T)}$$

[which is an easy consequence of the fact that on $B_1^{(T)}$ all the blocks (44) appear].

Now, let $x = b^0 \times_k b^1 \times_k \dots$ be a sequence and $B = (b_0, \dots, b_{sk-1})$ be any k -block.

For $j=0, 1$ we put

$$\begin{aligned} \text{fr}_i(j, B) &= \text{fr}(j : (b_i, b_{k+i}, \dots, b_{(s-1)k+i})) \\ & \quad i=0, 1, \dots, k-1. \end{aligned} \tag{45}$$

We define some condition (★)

$$(\forall t \geq 0) \text{fr}_i(0, (b^t \times_k \dots \times_k b^{t+s})_j) - \text{fr}_i(1, (b^t \times_k \dots \times_k b^{t+s})_j) \rightarrow 0, \quad (\star)$$

$$i=0, \dots, k-1$$

and

$$\begin{aligned} b^t \times_k \dots \times_k b^{t+s} &= (b^t \times_k \dots \times_k b^{t+s})_1 \dots (b^t \times_k \dots \times_k b^{t+s})_{k'}, \\ & \quad j=1, \dots, k. \end{aligned}$$

LEMMA 2. — If (★) holds and $k^2 || b^t |$, $t \geq 0$, then x is strictly transitive.

Proof. — For a purpose of this proof we employ the symbol $\text{fr}(B, C)$ to indicate the average frequency $\frac{1}{|C|} \text{fr}(B, C)$ and also we employ $\text{fr}_i(j, B)$

to indicate $\frac{1}{|B|} \text{fr}_i(j, B)$.

A. Suppose, first, that we consider $\text{fr}(d, b \times_k c)$ but we assume $|d| \ll |b|$. If $b = B_1 \dots B_k$, then we can write

$$\begin{aligned} \text{fr}(d, b \times_k c) &\approx \text{fr}(d, B_1) \text{fr}_0(0, c) + \text{fr}(\vec{d}, B_1) \text{fr}_0(1, c) \\ & \quad + \text{fr}(d, B_2) \text{fr}_1(0, c) + \text{fr}(\vec{d}, B_2) \text{fr}_1(1, c) \\ & \quad + \dots \\ & \quad + \text{fr}(d, B_k) \text{fr}_{k-1}(0, c) + \text{fr}(\vec{d}, B_k) \text{fr}_{k-1}(1, c). \end{aligned}$$

since the average frequencies obtained by appearances of d on blocks of the form $B_i^\square B_{i+1}^\square$ or $B_k^\square B_1^\square$ are not essential.

Similarly

$$\begin{aligned} \text{fr}(\tilde{d}, b \times_k c) &\approx \text{fr}(d, \mathbf{B}_1) \text{fr}_0(1, c) + \text{fr}(\tilde{d}, \mathbf{B}_1) \text{fr}_0(0, c) \\ &\quad + \dots \\ &\quad + \text{fr}(d, \mathbf{B}_k) \text{fr}_{k-1}(1, c) + \text{fr}(\tilde{d}, \mathbf{B}_k) \text{fr}_{k-1}(0, c). \end{aligned}$$

This implies

$$\begin{aligned} |\text{fr}(d, b \times_k c) - \text{fr}(\tilde{d}, b \times_k c)| &\approx |\text{fr}(d, \mathbf{B}_1) - \text{fr}(\tilde{d}, \mathbf{B}_1)| \cdot |\text{fr}_0(0, c) - \text{fr}_0(1, c)| \\ &\quad + |\text{fr}(d, \mathbf{B}_2) - \text{fr}(\tilde{d}, \mathbf{B}_2)| |\text{fr}_1(0, c) - \text{fr}_1(1, c)| \\ &\quad + \dots \\ &\quad + |\text{fr}(d, \mathbf{B}_k) - \text{fr}(\tilde{d}, \mathbf{B}_k)| |\text{fr}_{k-1}(0, c) - \text{fr}_{k-1}(1, c)| \\ &= \sum_{i=0}^{k-1} |\text{fr}(d, \mathbf{B}_{i+1}) - \text{fr}(\tilde{d}, \mathbf{B}_{i+1})| |\text{fr}_i(0, c) - \text{fr}_i(1, c)|. \quad (46) \end{aligned}$$

Now, suppose that

$$\begin{aligned} b \times_k c &= \mathbf{D}_1 \dots \mathbf{D}_k, \quad |c| \gg |b|, \quad k^2 ||c|, \\ b &= \mathbf{B}_1 \dots \mathbf{B}_k, \\ c &= (c_0, \dots, c_{k-1})(c_k, \dots, c_{2k-1}) \dots (c_{(r-1)k}, \dots, c_{rk-1}). \end{aligned}$$

Thus

$$b \times_k c = \underbrace{\mathbf{B}_1^{c_0} \dots \mathbf{B}_k^{c_{k-1}} \mathbf{B}_1^{c_k} \dots \mathbf{B}_k^{c_{2k-1}} \dots \mathbf{B}_1^{c_{(r-1)k}} \dots \mathbf{B}_k^{c_{rk-1}}}_{\mathbf{D}_1} \dots \underbrace{\mathbf{B}_1^{c_{(r-1)k}} \dots \mathbf{B}_k^{c_{rk-1}}}_{\mathbf{D}_k}$$

We have

$$\begin{aligned} \text{fr}(d, \mathbf{D}_1) &\approx \text{fr}(d, \mathbf{B}_1) \text{fr}_0(0, c_0 \dots c_{r-1}) + \text{fr}(\tilde{d}, \mathbf{B}_1) \text{fr}_0(1, c_0 \dots c_{r-1}) \\ &\quad + \text{fr}(d, \mathbf{B}_2) \text{fr}_1(0, c_0 \dots c_{r-1}) + \text{fr}(\tilde{d}, \mathbf{B}_2) \text{fr}_1(1, c_0 \dots c_{r-1}) \\ &\quad + \dots \\ &\quad + \text{fr}(d, \mathbf{B}_k) \text{fr}_{k-1}(0, c_0 \dots c_{r-1}) + \text{fr}(\tilde{d}, \mathbf{B}_k) \text{fr}_{k-1}(1, c_0 \dots c_{r-1}) \end{aligned}$$

and generally

$$\begin{aligned} \text{fr}(d, \mathbf{D}_i) &\approx \text{fr}(d, \mathbf{B}_1) \text{fr}_0(0, c_{(i-1)r} \dots c_{ir-1}) \\ &\quad + \text{fr}(\tilde{d}, \mathbf{B}_1) \text{fr}_0(1, c_{(i-1)r} \dots c_{ir-1}) \\ &\quad + \dots \end{aligned}$$

$$\begin{aligned}
 &+ \text{fr}(d, \mathbf{B}_k) \text{fr}_{k-1}(0, c_{(i-1)r} \dots c_{ir-1}) \\
 &+ \text{fr}(\tilde{d}, \mathbf{B}_k) \text{fr}_{k-1}(1, c_{(i-1)r} \dots c_{ir-1}) \quad (47)
 \end{aligned}$$

Establishing the analogous formula for $\text{fr}(\tilde{d}, D_i)$ we get

$$\begin{aligned}
 |\text{fr}(d, D_i) - \text{fr}(\tilde{d}, D_i)| &\approx \sum_{j=1}^k |\text{fr}(d, \mathbf{B}_j) - \text{fr}(\tilde{d}, \mathbf{B}_j)| \\
 &\times |\text{fr}_{j-1}(0, c_{(i-1)r} \dots c_{ir-1}) - \text{fr}_{j-1}(1, c_{(i-1)r} \dots c_{ir-1})| \quad (48)
 \end{aligned}$$

B. $\lim_{\mathbf{T}} (\text{fr}(d, c_{\mathbf{T}}) - \text{fr}(\tilde{d}, c_{\mathbf{T}})) = 0$ and also

$$\begin{aligned}
 \lim_{\mathbf{T}} (\text{fr}(d, \mathbf{B}_i^{(\mathbf{T})}) - \text{fr}(\tilde{d}, \mathbf{B}_i^{(\mathbf{T})})) &= 0, \quad c_{\mathbf{T}} = \mathbf{B}_1^{(\mathbf{T})} \dots \mathbf{B}_k^{(\mathbf{T})}, \\
 &i = 1, \dots, k.
 \end{aligned}$$

It is enough to prove the latter statement.

From (47) we have the following

$$\begin{aligned}
 &|\text{fr}(d, \mathbf{B}_i^{(\mathbf{T})}) - \text{fr}(\tilde{d}, \mathbf{B}_i^{(\mathbf{T})})| - \sum_{j=1}^k |\text{fr}(d, \mathbf{B}_j^{(t)}) - \text{fr}(\tilde{d}, \mathbf{B}_j^{(t)})| \\
 &|\text{fr}_{j-1}(0, (b^t \times_k \dots \times_k b^{\mathbf{T}})_i) - \text{fr}_{j-1}(1, (b^t \times_k \dots \times_k b^{\mathbf{T}})_i)| \Big| \rightarrow 0_{\mathbf{T}}
 \end{aligned}$$

From (★) we obtain

$$\begin{aligned}
 &\sum_{j=1}^k |\text{fr}(d, \mathbf{B}_j^{(t)}) - \text{fr}(\tilde{d}, \mathbf{B}_j^{(t)})| \\
 &\times |\text{fr}_{j-1}(0, (b^t \times_k \dots \times_k b^{\mathbf{T}})_i) - \text{fr}_{j-1}(1, (b^t \times_k \dots \times_k b^{\mathbf{T}})_i)| \Big| \rightarrow 0_{\mathbf{T}}
 \end{aligned}$$

which implies

$$|\text{fr}(d, \mathbf{B}_i^{(\mathbf{T})}) - \text{fr}(\tilde{d}, \mathbf{B}_i^{(\mathbf{T})})| \rightarrow 0_{\mathbf{T}}$$

C. $\lim_{\mathbf{T}} \text{fr}(d, c_{\mathbf{T}})$ exists and also $\lim_{\mathbf{T}} \text{fr}(d, \mathbf{B}_i^{(\mathbf{T})})$ exists for $i = 1, \dots, k$.

First, take t_0 such that

$$|\text{fr}(d, \mathbf{B}_j^{(t_0)}) - \text{fr}(\tilde{d}, \mathbf{B}_j^{(t_0)})| < \varepsilon, \quad j = 1, \dots, k.$$

Then, applying (47)

$$\left| \text{fr}(d, \mathbf{B}_i^{(T)}) - \sum_{j=1}^k [\text{fr}(d, \mathbf{B}_j^{(t_0)}) \text{fr}_{j-1}(0, (b^{t_0} \times_k \dots \times_k b^T)_i) + \text{fr}(\tilde{d}, \mathbf{B}_j^{(t_0)}) \text{fr}_{j-1}(1, (b^{t_0} \times_k \dots \times_k b^T)_i)] \right| < \varepsilon$$

for $T \geq T_0$.

But

$$\begin{aligned} & \sum_{j=1}^k [\text{fr}(d, \mathbf{B}_j^{(t_0)}) \text{fr}_{j-1}(\dots) + \text{fr}(\tilde{d}, \mathbf{B}_j^{(t_0)}) \text{fr}_{j-1}(\dots)] \\ &= \sum_{j=1}^k [\text{fr}(d, \mathbf{B}_j^{(t_0)}) \text{fr}_{j-1}(\dots) + \text{fr}(\tilde{d}, \mathbf{B}_j^{(t_0)}) (1 - \text{fr}_{j-1}(0, \dots))] \\ &= \sum_{j=1}^k \text{fr}_{j-1}(0, (b^{t_0} \times_k \dots \times_k b^T)_i) |\text{fr}(d, \mathbf{B}_j^{(t_0)}) - \text{fr}(\tilde{d}, \mathbf{B}_j^{(t_0)})| \\ & \qquad \qquad \qquad + \sum_{j=1}^k \text{fr}(\tilde{d}, \mathbf{B}_j^{(t_0)}). \end{aligned}$$

The first sum is estimated by $k \varepsilon$, so

$$\begin{aligned} \left| \text{fr}(d, \mathbf{B}_i^{(T)}) - \sum_{j=1}^k \text{fr}(d, \mathbf{B}_j^{(t_0)}) \right| &< 100 k^2 \varepsilon \\ &\qquad \qquad \qquad \approx \\ & \sum_{j=1}^k \text{fr}(\tilde{d}, \mathbf{B}_j^{(t_0)}) \end{aligned}$$

so

$$|\text{fr}(d, \mathbf{B}_i^{(T)}) - \text{fr}(d, \mathbf{B}_i^{(T')})| < 200 k^2 \varepsilon$$

whenever $T, T' > T_0$. In other words $\{\text{fr}(d, \mathbf{B}_i^{(T)})\}$ is a Cauchy sequence. $D.x$ is strictly transitive.

If we take a long sector $x[k, n]$, then it is a juxtaposition $c_T^{i_1, \dots, i_k}$ (without some ends) and also it is a juxtaposition

$$\mathbf{B}_1^{(T) \square} \mathbf{B}_2^{(T) \square} \dots \mathbf{B}_k^{(T) \square} \dots$$

On any $\mathbf{B}_j^{(T)}$ ($\tilde{\mathbf{B}}_j^{(T)}$) there are frequencies of all blocks.

Remark. — The proof of Lemma 2 stems from the proof of Lemma 3 in [8].

where $c^{(p)} = c_{(p-1)r} \cdots c_{pr-1}$.

Similarly

$$\frac{1}{tr} \text{fr}_j(1, (b \times_k c)_p) = \sum_{s=0}^{k-1} \left[\frac{1}{t} \text{fr}_j(0, \mathbf{B}_{s+1}) \frac{1}{r} \text{fr}_s(1, c^{(p)}) + \frac{1}{t} \text{fr}_j(1, \mathbf{B}_{s+1}) \frac{1}{r} \text{fr}_s(0, c^{(p)}) \right] \quad (51)$$

Combining (50) and (51) we get

$$\begin{aligned} \left| \frac{1}{tr} \text{fr}_j(0, (b \times_k c)_p) - \frac{1}{tr} \text{fr}_j(1, (b \times_k c)_p) \right| &= \sum_{s=0}^{k-1} \left| \frac{1}{t} \text{fr}_j(0, \mathbf{B}_{s+1}) - \frac{1}{t} \text{fr}_j(1, \mathbf{B}_{s+1}) \right| \\ &\times \left| \frac{1}{r} \text{fr}_s(0, c^{(p)}) - \frac{1}{r} \text{fr}_s(1, c^{(p)}) \right| \leq \sum_{s=0}^{k-1} \left| \frac{1}{t} \text{fr}_j(0, \mathbf{B}_{s+1}) - \frac{1}{t} \text{fr}_j(1, \mathbf{B}_{s+1}) \right| \\ &\times \left| \frac{1}{r} \text{fr}_s(0, c^{(p)}) - \frac{1}{r} \text{fr}_s(1, c^{(p)}) \right| \end{aligned}$$

By an easy induction

$$\begin{aligned} &\left| \frac{1}{n_{t-1}} \text{fr}_j(0, (b^0 \times_k b^1 \times_k \dots \times_k b^{t-1})_p) \right. \\ &\quad \left. - \frac{1}{n_{t-1}} \text{fr}_j(1, (b^0 \times_k b^1 \times_k \dots \times_k b^{t-1})_p) \right| \\ &\leq \sum_{s_1=0}^{k-1} \sum_{s_2=0}^{k-1} \dots \sum_{s_{t-1}=0}^{k-1} \left| \frac{1}{\lambda_{t-1}} \text{fr}_{s_{t-1}}(0, b^{t-1(p)}) \right. \\ &\quad \left. - \frac{1}{\lambda_{t-1}} \text{fr}_{s_{t-1}}(1, b^{t-1(p)}) \right| \\ &\times \left| \frac{1}{\lambda_{t-2}} \text{fr}_{s_{t-2}}(0, b^{t-2(p)}) - \frac{1}{\lambda_{t-2}} \text{fr}_{s_{t-2}}(1, b^{t-2(p)}) \right| \\ &\quad \times \dots \\ &\quad \times \left| \frac{1}{\lambda_1} \text{fr}_{s_1}(0, b^1(p)) - \frac{1}{\lambda_1} \text{fr}_{s_1}(1, b^1(p)) \right| \\ &\quad \times \left| \frac{1}{\lambda_0} \text{fr}_j(0, b^{0(s_1+1)}) - \frac{1}{\lambda_0} \text{fr}_j(1, b^{0(s_1+1)}) \right| \end{aligned}$$

$$\begin{aligned} &\leq \sum_{s_{t-1}=0}^{k-1} \left| \frac{1}{\lambda_{t-1}} \text{fr}_{s_{t-1}}(0, b^{t-1(p)}) - \frac{1}{\lambda_{t-1}} \text{fr}_{s_{t-1}}(1, b^{t-1(p)}) \right| \\ &\quad \times \sum_{s_{t-2}=0}^{k-1} \left| \frac{1}{\lambda_{t-2}} \text{fr}_{s_{t-2}}(0, b^{t-2(p)}) \right. \\ &\quad \quad \left. - \frac{1}{\lambda_{t-2}} \text{fr}_{s_{t-2}}(1, b^{t-2(p)}) \right| \dots \\ &\quad \times \sum_{s_1=0}^{k-1} \left| \frac{1}{\lambda_0} \text{fr}_j(0, b^{0(s_1+1)}) - \frac{1}{\lambda_0} \text{fr}_j(1, b^{0(s_1+1)}) \right| \\ &\quad \quad \quad \left| \frac{1}{\lambda_1} \text{fr}_{s_1}(0, b^1(p)) - \frac{1}{\lambda_1} \text{fr}_{s_1}(1, b^1(p)) \right| \end{aligned}$$

It is well-known [8] that if the series (49) is divergent then the infinite product obtained converges to zero. Therefore (★) is satisfied (this reasoning does not require us to start with b^0).

We can put Lemma 2 and 3 together to obtain

THEOREM 8. — *Let $x = b^0 \times_k b^1 \times_k \dots$ and $|b^i| = k \lambda_i$, $k | \lambda_i$, $i \geq 0$ and*

$$\sum_{t=0}^{\infty} \left(\sum_{i=0}^{k-1} \min \left(\frac{1}{\lambda_t} \text{fr}_i(0, b^{t(i+1)}), \frac{1}{\lambda_t} \text{fr}_i(1, b^{t(i+1)}) \right) \right) = \infty.$$

then x is strictly transitive.

IV. ON METRIC THEORY OF k -MORSE SEQUENCES

In this section we develop metric theory of k -Morse sequences. Our main aim is to show how to distinguish between some k -Morse sequences.

Let $x = b^0 \times_k b^1 \times_k \dots$ be a k -Morse sequence,

$$\begin{aligned} c_t &= b^0 \times_k \dots \times_k b^t = B_1^{(t)} \dots B_k^{(t)}, \\ |b^i| &= k \lambda_i, \quad |c_t| = kn_t = k \lambda_0 \dots \lambda_t. \end{aligned}$$

In addition, we assume that for every $t \geq 0$

$$x_t = b^t \times_k b^{t+1} \times_k \dots = (a_{11}^{(t)} \dots a_{1k}^{(t)}) (a_{21}^{(t)} \dots a_{2k}^{(t)}) \dots$$

has the property:

$$\text{for every } B \in \{0, 1\}^k \text{ there is } j \text{ such that } B = (a_{j1}^{(t)} \dots a_{jk}^{(t)}). \quad (52)$$

Thus, x is a juxtaposition of the blocks

$$c_t^{i_1 \dots i_k} = (B_1^{(t)})^{i_1} (B_2^{(t)})^{i_2} \dots (B_k^{(t)})^{i_k}, \quad (53)$$

$$i_s \in \{0, 1\}.$$

Any block $c_t^{i_1 \dots i_k}$ will be said to be a t -symbol. Any $B_j^{(t)}$ (and its mirror image) we will call a sub- t -symbol, $j = 1, \dots, k$.

We treat x as two-sided sequence (see [8]) and by $X(x)$ we mean the closure of trajectory of x via the shift $\tau: \{0, 1\}^{\mathbb{Z}} \ni$, i. e.

$$X(x) = \overline{\{\tau^i x : i \in \mathbb{Z}\}}.$$

We have got then a strictly ergodic dynamical system $(X(x), \tau, \mu_x)$ called a k -Morse dynamical system.

LEMMA 4 [the structure of $X(x)$]. — Let $y \in X(x)$. Then for any $t \geq 0$ there is a unique $j_t, 0 \leq j_t \leq n_t - 1$ such that


$$y[-j_t + rn_t - j_t + (r+1)n_t - 1] = (B_{k_r}^{(t)})^r, \quad (54)$$

$$r = 0, \pm 1, \dots$$

Moreover, for a. e. $y \in X(x)$

$$j_t \nearrow_{t} \infty, \quad k_t = n_t - j_t \nearrow_{t} \infty. \quad (5)$$

Proof. — It is sufficient to prove (54) for x (see Picture 7).



Picture 7

The proof of (54) is now obvious from Picture 7 because we have assumed (52).

On the other hand any $y \in \{0, 1\}^Z$ satisfying (54) and (55) must belong to $X(x)$. So, by denoting

$$C_{j;0}^{n_t}(r) = \{y \in X(x) : y[rn_t, (r+1)n_t - 1] = (B_{k_r}^{(t)})^i, y[0, n_t - 1] = (B_r^{(t)})^j\}$$

$$j=0, 1, \quad r=1, \dots, k$$

we see that $C_{j;0}^{n_t}(r)$ may be considered to be the base of an n_t - τ -stack. In this way we obtain $2k$ of n_t - τ -stacks (mutually disjoint) and moreover the corresponding sequence of partitions will tend to the point partition as soon as we prove (55).

To prove (55) we put

$$D_0^{n_t} = C_{0;0}^{n_t}(1) \cup C_{1;0}^{n_t}(1) \cup \dots \cup C_{0;0}^{n_t}(k) \cup C_{1;0}^{n_t}(k).$$

Then

$$D_0^{n_t}, \tau D_0^{n_t}, \dots, \tau^{n_t-1} D_0^{n_t} \text{ are pairwise disjoint and } \bigcup_{i=0}^{n_t-1} D_i^{n_t} = X(x). \quad (57)$$

Since there is a homomorphism h from $(X(x), \tau)$ to $(X(\hat{x}), \tau)$ (see Section I) and any n_t - τ -stack satisfying (57) is unique (due to the ergodicity property) $h^{-1} D_i^{n_t} = \bar{D}^i$, where \bar{D}^i is the n_t -stack in $X(\hat{x})$ described in (5). The claim (55) is true for regular Toeplitz sequences ([5], [18]), so also for x .

Remark. — Another proof of Lemma 4 can be done on the basis of Lemma 1.

Remark. — In fact we have proved that the rank of a k -Morse sequence is bounded by $2k$. It is well-known that in that case spectral multiplicity is bounded by $2k$, too. But Theorem 4 strengthens this estimation. The spectral multiplicity of any k -Morse sequence is bounded by k (Theorem 5).

Lemma 4 says that a k -Morse sequence $x = b^0 \times_k b^1 \times_k \dots$ has a unique decomposition into a juxtaposition of sub- t -symbols. However, to develop metric theory of k -Morse sequences we will require x to have a unique decomposition into a juxtaposition of t -symbols, $t \geq 0$.

We start with a very simple observation.

LEMMA 5. — *Let $x = b^0 \times_k b^1 \times_k \dots$ be a k -Morse sequence. Then*

$$d(c_i^{i_1 \dots i_k}, c_i^{j_1 \dots j_k}) = \begin{cases} 0 & \text{if } (i_1 \dots i_k) = (j_1 \dots j_k) \\ \geq 1/k & \text{otherwise} \end{cases} \quad (58)$$

Let $x = b^0 \times_k b^1 \times_k \dots$ be a k -Morse sequence such that for each t , if

$$x[j + skn_t, j + (s+1)kn_t - 1] = c_t^{i_t^{(s)}} \cdots i_k^{(s)} \quad \text{for } s=0, 1, \dots,$$

then $kn_t \mid j$.

Hence, it is clear that for every $y \in X(x)$ and every t there is a unique j_t , $0 \leq j_t \leq kn_t - 1$ such that

$$y[-j_t + skn_t, -j_t + (s+1)kn_t - 1] = c_t^{i_t^{(s)}} \cdots i_k^{(s)}, \\ s=0, \pm 1, \dots$$

In this case one can define $2^k kn_t$ -stacks with the bases

$$E_{\mathcal{I}}^{i_1 \cdots i_k, t} = \{y \in X(x) : y[skn_t, (s+1)kn_t - 1] \\ = c_t^{i_t^{(s)}} \cdots i_k^{(s)} \text{ and } y[0, kn_t - 1] = c_t^{i_1} \cdots i_k\}. \quad (59)$$

$(i_1, \dots, i_k) \in \{0, 1\}^k$ and the corresponding sequence of partitions tends to the unit partition.

In others words we will assume that

for every t there is L_t such that if a

$$\text{string } \eta_1^{(t)} \dots \eta_{L_t}^{(t)} \text{ appears on } x \text{ at a place } j \\ \text{and } \eta_i^{(t)} \text{ are } t\text{-symbols, then } kn_t \mid j. \quad (60)$$

An extremely interesting case is when the sequence $\{L_t\}$ is bounded, i. e.

there is $L = L(x)$ such that for every t

$$\text{if a string } \eta_1^{(t)} \dots \eta_L^{(t)} \text{ appears on } x \\ \text{at a place } j \text{ and } \eta_i^{(t)} \text{ are } t\text{-symbols, then } kn_t \mid j. \quad (61)$$

The next lemma is an essential strengthener of the property (61) for the number $L(x)$.

LEMMA 6. — *If $x = b^0 \times_k b^1 \times_k \dots$ is a k -Morse sequence such that $k \mid \lambda_t$, $\sup k \lambda_t = k \lambda < \infty$ and $L = L(x)$ satisfies (61), then*

$$(\exists \delta > 0) \quad (\forall t \geq 0) \quad [\text{if } \eta = \eta_1^{(t)} \dots \eta_{L(x)}^{(t)}, \\ \eta_i^{(t)} \text{ is a } t\text{-symbol and } d(\eta, x[j, j + |\eta| - 1]) < \delta \\ \text{then } d(\eta, x[j, j + |\eta| - 1]) = 0 \text{ and } kn_t \mid j].$$

Proof. — The proof goes by induction.

We put $\delta = 1/k \lambda L$.

$1^0 t = 0$. If $\eta = \eta_1^{(0)} \dots \eta_L^{(0)}$ appears on x at a place j then $|\eta| = L k \lambda_0 \leq k \lambda L$, so if $d(\eta, x[j, j + L k \lambda_0 - 1]) < \delta$, then it is equal to 0 and from (61) $k \lambda_0 |j$.

2^0 Let

$$\eta^{(t+1)} = \eta_1^{(t+1)} \dots \eta_L^{(t+1)} \quad \text{and} \quad d(\eta^{(t+1)}, x[j, j + L k n_{t+1} - 1]) < \delta.$$

Since $\eta_i^{(t+1)}$ is a $(t+1)$ -symbol,

$$\eta_i^{(t+1)} = \eta_{i,1}^{(t)} \dots \eta_{i,\lambda_{t+1}}^{(t)} \quad \text{and} \quad \eta_{i,k}^{(t)} \text{ are } t\text{-symbols.} \quad (62)$$

Indeed, due to the condition $k | \lambda_{t+1}$ we have

$$\begin{aligned} c_{t+1} &= c_t \times_k b^{t+1} = (B_1^{(t)})^{b^{t+1}[0]} (B_2^{(t)})^{b^{t+1}[1]} \dots \\ &\quad (B_k^{(t)})^{b^{t+1}[k-1]} (B_1^{(t)})^{b^{t+1}[k]} = \dots \\ &= c_t^{b^{t+1}[0], \dots, b^{t+1}[k-1]} c_t^{b^{t+1}[k], \dots, b^{t+1}[2k-1]} = \dots \\ &= \underbrace{c_t^{b^{t+1}[0], \dots, b^{t+1}[k-1]} \dots c_t^{b^{t+1}[s_1 k], \dots, c_t^{b^{t+1}[(s_1+1)k-1]}}}_{B_1^{(t+1)}} \dots \underbrace{\dots}_{B_k^{(t+1)}} \end{aligned}$$

Therefore $c_{t+1}^{i_1 \dots i_k}$ is a juxtaposition of t -symbols since any $B_i^{(t+1)}$ was a juxtaposition of t -symbols.

Since $|B_i^{(t+1)}| = n_{t+1}$ and $|c_t^{j_1 \dots j_k}| = k n_t$, the number of t -symbols in $c_{t+1}^{i_1 \dots i_k}$ is equal to $(n_{t+1}/k n_t) k = \lambda_{t+1}$.

We write down $\eta^{(t+1)}$ as follows

$$\eta^{(t+1)} = \underbrace{\eta_{1,1}^{(t)} \dots \eta_{1,\lambda_{t+1}}^{(t)} \eta_{2,1}^{(t)} \dots \eta_{2,\lambda_{t+1}}^{(t)} \dots \eta_{L,1}^{(t)} \dots \eta_{L,\lambda_{t+1}}^{(t)}}_L$$

i.e. as λ_{t+1} groups of L consecutive t -symbols. Then, there must exist such a group that it appears on x within δ . Applying the induction hypothesis we come that

$$k n_t | j. \quad (63)$$

In view of this fact, in the representation

$$d(\eta^{(t+1)}, x[j, j + L k n_{t+1} - 1]) < \delta$$

in opposite of t -symbols from $\eta^{(t+1)}$ there are t -symbols from x . So, if the distance (64) is positive, then by virtue of Lemma 5

$$d(\eta^{(t+1)}, x[j, j + Lkn_{t+1} - 1]) \geq \frac{1}{k} |c_t / L | c_{t+1} | = 1/L k \lambda_{t+1} \geq \delta$$

and we get a contradiction. In order to obtain $kn_{t+1} | j$ it is enough to apply (61).

Remark. — The transparent example of a 2-Morse sequence satisfying (61) is Rudin-Shapiro sequence $0001 \times_2 0001 \times_2 \dots$. Therefore Lemma 6 is true for it. The significance of the lemma was presented (in case of substitutions) in [13]. In particular by the methods developed in [13] we are able to prove the following property of Rudin-Shapiro dynamical system:

Let (X, T, ν) be any ergodic automorphism of a Lebesgue space. Suppose Rudin-Shapiro dynamical system to be a factor of T and let U be another factor of T . If T is of local rank one and $\tau \times U$ is ergodic, then U is a rotation on a finite group.

Now, we will examine some class \mathcal{R} of 2-Morse sequences including Rudin-Shapiro sequence.

We put

$$x = b^0 \times_2 b^1 \times_2 \dots \in \mathcal{R} \quad \text{iff } b^i = 0001 \text{ or } 0010. \quad (65)$$

Arguing as in II.4 (proof of Theorem 6) we see that any such x determines a 2-Morse shift with twofold Lebesgue spectrum (in the orthocomplement of the eigenfunctions space).

We intend to prove

THEOREM 9. — *If $x = b^0 \times_2 b^1 \times_2 \dots, y = \beta^0 \times_2 \beta^1 \times_2 \dots \in \mathcal{R}$ and x and y are measure-theoretically isomorphic, then*

$$b^t = \beta^t \quad (66)$$

for t large enough.

As a simple consequence of this theorem we get a continuum pairwise nonisomorphic dynamical systems with twofold nonsingular spectra.

Before the proof of Theorem 9 we will settle down some auxiliary informations.

Let us suppose $\varphi: (X(x), \tau) \rightarrow (X(y), \tau)$ to establish an isomorphism between

$$x = b^0 \times_2 b^1 \times_2 \dots \in \mathcal{R} \quad \text{and} \quad y = \beta^0 \times_2 \beta^1 \times_2 \dots \in \mathcal{R}.$$

Given $\varepsilon > 0$ and using ergodic theorem we can find a finite code $\varphi_\varepsilon: X(x) \rightarrow \{0, 1\}^Z$ (i. e. a measurable map, $\varphi_\varepsilon \tau = \tau \varphi_\varepsilon$), such that

$$d(\varphi_\varepsilon(z), \varphi(z)) = \lim_m d(\varphi_\varepsilon(z)[-m, m], \varphi(z)[-m, m]) < \varepsilon$$

a. e. $z \in X(x)$ (67)

and the length of φ_ε is equal to n ($|\varphi_\varepsilon| = n$), i. e. $\varphi_\varepsilon(z)[0]$ depends only on $z[-n, n]$ and n is the smallest number with this property (for more details see [13]).

Now, we will establish some properties of finite codes.

$$\liminf_m d(\varphi_\varepsilon(z)[-m, m], \varphi_\varepsilon(\tilde{z})[-m, m]) > 1 - 2\varepsilon$$

for a. e. $z \in X(x)$ (68)

Indeed, it follows from (67) and from the simple fact that


$$\varphi(\tilde{z}) = \varphi(z) \quad \text{a. e. } z \in X(x).$$

(for every Z_2 -extension T_θ the map $(x, i) \mapsto (x, i + 1)$ belongs to the second centralizer of T_θ [13]).

Denote by $B_1^{(t)}, \tilde{B}_1^{(t)}, B_2^{(t)}, \tilde{B}_2^{(t)}$ ($E_1^{(t)}, \tilde{E}_1^{(t)}, E_2^{(t)}, \tilde{E}_2^{(t)}$) all sub- t -symbols of $x(y)$ and by

$$D_1^{(t)}, \bar{D}_1^{(t)}, D_2^{(t)}, \bar{D}_2^{(t)} \quad (|D_i^{(t)}| = |B_i^{(t)}| - 2n)$$

their codes via φ_ε (see *Picture 8*)



Picture 8

we will consider $t \in \mathbb{N}$ such that

$$n/n_t < \varepsilon. \tag{69}$$


In view of (68) we may assume

$$\left. \begin{aligned} d(D_i^{(t)}, \bar{D}_i^{(t)}) &> 1 - 3\varepsilon \\ i &= 1, 2, \text{ for } t \text{ large enough.} \end{aligned} \right\} \quad (70)$$

Let us fix t satisfying (69), (70) and denote by s_t the shift of sub- t -symbols of x with respect to sub- t -symbols of y (Picture 9).

We assert that

$$s_t/n_t < 1/8 \quad \text{or} \quad (n_t - s_t)/n_t < 1/8. \quad (71)$$



Picture 9

If (71) were false then from (67), (69) and (70) we would have that there are two functions $f_i: \{0, 1\} \rightarrow \{0, 1\}^2$ such that

$$f_i(1) = \widetilde{f_i(0)}, \quad i = 1, 2, \quad (72)$$

$$d(D_1^{(t)}, (E_1^{(t)})^{f_1(0)[0]}(E_2^{(t)})^{f_1(0)[1]}) < 100\varepsilon, \quad f_i(0) = (f_i(0)[0], f_i(0)[1]), \quad (73)$$

$$d(D_2^{(t)}, (E_2^{(t)})^{f_2(0)[0]}(E_1^{(t)})^{f_2(0)[1]}) < 100\varepsilon, \quad (74)$$

and

$$\left. \begin{aligned} d(D_1^{(t)}, (E_1^{(t)})^{k_1}(E_2^{(t)})^{k_2}) &\geq 1 - 100\varepsilon, \\ d(D_2^{(t)}, (E_2^{(t)})^{l_1}(E_1^{(t)})^{l_2}) &\geq 1 - 100\varepsilon \end{aligned} \right\} \quad (75)$$

for any pairs

$$(k_1, k_2) \neq (f_1(0)[0], f_1(0)[1]), \quad (l_1, l_2) \neq (f_2(0)[0], f_2(0)[1])$$

(for a suitable choice of ε).

For ε as small as we need one can assume there are $i_1^{(k)}, i_2^{(k)}, k = 1, \dots, 8$ such that (Picture 9)

$$f_1(i_1^{(1)}) = (i_1^{\prime(1)}, i_2^{\prime(1)}), \quad f_2(i_2^{(1)}) = i_2^{\prime(1)}, i_1^{\prime(2)}, \dots, f_2(i_2^{(8)}) = (i_2^{\prime(8)}, i_1^{\prime(9)}) \quad (76)$$

Put $f_1(0) = (0, 0)$, $f_2(0) = (0, 1)$ and $i_1^{(0)} = 0$. Then applying (76) we compute that

$$i_1^{(1)} \dots i_1^{(9)} = 001100110. \tag{77}$$


But $i_1^{(1)} \dots i_1^{(9)}$ appears on $x_{t+1} \in \mathcal{A}$ and

$$x_{t+1} = BC \times_2 x_{t+2} = B^\square C^\square B^\square C^\square \dots$$

and $B \neq C, \check{C}$, a contradiction.

The remaining cases are left to the reader. The proof of (71) is now complete.

Remark. — The proof of (71) does not exclude the following situation (Picture 10).




Picture 10

This case reduces to the preceding by passing from t into $t + 1$.


Now, we assert that

$$s_t \text{ is constant for } t \text{ large enough.} \tag{78}$$

An easy proof of it follows from Picture 11 and Picture 12 and the proof of the remark above.



Picture 11



Picture 12

So, we can assume $s = s_t$, $t \geq 0$ and by considering

$$\varphi\tau^{-s} \text{ instead of } \varphi$$

we may assume that the numbers j_t , k_t , $t \geq 0$ defined in (55) are the same for z and $\varphi(z)$, i. e.

$$j_t(z) = j_t(\varphi(z)), \quad k_t(z) = k_t(\varphi(z)). \quad (79)$$


Proof of Theorem 9. — Let t be large enough (i. e. determined by the condition $s/n_t < \varepsilon$).

Due to (71) we can define two functions $g_i: \{0, 1\} \rightarrow \{0, 1\}$, $i = 1, 2$ such that

$$g_i(1) = g_i(0), \quad i = 1, 2 \quad (80)$$

$$\left. \begin{aligned} d(D_1^{(t)}, (E_1^{(t)})^{g_1(0)} E_2^{(t)}) < 100\varepsilon, \\ d(D_2^{(t)}, (E_2^{(t)})^{g_2(0)} E_1^{(t)}) < 100\varepsilon \end{aligned} \right\} \text{ (Picture 13)} \quad (81)$$

$$\left. \begin{aligned} d(D_1^{(t)}, (E_1^{(t)})^{g_1(1)} E_2^{(t)}) &\geq 1 - 100\varepsilon, \\ d(D_2^{(t)}, (E_2^{(t)})^{g_2(1)} E_1^{(t)}) &\geq 1 - 100\varepsilon. \end{aligned} \right\} \quad (82)$$



Picture 13

Like in (76) we may assume

$$i_1^{(1)} = g_1(i_1^{(1)}), \dots, i_8^{(2)} = g_2(i_8^{(2)}), \quad (83)$$


and moreover

$$\{i_1^{(1)}, i_2^{(1)}, \dots, i_8^{(1)}\} = \{0, 1\}. \quad \{i_1^{(2)}, i_2^{(2)}, \dots, i_8^{(2)}\} = \{0, 1\}. \quad (84)$$

we assert that


$$g_1(i) = g_2(i), \quad i = 1, 2. \quad (85)$$

This follows from the following simple observation (*Picture 14*)



Picture 14

i. e. any pair (i, i) implies that $(t + 1)$ -symbol arises by joining the t -symbol considered $[(B_1^{(t)})^i (B_2^{(t)})^i$ on *Picture 14*] with the another from the right. Moreover, any pair $(j, j + 1)$ implies that $(t + 1)$ -symbol arises by joining the t -symbol considered $[(E_1^{(t)})^j (E_2^{(t)})^{j+1}$ on *Picture 14*] with the another from the left (this rule is valid for *any* $x \in \mathcal{O}$). Hence (85) is true because of (79).



Picture 15

Now, suppose that $b^{t+1} = 0001$ and $\beta^{t+1} = 0010$.

Then one of the following situations has to appear (Picture 15). This leads to an obvious contradiction since from the rule above $(t+2)$ -symbols on z are obtained by joining those from the right and on $\varphi(z)$ those from the left,

$$j_{t+2}(z) \neq j_{t+2}(\tau^{-s} \varphi z).$$

If we assume $y=x$, then any $\varphi: (X(x), \tau, \mu_x)$, $\varphi\tau = \tau\varphi$ has the property $\tau^{-s}\varphi$ acts inside of the atoms of the partition η [it follows from (79)], where $A \in \eta$ iff $A = \{z, \tilde{z}\}$, $z \in X(x)$. Then, immediately from the ergodicity of τ on $X(x)$ we obtain.

THEOREM 10. — *Let $x \in \mathcal{R}$ then the centralizer $C(\tau)$ of τ on $X(x)$. i. e.*

$$C(\tau) = \{ \varphi: \varphi: (X(x), \mu_x) \rightarrow \mathcal{R}, \varphi\tau = \tau\varphi \}$$

is of the form

$$C(\tau) = \{ \tau^i \sigma^j: i \in \mathbb{Z}, j = 0, 1 \}$$

where σ is the mirror map.

Two problems. First, observe that all Toeplitz Z_2 -extensions with 2^k -fold Lebesgue spectrum we have considered in III. 2 possess the point spectrum equal to $G\{2^t: t \geq 0\}$. Therefore for any odd number s , τ_s^s is ergodic and has 2^k -fold Lebesgue spectrum (in the orthocomplement of proper functions), $k \geq 1$. Hence for any even number r we get a class of Toeplitz Z_2 -extensions with Lebesgue spectral multiplicity equal to r . The problem is, when are odd numbers the values of Lebesgue spectral multiplicity of Toeplitz Z_2 -extensions (the most interesting case, $r = 1$).

As we have seen in III. 4 (Theorem 5) the spectral multiplicity of any k -Morse sequence is at most k . In what cases is it equal to k ?

REFERENCES

- [1] G. CHRISTOL, T. KAMAE, M. MENDÈS-FRANCE and G. RAUZY, Suites algébriques, automates et substitutions, *Bull. Soc. Math. France*, T. **108**, 1980, pp. 401-419 (in French).
- [2] P. COLLET and J. P. ECKMANN, *Iterated Maps on the Interval as Dynamical Systems*, Birkhäuser, Boston-Basel-Stuttgart, (1980).
- [3] H. HELSON and W. PARRY, Cocycles and Spectra, *Arkiv för Mat.*, Vol. **16**, 1978, pp. 195-206.
- [4] K. JACOBS, Ergodic Theory and Combinatorics, *Contemp. Math.*, Vol. **26**, 1984, pp. 171-187.
- [5] K. JACOBS and M. KEANE, 0-1 Sequences of Toeplitz Type, *Z. Wahr. verw. Geb.*, 1969, pp. 123-131.
- [6] T. KAMAE, *Spectral Properties of Automaton-Generating Sequences*, preprint (unpublished).
- [7] A. B. KATOK and A. M. STEPIN, Approximation in Ergodic Theory, *Usp. Mat. Nauk*, Vol. **22**, (137), 1967, pp. 81-105 (in Russian).
- [8] M. KEANE, Generalized Morse Sequences, *Z. Wahr. verw. Geb.*, Vol. **10**, 1968, pp. 335-353.
- [9] J. KWIATKOWSKI, Isomorphism of Regular Morse Dynamical Systems, *Studia Math.*, Vol. **62**, 1982, pp. 59-89.
- [10] M. LEMAŃCZYK, The Rank of Regular Morse Dynamical Systems, *Z. Wahr. verw. Geb.*, Vol. **70**, 1985, pp. 33-48.
- [11] M. LEMAŃCZYK, Ergodic Properties of Morse Sequences, *Thesis*, Toruń, 1985.
- [12] M. LEMAŃCZYK, Ergodic Z_2 -Extensions Over Rational Pure Point Spectrum, Category and Homomorphisms, *Compositio Math.*, Vol. **63**, 1987, pp. 63-81.
- [13] M. LEMAŃCZYK, M. K. MENTZEN, On Metric Properties of Substitutions, *Compositio Math.* (to appear).
- [14] J. MATHEW and M. G. NADKARNI, A Measure-Preserving Transformation Whose Spectrum has Lebesgue Component of Multiplicity Two, *Bull. Lon. Math. Soc.*, Vol. **16**, 1984, pp. 402-406.
- [15] D. NEWTON, On Canonical Factors of Ergodic Dynamical Systems, *J. Lon. Math. Soc.*, Vol. **19**, 1978, pp. 129-136.
- [16] M. QUEFFELEC, Contribution à l'étude spectrale de suites arithmétiques, *Thèse*, 1984 (in French).
- [17] W. PARRY, Compact Abelian Group Extensions of Discrete Dynamical Systems, *Z. Wahr. verw. Geb.*, Vol. **13**, 1969, pp. 95-113.
- [18] S. WILLIAMS, Toeplitz Minimal Flows Which Are Not Uniquely Ergodic, *Z. Wahr. verw. Geb.*, Vol. **67**, 1984, pp. 95-107.

(Manuscrit reçu le 6 mai 1986.)