

ANNALES DE L'I. H. P., SECTION B

PAVEL M. BLEHER

JOEL L. LEBOWITZ

Variance of number of lattice points in random narrow elliptic strip

Annales de l'I. H. P., section B, tome 31, n° 1 (1995), p. 27-58

http://www.numdam.org/item?id=AIHPB_1995__31_1_27_0

© Gauthier-Villars, 1995, tous droits réservés.

L'accès aux archives de la revue « Annales de l'I. H. P., section B » (<http://www.elsevier.com/locate/anihpb>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

Variance of number of lattice points in random narrow elliptic strip

by

Pavel M. BLEHER and Joel L. LEBOWITZ

Department of Mathematics and Physics,
Rutgers University,
New Brunswick, NJ 08540, U.S.A.

Dedicated to the memory of Claude Kipnis

ABSTRACT. – Let $N(x; s)$ be the number of integral points inside an elliptic strip of area s , bounded by the ellipses $E(x)$ and $E(x + s)$, where

$$E(x) = \{ (q_1, q_2) : (q_1^2 + \mu q_2^2) \pi \mu^{-1/2} = x \}.$$

We prove that if $\mu > 0$ is a diophantine number and $s = s(T) = \text{const } T^\gamma$, with $0 < \gamma < 1/2$, then

$$\lim_{T \rightarrow \infty} \frac{1}{T} \int_T^{2T} \frac{|N(x; s) - s|^2}{s} dx = 4$$

(with the factor 4 coming from symmetry considerations). Contrariwise, if μ is rational then

$$\lim_{T \rightarrow \infty} \frac{1}{T \log T} \int_T^{2T} \frac{|N(x; s) - s|^2}{s} dx = \left(\frac{1}{2} - \gamma \right) c(\mu)$$

with some $c(\mu) > 0$, and if μ is a Liouville number then

$$\limsup_{T \rightarrow \infty} \frac{1}{T} \int_T^{2T} \frac{|N(x; s) - s|^2}{s} dx = \infty.$$

Key words: Lattice points, random strip.

Classification A.M.S. : 10 J 25, 10 E 35.

Annales de l'Institut Henri Poincaré - Probabilités et Statistiques - 0246-0203

Vol. 31/95/01/\$ 4.00/© Gauthier-Villars

RÉSUMÉ. — Soit $N(x; s)$ le nombre de points entiers dans une bande elliptique d'aire s , comprise entre les ellipses $E(x)$ et $E(x + s)$, où

$$E(x) = \{q_1, q_2\} : (q_1^2 + \mu q_2^2) \pi \mu^{-1/2} = x \}$$

Nous montrons que, si $\mu > 0$ est un nombre diophantien et $s = s(T) = \text{const } T^\gamma$, with $0 < \gamma < 1/2$, alors

$$\lim_{T \rightarrow \infty} \frac{1}{T} \int_T^{2T} \frac{|N(x; s) - s|^2}{s} dx = 4$$

(le facteur 4 vient de considérations de symmétrie). Au contraire si μ est rationnel alors

$$\lim_{T \rightarrow \infty} \frac{1}{T \log T} \int_T^{2T} \frac{|N(x; s) - s|^2}{s} dx = \left(\frac{1}{2} - \gamma\right) c(\mu)$$

où $c(\mu) > 0$, et si μ est un nombre de Liouville, alors,

$$\limsup_{T \rightarrow \infty} \frac{1}{T} \int_T^{2T} \frac{|N(x; s) - s|^2}{s} dx = \infty.$$

1. INTRODUCTION

The statistical properties of the eigenvalues

$$0 = E_0 < E_1 \leq E_2 \leq \dots, \quad \lim_{n \rightarrow \infty} E_n = \infty,$$

of the Laplace-Beltrami operator $(-\Delta)$ on a compact d -dimensional Riemannian manifold M^d , $d \geq 2$, is a problem of both physical and mathematical interest. According to the Weyl law,

$$\left. \begin{aligned} N(E) &\equiv \#\{E_n \leq E\} \\ &= CE^{d/2} + n(E), \quad n(E) = o(E^{d/2}), \\ C &= \frac{(\text{Vol } M^d) \omega_d}{(2\pi)^d}, \quad \omega_d = \text{Vol}\{x \in \mathbb{R}^d : |x| \leq 1\}. \end{aligned} \right\} \quad (1.1)$$

Defining rescaled eigenvalues by

$$x_n = (C^{-1} E_n)^{2/d}, \quad 0 = x_0 < x_1 \leq x_2 \leq \dots,$$

(1.1) is equivalent to

$$N(x) \equiv \#\{x_n \leq x\} = x + n(x), \quad n(x) = o(x),$$

so that the density of the rescaled eigenvalues x_n is equal to 1,

$$\lim_{x \rightarrow \infty} x^{-1} N(x) = 1.$$

We are interested in statistical properties of the sequence $x_0 < x_1 \leq x_2 \dots$

To characterize the statistical properties of $\{x_n\}$ consider the asymptotic behavior as $T \rightarrow \infty$ of the “temporal” moments

$$M_k(T; s) = \frac{1}{T} \int_T^{2T} N^k(x; s) dx, \quad k = 0, 1, 2, \dots,$$

of $N(x; s) = N(x + s) - N(x)$, the number of x_n 's lying in the window $[x, x + s]$. A striking conjecture proposed by Berry and Tabor [BT] is that for a “generic” Riemannian manifold with completely integrable geodesic flow the limiting moments of $N(x; s)$ coincide with those of the Poisson process, in which points are thrown randomly and independently on the line with density one,

$$\lim_{T \rightarrow \infty} M_k(T; s) = \left. \frac{d^k}{d\xi^k} e^{s(e^\xi - 1)} \right|_{\xi=0} \equiv \gamma_k(s). \tag{1.2}$$

Numerical evidence and impressive analytic arguments in the favor of this conjecture were given in [BT] and also in subsequent papers of Berry [Ber], Casati, Chirikov and Guarneri [CCG], Sinai [Sin], Major [Maj], Cheng and Lebowitz [CL], Cheng, Lebowitz and Major [CLM] and others; *see* also monographs of Gutzwiller [Gu], Ozorio de Almeida [OdA], and Tabor [Tab]. By contrast, for a generic Riemannian manifold of negative curvature the limiting moments of $N(x; s)$ are expected to coincide with those of the Wigner-Dyson gaussian orthogonal ensemble of random matrices. The proof of this as well as of (1.2) remains a challenging open problem.

Keeping s fixed while $T \rightarrow \infty$ in $M_k(T; s)$ measures only local correlations among the x_n 's. To characterize long range correlations between different x_n consider windows of size $s = s(T)$ which grow with T . The question is then, for what $s(T)$ does (1.2) still hold? Casati, Chirikov and Guarneri [CCG] studied this problem for a simple model system of a torus \mathbb{T}^2 with the metric

$$dq^2 = dq_1^2 + \mu^{-1} dq_2^2, \tag{1.3}$$

and they discovered a saturation of the Poisson asymptotics. Berry [Ber] gave an analytic explanation of the phenomenon of saturation using a heuristic trace formula for $N(E)$ and a diagonal approximation for the second moment. As a matter of fact, Berry computed the saturation of rigidity, a more complicated statistical mean of the second order, but this does not matter for our discussion.

In the present paper we continue the rigorous study of the phenomenon of saturation which was begun in our paper [BL] (see also [BDL]). To describe the phenomenon of saturation we consider the variance

$$D(T; s) = \frac{1}{T} \int_T^{2T} |N(x; s) - s|^2 dx. \quad (1.4)$$

Then the claim is that for a generic integrable system,

$$D(T; s) \sim \begin{cases} s & \text{if } s \ll T^{1/2}, \\ cT^{1/2} & \text{if } s \gg T^{1/2}, \end{cases} \quad (1.5)$$

or more precisely,

$$\lim_{T \rightarrow \infty; s/T^{1/2} \rightarrow 0} s^{-1} D(T; s) = 1, \quad (1.6)$$

$$\lim_{T \rightarrow \infty; s/T^{1/2} \rightarrow \infty} (cT^{1/2})^{-1} D(T; s) = 1. \quad (1.7)$$

In other words, $D(T; s)$ grows linearly, as a function of s , for $s \ll T^{1/2}$, like for the Poisson process, and then it saturates at $cT^{1/2}$ for $s \gg T^{1/2}$. The crossover behavior of $D(T; s)$ is described by a scaling function $V(\lambda)$, so that

$$\lim_{T \rightarrow \infty; s/T^{1/2} \rightarrow 0} T^{-1/2} D(T; s) = V(\lambda). \quad (1.8)$$

The validity of (1.7), (1.8) was established in [BL] for a number of systems. In the present work we study (1.6).

We will consider the CCG (Casati-Chirikov-Guarnieri) model system of a torus \mathbb{T}^2 with the metric (1.3), i.e., we have a periodic box with sides of length 1 and $\mu^{-1/2}$. The eigenvalues of the Laplace operator are then

$$\{(2\pi)^2 (n_1^2 + \mu n_2^2), n_1, n_2 \in \mathbb{Z}\},$$

so that the scaled eigenvalues $x_0 < x_1 \leq x_2 \leq \dots$ are elements of the set

$$X = \{(n_1^2 + \mu n_2^2) \mu^{-1/2}, n_1, n_2 \in \mathbb{Z}\},$$

ordered in increasing order. Hence

$$\begin{aligned} N(x; s) &= N(x + S) - N(x) \\ &= \#\{(n_1, n_2) \in \mathbb{Z}^2 : x < (n_1^2 + \mu n_2^2) \pi \mu^{-1/2} \leq x + s\}. \end{aligned}$$

$N(x)$ and $N(x; s)$ have the obvious geometric interpretation of the number of integral points inside of the ellipse $E(s) = \{(q_1^2 + \mu q_2^2) \pi \mu^{-1/2} = x\}$ and inside of the elliptic strip between $E(x)$ and $E(x + s)$, respectively.

A number $\mu > 0$ is called diophantine if $\exists C, M > 0$ such that for every rational p/q ,

$$\left| \mu - \frac{p}{q} \right| \geq \frac{C}{q^M}.$$

μ is a Liouville number if there exist a sequence of rationals $\frac{p_i}{q_i}$ and a sequence of positive numbers $c_i, i = 1, 2, \dots$, such that $\lim_{i \rightarrow \infty} c_i = \infty$ and

$$\left| \mu - \frac{p_i}{q_i} \right| \leq e^{-c_i q_i}, \quad i = 1, 2, \dots$$

In what follows we will assume that $s = s(T)$ is a function which satisfies

$$\exists \omega > 0 : \lim_{T \rightarrow \infty} T^{-\omega} s = \infty; \quad \lim_{T \rightarrow \infty} T^{-1/2} s = 0. \quad (1.9)$$

For the sake of definiteness, one may take $s = \text{const } T^\gamma$ with an arbitrary $0 < \gamma < 1/2$. Let

$$d = \frac{s}{2T^{1/2}};$$

then d characterizes the width the elliptic strip between $E(x)$ and $E(x+s)$, $T \leq x \leq 2T$, and the second condition in (1.9) is equivalent to $d \rightarrow 0$ as $T \rightarrow \infty$. We prove the following theorems.

THEOREM 1.1. – *If μ is diophantine and s satisfies (1.9) then*

$$\lim_{T \rightarrow \infty} \frac{1}{T} \int_T^{2T} \frac{|N(x; s) - s|^2}{s} dx = 4. \quad (1.10)$$

THEOREM 1.2. – *If $\mu = 1$ and s satisfies (1.9) then*

$$\lim_{T \rightarrow \infty} \frac{1}{T} \int_T^{2T} \frac{|N(x; s) - s|^2}{s |\log d|} dx = \frac{4}{\pi}. \quad (1.11)$$

THEOREM 1.3. – *If μ is rational and s satisfies (1.9) then $\exists c(\mu) > 0$ such that*

$$\lim_{T \rightarrow \infty} \frac{1}{T} \int_T^{2T} \frac{|N(x; s) - s|^2}{s |\log d|} dx = c(\mu). \quad (1.12)$$

Remarks. – 1. An exact value of $c(\mu)$ follows from the formulae (4.8) and (A.2) below.

2. For $s = \text{const } T^\gamma$ (1.12) reduces to

$$\lim_{T \rightarrow \infty} \frac{1}{T \log T} \int_T^{2T} \frac{|N(x; s) - s|^2}{s} dx = c(\mu) \left(\frac{1}{2} - \gamma \right).$$

In our last theorem we will need a more restrictive condition on $s = s(T)$ than (1.9):

$$\exists \omega > 0 : \lim_{T \rightarrow \infty} T^{-\omega} s = \infty, \quad \lim_{T \rightarrow \infty} T^{-(1/2)+\omega} s = 0. \quad (1.9')$$

THEOREM 1.4. – *If μ is a Liouville number and s satisfies (1.9') then*

$$\lim_{T \rightarrow \infty} \frac{1}{T} \int_T^{2T} \frac{|N(x; s) - s|^2}{s} dx = \infty. \quad (1.13)$$

Before passing to the proof of the theorems we make some comments. The “temporal” averaging $\frac{1}{T} \int_T^{2T} \dots dx$ in Theorems 1.1-1.4 can be replaced by more general averaging

$$\frac{1}{T} \int_{Ta}^{Tb} \dots p(x/T) dx, \quad (1.14)$$

with arbitrary $b > a > 0$ and arbitrary bounded probability density $p(t)$ on $[a, b]$. The proof remains the same for the uniform density $p(t) = (b-a)^{-1}$, and then by linearity it can be extended to stepwise densities. Since any bounded probability density is approximated by a sequence of stepwise densities the general case follows; for details see [BCDL] and [Ble] (see also Theorem 2.1 below). The advantage of having $p(t)$ in (1.14) is that this allows to make change of variable in averaging. For instance, when $p(t) = (1/2)(b^{1/2} - a^{1/2})^{-1} t^{-1/2}$, (1.14) reduces to averaging with respect to dR where $R = x^{1/2}$ (cf. [BL]).

Note that if $\mu > 0$ is irrational then every rescaled eigenvalue

$$x_{n_1 n_2} = (n_1^2 + \mu n_2^2) \mu^{-1/2} \quad (1.15)$$

with $n_1, n_2 > 0$ is fourfold degenerate since in this case $x_{m_1 m_2} = x_{n_1 n_2}$ if and only if $m_1 = \pm n_1, m_2 = \pm n_2$. This fourfold degeneracy of eigenvalues is the origin of the 4 in the RHS of (1.10). More precisely, if we get rid of the degeneracy by defining

$$N_+(x; s) = \# \{ (n_1, n_2) \in \mathbb{Z}^2 : n_1, n_2 > 0,$$

$$x < (n_1^2 + \mu n_2^2)^{-1/2} \leq x + s \},$$

then (1.10) is equivalent to

$$\lim_{T \rightarrow \infty} \frac{1}{T} \int_T^{2T} \frac{|N(x; s) - (1/4)s|^2}{(1/4)s} dx = 1.$$

If μ is rational then the degeneracy can be higher than 4, since for some $k \in \mathbb{Z}$ there exist many solutions of the equation

$$qn_1^2 + pn_2^2 = k. \quad (1.16)$$

On the average the number of solutions of (1.16) grows like $c \log k$ (see appendix), which gives rise to the logarithmic multiplier in the denominator in the LHS of (1.11) and (1.12). Finally, if μ is a Liouville number then it is well approximated by rationals p_i/q_i . We will show that this implies that we can choose T_i such that $|N(x; s) - s|^2$ is on the average of order $c(p_i/q_i) s |\log d|$ on the interval $T_i < x < 2T_i$ and $c(p_i/q_i) |\log d| \rightarrow \infty$, hence (1.13) follows.

As we pointed out above, the second condition in (1.9) is equivalent to $d \rightarrow 0$ as $T \rightarrow \infty$, where $d = (2T^{1/2})^{-1} s$ characterizes the width of the elliptic strip between $E(x)$ and $E(x + s)$, $T \leq x \leq 2T$. This condition has a simple geometric origin. Namely, if d is bounded from below by a positive constant, then $N(x; s)$ “feels” the rigidity of the lattice and for typical (diophantine) μ this leads to deviation from the Poisson-like behavior of $D(T; s)$ as $T \rightarrow \infty$ and this leads also to saturation when $d \rightarrow \infty$ as $T \rightarrow \infty$ [cf. (1.7), (1.8)]. Theorem 1.1 can be interpreted then as an indication of the fact that if $d \rightarrow 0$ as $T \rightarrow \infty$ then for typical μ , $N(x; s)$ does not feel the rigidity of the lattice (modulo obvious symmetry with respect to the axes), and $D(T; s)$ behaves as if the lattice points were randomly and independently thrown on the positive quadrant with density one and then every point were symmetrically reflected with respect to the axes. An interesting open problem remains to prove (or to disprove) Theorem 1.1 for s fixed.

Theorem 1.1 can be extended to the case when the elliptic strip is bounded by ellipses shifted in a fixed vector $\alpha = (\alpha_1, \alpha_2)$ in the plane. Namely, let $N_\alpha(x; s)$ be the number of lattice points between ellipses $\alpha + E(x)$ and $\alpha + E(x + s)$. A minor modification of the proof of Theorem 1.1 below allows to prove the following extension of this theorem:

THEOREM 1.1'. – *If μ is diophantine and s satisfies (1.9) then*

$$\lim_{T \rightarrow \infty} \frac{1}{T} \int_T^{2T} \frac{|N_\alpha(x; s) - s|^2}{s} dx = m_\alpha,$$

where m_α is the symmetry factor,

$$m_\alpha = \begin{cases} 1 & \text{if both } 2\alpha_1 \text{ and } 2\alpha_2 \text{ are not integers,} \\ 2 & \text{if only one of the numbers } 2\alpha_1, 2\alpha_2 \text{ is an integer,} \\ 4 & \text{if both } 2\alpha_1 \text{ and } 2\alpha_2 \text{ are integers.} \end{cases}$$

The case of rational μ and nonzero α can be also treated by our methods; in this case the asymptotic behavior of the variance of $N_\alpha(x; s)$ depends

on arithmetic properties of α : for α diophantine,

$$\lim_{T \rightarrow \infty} \frac{1}{T} \int_T^{2T} \frac{|N_\alpha(x; s) - s|^2}{s} dx = 1,$$

while for α rational a log T -multiplier appears in the normalization factor, like in Theorem 1.3 (cf. [BL]).

We would like to mention here an interesting paper of Luo and Sarnak [LS] who proved some estimates on the variance for the eigenvalues of the Laplace-Beltrami operator on arithmetic hyperbolic surfaces. The situation they study is certainly very different from ours (hyperbolic geodesic flow versus integrable) but because of an arithmetic degeneracy, the behavior of the variance turns out to be somewhat similar to the integrable case.

Theorem 1.1 is proved in Section 2. The proof utilizes a uniform estimate on the average number of integral points in a narrow elliptic strip, which is proved in Section 3. In Section 4 we prove Theorems 1.2-1.4 and in the appendix we evaluate the average squared number of solutions of the equation (1.16).

2. PROOF OF THEOREM 1.1

By (1.4) the equation (1.10) reads

$$\lim_{T \rightarrow \infty} s^{-1} D(T; s) = 4. \quad (2.1)$$

It is useful to generalize this to

$$\lim_{T, V \rightarrow \infty} s^{-1} D(T, V; s) = 4, \quad (2.2)$$

where

$$D(T, V; s) = \frac{1}{V} \int_T^{T+V} |N(x; s) - s|^2 dx. \quad (2.3)$$

We will prove (2.2) under the assumption that

$$(1/4)T^{1-\xi} \leq V \leq 4T^{1-\xi}, \quad (2.4)$$

where $\xi > 0$ is a fixed small number which will be chosen later. Then (2.1) follows. Indeed, let L be the integral part of T^ξ , $L = [T^\xi]$. Divide the interval $T \leq x \leq 2T$ into L subintervals $[T_j, T_{j+1}]$, where

$$T_j = T + jL^{-1}T, \quad j = 0, 1, \dots, L-1.$$

Then

$$V = \Delta T = T_{j+1} - T_j = L^{-1} T = [T^\epsilon]^{-1} T$$

satisfies (2.4), hence (2.2) holds which implies that $\forall \epsilon > 0, \exists T(\epsilon) > 0$ such that if $T \geq T(\epsilon)$ then

$$|s^{-1} D(T_j, \Delta T; s) - 4| \leq \epsilon, \quad j = 0, 1, \dots, L - 1.$$

Now,

$$\begin{aligned} |s^{-1} D(T; s) - 4| &= \left| L^{-1} s^{-1} \sum_{j=0}^{L-1} D(T_j, \Delta T; s) - 4 \right| \\ &\leq L^{-1} \sum_{j=0}^{L-1} |s^{-1} D(T_j, \Delta T; s) - 4| \leq \epsilon, \end{aligned}$$

so that (2.1) indeed follows from (2.2) with the restriction (2.4).

The advantage of the restriction (2.4) is that it makes V much smaller than T (for large T) and this enables us to pass from averaging over x to averaging over $R = x^{1/2}$, which is useful for the application of the Hardy-Voronoi summation formula and for all subsequent considerations. So we can rewrite $D(T, V; s)$ as

$$D(T, V; s) = \frac{1}{V_0} \int_{T_0}^{T_0+V_0} |N_0(R+d(R)) - N_0(R) - s|^2 \frac{2RV_0}{V} dR \tag{2.5}$$

where $N_0(R) = N(R^2)$ and

$$T_0 = T^{1/2}, \quad T_0 + V_0 = (T + V)^{1/2}, \quad (R + d(R))^2 = R^2 + s.$$

Observe that

$$V_0 = (T + V)^{1/2} - T^{1/2} = \frac{V}{2T^{1/2}} (1 + O(V/T)), \tag{2.6}$$

so

$$\begin{aligned} \frac{2RV_0}{V} &= \frac{2T_0V_0}{V} (1 + O(V_0/T_0)) = \frac{2T^{1/2}V(2T^{1/2})^{-1}}{V} (1 + O(V/T)) \\ &= 1 + O(V/T), \end{aligned}$$

which implies that we can drop the multiplier $\frac{2RV_0}{V}$ in the RHS of (2.5), in the limit when $T \rightarrow \infty$ and $V/T \rightarrow 0$ [here and in what follows $f = O(g)$ means $|f| \leq \text{const.} \cdot g$]. More precisely, define

$$D_0(T_0, V_0; s) = \frac{1}{V_0} \int_{T_0}^{T_0+V_0} |N_0(R+d(R)) - N_0(R) - s|^2 dR \quad (2.7)$$

Then

$$D(T, V; s) = D_0(T_0, V_0; s)(1 + O(V/T)). \quad (2.8)$$

By (2.4) and (2.6), when T is large,

$$T_0^{1-2\xi} \leq V_0 \leq 2T_0^{1-2\xi}. \quad (2.9)$$

Put now

$$d = d(T_0) = (T_0^2 + s)^{1/2} - T_0 = \frac{2}{2T_0} (1 + O(s/T_0^2)). \quad (2.10)$$

We want to replace $d(R)$ in (2.7) by d . Define to that end

$$D_1(T_0, V_0; d) = \frac{1}{V_0} \int_{T_0}^{T_0+V_0} |N_0(R+d) - N_0(R) - 2Rd - d^2|^2 dR.$$

We will prove

THEOREM 2.1. – *Assume that μ is diophantine, $d = d(T_0)$ satisfies*

$$\exists \omega > 0 : T_0^{\omega-1} < d, \quad \lim_{T_0 \rightarrow \infty} d = 0 \quad (2.11)$$

and $V_0 = V_0(T_0)$ satisfies (2.9) with $0 < 2\xi < \omega$. Then

$$\lim_{T_0 \rightarrow \infty} (2T_0 d)^{-1} D_1(T_0, V_0; d) = 4. \quad (2.12)$$

Theorems 1.1 and 2.1 are very close and the difference between these two theorems is that in Theorem 1.1 we fix the area s of the elliptic strip while in Theorem 2.1 we fix its width d . To reduce Theorem 1.1 to Theorem 2.1 we prove also

LEMMA 2.2.

$$(T_0 d)^{-1} [D_1(T_0, V_0; d(T_0)) - D_0(T_0, V_0; s)] = o(1), \quad T \rightarrow \infty. \quad (2.13)$$

Theorem 2.1, Lemma 2.2 and (2.8) imply obviously Theorem 1.1.

Proof of Theorem 2.1. – To simplify notations we redenote T_0 , V_0 and $N_0(R)$ by T , V and $N(R)$, respectively, so that now

$$N(R) = \# \{ (n_1, n_2) \in \mathbb{Z}^2 : (n_1^2 + \mu n_2^2) \pi \mu^{-1/2} \leq R^2 \},$$

and (2.12) reads

$$\lim_{T \rightarrow \infty} \frac{1}{V} \int_T^{T+V} \frac{|N(R+d) - N(R) - 2Rd - d^2|^2}{2Td} dR = 4; \quad (2.14)$$

(2.9) and (2.11) read

$$T^{1-2\xi} \leq V \leq 2T^{1-2\xi}, \quad T^{\omega-1} < d. \quad (2.15)$$

Note that

$$N(R) = \sum_{n \in \mathbb{Z}^2} \chi(n; R),$$

where

$$\chi(x; R) = \begin{cases} 1 & \text{if } |x|_\mu \leq R, \\ 0 & \text{if } |x|_\mu > R, \end{cases}$$

and

$$|x|_\mu^2 = (x_1^2 + \mu x_2^2) \pi \mu^{-1/2}.$$

Let

$$\chi_\delta(x; R) = \int_{|y|_\mu \leq R} \frac{1}{\delta^2} \varphi\left(\frac{x-y}{\delta}\right) dy,$$

where $\varphi(x) \geq 0$ is a C^∞ function with $\int \varphi(x) dx = 1$ and $\varphi(x) = 0$ when $|x|_\mu > 1$. For what follows we put

$$\delta = T^{-\zeta}$$

where ζ satisfies

$$1 \geq \zeta > 1 + \xi - \frac{\omega}{2}. \quad (2.16)$$

Note that (2.15) and (2.16) imply

$$\frac{\delta}{d} \leq T^{-\zeta+1-\omega} \leq T^{-\xi-(\omega/2)},$$

hence δ is much smaller than d when T is large. Define

$$N_\delta(R) = \sum_{n \in \mathbb{Z}^2} \chi_\delta(n; R).$$

LEMMA 2.3. – $\exists \varepsilon_0 > 0$ such that

$$\frac{1}{V} \int_T^{T+V} \frac{|N_\delta(R) - N(R)|^2}{Td} dR = O(T^{-\varepsilon_0}), \quad T \rightarrow \infty.$$

Proof. – By (2.15),

$$VTd \geq T^{1-2\xi+\omega}$$

hence it suffices to show that

$$\frac{1}{T^{1-2\xi+\omega}} \int_T^{T+V} |N_\delta(R) - N(R)|^2 dR = O(T^{-\varepsilon_0}).$$

Now,

$$N_\delta(R) - N(R) = \sum_n [\chi_\delta(n; R) - \chi(n; R)],$$

hence

$$I \equiv \int_T^{T+V} |N_\delta(R) - N(R)|^2 dR = \sum_{n, n'} I(n, n'),$$

where

$$I(n, n') = \int_T^{T+V} [\chi_\delta(n; R) - \chi(n; R)] [\chi_\delta(n'; R) - \chi(n'; R)] dR.$$

By symmetry we can assume $|n|_\mu \geq |n'|_\mu$. Observe that $\chi_\delta(x; R) - \chi(x; R)$ has support in the δ -neighborhood of the ellipse $E(R)$. This implies that $I(n, n') = 0$ if $|n|_\mu - |n'|_\mu > 2\delta = 2T^{-\zeta}$ or $|n|_\mu < T$ or $|n|_\mu > T + 2V$. In addition,

$$I(n, n') \leq \int_{|n|_\mu - \delta}^{|n|_\mu + \delta} dR = 2\delta = 2T^{-\zeta},$$

hence

$$I \leq cT^{-\zeta} \sum_{n: T \leq |n|_\mu \leq T+V} \sum_{n': 0 \leq |n|_\mu - |n'|_\mu \leq 2T^{-\zeta}} 1 \equiv cT^{-\zeta} J.$$

We will prove in the next section that

$$J = O(T^{3-\zeta+\varepsilon}), \quad \forall \varepsilon > 0 \tag{2.17}$$

(see Lemma 3.1). This gives

$$\int_T^{T+V} |N_\delta(R) - N(R)|^2 dR = O(T^{3-2\zeta+\varepsilon}), \quad \forall \varepsilon > 0.$$

Hence

$$\frac{1}{T^{1-2\xi+\omega}} \int_T^{2T} |N_\delta(R) - N(R)|^2 dR = O(T^{2-2\zeta+2\xi-\omega+\varepsilon}), \quad \forall \varepsilon > 0.$$

Since by (2.16) $2 - 2\zeta + 2\xi - \omega < 0$, Lemma 2.3 follows.

By the triangle inequality Lemma 2.3 implies

$$\lim_{T \rightarrow \infty} \left\{ \left(\frac{1}{V} \int_T^{T+V} \frac{|N_\delta(R+d) - N_\delta(R) - 2Rd - d^2|^2}{2Td} dR \right)^{1/2} - \left(\frac{1}{V} \int_T^{T+V} \frac{|N(R+d) - N(R) - 2Rd - d^2|^2}{2Td} dR \right)^{1/2} \right\} = 0,$$

hence it suffices to prove

$$\lim_{T \rightarrow \infty} \frac{1}{V} \int_T^{T+V} \frac{|N_\delta(R+d) - N_\delta(R) - 2Rd - d^2|^2}{2Td} dR = 4. \tag{2.18}$$

Let

$$\psi(\xi) = \int_{\mathbb{R}^2} e^{2\pi i x \xi} \varphi(x) dx$$

and

$$|n| = \mu^{1/4} \pi^{-1/2} (n_1^2 + \mu^{-1} n_2^2)^{1/2}, \tag{2.19}$$

which is, up to a multiplier, a dual norm to $|\cdot|_\mu$. The Hardy-Voronoi summation formula gives

$$\begin{aligned} \frac{N_\delta(R) - R^2}{R^{1/2}} &= \pi^{-2} \sum_{n \in \mathbb{Z}^2 \setminus \{0\}} |n|^{-3/2} \psi(n\delta) \\ &\times \cos\left(2\pi |n|R - \frac{3\pi}{4}\right) + O(R^{-1}), \end{aligned} \tag{2.20}$$

with the remainder $O(R^{-1})$ uniform in δ . From (2.20)

$$\begin{aligned} & \frac{N_\delta(R+d) - N_\delta(R) - 2Rd - d^2}{R^{1/2}} \\ &= -2\pi^{-2} F(R; T) + O(R^{-1}), \end{aligned} \quad (2.21)$$

where

$$\begin{aligned} F(R; T) &= \sum_{n \in \mathbb{Z}^2 \setminus \{0\}} |n|^{-3/2} \psi(n\delta) \sin(\pi|n|d) \\ &\quad \times \sin\left(2\pi|n|\left(R + \frac{d}{2}\right) - \frac{3\pi}{4}\right), \end{aligned} \quad (2.22)$$

so that (2.18) is equivalent to

$$\lim_{T \rightarrow \infty} \frac{2\pi^{-4}}{Vd} \int_T^{T+V} |F(R; T)|^2 dR = 4. \quad (2.23)$$

By symmetry we can rewrite $F(R; T)$ as

$$\begin{aligned} F(R; T) &= \sum_{n \in \Lambda_+} r(n) |n|^{-3/2} \psi(n\delta) \sin(\pi|n|d) \\ &\quad \times \sin\left(2\pi|n|\left(R + \frac{d}{2}\right) - \frac{3\pi}{4}\right), \end{aligned} \quad (2.24)$$

where

$$\Lambda_+ = \{n = (n_1, n_2) \in \mathbb{Z}^2 : n_1, n_2 \geq 0; n \neq 0\}$$

and

$$r(n) = \begin{cases} 4 & \text{if } n_1, n_2 \neq 0, \\ 2 & \text{if } n_1 n_2 = 0, \end{cases}$$

By (2.24),

$$\frac{2\pi^{-4}}{Vd} \int_T^{T+V} |F(R; T)|^2 dR = \sum_{n, n'} a(n) a(n') K(n, n'), \quad (2.25)$$

where

$$a(n) = r(n) |n|^{-3/2} \psi(n\delta) \sin(\pi|n|d) \quad (2.26)$$

and

$$\begin{aligned}
 K(n, n') &= \frac{2\pi^{-4}}{Vd} \int_T^{T+V} \sin\left(2\pi|n|\left(R + \frac{d}{2}\right) - \frac{3\pi}{4}\right) \\
 &\quad \times \sin\left(2\pi|n'|\left(R + \frac{d}{2}\right) - \frac{3\pi}{4}\right) dR. \tag{2.27}
 \end{aligned}$$

The idea to prove (2.23) is to show that the diagonal sum

$$I_d = \sum_n a(n)^2 K(n, n)$$

converges to 4 as $T \rightarrow \infty$, while the off-diagonal sum

$$I_o = \sum_{n \neq n'} a(n) a(n') K(n, n')$$

converges to 0. The difficult part is to show the convergence of the off-diagonal sum. First we consider the diagonal sum.

Observe that

$$K(n, n) = \pi^{-4} d^{-1} (1 + O(V^{-1}|n|^{-1})),$$

so

$$I_d = \pi^{-4} d^{-1} \sum_{n \in \Lambda_+} r(n)^2 |n|^{-3} \psi(n\delta)^2 \sin^2(\pi|n|d) (1 + O(V^{-1}|n|^{-1})),$$

which is an approximating sum of the integral

$$M = 16\pi^{-4} \int_{\mathbb{R}_+^2} |x|^{-3} \sin^2(\pi|x|) dx,$$

so that $\lim_{T \rightarrow \infty} I_d = M$. A direct computation gives $M = 4$ (see the appendix to [BL]), hence

$$\lim_{T \rightarrow \infty} I_d = 4.$$

We turn now to the evaluation of I_o .

From (2.27),

$$|K(n, n')| \leq cd^{-1} \min\{1, V^{-1}||n| - |n'|\}^{-1}. \tag{2.28}$$

Define

$$\begin{aligned}
 S &= \{(n, n') \in \Lambda_+ \oplus \Lambda_+ : 0 < ||n| - |n'|| \leq 1\}, \\
 S^c &= \{(n, n') \in \Lambda_+ \oplus \Lambda_+ : ||n| - |n'|| > 1\}.
 \end{aligned}$$

Then

$$\begin{aligned}
 I^c &= \sum_{(n, n') \in S^c} a(n) a(n') K(n, n') \\
 &\leq c(Vd)^{-1} \sum_{(n, n') \in S^c} |n|^{-3/2} |n'|^{-3/2} \\
 &\quad \times \psi(n\delta) \psi(n'\delta) ||n| - |n'||^{-1}. \tag{2.29}
 \end{aligned}$$

The sum

$$\sum_{(n, n') \in S^c} |n|^{-3/2} |n'|^{-3/2} ||n| - |n'||^{-1}$$

diverges as \log^2 , *i.e.*,

$$\sum_{(n, n') \in S^c : |n|, |n'| \leq L} |n|^{-3/2} |n'|^{-3/2} ||n| - |n'||^{-1} \sim c \log^2 L.$$

The function $\psi(n\delta) \psi(n'\delta)$ produces a smooth cut-off at the scale $\delta^{-1} = T^\zeta$ in the RHS of (2.29) hence we obtain

$$|I^c| \leq c(Vd)^{-1} T^\varepsilon \leq c_0 T^{2\xi - \omega + \varepsilon}, \quad \forall \varepsilon > 0.$$

This implies $\lim_{T \rightarrow \infty} I^c = 0$.

To estimate

$$I = \sum_{(n, n') \in S} a(n) a(n') K(n, n')$$

we slice S into layers

$$S_j = \{jT^{-1} \leq ||n| - |n'|| \leq (j+1)T^{-1}\}$$

and estimate

$$I_j(N) = \sum_{(n, n') \in S_j : (N/2) \leq |n| \leq N} a(n) a(n') K(n, n').$$

Let us fix some ζ' such that $\zeta < \zeta' < 1$ and assume first that $N < T^\zeta$. By (2.28), for any $(n, n') \in S_j$,

$$|K(n, n')| \leq c(j+1)^{-1} (Vd)^{-1} T,$$

and by (2.26),

$$|a(n)|, |a(n')| \leq c|n|^{-3/2}g(|n|d)$$

where

$$g(t) = \min \{ 1, t \},$$

hence

$$I_j(N) \leq c(j+1)^{-1} N^{-3} (Vd)^{-1} T g^2(Nd) \sum_{(n, n') \in S_j : (N/2) \leq |n| \leq N} 1.$$

In the next section we will prove the estimate

$$\sum_{(n, n') \in S_j : (N/2) \leq |n| \leq N} 1 \leq cN^\kappa, \quad \kappa < 2, \tag{2.30}$$

(see Lemma 3.2) which implies

$$I_j(N) \leq c(j+1)^{-1} N^{-3+\kappa} (Vd)^{-1} T g^2(Nd). \tag{2.31}$$

When $N \leq d^{-1}$ this gives

$$I_j(N) \leq c(j+1)^{-1} N^{-1+\kappa} V^{-1} T d.$$

Summing this last estimate over $j = 0, 1, \dots, T$ and $N = [2^{-i} d^{-1}]$, $i = 0, 1, \dots, [\log_2 d]$, we obtain

$$\begin{aligned} & \sum_{(n, n') \in S : |n| \leq d^{-1}} |a(n)a(n')K(n, n')| \\ & \leq cd^{2-\kappa} V^{-1} T \log T \\ & \leq c_0 T^{-(2-\kappa)\omega+2\xi} \log T \rightarrow 0 \quad \text{as } T \rightarrow \infty, \end{aligned} \tag{2.32}$$

if we take ξ such that

$$0 < \xi < \frac{(2-\kappa)\omega}{2}. \tag{2.33}$$

When $N > d^{-1}$, (2.31) gives

$$I_j(N) \leq c(j+1)^{-1} N^{-3+\kappa} (Vd)^{-1} T.$$

Summing this estimate over $j = 0, 1, \dots, T$ and $N = [2^i d^{-1}]$, $i = 0, 1, \dots, [\log_2 (T^{\zeta'} d)]$, we obtain

$$\begin{aligned} & \sum_{(n, n') \in S : d^{-1} \leq |n| \leq T^{\zeta'}} |a(n) a(n') K(n, n')| \\ & \leq cd^{2-\kappa} V^{-1} T \log T \\ & \leq c_0 T^{-(2-\kappa)\omega+2\xi} \log T \rightarrow 0 \quad \text{as } T \rightarrow \infty, \end{aligned}$$

if (2.33) holds. When $|n| \geq T^{\zeta'}$, $a(n)$ is very small due to the cut-off function $\psi(n\delta)$. To see this note that $\forall M > 0, \exists T_0 > 0$ such that if $T > T_0$ then

$$|a(n)| \leq \psi(nT^{-\zeta}) \leq (|n|T^{-\zeta})^{-M}, \quad \forall n : |n| \geq T^{\zeta'},$$

which implies

$$|a(n)| \leq T^{-3}|n|^{-3} \quad (\text{say}) \quad \forall n : |n| \geq T^{\zeta'}; \quad T > T_0.$$

Therefore

$$\begin{aligned} & \sum_{(n, n') \in S : |n| \geq T^{\zeta'}} |a(n) a(n') K(n, n')| \\ & \leq c T^{-6+\gamma} \sum_{(n, n') \in S : |n| \geq T^{\zeta'}} |n|^{-6} \rightarrow 0 \quad \text{as } T \rightarrow \infty. \end{aligned}$$

Thus we have proved that

$$I = \sum_{(n, n') \in S} a(n) a(n') K(n, n') \rightarrow 0 \quad \text{as } T \rightarrow \infty,$$

which finishes the proof of Theorem 2.1.

Proof of Lemma 2.2. – For any two positive functions $f(R)$ and $g(R)$ on $[T, T + V]$, define

$$\rho(f, g) = \left(\frac{1}{sV} \int_T^{T+V} |n(R + f(R)) - n(R + g(R))|^2 dR \right)^{1/2},$$

where

$$\begin{aligned} n(R) &= \overline{N(R) - R^2} \\ &= \#\{ (n_1, n_2) \in \mathbb{Z}^2 : (n_1^2 + \mu n_2^2) \pi \mu^{-1/2} \leq R^2 \} - R^2. \end{aligned} \quad (2.34)$$

Obviously,

$$\rho^2(f, h) \leq \rho^2(f, g) + \rho^2(g, h).$$

To prove (2.13) it suffices to show that

$$\rho^2(f, g) = o(1), \quad T \rightarrow \infty, \tag{2.35}$$

with

$$f(R) = d(R), \quad g(R) \equiv d = d(T); \quad d(R) = (R^2 + s)^{1/2} - R. \tag{2.36}$$

Consider the sequence of stepwise functions

$$f_j(R) = f(T + k 2^{-j} V) \quad \text{when } T + k 2^{-j} V \leq R < T + (k + 1) 2^{-j} V, \\ k = 0, 1, \dots, 2^j - 1.$$

Then $f_0(R) = g(R)$ and $f_j(R) \rightarrow f(R)$ as $j \rightarrow \infty$. In addition,

$$|f'(R)| = |d'(R)| = |R(R^2 + s)^{-1/2} - 1| \sim R^{-2} s \sim T^{-1} d,$$

where $f \sim g$ means $cf \leq g \leq c'f$, which implies

$$d_j \equiv \sup_{T \leq R \leq T+V} |f(R) - f_j(R)| \sim 2^j VT^{-1} d. \tag{2.37}$$

LEMMA 2.4. - If $d_j > T^{-1}$ then

$$\rho^2(f, f_j) = O(2^{-2j} VT^\epsilon d), \quad \forall \epsilon > 0.$$

Proof. - Let us show that

$$\left. \begin{aligned} \frac{1}{sV} \int_T^{T+V} |N(R + f(R)) - N(R + f_j(R))|^2 dR \\ \leq c_\epsilon 2^{-2j} VT^\epsilon d, \quad \forall \epsilon > 0. \end{aligned} \right\} \tag{2.38}$$

The proof is the same as the one of Lemma 2.3. Namely,

$$I = \int_T^{T+V} |N(R + f(R)) - N(R + f_j(R))|^2 dR = \sum_{n, n'} I(n, n'),$$

where

$$I(n, n') = \int_T^{T+V} \chi(n; R) \chi(n'; R) dR$$

where $\chi(x; R)$ is the characteristic function of the elliptic strip between $E(R + f(R))$ and $E(R + f_j(R))$. Observe that the width of this strip is $O(d_j)$. Now using the same arguments as we used in the proof of Lemma 2.3, we obtain the estimate

$$I \leq c_\varepsilon T^{3+\varepsilon} d_j^2, \quad \forall \varepsilon > 0,$$

which implies (2.38) [use (2.37)]. In addition to (2.38),

$$\frac{1}{sV} \int_T^{T+V} |(R + f(R))^2 - (R + f_j(R))^2|^2 dR \leq c 2^{-2j} V d, \quad (2.39)$$

[use again (2.37)], hence Lemma 2.4 follows.

Take

$$j_0 = \max \{ j : d_j > T^{-1} \}.$$

Then by (2.37),

$$2^{-j_0} V d \sim 1, \quad (2.40)$$

hence

$$\rho^2(f, f_{j_0}) \leq c(Vd)^{-1} T^\varepsilon = o(1), \quad T \rightarrow \infty. \quad (2.41)$$

By (2.40), $j_0 \sim \log T$. Let us estimate now $\rho^2(f_j, f_{j+1})$ when $j \leq j_0$. Define

$$\rho_\delta(f, g) = \left(\frac{1}{sV} \int_T^{T+V} |n_\delta(R + f(R)) - n_\delta(R + g(R))|^2 dR \right)^{1/2},$$

where $n_\delta(R) = N_\delta(R) - R^2$. Put $\delta = T^{-\zeta}$ with some ζ satisfying (2.16). Lemma 2.3 gives then

$$\rho_\delta^2(f_j, f_{j+1}) - \rho^2(f_j, f_{j+1}) = O(T^{-\varepsilon_0}). \quad (2.42)$$

LEMMA 2.5. — $\exists \varepsilon_0 > 0$ such that

$$\rho_\delta^2(f_j, f_{j+1}) \leq cT^{-\varepsilon_0}, \quad J = 0, 1, \dots, j_0.$$

Proof. — With the help of the Hardy-Voronoi summation formula we obtain (as in the proof of Theorem 2.1 above)

$$\rho_\delta^2(f_j, f_{j+1}) = \sum_{k=0}^{2^{j+1}-1} \sum_{n, n'} a_k(n) a_k(n') K_k(n, n') + O(T^{-\varepsilon_0}),$$

where

$$a_k(n) = r(n)|n|^{-3/2}\psi(n\delta)\sin(\pi|n|\Delta_k), \quad \Delta_k = f_{j+1}(R_k) - f_j(R_k),$$

$$R_k = T + k2^{-j-1}V,$$

and

$$K_k(n, n') = \frac{2\pi^{-3}}{Vd} \int_{R_k}^{R_{k+1}} \sin\left(2\pi|n|\left(R + \frac{\Delta_k}{2}\right) - \frac{3\pi}{4}\right) \times \sin\left(2\pi|n'|\left(R + \frac{\Delta_k}{2}\right) - \frac{3\pi}{4}\right) dR$$

[cf. (2.26), (2.27)]. Now with the help of the same arguments that we used in the proof of Theorem 1.2 we obtain that

$$\sum_{k=0}^{2^{j+1}-1} \sum_{n, n'} a_k(n) a_k(n') K_k(n, n') = O(T^{-(2-\kappa)\omega+2\xi} \log T)$$

[cf. (2.32)]. This proves Lemma 2.5

Lemma 2.5 combined with (2.41), (2.42) proves (2.35) and thus Lemma 2.2 is proved.

3. UNIFORM ESTIMATE OF THE NUMBER OF CLOSE PAIRS

In this section we put $|n| = (n_1^2 + \mu n_2^2)^{1/2}$. We call a pair $(n, n') \in \mathbb{Z}^2 \oplus \mathbb{Z}^2$ close if $||n| - |n'||$ is small. We are interested in evaluation of the number of close pairs in the annulus $T \leq |n|, |n'| \leq 2T$.

LEMMA 3.1. - $\forall \varepsilon > 0, \exists c_\varepsilon > 0$ such that $\forall T > 1$ and $\forall 1 > \delta > T^{-1}$,

$$J \equiv \sum_{n: T \leq |n| \leq 2T} \sum_{n': 0 < ||n| - |n'|| \leq \delta} 1 \leq c_\varepsilon T^{3+\varepsilon} \delta.$$

Proof. - By symmetry we may assume $|n| > |n'|$ and also that $n_1, n'_1, n_2, n'_2 \geq 0$ and estimate the sum J with these restrictions. We have

$$|n| - |n'| = \frac{|n|^2 - |n'|^2}{|n| + |n'|} \geq \frac{(n_1^2 - n_1'^2) + \mu(n_2^2 - n_2'^2)}{T},$$

hence

$$J \leq \sum_{n: T \leq |n| \leq 2T} \sum_{n': 0 < (n^2 - n_1'^2) - \mu(n_2^2 - n_2'^2) \leq T\delta} 1.$$

Let

$$n_1^2 - n_1'^2 = m_1, \quad n_2^2 - n_2'^2 = m_2.$$

Then

$$0 < m_1 - \mu m_2 \leq T\delta. \quad (3.1)$$

We can count all pairs n, n' first counting m_2 from $-4T^2$ to $4T^2$, then counting m_1 which satisfy (3.1) and then counting the divisors of m_1, m_2 . Assume first that $m_1 m_2 \neq 0$. This part of J is estimated as

$$\begin{aligned} J_{\{m_1 m_2 \neq 0\}} &\leq c \sum_{m_2=1}^{4T^2} \sum_{m_1} d(m_1) d(m_2) \leq c \sum_{m_2=1}^{4T^2} \sum_{m_1} [d(m_1)^2 + d(m_2)^2] \\ &\leq c_0 T \delta \sum_{m=1}^{c_1 T^2} d(m)^2 \leq c_\varepsilon T \delta T^{2+\varepsilon} = c_\varepsilon T^{3+\varepsilon} \delta. \end{aligned}$$

Assume second that $m_2 = 0, m_1 \neq 0$. This part is estimated as

$$J_{\{m_2=0\}} \leq c \sum_{n_2=0}^{2T} \sum_{m_1} d(m_1) \leq c_\varepsilon T^{2+\varepsilon} \delta.$$

Similarly we estimate the part with $m_1 = 0, m_2 \neq 0$. Lemma 3.1 is proved.

LEMMA 3.2. – If μ is diophantine and $\zeta > 1$ then $\exists \kappa < 2$ such that

$$J = O(T^\kappa), \quad T \rightarrow \infty.$$

Proof. – If μ is diophantine and $\zeta > 1$, then the number of $m_2, |m_2| \leq 4T^2$, which satisfy (3.1) for at least one m_1 , is $O(T^{\kappa'})$ with some $\kappa' < 2$. This implies $J = O(T^{\kappa'+\varepsilon}), \forall \varepsilon > 0$. Lemma 3.2 is proved.

LEMMA 3.3. – If μ is a rational and $\zeta > 1$ then for large T ,

$$J = \sum_{n: T \leq |n| \leq 2T} \sum_{n': 0 < ||n| - |n'||| \leq T^{-\zeta}} 1 = 0.$$

Proof. – Let $\mu = \frac{p}{q}$. Then

$$||n| - |n'||| = \frac{||n|^2 - |n'|^2|}{|n| + |n'|} \geq \frac{q^{-1}}{4T} > T^{-\zeta},$$

hence $J = 0$ for $T^{\zeta-1} > 4q$. Lemma 3.3 is proved.

4. PROOF OF THEOREMS 1.2-1.4

Proof of Theorem 1.2. – There are two places where the proof of Theorem 1.2 differs from the proof of Theorem 1.1: the first place concerns the computation of the diagonal sum and the second one concerns the estimation of the off-diagonal sum. Let us first discuss the diagonal sum.

When $\mu = 1$, (2.19) reduces to

$$|n| = \pi^{-1/2} (n_1^2 + n_2^2)^{1/2}.$$

we rewrite (2.22) as

$$F(R; T) = \sum_{k=1}^{\infty} r_2(k) \underline{k}^{-3/4} \psi(\underline{k}^{1/2} \delta) \sin(\pi \underline{k}^{1/2} d) \times \sin\left(2\pi \underline{k}^{1/2} \left(R + \frac{d}{2}\right) - \frac{3\pi}{4}\right), \tag{4.1}$$

where $r_2(k)$ is the number of representations of k as a sum of two squares, $k = n_1^2 + n_2^2$, and $\underline{k} = \pi^{-1} k = |n|^2$. As is well known, $r_2(k) \leq c_\varepsilon k^\varepsilon$, $\forall \varepsilon > 0$. To prove Theorem 1.2 we have to show that

$$\lim_{T \rightarrow \infty} \frac{2\pi^{-4}}{V d |\log d|} \int_T^{T+V} |F(R; T)|^2 dR = \frac{4}{\pi} \tag{4.2}$$

[cf. (2.23)]. By (4.1),

$$\frac{2\pi^{-4}}{V d |\log d|} \int_T^{T+V} |F(R; T)|^2 dR = \sum_{k, k'} a(k) a(k') K(k, k'),$$

where

$$a(k) = r_2(k) \underline{k}^{-3/4} \psi(\underline{k}^{1/2} \delta) \sin(\pi \underline{k}^{1/2} d) \tag{4.3}$$

and

$$K(k, k') = \frac{2\pi^{-4}}{V d |\log d|} \int_T^{T+V} \sin\left(2\pi \underline{k}^{1/2} \left(R + \frac{d}{2}\right) - \frac{3\pi}{4}\right) \times \sin\left(2\pi \underline{k}'^{1/2} \left(R + \frac{d}{2}\right) - \frac{3\pi}{4}\right) dR.$$

The diagonal sum reduces to

$$I_d = \frac{\pi^{-4}}{d|\log d|} \sum_{k=1}^{\infty} r_2(k)^2 \underline{k}^{-3/2} \psi(\underline{k}^{1/2} \delta) \\ \times \sin^2(\pi \underline{k}^{1/2} d) (1 + O(V^{-1} k^{-1/2})). \quad (4.4)$$

As shown in [BD] (see also the appendix to the present paper),

$$\lim_{T \rightarrow \infty} (N \log N)^{-1} \sum_{k=1}^N r_2(k)^2 = 4. \quad (4.5)$$

Substituting $x_k = \pi^{-1} k d^2$ we can reduce (4.4) to an approximating sum

$$I_d = \pi^{-3} |\log d|^{-1} \sum_{k=1}^{\infty} r_2(k)^2 x_k^{-3/2} \sin^2(\pi x_k^{1/2}) \Delta x (1 + O(V^{-1} k^{-1/2})), \\ \Delta x = x_{k+1} - x_k,$$

of the integral

$$M = 8 \pi^{-3} \int_0^{\infty} x^{-3/2} \sin^2(\pi x^{1/2}) dx = \frac{4}{\pi}.$$

Indeed, when x_k is order of 1, k is of order of d^{-2} , hence by (4.5) $r_2(k)^2$ is on the average of order of $4 \log d^{-2}$, so that

$$\lim_{T \rightarrow \infty} I_d = M = \frac{4}{\pi}.$$

The off-diagonal sum I_o is estimated in the same way as in the proof of Theorem 1.1, except instead of Lemma 3.2 we use Lemma 3.3. Theorem 1.2 is proved.

Proof of Theorem 1.3. – For a general $\mu = \frac{p}{q}$, (2.19) reduces to

$$|n| = \left(\frac{p}{q}\right)^{1/4} \pi^{-1/2} \left(n_1^2 + \frac{q}{p} n_2^2\right)^{1/2} = \pi^{-1/2} (pq)^{-1/4} (pn_1^2 + qn_2^2)^{1/2},$$

and (4.1) reads

$$F(R; T) = \sum_{k=1}^{\infty} r_2(k; p, q) \underline{k}^{-3/4} \psi(\underline{k}^{1/2} \delta) \sin(\pi \underline{k}^{1/2} d) \\ \times \sin\left(2\pi \underline{k}^{1/2} \left(R + \frac{d}{2}\right) - \frac{3\pi}{4}\right),$$

where $r_2(k; p, q)$ is the number of representation of k as

$$k = pn_1^2 + qn_2^2,$$

and

$$\underline{k} = \pi^{-1} (pq)^{-1/2} k = |n|^2.$$

We will prove in the appendix that

$$\lim_{N \rightarrow \infty} (N \log N)^{-1} \sum_{k=1}^N r_2(k; p, q)^2 = \sigma(p, q) > 0, \tag{4.6}$$

which is a generalization of (4.5). The diagonal sum (4.4) now reads

$$I_d = \frac{\pi^{-3}}{d|\log d|} \sum_{k=1}^{\infty} r_2(k; p, q)^2 \underline{k}^{-3/2} \psi(\underline{k}^{1/2} \delta) \times \sin^2(\pi \underline{k}^{1/2} d) (1 + O(V^{-1} \underline{k}^{-1/2})). \tag{4.7}$$

Substituting $x_k = \underline{k}d^2$ we can reduce (4.7) to an approximating sum of the integral

$$M = \sigma(p, q) (pq)^{1/2} \pi^{-3} 2 \int_0^{\infty} x^{-3/2} \sin^2(\pi x^{1/2}) dx = \frac{\sigma(p, q) (pq)^{1/2}}{\pi},$$

hence

$$\lim_{T \rightarrow \infty} I_d = c(\mu), \quad c(\mu) = \frac{\sigma(p, q) (pq)^{1/2}}{\pi}, \tag{4.8}$$

where $\sigma(p, q)$ is defined in (A.2).

The off-diagonal sum I_o is again estimated in the same way as in the proof of Theorem 1.1 except instead of Lemma 3.2 we use Lemma 3.3. Theorem 1.3 is proved.

Proof of Theorem 1.4. – Assume

$$\left| \mu - \frac{p_i}{q_i} \right| \leq e^{-c_i q_i}, \quad i = 1, 2, \dots; \quad \lim_{i \rightarrow \infty} c_i = \infty. \tag{4.9}$$

Put $T_i = e^{b_i q_i}$ with some b_i such that $\lim_{i \rightarrow \infty} b_i = \infty$. By Theorem 1.3 when T is large,

$$\frac{1}{T} \int_T^{2T} \frac{|N(x; s, \mu_i) - s|^2}{s} dx \geq \frac{c(\mu_i) \omega}{2} \log T, \quad \mu_i = \frac{p_i}{q_i}, \tag{4.10}$$

where $N(x; s, \mu)$ stands for $N(x; s)$ referred to an ellipse with the value of parameter μ . A careful inspection of the proof of Theorem 1.3 shows that (4.10) holds when $T = T_i$. In addition (4.8), (A.2) imply

$$c(\mu_i) \geq \frac{\text{const.}}{q_i},$$

so that

$$\frac{1}{T_i} \int_{T_i}^{2T_i} \frac{|N(x; s, \mu_i) - s|^2}{s} dx \geq \text{const. } b_i. \quad (4.11)$$

Now, a simple estimate gives

$$\frac{1}{T} \int_T^{2T} |N(x; s, \mu) - N(x; s, \mu_i)|^2 dx \leq \text{const. } T^4 |\mu - \mu_i|,$$

which implies

$$\begin{aligned} \frac{1}{T_i} \int_{T_i}^{2T_i} |N(x; s, \mu) - N(x; s, \mu_i)|^2 dx \\ \leq \text{const. } \exp(4b_i q_i - c_i q_i). \end{aligned} \quad (4.12)$$

If we take $b_i = (1/5) c_i$, then (4.12) combined with (4.11) give

$$\frac{1}{T_i} \int_{T_i}^{2T_i} \frac{|N(x; s, \mu) - s|^2}{s} dx \geq \text{const. } b_i \rightarrow \infty, \quad i \rightarrow \infty.$$

Theorem 1.4 is proved.

APPENDIX

Let $r_2(k; p, q)$ with $\text{gcd}(p, q) = 1$ be the number of representations of k as $k = qn_1^2 + pn_2^2$.

LEMMA A.1.

$$\lim_{N \rightarrow \infty} (N \log N)^{-1} \sum_{k=1}^N r_2(k; p, q)^2 = \sigma(p, q) \quad (\text{A.1})$$

with

$$\sigma(p, q) = \begin{cases} \frac{4 d(p) d(q)}{pq} & \text{if } p \equiv q \equiv 1 \pmod{2}; \\ \frac{(6l + 2) d(p') d(q)}{pq} & \text{if } p = 2^l p', \quad p' \equiv q \equiv 1 \pmod{2}, \end{cases} \tag{A.2}$$

where $d(p)$ is the number of divisors of p .

Proof. – We follow the appendix to [BCDL] (see also [BD1], [BD2]) where (A.1) was proved for $p = q = 1$. Consider the exponential sum

$$S(b) = \sum_{k=1}^{\infty} r_2(k; p, q)^2 e^{-k/b}. \tag{A.3}$$

We will show that

$$\lim_{b \rightarrow \infty} \frac{1}{b \log b} S(b) = \sigma(p, q). \tag{A.4}$$

By the tauberian theorem of Hardy and Littlewood [HL], (A.1) follows from (A.4). We can rewrite $S(b)$ as

$$S(b) = \sum_{m, m' \in \mathbb{Z}^2 \setminus \{0\}} e^{-|m|^2/b}, \tag{A.5}$$

summed over integral vectors m, m' with

$$|m|^2 = |m'|^2, \quad |m|^2 = qm_1^2 + pm_2^2. \tag{A.6}$$

We want to convert (A.5) into an unrestricted sum. To that end let us analyze (A.6). Assume first that

$$p \equiv q \equiv 1 \pmod{2}. \tag{A.7}$$

Obviously (A.6) is equivalent to

$$q(m_1 + m'_1)(m_1 - m'_1) = p(m_2 + m'_2)(-m_2 + m'_2). \tag{A.8}$$

This is satisfied when

$$\left. \begin{aligned} m_1 + m'_1 &= p_1 jk, & m_1 - m'_1 &= p_2 hl, \\ \text{or } k \equiv m_2 + m'_2 &= q_1 jl, & -m_2 + m'_2 &= q_2 hk, \end{aligned} \right\} \tag{A.9}$$

where

$$p_1 p_2 = p, \quad q_1 q_2 = q. \tag{A.10}$$

We would like to have a one-to-one correspondence between pairs $m, m' \in \mathbb{Z}^2$ which satisfy (A.8) and the numbers h, j, k, l, p_1, q_1 . From (A.9),

$$4|m|^2 = 4|m'|^2 = (p_1 q_1 j^2 + p_2 q_2 h^2)(p_1 q_2 k^2 + p_2 q_1 l^2) \quad (\text{A.11})$$

and

$$jk \equiv hl \pmod{2}, \quad jl \equiv hk \pmod{2} \quad (\text{A.12})$$

[use (A.7)]. This implies that

$$j^2 + h^2 \neq 0, \quad k^2 + l^2 \neq 0 \quad (\text{A.13})$$

(otherwise $|m| = |m'| = 0$) and

$$\left. \begin{array}{l} \text{either } j \equiv h \equiv 0, \\ l \equiv 0, \text{ or } j \equiv h \equiv k \equiv l \equiv 1 \pmod{2}. \end{array} \right\} \quad (\text{A.14})$$

To secure the uniqueness of the representation (A.9) we impose the following conditions:

$$\left. \begin{array}{l} p_1, q_1 > 0; \quad \gcd(k, l) = 1; \\ \text{either } k > 0 \text{ or } k = 0, l > 0. \end{array} \right\} \quad (\text{A.15})$$

PROPOSITION A.2. – *If (A.7) holds then (A.9) sets a one-to-one correspondence between pairs $m, m' \in \mathbb{Z}^2$ with $|m| = |m'| \neq 0$ and the numbers j, h, k, l, p_1, q_1 satisfying (A.13)-(A.15).*

Proof. – Observe that (A.9) and (A.15) determine uniquely the signs of j, h, k, l so we may assume that $m_1 \pm m'_1, \pm m_2 \pm m'_2 \geq 0$ and look for nonnegative j, h, k, l . Put

$$\begin{aligned} j &= \gcd(m_1 + m'_1, m_2 + m'_2), & h &= \gcd(m_1 - m'_1, -m_2 + m'_2); \\ x &= \frac{m_1 + m'_1}{j}, & y &= \frac{m_1 - m'_1}{h}, & z &= \frac{m_2 + m'_2}{j}, & w &= \frac{-m_2 + m'_2}{h}. \end{aligned}$$

Then (A.7) reduces to

$$qxy = pzw$$

with

$$\gcd(p, q) = \gcd(x, z) = \gcd(y, w) = 1.$$

Hence p divides xy and q divides zw so that

$$\begin{aligned} x &= p_1 k, & y &= p_2 l, & p_1 p_2 &= p, & p_1 &> 0; \\ z &= q_1 m, & w &= q_2 n, & q_1 q_2 &= q, & q_1 &> 0, \end{aligned}$$

and

$$kl = mn, \quad \gcd(k, m) = \gcd(l, n) = 1,$$

which implies that $k = n$ and $l = m$. Therefore (A.9) holds.

To prove uniqueness assume that we have another representation

$$z = q'_k k', \quad w = q'_2 l', \quad q'_1 q'_2 = q, \quad q'_1 > 0.$$

Then on the one hand as we showed before, $k = n$ and $l = m$, on the other hand the same computation gives $k' = n$ and $l' = m$. Hence $k = k'$ and $l = l'$ which proves the uniqueness. Proposition A.2 is proved.

Observe that if j, h, k, l satisfy (A.7) then $-j, -h, -k, -l$ satisfy (A.7) as well. So if we drop the condition on the sign of k, l in (A.15) then we will have in (A.7) a one-to-two correspondence between m, m' and j, h, k, l, p_1, q_1 . Therefore,

$$S(b) = (1/2) \sum_{j h k l p_1 q_1} \times \exp(-(p_1 q_1 j^2 + p_2 q_2 h^2)(p_1 q_2 k^2 + p_2 q_1 l^2)/(4 b)) \tag{A.16}$$

summed over all divisors p_1 of p and q_1 of q and over all (j, h, k, l) satisfying (A.13), (A.14) and

$$\gcd(k, l) = 1. \tag{A.17}$$

Let us consider a partial sum $S(b; p_1, q_1)$ in (A.16) with p_1, q_1 fixed. (A.17) reduces the possibilities allowed by (A.14) to two. Therefore

$$S(b; p_1, q_1) = S_e + S_o,$$

where the terms with even j, h are

$$S_e = (1/2) \sum_{kl} [f_1(k, l) f_2(k, l) - 1], \tag{A.18}$$

summed over (k, l) satisfying (A.17), and the terms with j and h odd are

$$S_o = (1/2) \sum_{kl} g_1(k, l) g_2(k, l), \tag{A.19}$$

summed over over odd integers (k, l) satisfying (A.17). The functions (f_i, g_i) are defined by

$$\sum_x \exp(-x^2 a_i) = f_i \quad \text{or} \quad g_i, \quad i = 1, 2, \tag{A.20}$$

where the sum is over integer x for f_i and over half-odd-integer x for g_i . In (A.20) we have used the abbreviation

$$a_i = p_i q_i (p_1 q_2 k^2 + p_2 q_1 l^2)/b. \quad (\text{A.21})$$

The (-1) in (A.18) takes account of the fact that the term ($j = h = 0$) was omitted from (A.16). By the Poisson summation formula, (A.20) gives

$$\left. \begin{aligned} f_i &= (\pi/a_i)^{1/2} \sum_r \exp(-(\pi^2/a_i) r^2), \\ g_i &= (\pi/a_i)^{1/2} \sum_r (-1)^r \exp(-(\pi^2/a_i) r^2). \end{aligned} \right\} \quad (\text{A.22})$$

For $a_i \geq 1$, (A.20) gives the asymptotics

$$f_i = 1 + O(\exp(-a_i)), \quad g_i = O(\exp(-a_i)). \quad (\text{A.23})$$

For $a_i \leq 1$, (A.22) gives

$$f_i, g_i = (\pi/a_i)^{1/2} (1 + O(\exp(-\pi^2/a_i))). \quad (\text{A.24})$$

(A.23) implies $[f_1(k, l) f_2(k, l) - 1] = O(\exp(-(k^2 + l^2)/b))$, hence

$$\sum_{k^2 + l^2 > b} [f_1(k, l) f_2(k, l) - 1] = O(b), \quad b \rightarrow \infty.$$

The density of (k, l) with $\text{gcd}(k, l) = 1$ is $6/\pi^2$, hence (A.24) implies

$$\begin{aligned} (1/2) & \sum_{k^2 + l^2 \leq b} [f_1(k, l) f_2(k, l) - 1] \\ &= \frac{3}{\pi^2} \int_{1 \leq k^2 + l^2 \leq b} \frac{\pi}{(a_1 a_2)^{1/2}} dk ld + O(b) \\ &= \frac{3b}{\pi (pq)^{1/2}} \int_{1 \leq k^2 + l^2 \leq b} \frac{dk ld}{p_1 q_2 k^2 + p_2 q_1 l^2} + O(b) \\ &= \frac{3b}{pq} \log b + O(b). \end{aligned}$$

Therefore

$$S_e = \frac{3b}{pq} \log b + O(b).$$

Similar computation gives

$$S_o = \frac{b}{pq} \log b + O(b)$$

[note that the density of odd (k, l) with $\gcd(k, l) = 1$ is equal to $2/\pi^2$]. Thus

$$S(b; p_1, q_1) = \frac{4b}{pq} \log b + O(b).$$

Summing over p_1, q_1 we obtain that

$$S(b) = \frac{4d(p)d(q)}{pq} b \log b + O(b).$$

For p, q odd Lemma A.1 is proved. For p even and q odd the proof is similar, with a somewhat more tedious arithmetics.

ACKNOWLEDGEMENTS

Research was supported in part by NSF Grant No. DMR 92-13424.

REFERENCES

- [Ber] M. V. BERRY, Semiclassical theory of spectral rigidity, *Proc. R. Soc. Lond.*, Vol. A **400**, 1985, pp. 229-251.
- [BT] M. V. BERRY and M. TABOR, Level clustering in the regular spectrum, *Proc. R. Soc. Lond.*, Vol. A **356**, 1977, pp. 375-394.
- [BCDL] P. M. BLEHER, Zh. CHENG, F. J. DYSON and J. L. LEBOWITZ, Distribution of the error term of number of lattice points inside a shifted circle, *Commun. Math. Phys.*, Vol. **154**, 1993, pp. 433-469.
- [BD1] P. M. BLEHER and F. J. DYSON, The variance of the error function in the shifted circle problem is a wild function of the shift, *Commun. Math. Phys.* (to appear).
- [BD2] P. M. BLEHER and F. J. DYSON, Mean square value of exponential sums related to representation of integers as sum of two squares, *Acta Arithm.* (to appear).
- [BDL] P. M. BLEHER, F. J. DYSON and J. L. LEBOWITZ, Non-gaussian energy level statistics for some integrable systems, *Phys. Rev. Lett.*, Vol. **71**, 1993, pp. 3047-3050.
- [BL] P. M. BLEHER and J. L. LEBOWITZ, Energy-level statistics of model quantum systems: universality and scaling in a lattice-point problem, *J. Stat. Phys.*, Vol. **74**, 1994, pp. 167-217.
- [Ble] P. M. BLEHER, On the distribution of the number of lattice points inside a family of convex ovals, *Duke Math. Journ.*, Vol. **67**, 1992, pp. 461-481.
- [CCG] G. CASATI, B. V. CHIRIKOV and I. GUARNERI, Energy-level statistics of integrable quantum systems, *Phys. Rev. Lett.*, Vol. **54**, 1985, pp. 1350-1353.
- [CL] Zh. CHENG and J. L. LEBOWITZ, Statistics of energy levels in integrable quantum systems, *Phys. Rev.*, Vol. **A44**, 1991, pp. R3399-3402.
- [CLM] Zh. CHENG, L. LEBOWITZ and P. MAJOR, On the number of lattice points between two enlarged and randomly shifted copies of an oval, Preprint, Rutgers University, 1993.
- [Gu] M. C. GUTZWILLER, *Chaos in classical and quantum mechanics*, Springer-Verlag, N. Y. e.a., 1990.
- [HL] G. H. HARDY and J. E. LITTLEWOOD, Tauberian theorems concerning power series and Dirichlet series whose coefficients are positive, *Proc. London Math. Soc.*, Vol. **13**, 1914, p. 174.

- [LS] W. LUO and P. SARNAK, Number variance for arithmetic hyperbolic surfaces, Preprint, Princeton University, 1993.
- [Maj] P. MAJOR, Poisson law for the number of lattice points in a random strip with finite area, *Prob. Theory Related Fields*, Vol. **92**, 1992, pp. 423-464.
- [OdA] A. M. OZORIO de ALMEIDA, *Hamiltonian systems: chaos and quantization*, Cambridge Univ. Press, New York, 1988.
- [Sin] Ya. G. SINAI, Poisson distribution in a geometric problem, in *Advances in Soviet Mathematics*, Vol. **3**, Ya G. SINAI Ed. (AMS, Providence, Rhode Island), 1991, pp. 199-214.
- [Tab] M. TABOR, *Chaos and integrability in nonlinear dynamics. An introduction*, Wiley, New York, 1989.

(Manuscript received March 28, 1994;
revised May 20, 1994.)