

ANNALES DE L'I. H. P., SECTION C

A. AMBROSETTI

V. COTI ZELATI

**Addendum to closed orbits of fixed energy for
a class of N-body problems**

Annales de l'I. H. P., section C, tome 9, n° 3 (1992), p. 337-338

<http://www.numdam.org/item?id=AIHPC_1992__9_3_337_0>

© Gauthier-Villars, 1992, tous droits réservés.

L'accès aux archives de la revue « Annales de l'I. H. P., section C » (<http://www.elsevier.com/locate/anihpc>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

*Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>*

**Addendum to
Closed orbits of fixed energy for a class of N-body
problems**

by

A. AMBROSETTI

Scuola Normale Superiore, Pisa

and

V. COTI ZELATI

University of Napoli, Faculty of Architecture, Napoli

In Lemma of 6 the paper *Closed Orbits of fixed energy for a class of N-body problems*, published on this Journal (Vol. 9, No. 2, 1992, pp. 187-200) it is claimed that critical points of f_ε are found by applying the Mountain-Pass Theorem. Actually, such a Theorem needs to be slightly changed, in order to make it applicable in Lemma 6. We indicate below the required modification.

Let $\Sigma_\rho = \{u \in \Lambda_0 : \|u\| \geq \rho\}$, $\Gamma_\rho = \{p \in C([0, 1], \Sigma_\rho) : \|p(0)\| = \rho, p(1) = u_1\}$, and

$$c = \inf_{p \in \Gamma_\rho} \max_{0 \leq \xi \leq 1} f_\varepsilon(p(\xi)).$$

Hereafter, $0 < \varepsilon \leq \varepsilon_0$. By Lemma 2 (i), $a \geq \beta > 0$. Suppose c is not a critical level for f_ε . Since (PS^+) holds, then there exist $m > 0$ and $k \in]0, \beta/2[$ such that $\|f'_\varepsilon\| \geq m$, for all $u \in \Lambda_0 \cap \{c - k \leq f_\varepsilon(u) \leq c + k\}$. Let $\eta(s, u)$, $\eta : [0, \tau_u] \times \Lambda_0 \rightarrow E$ denote the steepest descent flow satisfying

$$\frac{d\eta}{ds} = -X(\eta), \quad \eta(0, u) = u,$$

where X is, as usual, a pseudo-gradient vector field for f_ε , such that (i) $X(u) = 0$ if $f_\varepsilon(u) \leq c - 2k$ or $f_\varepsilon(u) \geq c + 2k$, (ii) $X(u) = f'_\varepsilon$ if $c - k \leq f_\varepsilon(u) = c + k$, and (iii) $f_\varepsilon(\eta(s, u)) \leq f_\varepsilon(u)$, $\forall 0 \leq s < \tau_u$ (see the Deformation lemma in [2]).

Since, as a consequence of (2.3), $f_\varepsilon(u) \rightarrow +\infty$ whenever $u \rightarrow v \in \partial\Lambda_0 - \{0\}$, then (iii) above readily implies that $\eta(s, u) \in \Lambda_0$ whenever $u \in \Lambda_0$ and $\eta(s, u) \neq 0$.

Let us show that $\tau_u \geq \rho/2$, for all $u \in \Sigma_\rho$. According to the preceding remark, this follows in the usual way if $\|\eta(s, u)\| > \rho/2$, $\forall s < \tau_u$ (indeed, in such a case $\tau_u = +\infty$). Otherwise, let $S \in]0, \tau_u[$ be such that $\|\eta(S, u)\| = \rho/2$, for some $u \in \Sigma_\rho$. Then

$$\frac{\rho}{2} \leq \|\eta(S, u) - u\| \leq \int_0^S \|X(\eta)\| \leq S,$$

proving the claim.

Note also that the same arguments used to prove Lemma 2 (i) yield that $(f'_\varepsilon(u)|u) > 0$, $\forall u \in \Lambda_0$, $\|u\| = \rho$. Therefore $\|\eta(s, u)\| < \rho$, whenever $s \geq 0$ and $u \in \Lambda_0$, $\|u\| = \rho$.

After these preliminaries, let $k' < \min(k, \rho/4m)$ and let $p \in \Gamma_\rho$ be such that $\max_\xi f_\varepsilon(p(\xi)) < c + k'$. Consider the path $p_1(\xi) = \eta\left(\frac{\rho}{2}, p(\xi)\right)$. Using the properties of the vector field X , one shows in the standard way that $\max_\xi f_\varepsilon(p_1(\xi)) < c - k'$. Moreover, $p_1(1) = u_1$, whereas, as noted before, $\|p_1(0)\| < \rho$. Let $\xi_0 = \min\{\chi \in [0, 1] : \|p_1(\xi)\| > \rho, \forall \xi \in [\chi, 1]\}$. Setting $q(\xi) = p_1(\xi_0 + \xi(1 - \xi_0))$, it follows that $q \in \Gamma_\rho$ as well as $\max_\xi f_\varepsilon(q(\xi)) < c - k'$, a contradiction which shows that c is a critical level for f_ε .