

S. V. LUDKOVSKY

Properties of quasi-invariant measures on topological groups and associated algebras

Annales mathématiques Blaise Pascal, tome 6, n° 1 (1999), p. 33-45

http://www.numdam.org/item?id=AMBP_1999__6_1_33_0

© Annales mathématiques Blaise Pascal, 1999, tous droits réservés.

L'accès aux archives de la revue « Annales mathématiques Blaise Pascal » (<http://math.univ-bpclermont.fr/ambp/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

Properties of quasi-invariant measures on topological groups and associated algebras.

S.V. Ludkovsky

Mathematics subject classification (1991 Revision) 22A10, 43A05.

Permanent address: Theoretical Department, Institute of General Physics, Str. Vavilov 38, Moscow, 117942, Russia.

Abstract

Properties of quasi-invariant measures relative to dense subgroups are considered on topological groups. Mainly non-locally compact groups are considered such as

(i) a group of diffeomorphisms $Diff(t, M)$ of real or non-Archimedean manifold M in cases of locally compact and nonlocally compact M , where t is a class of smoothness,

(ii) a Banach-Lie group over a classical or non-Archimedean field,

(iii) loop groups of real and non-Archimedean manifolds.

Recently quasi-invariant measures on a group of diffeomorphisms were constructed for real locally compact M in [8, 24] and for non-locally compact real or non-Archimedean manifolds M in [10, 12, 14, 18, 20, 21]. Such groups are also Banach manifolds or strict inductive limits of their sequences. Then on a real and non-Archimedean loop groups and semigroups of families of mappings from one manifold into another they were elaborated in [13, 15, 16, 17]. On real Banach-Lie groups quasi-invariant measures were constructed in [3].

This article is devoted to the investigation of properties of quasi-invariant measures that are important for analysis on topological groups and for construction irreducible representations [8, 23]. The following properties are investigated:

(1) convolutions of measures and functions,

(2) continuity of functions of measures,

(3) non-associative algebras generated with the help of quasi-invariant measures. The theorems given below show that many differences appear to be between locally compact and non-locally compact groups. The groups considered below are supposed to have structure of Banach manifolds over the corresponding fields.

1. Definitions. (a). Let G be a Hausdorff separable topological group. A real (or complex) Radon measure μ on $Af(G, \mu)$ is called left-quasi-invariant (or right) relative to a dense subgroup H of G , if μ_h (or μ^h) is equivalent to μ for each $h \in H$, where $Bf(G)$ is the Borel σ -field of G , $Af(G, \mu)$ is its completion by μ , $\mu_h(A) := \mu(h^{-1}A)$, $\mu^h(A) := \mu(Ah^{-1})$ for each $A \in Af(G, \mu)$, $d_\mu(h, g) := \mu_h(dg)/\mu(dg)$ (or $\bar{d}_\mu(h, g) := \mu^h(dg)/\mu(dg)$) denote a left (or right) quasi-invariance factor. We assume that the uniformity τ_G on G is such that $\tau_G|_H \subset \tau_H$, (G, τ_G) and (H, τ_H) are complete. We suppose also that there exists an open base in $e \in H$ such that their closures in G are compact (such pairs exist for loop groups and groups of diffeomorphisms and Banach-Lie groups). We denote by $M_l(G, H)$ (or $M_r(G, H)$) the set of left- (or right) quasi-invariant measures on G relative to H with a finite norm $\|\mu\| := \sup_{A \in Af(G, \mu)} |\mu(A)| < \infty$.

(b). Let $L_H^p(G, \mu, \mathbf{C})$ for $1 \leq p \leq \infty$ denotes the Banach space of functions $f : G \rightarrow \mathbf{C}$ such that $f_h(g) \in L^p(G, \mu, \mathbf{C})$ for each $h \in H$ and

$$\|f\|_{L_H^p(G, \mu, \mathbf{C})} := \sup_{h \in H} \|f_h\|_{L^p(G, \mu, \mathbf{C})} < \infty,$$

where $f_h(g) := f(h^{-1}g)$ for each $g \in G$. For $\mu \in M_l(G, H)$ and $\nu \in M(H)$ let

$$(\nu * \mu)(A) := \int_H \mu_h(A) \nu(dh) \text{ and } (q \bar{*} f)(g) := \int_H f(hg) q(h) \nu(dh)$$

be convolutions of measures and functions, where $M(H)$ is the space of Radon measures on H with a finite norm, $\nu \in M(H)$ and $q \in L^s(H, \nu, \mathbf{C})$, that is

$$\left(\int_H |q(h)|^s |\nu|(dh) \right)^{1/s} =: \|q\|_{L^s(H, \nu, \mathbf{C})} < \infty \text{ for } 1 \leq s < \infty.$$

2. Lemma. The convolutions

$$* : M(H) \times M_l(G, H) \rightarrow M_l(G, H) \text{ and}$$

$$\tilde{*} : L^1(H, \nu, \mathbf{C}) \times L^1_H(G, \mu, \mathbf{C}) \rightarrow L^1_H(G, \mu, \mathbf{C}).$$

are continuous \mathbf{C} -bilinear mappings

Proof. It follows immediately from the definitions, Fubini theorem and because $d_\mu(h, g) \in L^1(H \times G, \nu \times \mu, \mathbf{C})$. In fact one has,

$$\|\nu * \mu\| \leq \|\nu\| \times \|\mu\|, \quad \|q\tilde{*}f\|_{L^1_H(G, \mu, \mathbf{C})} \leq \|q\|_{L^1(H, \nu, \mathbf{C})} \times \|f\|_{L^1_H(G, \mu, \mathbf{C})}.$$

3. Definition. For $\mu \in M(G)$ its involution is given by the following formula: $\mu^*(A) := \overline{\mu(A^{-1})}$, where \bar{b} denotes complex conjugated $b \in \mathbf{C}$, $A \in Af(G, \mu)$.

4. Lemma. Let $\mu \in M_l(G, H)$ and G and H be non-locally compact with structures of Banach manifolds. Then μ^* is not equivalent to μ .

Proof. Let $T : G \rightarrow TG$ be the tangent mapping. Then μ induces quasi-invariant measure λ from an open neighbourhood W of the unit $e \in G$ on a neighbourhood of the zero section V in T_eG and then it has an extension onto the entire T_eG . Let at first T_eG be a Hilbert space. Put $Inv(g) = g^{-1}$ then $T \circ Inv \circ T^{-1} =: K$ on V is such that there is not any operator B of trace class on T_eG such that $\tilde{M}_\lambda \subset B^{1/2}T_eG$ and $KT_eG \subset \tilde{M}_\lambda$, where $Re(1 - \theta(z)) \rightarrow 0$ for $(Bz, z) \rightarrow 0$ and $z \in T_eG$, $\theta(z)$ is the characteristic functional of λ , \tilde{M}_λ is the set of all $x \in T_eG$ such that λ_x is equivalent to λ (see theorem 19.1 [25]). Then using theorems for induced measures from a Hilbert space on a Banach space [2, 9], we get the statement of lemma 4.

5. Lemma. For $\mu \in M_l(G, H)$ and $1 \leq p < \infty$ the translation map $(q, f) \rightarrow f_q(g)$ is continuous from $H \times L^p_H(G, \mu, \mathbf{C})$ into $L^p_H(G, \mu, \mathbf{C})$.

Proof. For metrizable G in view of the Lusin theorem (2.3.5 in [5]) and definitions of τ_G and τ_H for each $\epsilon > 0$ there are a neighbourhood $V \ni e$ in H and compacts K_1 and K in G such that the closure $cl_G VK_1 =: K_2$ is compact in G with $K_2 \subset K$, the restriction $f|_{K_2}$ is continuous and $(|\tilde{\mu}| + |\mu|)(G \setminus K_2) < \epsilon$, where $\tilde{\mu}(dg) := f(g)\mu(dg)$.

6. Proposition. For a probability measure $\mu \in M(G)$ there exists an approximate unit, that is a sequence of non-negative continuous functions $\psi_i : G \rightarrow \mathbf{R}$ such that $\int_G \psi_i(g)\mu(dg) = 1$ and for each neighbourhood $U \ni e$ in G there exists i_0 such that $supp(\psi_i) \subset U$ for each $i > i_0$.

Proof follows from the Radon property of μ and the existence of countable base of neighbourhoods in $e \in G$.

7. Proposition. If $(\psi_i : i \in \mathbf{N})$ is an approximate unit in H relative to a probability measure $\nu \in M(H)$, then $\lim_{i \rightarrow \infty} \psi_i * f = f$ in the $L^1_H(G, \mu, \mathbf{C})$ norm, where $\mu \in M_l(G, H)$, $f \in L^1_H(G, \mu, \mathbf{C})$.

Proof follows from lemma III.11.18 [6] and lemmas 2, 5.

8. Lemma. Suppose $g \in L^q_H(G, \mu, \mathbf{C})$ and $(g^x|_H) \in L^q(H, \nu, \mathbf{C})$ for each $x \in G$, $f \in L^p(H, \nu, \mathbf{C})$ with $1 < p < \infty$, $1/p + 1/q = 1$, where $g^x(y) := g(yx)$ for each x and $y \in G$. Let μ and ν be probability measures, $\mu \in M_l(G, H)$, $\nu \in M(H)$. Then $f \tilde{*} g \in L^1_H(G, \mu, \mathbf{C})$ and there exists a function $h : G \rightarrow \mathbf{C}$ such that $h|_H$ is continuous, $h = f \tilde{*} g$ μ -a.e. on G and h vanishes at ∞ on G .

Proof. In view of Fubini theorem and Hölder inequality we have

$$\begin{aligned} \|f \tilde{*} g\|_{L^1_H(G, \mu, \mathbf{C})} &= \sup_{s \in H} \int_G \int_H |f(y)| \times |g(z)| \nu(dy) \mu((ys)^{-1} dz) \leq \\ \sup_{s \in H} \left(\int_G \int_H |g(z)|^q \nu(dy) \mu((ys)^{-1} dz) \right)^{1/q} &\times \left(\int_G \int_H |f(y)|^p \nu(dy) \mu((ys)^{-1} dz) \right)^{1/p} \leq \\ \|f\|_{L^p(H, \nu, \mathbf{C})} \times \|g\|_{L^q_H(G, \mu, \mathbf{C})} &\times \nu(H) \mu(G). \end{aligned}$$

The equation $\alpha_f(\phi) := \int_H f(y) \overline{\phi(y)} \nu(dy)$ defines a continuous linear functional on $L^q(H, \nu, \mathbf{C})$. In view of lemma 5 the function $\alpha_f(g^{(sx)^{-1}}) =: \tilde{h}((sx)^{-1}) =: w(s, x)$ of two variables s and x is continuous on $H \times H$ for $s, x \in H$, since the mapping $(s, x) \mapsto (sx)^{-1}$ is continuous from $H \times H$ into H . By Fubini theorem (see §2.6.2 in [5])

$$\begin{aligned} \int_G h(y) \psi(y) \mu(dy) &= \int_G \int_H f(y) g(yx) \psi(x) \nu(dy) \mu(dx) = \\ \int_H f(y) \left[\int_G g(yx) \psi(x) \mu(dx) \right] \nu(dy) \end{aligned}$$

for each $\psi \in L^p(G, \mu, \mathbf{C})$, since

$$\int_G \int_H |f(y) g(yx) \psi(x)| |\nu|(dy) |\mu|(dx) < \infty,$$

where $|\nu|$ denotes the variation of the real-valued measure ν , $h(y) := \tilde{h}(y^{-1})$. Here ψ is arbitrary in $L^p(G, \mu, \mathbf{C})$, from this it follows, that $\mu(\{y : h(y) \neq (f \tilde{*} g)(y), y \in G\}) = 0$, since h and $(f \tilde{*} g)$ are μ -measurable functions due to Fubini theorem and the continuity of the composition and the inversion in a topological group. In view of Lusin theorem (see §2.3.5 in [5]) for each $\epsilon > 0$ there are compact subsets $C \subset H$ and $D \subset G$ and functions $f' \in L^p(H, \nu, \mathbf{C})$

and $g' \in L^q_H(G, \mu, \mathbf{C})$ with closed supports $\text{supp}(f') \subset C$, $\text{supp}(g') \subset D$ such that $cl_G CD$ is compact in G ,

$$\|f' - f\|_{L^p(H, \nu, \mathbf{C})} < \epsilon \text{ and } \|g' - g\|_{L^q_H(G, \mu, \mathbf{C})} < \epsilon,$$

since by the supposition of §1 the group H has the base \mathbf{B}_H of its topology τ_H , such that the closures $cl_G V$ are compact in G for each $V \in \mathbf{B}_H$. From the inequality

$$|h'(x) - h(x)| \leq (\|f\|_{L^p(H, \nu, \mathbf{C})} + \epsilon)\epsilon + \epsilon\|g\|_{L^q_H(G, \mu, \mathbf{C})}$$

it follows that for each $\delta > 0$ there exists a compact subset $K \subset G$ with $|h(x)| < \delta$ for each $x \in G \setminus K$, where $h'(x^{-1}) := \alpha_{f'}(g^x)$.

9. Proposition. *Let $A, B \in Af(G, \mu)$, μ and ν be probability measures, $\mu \in M_l(G, H)$, $\nu \in M(H)$. Then the function $\zeta(x) := \mu(A \cap xB)$ is continuous on H and $\nu(yB^{-1} \cap H) \in L^1(H, \nu, \mathbf{C})$. Moreover, if $\mu(A)\mu(B) > 0$, $\mu(\{y \in G : yB^{-1} \cap H \in Af(H, \nu) \text{ and } \nu(yB^{-1} \cap H) > 0\}) > 0$, then $\zeta(x) \neq 0$ on H .*

Proof. Let $g_x(y) := ch_A(y)ch_B(x^{-1}y)$, then $g_x(y) \in L^q_H(G, \mu, \mathbf{C})$ for $1 < q < \infty$, where $ch_A(y)$ is the characteristic function of A . In view of propositions 6 and 7 there exists $\lim_{i \rightarrow \infty} \psi_i * g_x = g_x$ in $L^1_H(G, \mu, \mathbf{C})$. In view of lemma III.11.18 [6] and lemma 8, $\zeta(x)|_H$ is continuous. There is the following inequality:

$$1 \geq \int_H \mu(A \cap xB)\nu(dx) = \int_H \int_G ch_A(y)ch_B(x^{-1}y)\mu(dy)\nu(dx).$$

In view of Fubini theorem there exists

$$\int_H ch_B(x^{-1}y)\nu(dy) = \nu((yB^{-1}) \cap H) \in L^1(G, \mu, \mathbf{C}), \text{ hence}$$

$$\int_H \mu(A \cap xB)\nu(dx) = \int_G \nu(yB^{-1} \cap H)ch_A(y)\mu(dy).$$

10. Corollary. *Let $A, B \in Af(G, \mu)$, $\nu \in M(H)$ and $\mu \in M_l(G, H)$ be probability measures. Then denoting $\text{Int}_H V$ the interior of a subset V of H with respect to τ_H , one has*

(i) $\text{Int}_H(AB) \cap H \neq \emptyset$, when

$$\mu(\{y \in G : \nu(yB \cap H) > 0\}) > 0;$$

(ii) $\text{Int}_H(AA^{-1}) \ni e$, when

$$\mu(\{y \in G : \nu(yA^{-1} \cap H) > 0\}) > 0.$$

Proof. $AB \cap H \supset \{x \in H : \mu(A \cap xB^{-1}) > 0\}$.

11. Corollary. Let $G = H$. If $\mu \in M_1(G, H)$ is a probability measure, then G is a locally compact topological group.

Proof. Let us take $\nu = \mu$ and $A = C \cup C^{-1}$, where C is a compact subset of G with $\mu(C) > 0$, whence $\mu(yA) > 0$ for each $y \in G$ and inevitably $\text{Int}_G(AA^{-1}) \ni e$.

12. Lemma. Let $\mu \in M_1(G, H)$ be a probability measure and G be non-locally compact. Then $\mu(H) = 0$.

Proof. This follows from theorem 19.2 [25] and theorem 3.21 and lemma 3.26 [19] and the proof of lemma 4, since the embedding $T_e H \hookrightarrow T_e G$ is a compact operator in the non-Archimedean case and of trace class in the real case (see also the papers about construction of quasi-invariant measures on the groups considered here [3, 8, 10, 12, 13, 14, 16, 17, 18], [20, 21, 24]). Indeed, the measure μ on G is induced by the corresponding measure ν on a Banach space Z for which there exists a local diffeomorphism $A : W \rightarrow V$, where W is a neighbourhood of e in G and V is a neighbourhood of 0 in Z . The measure μ on G is quasi-invariant relative to H . Therefore, the measure ν on U is quasi-invariant relative to the action of elements $\psi \in W' \subset W \cap H$ due to the local diffeomorphism A , that is, ν_ϕ is equivalent to ν for each $\phi := A\psi A^{-1}$, where $AW'A^{-1}U \subset V$, W' is an open neighbourhood of e in H and U is an open neighbourhood of 0 in Z , $\nu_\phi(E) := \nu(\phi^{-1}E)$, ϕ is an operator on Z such that it may be non-linear. The quasi-invariance factor $\rho_\nu(\phi, \nu)$ has expression through $|\det(\phi')|$ and the quasi-invariance factor $q_\nu(z, x)$ relative to linear shifts $z \in Z'$ given by theorems from §26 [25] in the real case and theorem 3.28 [19] in the non-Archimedean case:

$$\nu_\phi(dx)/\nu(dx) = |\det\{\phi'(\phi^{-1}(x))\}|q_\nu(x - \phi^{-1}(x), x),$$

where $x \in U$, $\phi = A\psi A^{-1}$, $\psi \in W'$. Then $(A\psi A^{-1}v - v) \in Z'$ for each $v \in V$ and $\psi \in W'$, where ν on Z is quasi-invariant relative to shifts on vectors $z \in Z'$ and there exists a compact operator in the non-Archimedean case and an operator of trace class in the real case of embedding $\theta : Z' \hookrightarrow Z$ such that $\nu(Z') = 0$.

13. Theorem. *Let (G, τ_G) and (H, τ_H) be a pair of topological non-locally compact groups G, H (Banach-Lie, Frechet-Lie or groups of diffeomorphisms or loop groups) with uniformities τ_G, τ_H such that H is dense in (G, τ_G) and there is a probability measure $\mu \in M_1(G, H)$ with continuous $d_\mu(z, g)$ on $H \times G$. Also let X be a Hilbert space over \mathbb{C} and $U(X)$ be the unitary group. Then (1) if $T : G \rightarrow U(X)$ is a weakly continuous representation, then there exists $T' : G \rightarrow U(X)$ equal μ -a.e. to T and $T'|_{(H, \tau_H)}$ is strongly continuous; (2) if $T : G \rightarrow U(X)$ is a weakly measurable representation and X is separable, then there exists $T' : G \rightarrow U(X)$ equal to T μ -a.e. and $T'|_{(H, \tau_H)}$ is strongly continuous.*

Proof. Let $R(G) := (I) \cup L^1(G, \mu)$, where I is the unit operator on L^1 . Then we can define

$$A_{(\lambda e+a)_h} := \lambda I + \int_G a_h(g)[d(h^{-1}, g)]T_g\mu(dg),$$

where $a_h(g) := a(h^{-1}g)$. Then

$$|(A_{(\lambda e+a)_h} - A_{\lambda e+a}\xi, \eta)| \leq \int_G |a_h(g)d_\mu(h, g) - a(g)| |(T_g\xi, \eta)|\mu(dg),$$

hence A_{a_h} is strongly continuous with respect to $h \in H$, that is,

$$\lim_{h \rightarrow e} |A_{a_h}\xi - A_a\xi| = 0.$$

Denote $A_{a_h} = T'_h A_a$ as in §29 [22], so $T'_h\xi = A_{a_h}\xi$, where $\xi = A_a\xi_0$, $a \in L^1$. Whence

$$\begin{aligned} (T'_h\xi, T'_h\xi) &= (A_{a_h}\xi_0, A_{a_h}\xi_0) = \\ &= \int_G \bar{a}_h(g)(T_g\xi_0, T_{g'}\xi_0)d_\mu(h^{-1}, g)d_\mu(h^{-1}, g')a_h(g')\mu(dg)\mu(dg') \\ &= \int_G \bar{a}(z)a(z')(U_z\xi_0, U_{z'}\xi_0)\mu(dz)\mu(dz') = (\xi, \xi). \end{aligned}$$

Therefore, T'_h is uniquely extended to a unitary operator in the Hilbert space $X' \subset X$. In view of lemma 12, $\mu(H) = 0$. Hence T' may be considered equal to T μ -a.e. Then the space $\text{span}_{\mathbb{C}}[A_{a_h} : h \in H]$ is evidently dense in X , since

$$\begin{aligned} (A_{a_h}\xi_1, A_{x_q}\xi_0) &= \left(\int_G a_h(g)T_g d(h^{-1}, g)\mu(dg)\xi_1, \int_G x_q(g')T_{g'} d(q^{-1}, g')\mu(dg')\xi_0 \right) = \\ &= (T_h \int_G a(g)T_g\mu(dg)\xi_1, T_q \int_G x(g')T_{g'}\mu(dg')) = (T_{q^{-1}h}A_a\xi_1, A_x\xi_0). \end{aligned}$$

For proving the second statement let

$R := [\xi : A_a \xi = 0 \text{ for each } a \in L^1(G, \mu)]$. If

$$(A_a \xi, \eta) = \int_G a(g)(T_g \xi, \eta) \mu(dg) = \int_G a(g)(T'_g \xi, \eta) \mu(dg)$$

for each $a(g) \in L^1(G, \mu, \mathbb{C})$, then $(T_g \xi, \eta) = (T'_g \xi, \eta)$ for μ -almost all $g \in G$. Suppose that $\{\xi_n : n \in \mathbb{N}\}$ is a complete orthonormal system in X . If $\xi \in X$, then

$$\int_G a(g)(T_g \xi, \xi_m) \mu(dg) = 0$$

for each $g \in G \setminus S_m$, where $\mu(S_m) = 0$. Therefore, $(T_g \xi, \xi_m) = 0$ for each $m \in \mathbb{N}$, if $g \in G \setminus S$, where $S := \bigcup_{m=1}^{\infty} S_m$. Hence $T_g \xi = 0$ for each $g \in G \setminus S$, consequently, $\xi = 0$. Then $(T_g \xi_n, \xi_m) = (T'_g \xi_n, \xi_m)$ for each $g \in G \setminus \gamma_{n,m}$, where $\mu(\gamma_{n,m}) = 0$. Hence $(T_g \xi_n, \xi_m) = (T'_g \xi_n, \xi_m)$ for each $n, m \in \mathbb{N}$ and each $g \in G \setminus \gamma$, where $\gamma := \bigcup_{n,m} \gamma_{n,m}$ and $\mu(\gamma) = 0$. Therefore, $R = 0$.

14. Definition and note. Let $\{G_i : i \in \mathbb{N}_o\}$ be a sequence of topological groups such that $G = G_0$, $G_{i+1} \subset G_i$ and G_{i+1} is dense in G_i for each $i \in \mathbb{N}_o$ and their topologies are denoted τ_i , $\tau_i|_{G_{i+1}} \subset \tau_{i+1}$ for each i , where $\mathbb{N}_o := \{0, 1, 2, \dots\}$. Suppose that these groups are supplied with real probability quasi-invariant measures μ^i on G_i relative to G_{i+1} . For example, such sequences exist for groups of diffeomorphisms or loop groups considered in previous papers [10, 12, 13, 15, 16, 17, 18], [20, 21]. Let $L^2_{G_{i+1}}(G_i, \mu^i, \mathbb{C})$ denotes a subspace of $L^2(G_i, \mu^i, \mathbb{C})$ as in §1(b). Such spaces are Banach and not Hilbert in general. Let $\tilde{L}^2(G_{i+1}, \mu^{i+1}, L^2(G_i, \mu^i, \mathbb{C})) := H_i$ denotes the subspace of $L^2(G_i, \mu^i, \mathbb{C})$ of elements f such that

$$\|f\|_i^2 := [\|f\|_{L^2(G_i, \mu^i, \mathbb{C})}^2 + \|f\|_i'^2]/2 < \infty, \text{ where}$$

$$\|f\|_i'^2 := \int_{G_{i+1}} \int_{G_i} |f(y^{-1}x)|^2 \mu^i(dx) \mu^{i+1}(dy).$$

Evidently H_i are Hilbert spaces due to the parallelogram identity. Let

$$f^{i+1} * f^i(x) := \int_{G_{i+1}} f^{i+1}(y) f^i(y^{-1}x) \mu^{i+1}(dy)$$

denotes the convolution of $f^i \in H_i$.

15. Lemma. *The convolution $*$: $H_{i+1} \times H_i \rightarrow H_i$ is a continuous bilinear mapping.*

Proof. In view of Fubini theorem and Cauchy inequality:

$$\begin{aligned}
 & \int_{G_{i+1}} \int_{G_i} |f^{i+1} * f^i(z^{-1}x)|^2 \mu^i(dx) \mu^{i+1}(dz) = \\
 & \int_{G_{i+1}} \int_{G_i} \int_{G_{i+1}} f^{i+1}(y) f^i(y^{-1}z^{-1}x) \mu^{i+1}(dy) \int_{G_{i+1}} \bar{f}^{i+1}(q) \bar{f}^i(q^{-1}z^{-1}x) \mu^{i+1}(dq) \mu^i(dx) \mu^{i+1}(dz) \\
 & \leq \int_{G_i} \int_{G_{i+1}} \left(\int_{G_{i+1}} |f^{i+1}(y)|^2 \mu^{i+1}(dy) \right)^{1/2} \left(\int_{G_{i+1}} |f^{i+1}(q)|^2 \mu^{i+1}(dq) \right)^{1/2} \\
 & \left(\int_{G_{i+1}} |f^i(y^{-1}z^{-1}x)|^2 \mu^{i+1}(dy) \right)^{1/2} \left(\int_{G_{i+1}} |f^i(q^{-1}z^{-1}x)|^2 \mu^{i+1}(dq) \right)^{1/2} \mu^i(dx) \mu^{i+1}(dz) \leq \\
 & \|f^{i+1}\|_{L^2(G_{i+1}, \mu^{i+1}, \mathbb{C})}^2 \int_{G_i} \left[\int_{G_{i+1}} \int_{G_{i+1}} |f^i(y^{-1}z^{-1}x)|^2 \mu^{i+1}(dy) \mu^{i+1}(dz) \right]^2 \\
 & \left[\int_{G_{i+1}} \int_{G_{i+1}} |f^i(q^{-1}z^{-1}x)|^2 \mu^{i+1}(dq) \mu^{i+1}(dz) \right]^2 \mu^i(dx) \leq \\
 & \|f^{i+1}\|_{L^2(G_{i+1}, \mu^{i+1}, \mathbb{C})}^2 \int_{G_{i+1}} \int_{G_i} \int_{G_{i+1}} |f^i(y^{-1}z^{-1}x)|^2 \mu^{i+1}(dy) \mu^{i+1}(dz) \mu^i(dx) \\
 & = \|f^{i+1}\|_{L^2(G_{i+1}, \mu^{i+1}, \mathbb{C})}^2 \left(\int_{G_i} \int_{G_{i+1}} \int_{G_{i+1}} |f^i(y^{-1}\gamma)|^2 \mu^{i+1}(dy) \mu^{i+1}(dz) d_{\mu^i}(z^{-1}, \gamma) \mu^i(d\gamma) \right) \leq \\
 & \|f^{i+1}\|_{L^2(G_{i+1}, \mu^{i+1}, \mathbb{C})}^2 \int_{G_i} \int_{G_{i+1}} |f^i(z^{-1}x)|^2 \mu^{i+1}(dz) \mu^i(dx), \text{ since} \\
 & \int_{G_i} \int_{G_{i+1}} d_{\mu^i}(z^{-1}, \gamma) \mu^i(d\gamma) \mu^{i+1}(dz) = \int_{G_{i+1}} \mu^{i+1}(dz) \int_{G_i} \mu^i(zd\gamma) = 1. \text{ Then} \\
 & \|f^{i+1} * f^i\|_{L^2(G_i, \mu^i, \mathbb{C})}^2 = \int_{G_i} \left| \int_{G_{i+1}} f^{i+1}(y) f^i(y^{-1}x) \mu^{i+1}(dy) \right|^2 \mu^i(dx) \\
 & \leq \|f^{i+1}\|_{L^2(G_{i+1}, \mu^{i+1}, \mathbb{C})}^2 \int_{G_i} \int_{G_{i+1}} |f^i(z^{-1}x)|^2 \mu^{i+1}(dz) \mu^i(dx). \text{ Therefore,} \\
 & \|f^{i+1} * f^i\|_i \leq \|f^{i+1}\|_{L^2(G_{i+1}, \mu^{i+1}, \mathbb{C})} \|f^i\|_i.
 \end{aligned}$$

16. Definition. Let $l_2(\{H_i : i \in \mathbf{N}_o\}) =: H$ be the Hilbert space consisting of elements $f = (f^i : f^i \in H_i, i \in \mathbf{N}_o)$, for which

$$\|f\|^2 := \sum_{i=0}^{\infty} \|f^i\|_i^2 < \infty.$$

For elements f and $g \in H$ their convolution is defined by the formula: $f \star g := h$ with $h^i := f^{i+1} * g^i$ for each $i \in \mathbf{N}_o$. Let $*$: $H \rightarrow H$ be an involution

such that $f^* := (\bar{f}^{j^\wedge} : j \in \mathbf{N}_\circ)$, where $f^{j^\wedge}(y_j) := f^j(y_j^{-1})$ for each $y_j \in G_j$, $f := (f^j : j \in \mathbf{N}_\circ)$, \bar{z} denotes the complex conjugated $z \in \mathbf{C}$.

17. Lemma. *H is a non-associative non-commutative Hilbert algebra with involution $*$, that is $*$ is conjugate-linear and $f^{**} = f$ for each $f \in H$.*

Proof. In view of Lemma 15 the convolution $h = f * g$ in the Hilbert space H has the norm $\|h\| \leq \|f\| \|g\|$, hence is a continuous mapping from $H \times H$ into H . From its definition it follows that the convolution is bilinear. It is non-associative as follows from the computation of i -th terms of $(f * g) * q$ and $f * (g * q)$, which are $(f^{i+2} * g^{i+1}) * q^i$ and $f^{i+1} * (g^{i+1} * q^i)$ respectively, where f, g and $q \in H$. It is non-commutative, since there are f and $g \in H$ for which $f^{i+1} * g^i$ are not equal to $g^{i+1} * f^i$. Since $f^{j^\wedge}(y_j) = f^j(y_j)$ and $\bar{\bar{z}} = z$, one has $f^{**} = (f^*)^* = f$.

18. Note. In general $(f * g^*)^* \neq g * f^*$ for f and $g \in H$, since there exist f^j and g^j such that $g^{j+1} * (f^j)^* \neq (f^{j+1} * (g^j)^*)^*$. If $f \in H$ is such that $f^j|_{G_{j-i}} = f^{j+1}$, then

$$((f^{j-1})^* * f^j)(e) = \int_{G_{j+1}} \bar{f}^{j+1}(y^{-1}) f^{j+1}(y) \mu^{j+1}(dy) = \|f^{j+1}\|_{L^2(G_{j+1}, \mu^{j+1}, \mathbf{C})}^2,$$

where $j \in \mathbf{N}_\circ$.

19. Definition. Let $l_2(\mathbf{C})$ be the standard Hilbert space over the field \mathbf{C} be considered as a Hilbert algebra with the convolution $\alpha * \beta = \gamma$ such that $\gamma^i := \alpha^{i+1} \beta^i$, where $\alpha := (\alpha^i : \alpha^i \in \mathbf{C}, i \in \mathbf{N}_\circ)$, α, β and $\gamma \in l_2(\mathbf{C})$.

20. Note. The algebra $l_2(\mathbf{C})$ has two-sided ideals $J_i := \{\alpha \in l_2(\mathbf{C}) : \alpha^j = 0 \text{ for each } j > i\}$, where $i \in \mathbf{N}_\circ$. That is, $J * l_2(\mathbf{C}) \subset J$ and $l_2(\mathbf{C}) * J = J$ and J is the \mathbf{C} -linear subspace of $l_2(\mathbf{C})$, but $J * l_2(\mathbf{C}) \neq J$. There are also right ideals, which are not left ideals: $K_i := \{\alpha \in l_2(\mathbf{C}) : \alpha^j = 0 \text{ for each } j = 0, \dots, i\}$, where $j \in \mathbf{N}_\circ$. That is, $l_2(\mathbf{C}) * K_i = K_i$, but $K_i * l_2(\mathbf{C}) = K_{i-1}$ for each $i \in \mathbf{N}_\circ$, where $K_{-1} := l_2(\mathbf{C})$. The algebra $l_2(\mathbf{C})$ is the particular case of H , when $G_j = \{e\}$ for each $j \in \mathbf{N}_\circ$. We consider further H for non-trivial topological groups outlined above.

21. Theorem. *If F is a maximal proper left or right ideal in H , then H/F is isomorphic as the nonassociative noncommutative algebra over \mathbf{C} with $l_2(\mathbf{C})$.*

Proof. Since F is the ideal, it is the \mathbf{C} -linear subspace of H . Suppose, that there exists $j \in \mathbf{N}_\circ$ such that $f^j = 0$ for each $f \in F$, then $f^i = 0$ for each $i \in \mathbf{N}_\circ$, since the space of bounded complex-valued continuous functions $C_b^0(G_\infty, \mathbf{C})$ on $G_\infty := \bigcap_{j=0}^\infty G_j$ is dense in each $H_j := \{f^j : f \in H\}$

and $C_b^0(G_\infty, \mathbb{C}) \cap F_j = \{0\}$ and $C_b^0(G_j, \mathbb{C})|_{G_{j+1}} \supset C_b^0(G_{j+1}, \mathbb{C})$. Therefore, $F_j \neq \{0\}$ for each $j \in \mathbb{N}_o$, consequently, $\mathbb{C} \hookrightarrow F_j$ for each $j \in \mathbb{N}_o$. Since \mathbb{C} is embeddable into each F_j , then there exists the embedding of $l_2(\mathbb{C})$ into F , where $H_j := \{f^j : f \in H\}$, $\pi_j : H \rightarrow H_j$ are the natural projections.

The subalgebra F is closed in H , since H is a topological algebra and F is a maximal proper subalgebra. The space $H_\infty := \bigcap_{j \in \mathbb{N}_o} H_j$ is dense in each H_j and the group $G_\infty := \bigcap_{j \in \mathbb{N}_o} G_j$ is dense in each G_j .

Suppose that $F_i = H_i$ for some $i \in \mathbb{N}_o$, then $F_j = H_j$ for each $j \in \mathbb{N}_o$, since $C_b^0(G_\infty, \mathbb{C})$ is dense in each H_j and $C_b^0(G_j, \mathbb{C})|_{G_{j+1}} \supset C_b^0(G_{j+1}, \mathbb{C})$. The ideal F is proper, consequently, $F_j \neq H_j$ as the \mathbb{C} -linear subspace for each $j \in \mathbb{N}_o$, where $F_j = \pi_j(F)$.

There are linear continuous operators from $l_2(\mathbb{C})$ into $l_2(\mathbb{C})$ given by the following formulas: $x \mapsto (0, \dots, 0, x^0, x^1, x^2, \dots)$ with 0 as n coordinates at the beginning, $x \mapsto (x^n, x^{n+1}, x^{n+2}, \dots)$ for $n \in \mathbb{N}$; $x \mapsto (x^{k_l + \sigma_k(i)} : k \in \mathbb{N}_o, i \in (0, 1, \dots, l-1))$, where $\mathbb{N} \ni l \geq 2$, $\sigma_k \in S_l$ are elements of the symmetric group S_l of the set $(0, 1, \dots, l-1)$. Then $f \star (g \star h) + l_2(\mathbb{C})$ and $(f \star g) \star h + l_2(\mathbb{C})$ are considered as the same class, also $f \star g + l_2(\mathbb{C}) = g \star f + l_2(\mathbb{C})$ in $H/l_2(\mathbb{C})$, since $(f + l_2(\mathbb{C})) \star (g + l_2(\mathbb{C})) = f \star g + l_2(\mathbb{C})$ for each f, g and $h \in H$. For each $f, g, h \in F$: $f \star (g \star h) + l_2(\mathbb{C})$ and $(f \star g) \star h + l_2(\mathbb{C})$ are considered as the same class, also $f \star g + l_2(\mathbb{C}) = g \star f + l_2(\mathbb{C})$ in $F/l_2(\mathbb{C})$, since $(f + l_2(\mathbb{C})) \star (g + l_2(\mathbb{C})) = f \star g + l_2(\mathbb{C}) \subset F$ for each f and $g \in F$. Therefore, the quotient algebras $H/l_2(\mathbb{C})$ and $F/l_2(\mathbb{C})$ are associative commutative Banach algebras.

Let us adjoin a unit to $H/l_2(\mathbb{C})$ and $F/l_2(\mathbb{C})$. As a consequence of the Gelfand and Mazur theorem we have, that $(H/l_2(\mathbb{C})) / (F/l_2(\mathbb{C}))$ is isomorphic with \mathbb{C} (see theorem V.6.12 [6] and theorem III.11.1 [22]). On the other hand, as it was proved above $F_j \neq H_j$ for each $j \in \mathbb{N}_o$, hence there exists the following embedding $l_2(\mathbb{C}) \hookrightarrow (H/F)$ and $(H/F) / l_2(\mathbb{C})$ is isomorphic with $(H/l_2(\mathbb{C})) / (F/l_2(\mathbb{C}))$. Therefore, H/F is isomorphic with $l_2(\mathbb{C})$.

References

- [1] N. Bourbaki. Lie groups and algebras (Moscow: Mir, 1976).
- [2] Yu.L. Dalecky, S.V. Fomin. Measures and differential equations in infinite-dimensional space (Kluwer: Dordrecht, The Netherlands, 1991).
- [3] Yu.L. Daletskii, Ya.I. Shnaiderman. Diffusion and quasi-invariant measures on infinite-dimensional Lie groups. *Funct. Anal. and its Applications*. **3** (1969), 156-158.
- [4] R. Engelking. General topology (Moscow: Mir, 1986).
- [5] H. Federer. Geometric measure theory (Berlin: Springer-Verlag, 1969).
- [6] J.M.G. Fell, R.S. Doran. Representations of $*$ -algebras, locally compact groups, and Banach $*$ -algebraic bundles (Acad. Pr.: Boston, 1988).
- [7] E. Hewitt and K.A. Ross. Abstract harmonic analysis. Second edition (Berlin: Springer-Verlag, 1979).
- [8] A.V. Kosyak. Irreducible Gaussian representations of the group of the interval and circle diffeomorphisms. *J. Funct. Anal.* **125**(1994), 493-547.
- [9] H.-H. Kuo. Gaussian measures in Banach spaces (Springer, Berlin, 1975).
- [10] S.V. Ludkovsky. Measures on groups of diffeomorphisms of non-Archimedean Banach manifolds, *Usp. Mat. Nauk.* **51**(1996), 169-170 (*N^o 2*).
- [11] S.V. Ludkovsky. Measurability of representations of locally compact groups. *Math. Sb.* **186** 1995, 83-92 (*N^o 2*).
- [12] S.V. Ludkovsky. Measures on groups of diffeomorphisms of non-Archimedean manifolds, representations of groups and their applications. *Theoret. i Mathem. Phys.*, 1999.
- [13] S.V. Ludkovsky. Quasi-invariant measures on non-Archimedean semi-groups of loops. *Usp. Mat. Nauk.* **53** (1998), 203-204 (*N^o 3*).

- [14] S.V. Ludkovsky. Quasi-invariant measures on a group of diffeomorphisms of an infinite-dimensional real manifold and induced irreducible unitary representations. *Rendiconti dell'Istituto di Matematica dell'Università di Trieste. Nuova Serie.* 26 pages, is accepted for publication, 1999.
- [15] S.V. Ludkovsky. Quasi-invariant measures on loop groups of Riemann manifolds. *Dokl. Ross. Acad. Nauk*, to appear.
- [16] S.V. Ludkovsky. Gaussian quasi-invariant measures on loop groups and semigroups of real manifolds and their representations. I.H.E.S., Bures, France, preprint **IHES/M/97/95**.
- [17] S.V. Ludkovsky. Quasi-invariant measures on non-Archimedean groups and semigroups of loops and paths, their representations. **IHES/M/98/36**.
- [18] S.V. Ludkovsky. Quasi-invariant measures on groups of diffeomorphisms of Schwarz class of smoothness for real manifolds. **IHES/M/97/96**.
- [19] S.V. Ludkovsky. Quasi-invariant and pseudo-differentiable measures on a non-Archimedean Banach space. Intern. Centre for Theoret. Phys. Trieste, Italy. Preprint (<http://www.ictp.trieste.it>) **IC/96/210**, October 1996.
- [20] S.V. Ludkovsky. Quasi-invariant measures on a non-Archimedean group of diffeomorphisms and on a Banach manifold. ICTP. **IC/96/215**, October, 1996.
- [21] S.V. Ludkovsky. Quasi-invariant measures on groups of diffeomorphisms of real Banach manifolds. ICTP. **IC/96/218**, October, 1996.
- [22] M.A. Naimark. *Normed rings* (Moscow: Nauka, 1968).
- [23] Yu.A. Neretin. Representations of the Virasoro algebra and affine algebras. in: *Itogi Nauki i Tech. Ser. Sovr. Probl. Math. Fund. Napravl*(Moscow: Nauka) **22**(1988), 163-230.
- [24] E.T. Shavgulidze. About one measure quasi-invariant relative to an action of a diffeomorphisms group of a finite-dimensional manifold. *Dokl. Akad. Nauk SSSR.* **303**(1988), 811-814.
- [25] A.V. Skorohod. *Integration in the Hilbert space* (Moscow: Nauka, 1975).