

ANNALI DELLA
SCUOLA NORMALE SUPERIORE DI PISA
Classe di Scienze

L. FUCHS

On quasi-injective modules

Annali della Scuola Normale Superiore di Pisa, Classe di Scienze 3^e série, tome 23, n° 4 (1969), p. 541-546

http://www.numdam.org/item?id=ASNSP_1969_3_23_4_541_0

© Scuola Normale Superiore, Pisa, 1969, tous droits réservés.

L'accès aux archives de la revue « Annali della Scuola Normale Superiore di Pisa, Classe di Scienze » (<http://www.sns.it/it/edizioni/riviste/annaliscienze/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>

ON QUASI-INJECTIVE MODULES

By L. FUCHS

Dedicated to B. H. NEUMANN on his 60 th birthday

The purpose of this note is to point out certain analogies between injective and quasi-injective left modules over an arbitrary ring R with identity.

We shall show that quasi-injective modules can be characterized in the same way as injective modules M by the extensibility of homomorphisms $L \rightarrow M$ (where L is a left ideal of R) to $R \rightarrow M$, but in the quasi-injective case only homomorphisms are admitted whose kernels contain the annihilator left ideal of some $a \in M$.

The notion of K -bounded module (with K an ideal of R) is introduced as a module M which is annihilated by K and which contains an element whose annihilator is exactly K . For K -bounded modules quasi-injectivity turns out to be equivalent to R/K -injectivity. A K -bounded module is a direct summand of every module containing it as a pure submodule where purity can be taken in two, inequivalent ways.

Finally, the so-called exchange property will be proved for quasi-injective modules.

1. By a ring we mean an associative ring with 1 and by a module a unital left module over a ring R .

An R module M is said to be *quasi-injective* ⁽¹⁾ if every R -homomorphism of every R submodule of M into M is induced by an R -endomorphism of M . A module is quasi-injective exactly if it is a fully invariant submodule of its injective hull.

For an R -module M , we denote by $\Omega(M)$ the set of all left ideals L of R such that L contains $\text{Ann } a = \{r \in R \mid ra = 0\}$ for some $a \in M$.

Pervenuto alla Redazione il 24 Dicembre 1968.

⁽¹⁾ For properties of quasi-injective modules we refer e.g. to Faith [5].

LEMMA 1. *The following conditions on an R -module M are equivalent:*

- (i) M is quasi-injective;
- (ii) if B is a submodule of a cyclic submodule $A = Ra$ of M and if $\beta: B \rightarrow M$ is any R -homomorphism, then there is an extension $\alpha: A \rightarrow M$ of β ;
- (iii) if B is a submodule of any R -module A with $\Omega(A) \subseteq \Omega(M)$, then every R -homomorphism $\beta: B \rightarrow M$ can be extended to an R -homomorphism $\alpha: A \rightarrow M$.

The implication (i) \implies (ii) is trivial. To prove (ii) \implies (iii), assume (ii) and let A, B, β be given as in (iii). We use the standard argument and consider submodules C of A and homomorphisms $\gamma: C \rightarrow M$ such that $B \subseteq C \subseteq A$ and $\gamma|_B = \beta$. If the pairs (C, γ) are ordered in the obvious way, then we can pick out a maximal pair (C_0, γ_0) in the set of pairs (C, γ) . By way of contradiction, suppose there is an $a \in A$ not in C_0 .

Clearly, $L = \{r \in R \mid ra \in C_0\}$ is a left ideal of R contained in $\Omega(A)$, and hence in $\Omega(M)$. Choose some $x \in M$ such that $L \supseteq \text{Ann } a \supseteq \text{Ann } x$, and consider the submodule $N = Lx$ of M . The correspondence $rx \mapsto \gamma_0(ra)$ with $r \in L$ defines a homomorphism $\varphi': N \rightarrow M$ which can be extended, in view of our hypothesis (ii), to a homomorphism $\varphi: Rx \rightarrow M$. Now let $C' = C_0 + Ra$ and let $\gamma': C' \rightarrow M$ be defined as $\gamma': c + ra \mapsto \gamma_0(c) + \varphi(rx)$ for $c \in C_0$, $r \in R$. It is easy to check that γ' is a well-defined homomorphism, so $(C_0, \gamma_0) < (C', \gamma')$ contradicts the maximal choice of (C_0, γ_0) . Hence $C_0 = A$ and $\gamma_0 = \alpha$ is an extension of β .

The choice $A = M$ in (iii) yields (i). This completes the proof.

Condition (ii) may be reformulated to give a characterization of quasi-injectivity which is similar to a well-known characterization of injectivity [1].

LEMMA 2. *An R -module M is quasi-injective if and only if for every left ideal L of R and for every R -homomorphism $\eta: L \rightarrow M$ with $\text{Ker } \eta \in \Omega(M)$ there exists an R -homomorphism $\psi: R \rightarrow M$ that extends η ⁽²⁾.*

If $a \in M$ is such that $\text{Ann } a \subseteq \text{Ker } \eta$, then η induces an R -homomorphism $\beta: La \rightarrow M$, and the equivalence with (ii) becomes evident.

In connection with Lemma 2 let us notice that $\Omega(M)$ can be replaced by the filter (i.e. the dual ideal) $\bar{\Omega}(M)$ generated by $\Omega(M)$ in the lattice of all submodules of M . In fact, if M is quasi-injective, then so is $M \oplus \dots \oplus M$ with a finite number n of summands and $\Omega(M \oplus \dots \oplus M)$ contains all $L_1 \cap \dots \cap L_n$ with $L_j \in \Omega(M)$. Together with $M \oplus \dots \oplus M$ also M must have the property stated in Lemma 2 for the finite intersections $L_1 \cap \dots \cap L_n$.

⁽²⁾ Notice that the stated condition makes sense only for $L \in \Omega(M)$.

2. Next we introduce the notion of bounded modules.

Let K be a left ideal of R . An R -module M will be called K -bounded if $\Omega(M)$ consists exactly of the left ideals of R which contain K . That is, $K \in \Omega(M)$ is the unique minimal element of $\Omega(M)$, or, in other words, $\Omega(M)$ is the filter generated by K ⁽³⁾.

It follows at once that K must be two-sided, since it is the annihilator of M . We can thus form the factor ring R/K and may consider M as an R/K -module in the obvious way. If we do so, then we are led to

THEOREM 1. *A K -bounded R -module M is quasi-injective if and only if it is injective as an R/K -module.*

In the K -bounded case, the condition in Lemma 2 amounts to R/K -injectivity. Hence Theorem 1 holds.

For $K = 0$, we have: a 0-bounded quasi-injective is injective.

If we drop the hypothesis of K -boundedness, then — under rather restrictive conditions — a similar result can be established with R/K replaced by a topological ring which is constructed as an inverse limit [4].

3. Following Cohn [2], we call a submodule N of the R -module M *pure* if for all right R modules U , the homomorphism $U \otimes_R N \rightarrow U \otimes_R M$ [induced by the inclusion $N \rightarrow M$] is monic. This is equivalent to the following condition which is more suitable for our purposes: if

$$(1) \quad \sum_{j=1}^n r_{ij} x_j = a_i \in N \quad (i = 1, 2, \dots, m)$$

is a finite set of equations in the unknowns x_1, \dots, x_n where $r_{ij} \in R$, and if this system has a solution in M , then it has a solution in N too.

An R -module A is called *algebraically compact* (see [6], [8]) if it is a direct summand in every R -module in which it is a pure submodule. Or, equivalently, if

$$(2) \quad \sum_j r_{ij} x_j = a_i \in A \quad (i \in I)$$

is an arbitrary set of equations in the unknowns x_j ($j \in J$) [where I and J are arbitrary index sets and each equation contains but a finite number of

⁽³⁾ For Z -modules, i.e. for abelian groups, K -boundedness means that the group is a direct sum of cyclic groups of order n and orders m dividing n or it contains an element of infinite order, according as $K = (n)$ or $K = (0)$.

non-zero $r_{ij} \in R$], and if every finite set of equations in (2) has a solution in A , then the entire system (2) is solvable in A .

THEOREM 2. *A K -bounded quasi-injective R -module is algebraically compact.*

Let A be a K -bounded quasi-injective R -module and (2) a system of equations which is finitely solvable in A . If we consider A as an R/K -module and replace r_{ij} by $r_{ij} + K = \overline{r_{ij}}$, then (2) may be viewed as a system of equations over the R/K -module A . Finite solvability implies that this system is compatible in the sense of Kertész [7], thus it has a solution in the injective R/K -module A . This is evidently a solution of the original form (2), hence the algebraic compactness of A follows.

Algebraic compactness is preserved under direct products and direct summands, hence

COROLLARY 1. *Let M_i ($i \in I$) be K_i -bounded quasi-injective R -modules and M a direct summand of their direct product $\prod M_i$. Then M is algebraically compact⁽⁴⁾.*

4. There are various definitions of purity for modules which all reduce to ordinary purity for abelian groups. We are going to show that Theorem 2 holds if we replace purity in the sense of P. M. Cohn by the following one.

A submodule N of the R -module M is now called *pure* if

$$LN = N \cap LM$$

for all two-sided ideals L of R . Algebraic compactness can be defined in the same way as in 3 by using this definition of purity.

Next we prove Theorem 2 for this algebraic compactness. Let A be a K -bounded quasi-injective R -module and let M contain A as a pure submodule. The module M/KM is annihilated by K , thus $\Omega(M/KM)$ contains only left ideals containing K . In view of $0 = KA = A \cap KM$, the natural homomorphism $\varphi: M \rightarrow M/KM$ maps A isomorphically upon φA which is thus quasi-injective. The two last sentences imply, by Lemma 1, that the identity map of φA extends to a homomorphism $M/KM \rightarrow \varphi A$ showing that $M/KM = \varphi A \oplus N/KM$ for a submodule N of M . Hence $M = A \oplus N$, and A is algebraically compact.

⁽⁴⁾ Notice that for abelian groups the converse also holds: every algebraically compact group is a summand of a direct product of K -bounded quasi-injectives.

5. Next we turn our attention to the so-called exchange property which was systematically discussed by Crawley and Jónsson [3].

Recall that an R -module M is said to have the *exchange property* if for every R -module A containing M and for submodules N and $A_i (i \in I)$ of A , the direct decomposition

$$(3) \quad A = M \oplus N = \bigoplus_{i \in I} A_i \quad (I = \text{arbitrary index set})$$

implies the existence of R -submodules B_i of $A_i (i \in I)$ satisfying

$$(4) \quad A = M \oplus \left(\bigoplus_{i \in I} B_i \right).$$

It is known [3] that M has the exchange property if it has the stated property with the A_i subject to the condition that each A_i is isomorphic to a submodule of M . The following result generalizes a theorem of Warfield [9] from injectives to quasi-injectives.

THEOREM 3. *A quasi-injective module has the exchange property.*

Let M be a quasi-injective R -module and assume (3) holds for R -modules $N, A_i (i \in I)$ with A_i isomorphic to submodules of M . Select a submodule B of A which is maximal with respect to the properties: (i) $B = \bigoplus_i B_i$ with $B_i \subseteq A_i$, and (ii) $M \cap B = 0$. We claim (4) holds with these B_i .

Denote by φ the natural homomorphism $A \rightarrow A/B$. Because of (ii), $\varphi|_M$ is monic, so $\varphi(M)$ is a quasi-injective submodule of the R -module $A/B = \bigoplus_i A_i/B_i$ where A_i/B_i has been identified with $(A_i + B)/B$ under the canonical isomorphism. The maximal choice of B guarantees that no A_i/B_i has a non-zero submodule with 0 intersection with $\varphi(M)$, that is, $\varphi(M) \cap (A_i/B_i)$ is essential in A_i/B_i , and so $\bigoplus_i [\varphi(M) \cap (A_i/B_i)]$ and a fortiori $\varphi(M)$ is essential in A/B . Now φ maps A_i into A/B , but since A_i is isomorphic to a submodule of $\varphi(M)$ and $\varphi(M)$ is fully invariant in its injective hull (which contains A/B), we see that $\varphi(A_i)$ must be contained in $\varphi(M)$. Consequently, φ maps the whole A into $\varphi(M)$, i. e. $\varphi(M) = A/B$, so M and B generate A . This proves $A = M \oplus B$.

An immediate consequence is:

COROLLARY 2. *Assume that*

$$A = M_1 \oplus \dots \oplus M_m = \bigoplus_{i \in I} N_i$$

are two direct decompositions of an R -module A where every M_j and every

N_i is a quasi-injective R -module, and I is an arbitrary index set. Then they have isomorphic refinements, i. e. there exist R -modules A_{ji} ($j = 1, \dots, m; i \in I$) such that

$$M_j \cong \bigoplus_{i \in I} A_{ji} \quad \text{and} \quad N_i \cong A_{1i} \oplus \dots \oplus A_{mi}$$

for every j and i , respectively.

An application of Theorem 3 yields $A = M_1 \oplus (\bigoplus_i N_i')$ for submodules N_i' of N_i . Write $N_i = N_i' \oplus A_{1i}$ to get $M_1 \cong \bigoplus_{i \in I} A_{1i}$ and $M_2 \oplus \dots \oplus M_m \cong \bigoplus_i N_i'$. A simple induction completes the proof.

It is an open problem whether or not two infinite decompositions have isomorphic refinements ⁽⁵⁾.

⁽⁵⁾ This holds for injectives as was shown by Warfield [9].

REFERENCES

- [1] H. CARTAN and S. EILENBERG, *Homological algebra* (Princeton, 1956).
- [2] P. M. COHN, *On the free product of associative rings*, Math. Z. 71 (1959), 380-398.
- [3] P. CRAWLEY and B. JÓNSSON, *Refinements for infinite direct decompositions of algebraic system*, Pac. Journ. Math. 14 (1964), 797-855.
- [4] S. EILENBERG and N. STEENROD, *Foundations of algebraic topology* (Princeton, 1952).
- [5] C. FAITH, *Lectures on injective modules and quotient rings*, Lecture Notes in Mathematics, vol. 49 (1967).
- [6] L. FUCHS, *Algebraically compact modules over Noetherian rings*, Indian Journ. Math. 9 (1967), 357-374.
- [7] A. KERTÉSZ, *Systems of equations over modules*, Acta Sci. Math. Szeged 18 (1957), 207-234.
- [8] R. B. WARFIELD, JR., *Purity and algebraic compactness for modules*. Pac. Journ. Math. 28 (1969), 699-720.
- [9] R. B. WARFIELD, JR., *Decompositions of injective modules*.