

Astérisque

ALAIN ESCASSUT

PHILIPPE ROBBA

Summary

Astérisque, tome 10 (1973), p. 219-220

http://www.numdam.org/item?id=AST_1973__10__219_0

© Société mathématique de France, 1973, tous droits réservés.

L'accès aux archives de la collection « Astérisque » (<http://smf4.emath.fr/Publications/Asterisque/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

SUMMARY

Alain ESCASSUT

In particular, we shall show that the integral algebras of Krasner-Tate are Tate algebras of the form $\frac{K\{T, X\}}{P(x) - T Q(x)}$ where P and Q are 2 unitary and relatively prime polynoms with coefficients of absolute value ≤ 1 and that these algebras are characterised among ultrametric Banach algebras by five algebraic and topologic properties. We shall prove that if D is the spectrum of an element of an ultrametric Banach algebra and if the set of the infraconnected components is finite, then every infraconnected component has an associated idempotent in A . At last, we shall characterise the reduced Tate algebras of degree 1 among the ultrametric Banach algebras with the help of their algebraic and topologic properties.

Philippe ROBBA

The uniform limit of rational functions on a subset A of a non-archimedean field K is called an analytic element on A if the functions have no poles inside A (the idea is due to Krasner). Given a family \mathcal{A} of subsets of K , if a function f is defined on a chained union $\bigcup_i A_i$ of members of \mathcal{A} and $f|_{A_i}$ is an analytic element for each i , then f is said to be a \mathcal{A} -analytic function.

Let \mathcal{Q} be the most general class of subsets of K for which \mathcal{Q} -analytic functions verify the principle of analytic continuation. Then \mathcal{Q} has been completely determined by Escassut, Motzkin and the author [6].

But the class of all \mathcal{Q} -analytic functions is not stable under the elementary operations of algebra and analysis. We show that such a stability can be obtained either by restricting the class \mathcal{Q} (in various ways : see §§ 8, II) or by

imposing stronger conditions on K (e.g., K must be maximally complete if we want an analytic function to be representable by Laurent series in an annulus (§ 10) , or if we want a uniform limit of analytic functions to be analytic (§ II)).

An essential tool in this discussion will be the Theorem of representation of an analytic element as a sum of its singular parts (§ 4).

We obtain necessary conditions for the analytic continuation of a Taylor series outside its disk of convergence, and we give a general, if not very practical, constructive procedure for obtaining the continuation (§ 15).

We also obtain a factorization of meromorphic functions according to their zeros and poles analogous to Hensel's Theorem for analytic elements (§ 13).