

# Astérisque

EIRA J. SCOURFIELD

**On the property  $(f(n), g(n)) = 1$  for certain  
multiplicative functions**

*Astérisque*, tome 41-42 (1977), p. 273-276

[http://www.numdam.org/item?id=AST\\_1977\\_\\_41-42\\_\\_273\\_0](http://www.numdam.org/item?id=AST_1977__41-42__273_0)

© Société mathématique de France, 1977, tous droits réservés.

L'accès aux archives de la collection « Astérisque » (<http://smf4.emath.fr/Publications/Asterisque/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme  
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

ON THE PROPERTY  $(f(n), g(n)) = 1$  FOR CERTAIN MULTIPLICATIVE FUNCTIONS

by

Eira J. SCOURFIELD

The problem of investigating the sum

$$\Sigma_h(x) = \sum_{\substack{n \leq x \\ (n, h(n))=1}} 1 \quad (1)$$

for certain integer-valued arithmetic functions  $h$  has been considered by several authors in cases when the arithmetic properties of  $n$  and  $h(n)$  are not too closely related, and the expected result

$$\Sigma_h(x) \sim 6x/\pi^2$$

established ; for references, see this author's paper [2] . Multiplicative functions, however, present a rather different problem, and in 1948 [1], Erdős obtained the result

$$\Sigma_{\varphi}(x) \sim e^{-\gamma} x / \log \log \log x \quad \text{as } x \rightarrow \infty$$

for Euler's function  $\varphi$  . In [2], we considered the sum (1) for a class of integer-valued multiplicative functions, called polynomial-like, that includes  $\varphi$  and the divisor functions  $\sigma_{\nu}$  ( $\nu \geq 0$ ) ;  $f$  is polynomial-like if there exists a polynomial  $W \in \mathbb{Z}[x]$  such that

$$f(p) = W(p) \quad \text{for all primes } p. \quad (2)$$

For these functions, we proved in [2] :

THEOREM 1. If the polynomial  $W$  of (2) satisfies  $\deg W > 0$  ,  $W(0) \neq 0$  , then there exist constants  $C > 0$  ,  $\lambda$  ( $0 < \lambda \leq 1$  ,  $\lambda$  rational), depending on  $f$  , such that

$$\Sigma_f(x) \sim C x (\log \log \log x)^{-\lambda} \quad \text{as } x \rightarrow \infty .$$

If  $W$  is a non-zero constant, then there exists a constant  $C$  ( $0 < C \leq 1$ ) such that

$$\Sigma_f(x) \sim C x \quad \text{as } x \rightarrow \infty .$$

If  $W(0) = 0$  ,

$$\Sigma_f(x) = O(x^{\frac{1}{2}}) .$$

Example.  $f = \sigma_\nu$  ( $\nu > 0$ ) . For  $\nu$  odd,  $\lambda = 1$  ,  $C = e^{-\gamma}$  , whilst for  $\nu$  even,  $\lambda = 2^{-\beta}$  , where  $2^\beta \parallel \nu$  .

We obtain a generalization of the sum in theorem 1 by noting that  $n$  itself is a polynomial-like multiplicative function. Let  $f$  ,  $g$  be multiplicative polynomial-like functions, and let  $W_1$  ,  $W_2 \in \mathbb{Z}[x]$  be the polynomials such that

$$f(p) = W_1(p) , \quad g(p) = W_2(p) \quad \text{for all primes } p .$$

Suppose that the following conditions hold :

- (i)  $\deg W_i > 0$  ( $i = 1, 2$ ) ;
- (ii)  $W_1(x) = x^\alpha W_1^*(x)$  where  $\alpha \geq 0$  ,  $W_1^*(0) \neq 0$  ,  $\deg W_1^* > 0$  , and  $W_2(0) \neq 0$  ;
- (iii)  $W_1$  ,  $W_2$  are coprime polynomials.

It follows from (iii) that the set

MULTIPLICATIVE FUNCTIONS

$$S_0 = \{p : p | (f(q), g(q)) \text{ for all primes } q \neq p\}$$

of primes is finite (possibly empty). If  $p \in S_0$ ,  $p | (f(n), g(n))$  whenever there exists a prime  $q \neq p$  with  $q | n$ , and hence for "most"  $n$ . This suggests that for our generalization of the sum  $\Sigma_f(x)$ , we consider

$$\Sigma_{f,g}(x) = \sum_{\substack{n \leq x \\ p | (f(n), g(n)) \forall p \notin S_0}} 1.$$

Using results from sieve theory, we can prove

THEOREM 2. If conditions (i), (ii), (iii) above hold, there exist constants  $C > 0$ ,  $\lambda$  ( $0 < \lambda \leq 1$ ,  $\lambda$  rational) such that

$$\frac{x}{\log x} \log \log x \ll \Sigma_{f,g}(x) \ll \frac{x}{\log x} \exp\left(\frac{C \log \log x}{(\log \log \log x)^\lambda}\right).$$

Conditions (i), (ii), (iii) ensure that the sum  $\Sigma_{f,g}(x)$  is not too small and does not reduce to the sum considered in theorem 1 or in other published papers.

Examples. (i)  $f = \varphi$ ,  $g = \sigma_\nu$  ( $\nu > 0$ ), when  $S_0 = \{2\}$ ,  $\lambda = 2^{-\beta}$  where  $2^\beta || \nu$ .  
(ii)  $f = \sigma_\nu$ ,  $g = \sigma_\kappa$  ( $\nu, \kappa > 0$ ,  $\beta > \gamma$ ), where  $2^\beta || \nu$ ,  $2^\gamma || \kappa$ , when  $S_0 = \{2\}$ ,  $\lambda = 2^{-\beta}$ .

The method used to prove the upper bound in theorem 2 also establishes

THEOREM 3. The number of positive integers  $n \leq x$  with the property that  $n$  does not have a prime divisor in every residue class (mod  $p$ ) coprime to  $p$  for any odd prime  $p$  is

$$\ll \frac{x}{\log x} \exp\left(\frac{B \log \log x}{\log \log \log x}\right),$$

where  $B > 0$  is constant.

--:--:--

BIBLIOGRAPHY

- [1] P. ERDÖS.- Some asymptotic formulas in number theory. *J. Indian Math. Soc.* (N.S.), 12 (1948), pp. 75-78.
- [2] E.J. SCOURFIELD.- An asymptotic formula for the property  $(n, f(n)) = 1$  for a class of multiplicative functions. *Acta Arithmetica*, 29 (1976), pp. 401-423.

Eira J. SCOURFIELD  
Westfield College  
Department of Mathematics  
LONDON NW3 7ST, England U.K.