

Astérisque

AST Summary

Astérisque, tome 215 (1993), p. 205

<http://www.numdam.org/item?id=AST_1993__215__205_0>

© Société mathématique de France, 1993, tous droits réservés.

L'accès aux archives de la collection « Astérisque » ([http://smf4.emath.fr/
Publications/Asterisque/](http://smf4.emath.fr/Publications/Asterisque/)) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

*Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>*

Summary

Let G be a real Lie group acting on a manifold M . H. Cartan introduced the equivariant de Rham complex and its cohomology $H_G^*(M)$. For a free action, $H_G^*(M)$ is the cohomology $H^*(G \backslash M)$ of the orbit space and if M is the point \bullet , $H_G^*(\bullet)$ is the algebra of invariant polynomial functions on the Lie algebra \mathfrak{g} of G . To G -equivariant vector bundles on M , equipped with a G -invariant connection, are associated equivariant Chern classes. It became necessary to consider more general cohomological objects, like the algebra $H_G^\infty(M)$ of equivariant cohomology with C^∞ -coefficients, which is an algebra over $H_G^\infty(\bullet) = C^\infty(\mathfrak{g})^G$, and also the space $H_G^{-\infty}(M)$ of equivariant cohomology with $C^{-\infty}$ -coefficients, which is a module for $H_G^\infty(M)$, and for which $H_G^{-\infty}(\bullet)$ is the space of invariant generalized functions $C^{-\infty}(\mathfrak{g})^G$.

The first of the two articles of this volume, *Cohomologie équivariante et descente*, by Michel Duflo and Michèle Vergne, studies a generalization, denoted by $\mathcal{K}_G(M)$, of the cohomology $H_G^\infty(M)$. It is an algebra over $\mathcal{K}_G(\bullet) = C^\infty(G)^G$. It can be considered as a global analog of $H_G^\infty(M)$, and also as a de Rham version of the equivariant K -theory of M . The construction of $\mathcal{K}_G(M)$ uses the fixed point sets in M of the elements of G which are contained in compact subgroups of G . At least when G is compact, and under certain orientation assumptions, the “integral” over M of an element of $\mathcal{K}_G(M)$ is a G -invariant function on G .

The second article, *Equivariant cohomology with generalized coefficients*, by Shrawan Kumar and Michèle Vergne, undertakes a systematic study of the spaces $H_G^{-\infty}(M)$. Remarkable classes, with no analogs in the C^∞ -theory, are discovered. In particular, for a free action, the integral over M of an element of $H_G^{-\infty}(M)$ is a generalized function on \mathfrak{g} with support 0. When G is compact, a spectral sequence allows a comparison of $H_G^{-\infty}(M)$ with the equivariant cohomology $H_G^*(M)$.

The two articles have common motivations, but they can be read independently.