

# COMPOSITIO MATHEMATICA

MICHAEL P. ANDERSON

**Some finiteness properties of the fundamental  
group of a smooth variety**

*Compositio Mathematica*, tome 31, n° 3 (1975), p. 303-308

<[http://www.numdam.org/item?id=CM\\_1975\\_\\_31\\_3\\_303\\_0](http://www.numdam.org/item?id=CM_1975__31_3_303_0)>

© Foundation Compositio Mathematica, 1975, tous droits réservés.

L'accès aux archives de la revue « Compositio Mathematica » (<http://http://www.compositio.nl/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme  
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

**SOME FINITENESS PROPERTIES OF  
THE FUNDAMENTAL GROUP OF A SMOOTH VARIETY**

Michael P. Anderson

In this paper we prove that for any smooth variety  $X$  over an algebraically closed field of characteristic  $p \neq 2, 3, 5$  the group  $\prod_1^{(p)}(X)$  is a finitely presented pro- $(p)$ -group. We recall that  $\prod_1^{(p)}(X)$  denotes the maximal quotient of  $\prod_1(X)$  of order prime to  $p$ . In [8] Exposé II this result is demonstrated for smooth  $X$  provided there exists a projective smooth compactification  $\bar{X}$  of  $X$  such that  $\bar{X} \setminus X$  is a divisor with normal crossings on  $\bar{X}$  and for all  $X$  provided we assume strong resolution of singularities for all varieties of dimension  $\leq n$ . Thus the result was previously known for  $X$  of dimension  $\leq 2$ .

The essential new step is Lemma 1 which allows us to reduce to the case of dimension 2. The proof of this lemma uses Abhyankar's work on resolution of singularities [1] together with the technique of fibering by curves. We follow the notation of [7] Exposé XIII and [8] Exposé II.

Let us now state our proposition.

**PROPOSITION 1:** *Let  $X/k$  be a connected smooth variety over the algebraically closed field  $k$  of characteristic  $p \neq 2, 3, 5$ . Then  $\prod_1^{(p)}(X)$  is a finitely presented pro- $(p)$ -group.*

**PROOF:** By [7] Exposé IX it is sufficient to prove the result for the elements of a Zariski covering of  $X$ . Thus the result follows by induction on dimension from the result in dimension 2, [8] Exposé II Theorem 2.3.1, and the following lemma:

**LEMMA 1:** *Let  $X$  be a smooth variety of dimension  $n \geq 3$  over the algebraically closed ground field  $k$  and  $x$  a point of  $X$ . Then  $x$  has a Zariski neighborhood  $U$  such that there exists an algebraically closed extension  $\Omega/k$  and a smooth variety  $V$  over  $\Omega$  of dimension  $n-1$  and a morphism  $f: V \rightarrow U$  such that  $f$  induces a surjection  $\prod_1(V) \rightarrow \prod_1(U)$  and an isomorphism  $\prod_1^{(p)}(V) \rightarrow \prod_1^{(p)}(U)$ .*

PROOF OF LEMMA 1: We proceed by induction on the dimension of  $X$ .

Let  $U$  be an affine neighborhood of  $x$ . By [1] *Birational Resolution* there exists a smooth projective model of the function field  $k(U)$ . Let  $\bar{U}$  be a projective compactification of  $U$ . By [1] *Dominance* there exists a smooth projective variety  $X'$  together with a birational morphism  $X' \rightarrow \bar{U}$ . By [1] *Global Resolution* there exists a smooth projective variety  $X''$  together with a birational morphism  $X'' \rightarrow \bar{U}$  and such that the inverse image of  $\bar{U} \setminus U$  is a divisor with normal crossings on  $X''$ . Let  $U''$  be the complement of this divisor. Then the map  $g : U'' \rightarrow U$  is a proper birational mapping of smooth varieties. The subvariety of points of  $U$  where  $g$  is not an isomorphism is of codimension  $\leq 2$ . Thus by the Purity Theorem [7] Exposé X.3,  $g$  induces an isomorphism

$$\prod_1(U'') \rightarrow \prod_1(U).$$

By [9], [5], or [10], a general hyperplane section of  $U''$ , call it  $V$ , gives a smooth surface in  $U''$  such that

$$\prod_1^{(p)}(V) \simeq \prod_1^{(p)}(U'') \simeq \prod_1^{(p)}(U).$$

Thus the lemma is proved for  $n = 3$ .

Now assume  $n > 3$ . By [4] Exposé XI,  $x$  has a Zariski neighborhood  $W$  which admits an elementary fibration  $g : W \rightarrow W'$  with  $W'$  smooth of dimension  $n - 1$ . Moreover, by [6] Proposition 2.8 we may assume that  $g$  admits a finite etale multisection i.e. there exists a finite etale map  $s : S \rightarrow W'$  together with a closed immersion  $i : S \rightarrow W$  such that  $gi = s$ . Let  $y = g(x)$ . By induction  $y$  admits a Zariski neighborhood  $U'$  in  $W'$  such that there exists a smooth variety  $V'$  of dimension  $n - 2$  and a morphism  $f' : V' \rightarrow U'$  such that  $f'$  induces an isomorphism of the  $(p)$ -completions of the fundamental groups of  $V'$  and  $U'$ . Let  $U = g^{-1}(U')$  and  $V = V' \times_{V'} U$  with projections  $f : V \rightarrow U$  and  $g' : V \rightarrow V'$ . Then  $g'$  is an elementary fibration admitting an etale multisection. Letting  $C$  be a geometric fiber of  $g'$ , we have, by [7] Exposé XIII Proposition 4.3, exact sequences

$$\begin{array}{ccccccc} e & \rightarrow & \prod_1^{(p)}(C) & \rightarrow & \prod_1^{(p)}(V) & \rightarrow & \prod_1^{(p)}(V') \rightarrow e \\ & & \parallel & & \downarrow & & \downarrow \\ e & \rightarrow & \prod_1^{(p)}(C) & \rightarrow & \prod_1^{(p)}(U) & \rightarrow & \prod_1^{(p)}(U') \rightarrow e. \end{array}$$

Let  $K$  be the kernel of the homomorphism  $\prod_1'(U) \rightarrow \prod_1'(V)$  and  $K'$  the kernel of  $\prod_1(V') \rightarrow \prod_1(U')$ . Then the natural map  $K \rightarrow K'$  is an isomorphism. Moreover, by hypothesis  $K'$  is contained in the closed normal subgroup of  $\prod_1(V')$  generated by the Sylow  $p$  subgroups of  $\prod_1(V')$ . Since any Sylow  $p$  subgroup of  $\prod_1(V')$  is the image of a Sylow  $p$  subgroup of  $\prod_1'(V)$ ,  $K$  is also contained in the subgroup generated by the conjugates of the Sylow  $p$  subgroups. Thus  $K$  is contained in the kernel of  $\prod_1(V) \rightarrow \prod_1^{(p)}(V)$ . Therefore the homomorphism

$$\prod_1^{(p)}(V) \rightarrow \prod_1^{(p)}(U)$$

is injective and, by the five lemma, it is surjective. Thus the lemma and proposition are proved.

Using Proposition 1 and standard descent techniques we can weaken the resolution hypotheses required to prove finite presentation of  $\prod_1^{(p)}(X)$  for arbitrary  $X$ . We shall say that a point  $x$  of a variety  $X$  admits a ‘weak resolution of singularities’ if there exists a Zariski neighborhood  $U$  of  $x$  in  $X$  and a morphism of effective descent for the category of etale coverings  $f: U' \rightarrow U$  such that  $U'$  is a smooth variety. We have then the following:

**PROPOSITION 2:** *Let  $X$  be a variety over an algebraically closed field of characteristic  $p \neq 2, 3, 5$ . Assume that every point of  $X$  admits a weak resolution of singularities. Then  $\prod_1^{(p)}(X)$  is a finitely presented pro- $(p)$ -group.*

**COROLLARY:** *Let  $X$  be a variety of dimension 3 over an algebraically closed field of characteristic  $p \neq 2, 3, 5$ . Then  $\prod_1^{(p)}(X)$  is a finitely  $p$  presented pro- $(p)$ -group.*

**PROOF:** Proposition 2 is a straightforward application of [7] IX.5 together with Proposition 1. The Corollary follows from Proposition 2 and Abhyankar’s results on resolution [1].

As another application of the fibering by curves method we will outline a proof of the following result:

**PROPOSITION 3 (Kunneth Formula):** *Let  $X$  and  $Y$  be connected varieties over the algebraically closed field  $k$  of characteristic  $p$ . Then the natural homomorphism*

$$\prod_1^{(p)}(X \times Y) \rightarrow \prod_1^{(p)}(X) \times \prod_1^{(p)}(Y)$$

is an isomorphism.

In [7] Exposé XIII this proposition is demonstrated using the hypothesis of strong resolution of singularities. We avoid the use of resolution of singularities as follows:

First we consider the case where  $X$  and  $Y$  are normal varieties. Then it is sufficient to prove the formula for some non-trivial open subsets of  $X$  and  $Y$ . Choose  $U$  in  $X$  and  $V$  in  $Y$  such that  $U$  and  $V$  admit elementary fibrations  $f : U \rightarrow U'$  and  $g : V \rightarrow V'$  with étale multisections. By induction on the dimensions of  $U$  and  $V$  we may assume the proposition holds for  $U'$  and  $V'$ . Let  $C$  and  $D$  be geometric fibers of  $f$  and  $g$  respectively. Since  $f$  and  $g$  are elementary fibrations admitting étale multisections we have the following exact sequences

$$\begin{aligned} e \rightarrow \prod_1^{(p)}(C) &\rightarrow \prod_1'(U) \rightarrow \prod_1(U) \rightarrow e \\ e \rightarrow \prod_1^{(p)}(D) &\rightarrow \prod_1'(V) \rightarrow \prod_1(V) \rightarrow e \\ e \rightarrow \prod_1^{(p)}(C \times D) &\rightarrow \prod_1'(U \times V) \rightarrow \prod_1(U \times V) \rightarrow e. \end{aligned}$$

Arguing now as in the proof of Lemma 1, we see that the natural homomorphism

$$\prod_1'(U \times V) \rightarrow \prod_1'(U) \times \prod_1'(V)$$

induces an isomorphism on  $(p)$ -completions.

Consider now the case in which  $Y$  is assumed normal, and  $X$  is arbitrary. Let  $X' \rightarrow X$  be the normalization of  $X$ , and define

$$X'' = X' \times_X X', \quad X''' = X' \times_X X' \times_X X'.$$

Let  $X'_\alpha, \alpha \in \prod_0(X')$ , be the connected components of  $X'$ . Then by [7] IX Theorem 5.1,  $\prod_1(X)$  is the free product of the groups  $\prod_1(X_\alpha)$  and the

free group generated by the elements of the set  $\prod_0(X'')$  modulo certain relations defined by the projections:

$$X''' \rightrightarrows X'' \rightrightarrows X' \rightarrow X.$$

Thus the same description holds for  $\prod_1^{(p)}(X)$  after replacing all the groups involved by their prime to  $p$  completions. Moreover, the same result applies to  $X' \times Y \rightarrow X \times Y$ . This gives a description of  $\prod_1^{(p)}(X \times Y)$  as the free product (in the category of pro- $(p)$ -groups) of the groups  $\prod_1^{(p)}(X_\alpha \times Y)$  and the free pro- $(p)$ -group generated by the elements of the set  $\prod_0(X'' \times Y)$  modulo relations defined by the projections:

$$X''' \times Y \rightrightarrows X'' \times Y \rightrightarrows X' \times Y \rightarrow X \times Y.$$

It is long and tedious, but straightforward, to check that, since

$$\prod_1^{(p)}(X_\alpha \times Y) = \prod_1^{(p)}(X_\alpha) \times \prod_1^{(p)}(Y) \quad \text{and} \quad \prod_0(X'' \times Y) = \prod_0(X''),$$

the above relations force

$$\prod_1^{(p)}(X \times Y) = \prod_1^{(p)}(X) \times \prod_1^{(p)}(Y).$$

Now applying the same argument as above without the assumption that  $Y$  is normal (which is valid because we just verified that

$$\prod_1^{(p)}(X_\alpha \times Y) = \prod_1^{(p)}(X_\alpha) \times \prod_1^{(p)}(Y)$$

for  $X_\alpha$  normal and  $Y$  arbitrary) gives the result for  $X$  and  $Y$  arbitrary varieties.

#### BIBLIOGRAPHY

- [1] S. ABHYANKAR: *Resolution of Singularities of Embedded Algebraic Surfaces*. Academic Press, New York, 1966.
- [2] MICHAEL P. ANDERSON: *Profinite Groups and the Topological Invariants of Algebraic Varieties*. Princeton Ph.D. thesis, 1974.
- [3] MICHAEL P. ANDERSON: EXACTNESS PROPERTIES OF PROFINITE COMPLETION FUNCTORS. *Topology* 13 (1974) 229–239.
- [4] M. ARTIN, A. GROTHENDIECK, J. L. VERDIER: *Theorie des Topos et Cohomologie Etale des Schemas. Lecture Notes in Mathematics* 305 (1973).

- [5] WOUT DE BRUIN: Une forme algebrique du theoreme de Zariski pour  $\Pi_1$ . *C. R. Acad. Sci. Paris Ser. A-B* 272 (1971) A769–A771.
- [6] E. M. FRIEDLANDER: The Etale Homotopy Theory of a Geometric Fibration. *Manuscripta Mathematica* 10 (1973) 209–244.
- [7] A. GROTHENDIECK et al.: Revêtements Etales et Groupe Fondamental. *Lecture Notes in Mathematics* 224 (1971).
- [8] A. GROTHENDIECK et al.: Groupes de Monodromie en Geometrie Algebrique. *Lecture Notes in Mathematics* 288 (1972).
- [9] HERBERT POPP: Ein Satz vom Lefschetzschen Typ Uber die Fundamentalgruppe quasi-projectiver Schemata. *Math. Z.* 116 (1970) 143–152.
- [10] M. RAYNAUD: Theoreme de Lefschetz en Cohomologie des Faisceaux coherents et en cohomologie etale. *Ann. E. N. S.* 4 (1974) 29–52.

(Oblatum 31–I–1975 & 26–VI–1975)

Department of Mathematics  
Yale University  
New Haven, Connecticut 06520  
U.S.A.